



Centralisers, complex reflection groups and actions in the Weyl group E_6

Graham A. Niblo¹ · Roger Plymen¹ · Nick Wright¹

Received: 18 October 2022 / Accepted: 3 April 2023 / Published online: 8 June 2023
© The Author(s) 2023, corrected publication 2023

Abstract

The compact, connected Lie group E_6 admits two forms: simply connected and adjoint type. As we previously established, the Baum–Connes isomorphism relates the two Langlands dual forms, giving a duality between the equivariant K-theory of the Weyl group acting on the corresponding maximal tori. Our study of the A_n case showed that this duality persists at the level of homotopy, not just homology. In this paper we compute the extended quotients of maximal tori for the two forms of E_6 , showing that the homotopy equivalences of sectors established in the A_n case also exist here, leading to a conjecture that the homotopy equivalences always exist for Langlands dual pairs. In computing these sectors we show that centralisers in the E_6 Weyl group decompose as direct products of reflection groups, generalising Springer’s results for regular elements, and we develop a pairing between the component groups of fixed sets generalising Reeder’s results. As a further application we compute the K -theory of the reduced Iwahori-spherical C^* -algebra of the p -adic group E_6 , which may be of adjoint type or simply connected.

Keywords Weyl groups · Complex reflection groups · Exceptional Lie groups · Centralisers · Duality · K -theory

Mathematics Subject Classification Primary 20G41 · 20G07; Secondary 20G05 · 20G10

Communicated by Christian Voigt.

✉ Nick Wright
wright@soton.ac.uk

Graham A. Niblo
g.a.niblo@soton.ac.uk

Roger Plymen
r.j.plymen@soton.ac.uk

¹ Mathematical Sciences, University of Southampton, SO17 1BJ, England

Contents

Introduction	220
1 Component groups of fixed sets	222
2 Elements, conjugacy classes and centralisers	230
2.1 The special elements	231
2.2 The elementary part of the centralisers	233
2.3 The full centralisers	237
3 Fixed sets and their quotients: the simply connected type	241
3.1 Alternative coordinate systems	244
3.2 The fixed sets and their quotients	246
4 Fixed sets and their quotients: the group of adjoint type	256
4.1 Ramified torus	257
4.2 Ramified component groups	258
5 Applications	261
5.1 Extended affine Weyl groups of type E_6	261
5.2 p -adic groups of type E_6	262
References	264

Introduction

The structure of element centralisers in Weyl groups plays a key role in our understanding of the representation theory of reductive groups of p -adic type. For the narrow class of regular elements Springer showed that the centralisers have the structure of a complex reflection group inherited directly from the action on the corresponding regular eigenspace [12]. Building on the work of Springer, Brewer [3] classified the irreducible, rank n complex reflection groups in a rank $2n$ real reflection group, exhibiting these as subgroups of centralisers in the automorphism group of the root graph. In the case of elliptic elements Reeder studied the centralisers by constructing a symplectic form on the coinvariant representation [7].

As remarked by Reeder, there is no general theory for the structure of centralisers in Weyl groups, though the classical cases A_n , B_n , C_n , D_n are well understood. In this paper we determine the structure for all the centralisers in the Weyl group of type E_6 . We provide a description in terms of generators and relations and give a classification of each centraliser as a product of reflection groups. In the case of the seven conjugacy classes of regular elements this recovers Springer's identification of the centralisers as complex reflection groups, though our approach is somewhat different to Springer.

Theorem 1 *Let W be the Weyl group of type E_6 . For each $w \in W$ the centraliser $Z_W(w)$ decomposes as a product of reflection groups.*

For all the elements of order 1 or 2 in the Weyl group, the centralisers are real reflection groups. In all but one of the remaining cases they are truly complex reflection groups. The one exception is the centraliser of the element of type A_2 . This is the product of the cyclic group C_3 with the group $(S_3 \wr C_2)$: the latter of which is not a complex reflection group. This factor however is a reflection group over the finite field \mathbb{F}_3 . The structure as reflection groups is outlined in Fig. 3.

In the case of an elliptic element w in a Weyl group W , Reeder studied the centraliser via its actions on the (finite) group of coinvariants $\Gamma/(I - w)\Gamma$ where Γ is the root

lattice, see [7]. Reeder defined a pairing on $\Gamma/(I - w)\Gamma$, for an elliptic element w , and showed that for Γ a self-dual lattice the pairing is non-degenerate. Dual to the action on coinvariants one may consider the action of W on the invariant part of the Pontryagin dual of Γ , which is the maximal torus \mathcal{T}^\vee in the Langlands dual of the simply connected form of the Lie group. For non-elliptic elements, the coinvariants and invariants become infinite, leading us to study the finite component groups of the fixed sets in maximal tori \mathcal{T} and \mathcal{T}^\vee of Langlands dual groups. The centraliser of an element $w \in W$ can be represented by automorphisms of these two finite groups, and we show that these groups and actions are dual.

We will establish this duality in a more general context: let W be a finite group acting orthogonally on a Euclidean vector space \mathfrak{t} , preserving a lattice Γ in \mathfrak{t} . Let Γ^\vee denote the dual lattice in \mathfrak{t}^* and let $\mathcal{T} = \mathfrak{t}/\Gamma$, $\mathcal{T}^\vee = \mathfrak{t}^*/\Gamma^\vee$. The fixed sets \mathcal{T}^w and $(\mathcal{T}^\vee)^w$ each consist of a product of a sub-torus with a finite subgroup: $\mathcal{T}^w \cong \mathcal{T}_1^w \times F_w$ and $(\mathcal{T}^\vee)^w \cong (\mathcal{T}^\vee)_1^w \times F_w^\vee$ where \mathcal{T}_1^w and $(\mathcal{T}^\vee)_1^w$ denote the identity components of the w -fixed sets in \mathcal{T} and \mathcal{T}^\vee respectively. (In the case where W is a Weyl group, Γ the root lattice and $w \in W$ an elliptic element, the group F_w^\vee is the fixed set for the action of w on \mathcal{T}^\vee and the Pontryagin dual of F_w^\vee is identified with Reeder’s group of coinvariants.) We will construct a non-degenerate pairing between F_w , F_w^\vee for all elements of the group W , allowing us to prove:

Theorem 2 *Let W be a finite group acting orthogonally on a Euclidean vector space \mathfrak{t} , preserving a lattice Γ in \mathfrak{t} . Then the dual component group F_w^\vee is (canonically) isomorphic to the Pontryagin dual of F_w . Therefore F_w^\vee is (non-canonically) isomorphic to F_w .*

Theorem 3 *The actions of $Z_W(w)$ on F_w and F_w^\vee are dual.*

Moreover, by exploiting the relationship between invariants and coinvariants we establish the following formula for the cardinality of the component group.

Theorem 4 *Let $w \in W$ with $w \neq I$ and let r be the rank of $I - w$. Let g be the greatest common divisor of the $r \times r$ -minors of the matrix of $I - w$ expressed in coordinates with respect to a basis for the dual lattice Γ^\vee . Then*

$$g = |F_w|.$$

The above results allow us to determine the fixed sets and to show that \mathcal{T}^w and $(\mathcal{T}^\vee)^w$ agree:

Theorem 5 *For $w \in W$ the fixed sets \mathcal{T}^w and $(\mathcal{T}^\vee)^w$ are non-canonically isomorphic as topological groups. In the special case of Weyl groups acting on maximal tori, this gives an isomorphism between the w -fixed sets for the Langlands dual forms.*

For a Weyl group W and maximal torus $\mathcal{T} = \mathfrak{t}/\Gamma$, the extended affine Weyl group is defined to be $\Gamma \rtimes W$. The W -equivariant K -theory of the maximal torus \mathcal{T}^\vee in the Langlands dual is identified with the K -theory of $C_r^*(\Gamma \rtimes W)$, while the W -equivariant K -homology of \mathcal{T} is identified with $K_*^{\Gamma \rtimes W}(\mathfrak{t})$ (see [5]). Hence the Baum–Connes isomorphism for $\Gamma \rtimes W$ corresponds to a pairing between the equivariant K -theory

groups $K_W^*(\mathcal{T})$ and $K_W^*(\mathcal{T}^\vee)$ which we showed in [5] is given by a Poincaré duality. In particular the pairing makes these two groups isomorphic up to torsion.

Returning to the case of E_6 , our description of the centralisers and the fixed sets allows us to compute the action of the Weyl group W on the inertia space (cf. Theorem 8) associated to the action of W on the maximal torus, thereby obtaining a description of the extended quotient. Using the equivariant Chern character, we compute (up to torsion) the W -equivariant K -theory for the maximal tori of both the simply connected and adjoint-type Lie groups of type E_6 , exhibiting the isomorphism between these K -theory groups.

The isomorphism between these K -theory groups corresponds to a cohomological relation between the corresponding extended quotients for the actions of the Weyl group W on the maximal tori. We refine this in the case of E_6 by showing that the isomorphism in cohomology arises from a homotopy equivalence at the level of sectors.

Theorem 6 *Let W be the Weyl group of type E_6 and let \mathcal{T} and \mathcal{T}^\vee denote the corresponding (real) maximal tori. For each $w \in W$ there is a homotopy equivalence*

$$\mathcal{T}^w/Z_W(w) \sim (\mathcal{T}^\vee)^w/Z_W(w).$$

This stratification phenomenon was introduced and studied in the case of A_n in [6], and we conjecture that this demonstrates a general principle for any compact connected semisimple Lie group:

Conjecture (Comparison of sectors) *For any compact connected semisimple Lie group, the homeomorphisms between $\mathcal{T}^w, (\mathcal{T}^\vee)^w$ provided by Theorem 5 descend to homotopy equivalences*

$$\mathcal{T}^w/Z_W(w) \sim (\mathcal{T}^\vee)^w/Z_W(w).$$

In Sect. 5, we provide two main applications. The first is the computation of the K -theory for the group C^* -algebras of the two extended affine Weyl groups of type E_6 showing that these agree. The second is a geometric description of the set of tempered representations in the Iwahori-spherical block of the p -adic adjoint group E_6 . For each of these applications, full use is made of our results in Table 2.

The paper is organised as follows:

1 Component groups of fixed sets

In this section we will consider the general case of a finite group W acting orthogonally on a Euclidean vector space \mathfrak{t} of dimension n , preserving a lattice Γ . The action thus descends to the torus $\mathcal{T} = \mathfrak{t}/\Gamma$. Let w be an element of W and \mathcal{T}^w the fixed set of the action of w on \mathcal{T} . Since the action is by automorphisms of the group \mathcal{T} , in each case the fixed set is a (closed) subgroup of \mathcal{T} and is therefore isomorphic to the direct product of a finite abelian group with the identity component of \mathcal{T}^w , which we denote \mathcal{T}_1^w . We note that \mathcal{T}_1^w is the image under the exponential map of the 1-eigenspace of

w acting on the Lie algebra \mathfrak{t} . Let F_w denote the group of components of the w -fixed set, that is

$$F_w = \mathcal{T}^w / \mathcal{T}_1^w$$

so that $\mathcal{T}^w \cong \mathcal{T}_1^w \times F_w$.

We use the inner product to identify \mathfrak{t} with \mathfrak{t}^* . Note that this identifies the W -action on \mathfrak{t} with the dual action on \mathfrak{t}^* , preserving the dual lattice

$$\Gamma^\vee = \{x \in \mathfrak{t} : \langle x, y \rangle \in \mathbb{Z}, \forall y \in \Gamma\}.$$

This induces an action of W on the dual torus $\mathcal{T}^\vee = \mathfrak{t} / \Gamma^\vee$. Let F_w^\vee denote the component group for the action on the dual torus: $F_w^\vee = (\mathcal{T}^\vee)^w / (\mathcal{T}^\vee)_1^w$.

Theorem 2 *Let W be a finite group acting orthogonally on a Euclidean vector space \mathfrak{t} , preserving a lattice Γ in \mathfrak{t} . Then the dual component group F_w^\vee is (canonically) isomorphic to the Pontryagin dual of F_w . Therefore F_w^\vee is (non-canonically) isomorphic to F_w .*

To prove the theorem we will construct a \mathbb{T}^1 -valued pairing between the Pontryagin duals $\widehat{F_w}$ and $\widehat{F_w^\vee}$ thus establishing an isomorphism

$$\widehat{F_w} \cong \widehat{\widehat{F_w^\vee}} \cong F_w^\vee.$$

The Pontryagin dual $\widehat{F_w}$ is canonically isomorphic to the set of characters on \mathcal{T}^w which are trivial on the subgroup \mathcal{T}_1^w . The group of characters on \mathcal{T} is identified with the lattice Γ^\vee via

$$\chi_x(y + \Gamma) := e^{2\pi i \langle x, y \rangle}.$$

The characters on \mathcal{T}^w are given by the quotient:

$$\widehat{\mathcal{T}^w} = \Gamma^\vee / ((I - w)\Gamma^\vee)$$

Hence the set of characters on \mathcal{T}^w vanishing on \mathcal{T}_1^w is given by

$$\{x \in \Gamma^\vee : \chi_x \equiv 1 \text{ on } \mathcal{T}_1^w\} / ((I - w)\Gamma^\vee).$$

Note that $\chi_x \equiv 1$ on \mathcal{T}_1^w if and only if x is orthogonal to the fixed set \mathfrak{t}^w , so we obtain the identification:

$$\widehat{F_w} \cong \{x \in \Gamma^\vee : x \perp \mathfrak{t}^w\} / ((I - w)\Gamma^\vee).$$

Likewise for the dual F_w^\vee we have

$$\widehat{F_w^\vee} \cong \{y \in \Gamma : y \perp \mathfrak{t}^w\} / ((I - w)\Gamma).$$

We introduce the notation

$$\begin{aligned} \mathfrak{r}_w &= \{y \in \mathfrak{t} : y \perp \mathfrak{t}^w\} \\ \mathfrak{r}_w(\Gamma) &= \mathfrak{r}_w \cap \Gamma \\ \mathfrak{r}_w(\Gamma^\vee) &= \mathfrak{r}_w \cap \Gamma^\vee. \end{aligned}$$

Following Reeder, let m denote the minimal polynomial of w and let $\dot{m}(t) = \frac{m(t) - m(1)}{t - 1}$. Reeder restricts to the elliptic case, i.e. where $m(1) \neq 0$. In the non-elliptic case we will define $\ddot{m}(t) = \frac{\dot{m}(t) - \dot{m}(1)}{t - 1}$ and in general we set

$$\bar{m} = \begin{cases} \dot{m} & m(1) \neq 0 \\ \ddot{m} & m(1) = 0. \end{cases}$$

Definition 1.1 For $x \in \Gamma^\vee$ and $y \in \Gamma$ we define a twisted pairing

$$\langle x, y \rangle_w := \langle x, \bar{m}(w)y \rangle.$$

We will consider this integer valued pairing modulo μ where

$$\mu = \begin{cases} m(1) & \text{if } m(1) \neq 0 \\ \dot{m}(1) & \text{otherwise.} \end{cases}$$

Note that μ is always non-zero since m is the minimal polynomial, and w is a normal operator.

The following Lemma shows that $\bar{m}(w)$ provides an inverse in the elliptic case, and a partial inverse in general, to $I - w$.

Lemma 1.2 *The operator $\bar{m}(w)(I - w)$ is $\mu p_{\mathfrak{r}_w}$ where $p_{\mathfrak{r}_w}$ denotes the orthogonal projection onto \mathfrak{r}_w .*

Proof Clearly $\bar{m}(w)(I - w) = 0$ on the fixed set \mathfrak{t}^w which is the orthogonal complement of \mathfrak{r}_w , so it suffices to show that $\bar{m}(w)(I - w)y = \mu y$ for $y \in \mathfrak{r}_w$.

In the elliptic case the result is immediate from the fact that $\dot{m}(t)(1 - t) = \mu - m(t)$ and $m(w) = 0$.

In the non-elliptic case we have $\ddot{m}(t)(1 - t) = \mu - \dot{m}(t)$. The operator $\dot{m}(w)$ satisfies the equation

$$\dot{m}(w)(I - w) = m(1) - m(w) = 0.$$

The image of $I - w$ is precisely \mathfrak{r}_w since w is a normal operator and $\mathfrak{t}^w = \ker(I - w)$ hence for $y \in \mathfrak{r}_w$

$$\dot{m}(w)(I - w)y = \mu y - \dot{m}(w)y = \mu y.$$

□

Lemma 1.3 *The twisted pairing $\langle -, - \rangle_w$ descends to a well-defined pairing*

$$\widehat{F}_w \times \widehat{F}_w^\vee \rightarrow \mathbb{Z}/\mu\mathbb{Z}.$$

Proof Recall that \widehat{F}_w is identified with $\mathfrak{r}_w(\Gamma^\vee)/(I - w)\Gamma^\vee$ and \widehat{F}_w^\vee is identified with $\mathfrak{r}_w(\Gamma)/(I - w)\Gamma$.

Let $x \in \mathfrak{r}_w(\Gamma^\vee)$ and $y \in \Gamma$. Then

$$\begin{aligned} \langle x, (I - w)y \rangle_w &= \langle x, \overline{m}(w)(I - w)y \rangle \\ &= \langle x, \mu p_{\mathfrak{r}_w} y \rangle \\ &= \langle p_{\mathfrak{r}_w} x, \mu y \rangle \\ &= \mu \langle x, y \rangle \in \mu\mathbb{Z}. \end{aligned}$$

Similarly if $x \in \Gamma^\vee$ and $y \in \mathfrak{r}_w(\Gamma)$ then

$$\begin{aligned} \langle (I - w)x, y \rangle_w &= \langle (I - w)x, \overline{m}(w)y \rangle \\ &= \langle (w^{-1} - I)wx, \overline{m}(w)y \rangle \\ &= \langle wx, (w - I)\overline{m}(w)y \rangle \\ &= \langle wx, -\mu p_{\mathfrak{r}_w} y \rangle \\ &= -\mu \langle wx, y \rangle \in \mu\mathbb{Z}. \end{aligned}$$

Hence the pairing is well defined on the quotients. □

We note that since $\mathfrak{r}_w(\Gamma)$ contains $(I - w)\Gamma$ it spans the space $\mathfrak{r}_w = (I - w)t$. In particular $\mathfrak{r}_w(\Gamma)$ is a lattice in the space \mathfrak{r}_w , and similarly for $\mathfrak{r}_w(\Gamma^\vee)$.

We consider the lattices $(\mathfrak{r}_w(\Gamma))^\vee = \{x \in \mathfrak{r}_w : \langle x, \mathfrak{r}_w(\Gamma) \rangle \subseteq \mathbb{Z}\}$ and $(\mathfrak{r}_w(\Gamma^\vee))^\vee = \{y \in \mathfrak{r}_w : \langle \mathfrak{r}_w(\Gamma^\vee), y \rangle \subseteq \mathbb{Z}\}$. Since $\mathfrak{r}_w(\Gamma^\vee)$ lies in Γ^\vee it pairs integrally with Γ and hence $\mathfrak{r}_w(\Gamma^\vee) \subseteq (\mathfrak{r}_w(\Gamma))^\vee$. Similarly $\mathfrak{r}_w(\Gamma) \subseteq (\mathfrak{r}_w(\Gamma^\vee))^\vee$. Let

$$\begin{aligned} \mathfrak{t}^w(\Gamma) &= \mathfrak{t}^w \cap \Gamma, \\ \mathfrak{t}^w(\Gamma^\vee) &= \mathfrak{t}^w \cap \Gamma^\vee. \end{aligned}$$

Lemma 1.4 *The projection $p_{\mathfrak{r}_w}$ gives maps $\Gamma^\vee \rightarrow (\mathfrak{r}_w(\Gamma))^\vee$ and $\Gamma \rightarrow (\mathfrak{r}_w(\Gamma^\vee))^\vee$ which induce isomorphisms*

$$\begin{aligned} \Gamma^\vee/\mathfrak{t}^w(\Gamma^\vee) &\cong (\mathfrak{r}_w(\Gamma))^\vee, \\ \Gamma/\mathfrak{t}^w(\Gamma) &\cong (\mathfrak{r}_w(\Gamma^\vee))^\vee. \end{aligned}$$

Proof If $x \in \Gamma^\vee$ then $\langle x, y \rangle \in \mathbb{Z}$ for all $y \in \mathfrak{r}_w(\Gamma)$. Now

$$\langle p_{\mathfrak{r}_w} x, y \rangle = \langle x, p_{\mathfrak{r}_w} y \rangle = \langle x, y \rangle \in \mathbb{Z}$$

and $p_{\mathfrak{r}_w}x \in \mathfrak{r}_w$ so $p_{\mathfrak{r}_w}x \in (\mathfrak{r}_w(\Gamma))^\vee$. Clearly this map has kernel $\mathfrak{t}^w(\Gamma^\vee)$.

It remains to check surjectivity. Since $\mathfrak{r}_w(\Gamma)$ is a lattice in \mathfrak{r}_w and is the intersection of \mathfrak{r}_w with Γ it follows that \mathfrak{r}_w is complemented in Γ . Picking a complement for $\mathfrak{r}_w(\Gamma)$, let $\pi : \Gamma \rightarrow \mathfrak{r}_w(\Gamma)$ denote the retraction obtained by killing the complement.

Given any $x \in (\mathfrak{r}_w(\Gamma))^\vee$ consider the homomorphism from Γ to \mathbb{Z} defined by

$$y \mapsto \langle x, \pi(y) \rangle.$$

Since there is a perfect pairing between $\widehat{\Gamma}$ and Γ^\vee , there exists $x' \in \Gamma^\vee$ such that $\langle x, \pi(y) \rangle = \langle x', y \rangle$. Now for all $y \in \mathfrak{r}_w(\Gamma)$ we have

$$\langle p_{\mathfrak{r}_w}x', y \rangle = \langle x', p_{\mathfrak{r}_w}y \rangle = \langle x', y \rangle = \langle x, \pi(y) \rangle = \langle x, y \rangle.$$

Thus $x = p_{\mathfrak{r}_w}x'$ which is in the image.

Exchanging the roles of Γ and Γ^\vee gives the dual case. □

Proposition 1.5 *The pairing*

$$\widehat{F}_w \times \widehat{F}_w^\vee \rightarrow \mathbb{Z}/\mu\mathbb{Z}.$$

is left and right non-degenerate.

Proof Let $x \in \mathfrak{r}_w(\Gamma^\vee)$ and suppose that $\langle x, y \rangle_w \cong 0 \pmod{\mu}$ for all $y \in \mathfrak{r}_w(\Gamma)$, that is $\langle x, \frac{1}{\mu}\overline{m}(w)y \rangle \in \mathbb{Z}$ for all y in $\mathfrak{r}_w(\Gamma)$.

Hence $\frac{1}{\mu}\overline{m}(w)^*x$ pairs integrally with all y in $\mathfrak{r}_w(\Gamma)$, and lies in the space \mathfrak{r}_w , so $\frac{1}{\mu}\overline{m}(w)^*x \in (\mathfrak{r}_w(\Gamma))^\vee$.

By Lemma 1.4 $(\mathfrak{r}_w(\Gamma))^\vee = p_{\mathfrak{r}_w}\Gamma^\vee$ and by Lemma 1.2

$$\mu p_{\mathfrak{r}_w} = \overline{m}(w)(I - w) = (\overline{m}(w)(I - w))^*$$

so we have

$$\overline{m}(w)^*x \in \mu p_{\mathfrak{r}_w}\Gamma^\vee = \overline{m}(w)^*(I - w)^*\Gamma^\vee$$

since $\overline{m}(w)$ and $I - w$ commute.

Now $x, (I - w)^*\Gamma^\vee$ lie in \mathfrak{r}_w and $\overline{m}(w)^*$ is injective on this space (again by Lemma 1.2) giving

$$x \in (I - w)^*\Gamma^\vee = (I - w^{-1})\Gamma^\vee = (w - I)w^{-1}\Gamma^\vee = (I - w)\Gamma^\vee.$$

Thus the pairing is non-degenerate on the left.

Now let $y \in \mathfrak{r}_w(\Gamma)$ and suppose that $\langle x, y \rangle_w \cong 0 \pmod{\mu}$ for all $x \in \mathfrak{r}_w(\Gamma^\vee)$, that is $\langle x, \frac{1}{\mu}\overline{m}(w)y \rangle \in \mathbb{Z}$ for all x in $\mathfrak{r}_w(\Gamma^\vee)$.

Hence $\frac{1}{\mu}\overline{m}(w)y \in (\mathfrak{r}_w(\Gamma^\vee))^\vee = p_{\mathfrak{r}_w}\Gamma$. Then

$$\overline{m}(w)y \in \mu p_{\mathfrak{r}_w}\Gamma = \overline{m}(w)(I - w)\Gamma$$

and as $\overline{m}(w)$ is injective on \mathfrak{x}_w we have $y \in (I - w)\Gamma$. Hence the pairing is also non-degenerate on the right. \square

Proof of Theorem 2 As noted above it suffices to prove that \widehat{F}_w is the Pontryagin dual of \widehat{F}_w^\vee .

Define a pairing $\widehat{F}_w \times \widehat{F}_w^\vee \rightarrow \mathbb{T}^1$ by

$$\chi(x + (I - w)\Gamma^\vee, y + (I - w)\Gamma) := e^{\frac{2\pi i}{\mu} \langle x, y \rangle_w}.$$

Fixing an element of \widehat{F}_w this gives a character on \widehat{F}_w^\vee , hence we have a homomorphism from \widehat{F}_w to the Pontryagin dual of \widehat{F}_w^\vee . Left non-degeneracy of the pairing $\langle -, - \rangle_w$ implies injectivity of this homomorphism.

Likewise fixing an element of \widehat{F}_w^\vee gives a character on \widehat{F}_w , hence we have a homomorphism from \widehat{F}_w^\vee to the Pontryagin dual of \widehat{F}_w . Right non-degeneracy again implies injectivity.

Combining these two maps we deduce that

$$|\widehat{F}_w| \leq |\widehat{F}_w^\vee| = |\widehat{F}_w^\vee| \leq |\widehat{F}_w| = |\widehat{F}_w|.$$

Since the cardinalities are equal the injections must be isomorphisms.

The following result gives a tool for computing the cardinality of these groups, which is relevant in the computation of K -theory. It may be useful to bear in mind the following example.

Example 1.6 The conjugacy class representative $s_0s_1s_5s_3$ of type A_1^4 is represented by the matrix

$$M = \begin{pmatrix} -1 & 1 & 0 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 & 0 & -2 \\ 0 & 1 & -1 & 1 & 0 & -2 \\ 0 & 0 & 0 & 1 & 0 & -2 \\ 0 & 0 & 0 & 1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}$$

with respect to the lattice Γ and its transpose with respect to the lattice Γ^\vee . The rank of $I - M$ is 4 and the greatest common divisors of the 4×4 minors is easily seen to be 4 for both $I - M$ and $I - M^T$. As we will establish in the next theorem, this computes the order of the component group of the corresponding fixed set.

This example is simplified by the fact that the operator is self adjoint, ensuring that the dual matrices over Γ and Γ^\vee are transpose. In general, the matrices with respect to the dual bases are related by the formula $(I - M) \rightarrow (I - M^{-1})^T$, however the argument given in Theorem 5 below shows that the gcd of the corresponding minors is still the same in these two matrices.

Theorem 4 *Let $w \in W$ with $w \neq I$ and let r be the rank of $I - w$. Let g be the greatest common divisor of the $r \times r$ -minors of the matrix of $I - w$ expressed in coordinates with respect to a basis for the dual lattice Γ^\vee . Then*

$$g = |F_w|.$$

Proof Let $i = |F_w| = [\mathfrak{x}_w(\Gamma^\vee) : (I - w)\Gamma^\vee]$.

Let v_1, \dots, v_r be a basis for $(I - w)\Gamma^\vee$, and let w_1, \dots, w_r be a basis for $\mathfrak{x}_w(\Gamma^\vee)$.

Expressing these vectors in coordinates with respect to a basis for Γ^\vee , each element of the basis $\{v_i\}$ lies in the integer column span of $I - w$, so the greatest common divisor of the $r \times r$ -minors of the matrix $(v_1 | \dots | v_r)$ must be divisible by g . But conversely each column of $I - w$ can be written in terms of the basis, hence we deduce that g equals the greatest common divisor of the $r \times r$ -minors of the matrix $(v_1 | \dots | v_r)$.

The subgroup $\mathfrak{x}_w(\Gamma^\vee) = \mathfrak{x}_w \cap \Gamma^\vee$ has a complement Λ in Γ^\vee and the index i of $(I - w)\Gamma^\vee$ in the lattice $\mathfrak{x}_w(\Gamma^\vee)$ equals the index of $(I - w)\Gamma^\vee \oplus \Lambda$ in Γ^\vee . Let x_1, \dots, x_{n-r} be a basis of Λ , so $w_1, \dots, w_r, x_1, \dots, x_{n-r}$ gives a basis for Γ^\vee and hence the corresponding $n \times n$ matrix has determinant 1.

The elements $v_1, \dots, v_r, x_1, \dots, x_{n-r}$ form a basis for $(I - w)\Gamma^\vee \oplus \Lambda$ which has index i in Γ^\vee , hence the corresponding matrix has determinant i . But the determinant can be built from the minors of $(v_1 | \dots | v_r)$, so must be divisible by g . That is $g|i$.

Now consider exterior products: The determinants are given by $v_1 \wedge \dots \wedge v_r \wedge x_1 \wedge \dots \wedge x_{n-r}$ and by $w_1 \wedge \dots \wedge w_r \wedge x_1 \wedge \dots \wedge x_{n-r}$ thus

$$v_1 \wedge \dots \wedge v_r \wedge x_1 \wedge \dots \wedge x_{n-r} = i(w_1 \wedge \dots \wedge w_r \wedge x_1 \wedge \dots \wedge x_{n-r})$$

As v_1, \dots, v_r and w_1, \dots, w_r span the same r -dimensional subspace of \mathfrak{t}^* it follows that $v_1 \wedge \dots \wedge v_r$ is a multiple of $w_1 \wedge \dots \wedge w_r$, and from the above equation we see that the coefficient is i :

$$v_1 \wedge \dots \wedge v_r = i(w_1 \wedge \dots \wedge w_r).$$

The coefficients of $v_1 \wedge \dots \wedge v_r$ in the standard basis $\{e_{i_1} \wedge \dots \wedge e_{i_r}\}$ for the exterior algebra of Γ^\vee are precisely the $r \times r$ minors. But the above equation tells us that as $w_1 \wedge \dots \wedge w_r$ has integer coefficients, the coefficients of $v_1 \wedge \dots \wedge v_r$ must be divisible by i . Hence $i|g$.

Therefore $i = g$ as claimed. □

Remark 1.7 We note that since $|F_w| = |F_w^\vee|$ by Theorem 2 we may use the greatest common divisors of minors of the matrix of $I - w$ with respect to a basis for either the lattice Γ or the lattice Γ^\vee to compute the cardinality of the groups F_w, F_w^\vee .

As a consequence of Theorem 2 we have the following:

Theorem 5 *For $w \in W$ the fixed sets \mathcal{T}^w and $(\mathcal{T}^\vee)^w$ are non-canonically isomorphic as topological groups. In the special case of Weyl groups acting on maximal tori, this gives an isomorphism between the w -fixed sets for the Langlands dual forms.*

Proof The fixed sets are isomorphic to $T_1^w \times F_w, (T_1^\vee)^w \times F_w^\vee$ and we know that the component groups are isomorphic by 2 so we only need to show that the identity components are isomorphic. This follows from the fact that these are precisely the image of the fixed sets in $\mathfrak{t} \cong \mathfrak{t}^*$ under the respective quotient maps. \square

We now turn to the actions of the centraliser $Z_W(w)$ on the groups F_w and F_w^\vee , induced by the actions of $Z_W(w)$ on the w -fixed sets T^w and $(T^\vee)^w$. Although the groups F_w and F_w^\vee are isomorphic, there is in general no $Z_W(w)$ -equivariant isomorphism between the component groups. Instead we will show that the actions are dual. Nonetheless even though these dual actions can be very different we will see (cf. Section 4) that the numbers of orbits of the centraliser on F_w and F_w^\vee are the same. This is an instance of our conjecture and can be seen clearly in the dual actions of G_{25} on \mathbb{F}_3^3 in case (21) in Sect. 3.

Theorem 3 *The actions of $Z_W(w)$ on F_w and F_w^\vee are dual.*

Proof For $s \in F_w, t \in F_w^\vee \cong \widehat{F_w}$ denote the corresponding pairing by $\langle s, t \rangle^w$. We must show:

$$\langle gs, gt \rangle^w = \langle s, t \rangle^w \quad \forall s \in F_w, t \in F_w^\vee, g \in Z_W(w).$$

Recall that $\widehat{F_w} \cong \mathfrak{r}_w(\Gamma^\vee)/(I-w)\Gamma^\vee$ and $\widehat{F_w^\vee} \cong \mathfrak{r}_w(\Gamma)/(I-w)\Gamma$. By construction these isomorphisms are equivariant for the action of $Z_W(w)$.

The pairing of $\widehat{F_w}$ and $\widehat{F_w^\vee}$ is defined by

$$\chi(x + (I-w)\Gamma^\vee, y + (I-w)\Gamma) := e^{\frac{2\pi i}{\mu} \langle x, y \rangle_w}$$

and for $g \in Z_W(w)$ we have $\chi(gx + (I-w)\Gamma^\vee, gy + (I-w)\Gamma) := e^{2\pi i \langle gx, gy \rangle_w}$. Now note that

$$\langle gx, gy \rangle_w = \langle gx, \overline{m}(w)gy \rangle = \langle gx, g\overline{m}(w)y \rangle$$

since g is in the centraliser of w , so as W acts isometrically on \mathfrak{t} we have $\langle gx, gy \rangle_w = \langle x, y \rangle_w$.

Now for $s \in F_w, t \in F_w^\vee$ the pairing of gs with gt is defined to be the pairing of gs with the image of gt under the composition

$$F_w^\vee \cong \widehat{\widehat{F_w^\vee}} \cong \widehat{F_w}$$

The first isomorphism is tautologically equivariant while the second is equivariant by the above calculation. Letting ψ denote the image of t under this composition we have

$$\langle gs, gt \rangle^w = \langle g\psi, gs \rangle = \langle \psi, s \rangle = \langle s, t \rangle^w.$$

\square

Bearing in mind the example of E_6 we now suppose that Γ^\vee refines Γ with quotient $\mathcal{Z} = \Gamma^\vee / \Gamma$ cyclic of prime order. Moreover we assume that the induced action of W on the quotient \mathcal{Z} is trivial. It follows that \mathcal{Z} lies in each of the fixed groups \mathcal{T}^w .

The w -fixed set in the dual torus $\mathcal{T}^\vee = \mathfrak{t} / \Gamma^\vee = \mathcal{T} / \mathcal{Z}$ contains the image of \mathcal{T}^w under the quotient map $\mathcal{T} \rightarrow \mathcal{T}^\vee$. In particular $(\mathcal{T}^\vee)_1^w$ is the image of \mathcal{T}_1^w under the quotient map, since the identity component of the fixed set is precisely the image of the fixed set \mathfrak{t}^w under the respective exponential map. Hence the quotient map induces a map from $F_w = \mathcal{T}^w / \mathcal{T}_1^w$ to $F_w^\vee = (\mathcal{T}^\vee)^w / (\mathcal{T}^\vee)_1^w$.

We have two cases: either \mathcal{Z} lies in \mathcal{T}_1^w or $\mathcal{Z} \cap \mathcal{T}_1^w$ is trivial. In the former case the map from F_w to F_w^\vee is injective. It is thus an isomorphism since the groups have the same cardinality by Theorem 2. Hence in this case the fixed set $(\mathcal{T}^\vee)^w$ is precisely the image of the fixed set \mathcal{T}^w under the quotient by \mathcal{Z} , that is:

$$(\mathcal{T}^\vee)^w = (\mathcal{T}^\vee)_1^w \times F_w^\vee \cong \mathcal{T}_1^w / \mathcal{Z} \times F_w.$$

In the latter case the quotient map $\mathcal{T}^w \rightarrow (\mathcal{T}^\vee)^w$ gives an isomorphism of the identity components and takes F_w to a subgroup of F_w^\vee , which, again by Theorem 2, must have index $|\mathcal{Z}|$ in F_w^\vee . Hence in this case the image of \mathcal{T}^w in $(\mathcal{T}^\vee)^w$ is

$$\mathcal{T}_1^w \times F_w / \mathcal{Z}$$

which has index $|\mathcal{Z}|$ in the fixed set $(\mathcal{T}^\vee)^w$.

Remark 1.8 In the case that $\mathcal{Z} < F_w$ the $Z_W(w)$ -orbits in F_w include $|\mathcal{Z}|$ singletons. In this case, dually the orbits in F_w^\vee are partitioned into $|\mathcal{Z}|$ sets. To see this we note that the inclusion of \mathcal{Z} in F_w induces a quotient $\widehat{F}_w \rightarrow \widehat{\mathcal{Z}}$. Since $F_w^\vee \cong \widehat{F}_w$ we have a quotient map $F_w^\vee \rightarrow \widehat{\mathcal{Z}}$, given by the pairing of F_w^\vee with \mathcal{Z} . By Theorem 3 the pairing is equivariant but \mathcal{Z} is fixed by $Z_W(w)$, hence the map $F_w^\vee \rightarrow \widehat{\mathcal{Z}}$ is invariant under the action of $Z_W(w)$. Each $Z_W(w)$ -orbit in F_w^\vee thus lies in a coset of the kernel of the map $F_w^\vee \rightarrow \widehat{\mathcal{Z}}$.

2 Elements, conjugacy classes and centralisers


In this section we will provide a list of carefully selected representatives for the conjugacy classes in the Weyl group of type E_6 together with key properties of these elements. We will also introduce a number of special elements of the Weyl group that will play a key role in understanding the centraliser subgroups and their actions.

The elements s_1, \dots, s_6 denote the standard simple reflections generating the Weyl group W , with corresponding roots denoted r_1, \dots, r_6 . We define s_0 to be the reflection corresponding to the root

$$r_0 := -r_1 - 2r_2 - 3r_3 - 2r_4 - r_5 - 2r_6$$

which is the negation of the root of highest weight. This is at 120° to the simple root r_6 , so the group generated by s_0 and s_6 is isomorphic to the symmetric group S_3 . On the

Fig. 1 Coxeter–Dynkin Diagram of the roots r_0, \dots, r_6



other hand, r_0 is orthogonal to the roots r_1, \dots, r_5 , so s_0 commutes with the reflections s_1, \dots, s_5 .

These relationships can be summarised in the following extension of the Dynkin diagram for E_6 . We remark that the reflection s_0 is the linear part of the additional simple reflection in the affine Weyl group \tilde{E}_6 , and the extended diagram is isomorphic to the Dynkin diagram of this group (Fig. 1).

This diagram demonstrates additional symmetries of the set of roots for E_6 , which is the reason that including r_0 is so valuable. In choosing representatives for the conjugacy classes we also make use of the reflection $T = s_3^{s_2 s_4 s_6}$ corresponding to the root $r_T := -(r_2 + r_3 + r_4 + r_6) = 1/2(r_0 + r_3 + r_1 + r_5)$. The name is chosen to reflect the geometry of the roots r_2, r_3, r_4, r_6 in the Dynkin diagram, and the symmetry of this set in the extended diagram again motivates the inclusion of this element.

The roots r_0, r_1, r_5 and $-r_T$ form a Dynkin diagram of type D_4 with $-r_T$ as the central vertex, hence the group $\langle s_0, s_1, s_5, T \rangle$ is a Weyl group of type D_4 . We remark in passing that this group contains s_3 as the reflection with root $-2r_T + r_0 + r_1 + r_5$.

The centralisers will be built from an elementary part along with a small number of additional elements $u_1, u_2, u_3, T, T s_3$.


2.1 The special elements

We now define elements u_1, u_2 in W which generate a copy of S_3 acting (by conjugation) as a dihedral group on the reflections s_0, \dots, s_6 as illustrated in Fig. 2.

The element u_1 is defined as the product of four commuting reflections with roots

$$\{r_0, r_3, r_2 + r_3 + r_4, r_1 + r_2 + r_3 + r_4 + r_5\}.$$

Fig. 2 The action of the special elements u_i



It is easy to see that this element negates the roots r_0, r_3 , so $s_0^{u_1} = s_0$ and $s_3^{u_1} = s_3$. Acting with u_1 on the root r_6 gives the sum

$$r_6 + r_0 + r_3 + (r_2 + r_3 + r_4) + (r_1 + r_2 + r_3 + r_4 + r_5) = -r_6$$

hence $s_6^{u_1} = s_6$. Its action on r_2 takes it to

$$r_2 + r_3 - (r_2 + r_3 + r_4) = -r_4$$

so u_1 conjugates s_2 to s_4 and vice versa. Finally, acting on the root r_1 takes it to

$$r_1 + (r_2 + r_3 + r_4) - (r_1 + r_2 + r_3 + r_4 + r_5) = -r_5,$$

so conjugation by u_1 switches s_1 and s_5 as illustrated in the diagram. We note in passing that $-u_1$ is the so-called non-trivial pinned automorphism for E_6 , as described by Reeder [8].

Turning to the element u_2 , this is defined as the product of the four commuting reflections with roots

$$\{r_5, r_3, r_6 + r_3 + r_2, r_0 + r_6 + r_3 + r_2 + r_1\}.$$

A similar calculation shows that conjugation by this element preserves s_5, s_4, s_3 and switches the pairs s_6, s_2 and s_0, s_1 . Since the root representation is faithful we see that the product $u_1 u_2$ has order three, and therefore these elements generate a copy of the dihedral group \mathcal{D}_3 and the element $u_3 = u_1 u_2 u_1 = u_2 u_1 u_2$ is the product of four commuting reflections with roots

$$\{r_1, r_3, r_4 + r_3 + r_6, r_5 + r_4 + r_3 + r_6 + r_0\}.$$

In many cases we will see that the u_1, u_2 dihedral group acts on one or more two-dimensional subtori of the maximal torus of E_6 . There are two natural actions of the dihedral group \mathcal{D}_3 on a 2-torus: considering \mathcal{D}_3 as the Weyl group $W(A_2)$ we have the actions on the maximal tori in the Lie groups SU_3 and PSU_3 . The maximal tori are given respectively by triples $(\alpha, \beta, \gamma) \in \mathbb{T}^3$ such that $\alpha\beta\gamma = 1$, and the quotient of this by the group $C_3 = \{(\omega, \omega, \omega) : \omega^3 = 1\}$. The action of the dihedral group simply permutes the three coordinates. In both cases the maximal torus can be described as a hexagon with opposite sides identified, and the two actions correspond to the two conjugacy classes of \mathcal{D}_3 in \mathcal{D}_6 . The orbifold quotients are an equilateral triangle in the SU_3 case and a cone for PSU_3 .

Given that u_1, u_2 act by signed permutations on the roots $r_1 \dots, r_6, r_0$ we will see that the action of these on the 2-tori are again given by signed permutations:

$$\begin{aligned} u_1 &: (\alpha, \beta, \gamma) \mapsto (\beta^{-1}, \alpha^{-1}, \gamma^{-1}) \\ u_2 &: (\alpha, \beta, \gamma) \mapsto (\gamma^{-1}, \beta^{-1}, \alpha^{-1}). \end{aligned}$$

This has the effect of dualising the actions: the signed permutation action on the SU_3 torus is equivariantly isomorphic to the standard permutation action on the PSU_3 torus, and similarly the signed permutation action on the PSU_3 torus is equivariantly isomorphic to the standard permutation action on the SU_3 torus.

The two equivariant isomorphisms are both given by the map

$$(\alpha : \beta : \gamma) \mapsto (\beta\gamma^{-1}, \gamma\alpha^{-1}, \alpha\beta^{-1}).$$

2.2 The elementary part of the centralisers

Our approach is slightly different to that of Carter, in that we begin with an elementary part of the centraliser that can easily be read off from the extended Coxeter–Dynkin diagram, and then seek additional elements to complete the centraliser. Nonetheless our elementary part lies within Carter’s direct product $W_1 \times W_2$ for the conjugacy class representative.

Consider an element $w = w_1 w_2 \dots w_l \in W$ where w_1, \dots, w_l are root reflections and $l = \bar{l}(w)$ is the word length of an element $w \in W$ in terms of the generating set of all root reflections in the Weyl group. Associated to such a word there is a root diagram (cf. Carter [4]) where roots are connected by an edge when they are not orthogonal (here we only have single edges as we are in the E_6 case). We write our representatives in such a way that each connected component of the diagram corresponds to a subword. Such subwords g_1, \dots, g_k of course commute with the element w (and with one another) giving a direct product $\langle g_1 \rangle \times \dots \times \langle g_k \rangle$ in the centraliser of w .

Any roots which are orthogonal to all roots in the diagram will again give root reflections in the centraliser. The group these generate is precisely Carter’s group W_2 . The elementary part of the centraliser is $\langle g_1 \rangle \times \dots \times \langle g_k \rangle \times W_2$.

Carter determined the order of the centraliser subgroups, and for 11 of the conjugacy classes this shows that the elementary part is the entire centraliser and the structure is obvious. In the remaining 14 cases we are left to discover additional elements of the centraliser. In all cases we will show that these additional generators can be found in the subgroup $\langle u_1, u_2, T, Ts_3 \rangle \cong \mathcal{D}_3 \times \mathcal{D}_3$, as a consequence of our careful choice of representatives. In determining the centralisers and their actions we will also exploit the fact that the centraliser of an element g is contained in the centraliser of g^n , and have chosen conjugacy class representatives accordingly. This relationship between the centralisers is summarised in Fig. 3. Our choices of conjugacy class representatives are tabulated below (Table 1). We list the eigenvalues for each of these representatives for the standard six dimensional representation: these are distinct, confirming that our representatives give distinct conjugacy classes and that the conjugacy classes are characterised by their eigenvalues. We also describe the elementary part of the centraliser, and the index of this group in the full centraliser where it is not 1.

Table 1 The conjugacy class representatives and the elementary part of their centralisers

Conjugacy class		Centraliser			Elementary part of the centraliser		Index in full centraliser
Carter type	Representative	Eigenvalues	Order	Generators	Structure		
\emptyset	e	$1, 1, 1, 1, 1$	51,840	$\{e\} \times W$	W		
A_1	s_0	$1, 1, 1, 1, -1$	1440	$\langle s_0 \rangle \times \langle s_1, \dots, s_5 \rangle$	$C_2 \times S_6$		
A_1^2	$s_0 s_1$	$1, 1, 1, 1, -1, -1$	192	$\langle s_0, s_1 \rangle \times \langle s_3, \dots, s_5 \rangle$	$C_2 \times C_2 \times S_4$	2	
A_2	$s_0 s_6$	$1, 1, 1, 1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}$	216	$\langle s_0 s_6 \rangle \times \langle s_1, s_2, s_4, s_5 \rangle$	$C_3 \times S_3 \times S_3$	2	
A_1^3	$s_0 s_1 s_5$	$1, 1, 1, -1, -1, -1$	96	$\langle s_0, s_1, s_5 \rangle \times \langle s_3 \rangle$	$C_2 \times C_2 \times C_2 \times C_2$	6	
$A_2 \times A_1$	$s_0 s_6 s_1$	$1, 1, 1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, -1$	36	$\langle s_0 s_6, s_1 \rangle \times \langle s_4, s_5 \rangle$	$C_3 \times C_2 \times S_3$		
A_3	$s_0 s_6 s_3$	$1, 1, 1, i, -1, -i$	32	$\langle s_0 s_6 s_3 \rangle \times \langle s_1, s_5 \rangle$	$C_4 \times C_2 \times C_2$	2	
A_1^4	$s_0 s_1 s_5 s_3$	$1, 1, -1, -1, -1, -1$	1152	$\langle s_0, s_1, s_5, s_3 \rangle$	$C_2 \times C_2 \times C_2 \times C_2$	72	
$A_2 \times A_1^2$	$s_0 s_6 s_1 s_5$	$1, 1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, -1, -1$	24	$\langle s_0 s_6, s_1, s_5 \rangle$	$C_3 \times C_2 \times C_2$	2	
A_2^2	$s_0 s_6 s_1 s_2$	$1, 1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}$	108	$\langle s_0 s_6, s_1 s_2 \rangle \times \langle s_4, s_5 \rangle$	$C_3 \times C_3 \times S_3$	2	
$A_3 \times A_1$	$s_0 s_6 s_3 s_1$	$1, 1, i, -1, -i, -1$	16	$\langle s_0 s_6 s_3, s_1 \rangle \times \langle s_5 \rangle$	$C_4 \times C_2 \times C_2$		
A_4	$s_0 s_6 s_3 s_4$	$1, 1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, e^{\frac{6}{5}\pi i}, e^{\frac{8}{5}\pi i}$	10	$\langle s_0 s_6 s_3 s_4 \rangle \times \langle s_1 \rangle$	$C_5 \times C_2$		
D_4	$s_0 s_1 s_5 T$	$1, 1, -1, -1, e^{\frac{1}{3}\pi i}, e^{\frac{5}{3}\pi i}$	36	$\langle s_0 s_1 s_5 T \rangle$	C_6	6	
$D_4[\sigma_1]$	$s_1 T s_5 s_0^T$	$1, 1, i, -i, i, -i$	96	$\langle s_1 T s_5 s_0^T \rangle$	C_4	24	

Table 1 continued

Conjugacy class		Centraliser		Elementary part of the centraliser		Index in full centraliser
Carter type	Representative	Eigenvalues	Order	Generators	Structure	
$A_2^2 \times A_1$	$s_0s_6s_5s_1s_2$	$1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, -1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}$	36	$\langle s_0s_6, s_5, s_1s_2 \rangle$	$C_3 \times C_2 \times C_3$	2
$A_3 \times A_1^2$	$s_0s_6s_3s_1s_5$	$1, i, -1, -i, -1, -1$	96	$\langle s_0s_6s_3, s_1, s_5 \rangle$	$C_4 \times C_2 \times C_2$	6
$A_4 \times A_1$	$s_0s_6s_3s_4s_1$	$1, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, e^{\frac{6}{3}\pi i}, e^{\frac{8}{3}\pi i}, -1$	10	$\langle s_0s_6s_3s_4, s_1 \rangle$	$C_5 \times C_2$	
A_5	$s_0s_6s_3s_4s_5$	$1, e^{\frac{1}{3}\pi i}, e^{\frac{2}{3}\pi i}, -1, e^{\frac{4}{3}\pi i}, e^{\frac{5}{3}\pi i}$	12	$\langle s_0s_6s_3s_4s_5 \rangle \times \langle s_1 \rangle$	$C_6 \times C_2$	
D_5	$s_0s_6s_3s_4s_3^{s_2s_4}$	$1, -1, e^{\frac{1}{4}\pi i}, e^{\frac{3}{4}\pi i}, e^{\frac{5}{4}\pi i}, e^{\frac{7}{4}\pi i}$	8	$\langle s_0s_6s_3s_4s_3^{s_2s_4} \rangle$	C_8	
$D_5[a_1]$	$s_0s_6s_3s_4T$	$1, -1, i, -i, e^{\frac{1}{3}\pi i}, e^{\frac{5}{3}\pi i}$	12	$\langle s_0s_6s_3s_4T \rangle$	C_{12}	
A_2^3	$s_0s_6s_1s_2s_5s_4$	$e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}$	648	$\langle s_0s_6, s_1s_2, s_4s_5 \rangle$	$C_3 \times C_3 \times C_3$	24
$A_5 \times A_1$	$s_0s_6s_3s_4s_5s_1$	$e^{\frac{1}{3}\pi i}, e^{\frac{2}{3}\pi i}, -1, e^{\frac{4}{3}\pi i}, e^{\frac{5}{3}\pi i}, -1$	36	$\langle s_0s_6s_3s_4s_5, s_1 \rangle$	$C_6 \times C_2$	3
E_6	$s_1s_2s_3s_4s_5s_6$	$e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, e^{\frac{1}{6}\pi i}, e^{\frac{5}{6}\pi i}, e^{\frac{7}{6}\pi i}, e^{\frac{11}{6}\pi i}$	12	$\langle s_1s_2s_3s_4s_5s_6 \rangle$	C_{12}	
$E_6[a_1]$	$s_1s_2s_3s_4s_5s_6^{s_3}$	$e^{\frac{2}{9}\pi i}, e^{\frac{4}{9}\pi i}, e^{\frac{8}{9}\pi i}, e^{\frac{10}{9}\pi i}, e^{\frac{14}{9}\pi i}, e^{\frac{16}{9}\pi i}$	9	$\langle s_1s_2s_3s_4s_5s_6^{s_3} \rangle$	C_9	
$E_6[a_2]$	$s_6s_2s_0s_1s_4s_3$	$e^{\frac{1}{3}\pi i}, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}, e^{\frac{5}{3}\pi i}, e^{\frac{2}{3}\pi i}, e^{\frac{4}{3}\pi i}$	72	$\langle s_6s_2s_0s_1s_4s_3 \rangle$	C_6	12

2.3 The full centralisers

We begin by stating an elementary lemma which will help to identify elements of the centraliser.

Lemma 2.1 *Let A, B be diagonalisable matrices and let $V_1 \oplus \dots \oplus V_n$ be the eigenspace decomposition for B . If for all but one i , there is an eigenspace of A containing V_i then A, B commute.*

- In the cases $\emptyset, A_1, A_2 \times A_1, A_3 \times A_1, A_4, A_4 \times A_1, A_5, D_5, D_5[a_1], E_6, E_6[a_1]$ the entire centraliser is the elementary part described in the previous section.
- In cases $A_2, A_2 \times A_1^2, A_3$, the representatives are $s_0s_6, s_0s_6s_1s_5$ and $s_0s_6s_3$ respectively, which are all centralised by the element u_1 . In each case the elementary part of the centraliser has index 2 and it is easy to see that this does not contain u_1 . Moreover the action of the u_1 on the elementary part gives the structure of a wreath product as follows:

$$\begin{aligned} Z_W(s_0s_6) &= \langle s_0s_6 \rangle \wr \langle u_1 \rangle \cong C_3 \times (S_3 \wr C_2) \\ Z_W(s_0s_6s_1s_5) &= \langle s_0s_6 \rangle \wr \langle u_1 \rangle \cong C_3 \times (C_2 \wr C_2) \\ Z_W(s_0s_6s_3) &= \langle s_0s_6s_3 \rangle \wr \langle u_1 \rangle \cong C_4 \times (C_2 \wr C_2) \end{aligned}$$

The factors here are all complex reflection groups with the exception of $S_3 \wr C_2$. This group is isomorphic to the affine orthogonal group $O_2^-(\mathbb{F}_3) \times \mathbb{F}_3^2$ which can be embedded as a reflection group in $GL_3(\mathbb{F}_3)$, dual to the embedding as affine orthogonal transformations. Note that this group is also isomorphic to $O_4^+(\mathbb{F}_2)$ which is unique amongst orthogonal groups as it is not generated by reflections of \mathbb{F}_2^4 .

- Similarly in cases $A_1^2, A_2^2, A_2^2 \times A_1$, the representatives $s_0s_1, s_0s_6s_1s_2, s_0s_6s_5s_1s_2$ are centralised by u_2 and the structures are given by:

$$\begin{aligned} Z_W(s_0s_1) &= \langle s_0 \rangle \wr \langle u_2 \rangle \times \langle s_3, s_4, s_5 \rangle \cong (C_2 \wr C_2) \times S_4 \\ Z_W(s_0s_6s_1s_2) &= (\langle s_0s_6 \rangle \wr \langle u_2 \rangle) \times \langle s_4, s_5 \rangle \cong (C_3 \wr C_2) \times S_3 \\ Z_W(s_0s_6s_5s_1s_2) &= (\langle s_0s_6 \rangle \wr \langle u_2 \rangle) \times \langle s_5 \rangle \cong (C_3 \wr C_2) \times C_2 \end{aligned}$$

- In cases A_1^3 with representative $s_0s_1s_5$ and D_4 with representative $s_0s_1s_5T$, the elementary part of the centraliser has index 6. The elements s_0, s_1, s_5 are permuted and T is fixed by the group $\langle u_1, u_2 \rangle$, which therefore lies in both centralisers. In the first case, since $\langle u_1, u_2 \rangle$ also commutes with s_3 , the elementary part $\langle s_0 \rangle \times \langle s_1 \rangle \times \langle s_5 \rangle \times \langle s_3 \rangle$ is normal, and moreover faithfulness of the $\langle u_1, u_2 \rangle$ action implies these two groups have trivial intersection. Therefore

$$Z_W(s_0s_1s_5) = (\langle s_0 \rangle \wr_3 \langle u_1, u_2 \rangle) \times \langle s_3 \rangle \cong (C_2 \wr_3 D_3) \times C_2$$

where the notation $\wr_3 D_3$ indicates the permutation wreath product for the action of D_3 on the three factors.

For the element $s_0s_1s_5T$ the elementary part of the centraliser is simply the cyclic group $\langle s_0s_1s_5T \rangle \cong C_6$, so its intersection with $\langle u_1, u_2 \rangle$ must be central and therefore trivial. Hence we obtain the centraliser as

$$Z_W(s_0s_1s_5T) = \langle s_0s_1s_5T \rangle \times \langle u_1, u_2 \rangle \cong C_6 \times S_3.$$

- In the case A_1^4 with representative $s_0s_1s_5s_3$, the elementary part of the centraliser is the same as it was in the A_1^3 case above. The -1 eigenspace is the space spanned by r_0, r_1, r_5, r_3 , and in particular it contains $r_T = 1/2(r_0 + r_1 + r_5, +r_3)$. Hence the reflection T lies in the centraliser. As noted in Sect. 2, the group $\langle s_0, s_1, s_5, T \rangle$ is a Weyl group of type D_4 , which has index 6 in the centraliser. The group $\langle u_1, u_2 \rangle$ also centralises $s_0s_1s_5s_3$ and normalises $\langle s_0, s_1, s_5, T \rangle$.

It remains to consider the intersection which, since $\langle u_1, u_2 \rangle$ centralises T , must lie in the centraliser $Z_{W(D_4)}(T)$ which has order 16 (see [4]). Explicitly this is the Weyl group of type A_1^4 with roots $r_0 + r_1 - r_T, r_0 + r_5 - r_T, r_1 + r_5 - r_T, r_T$, and the first three of these are permuted by $\langle u_1, u_2 \rangle$, hence the intersection is trivial. We conclude that

$$Z_W(s_0s_1s_5s_3) = \langle s_0, s_1, s_5, T \rangle \times \langle u_1, u_2 \rangle \cong W(D_4) \times S_3.$$

- Case $A_3 \times A_1^2$, representative $s_0s_6s_3s_1s_5$: The elementary part of the centraliser is $\langle s_0s_6s_3 \rangle \times \langle s_1 \rangle \times \langle s_5 \rangle \cong C_4 \times C_2 \times C_2$. This can also be written as $\langle s_0s_6s_3s_1s_5 \rangle \times \langle s_1 \rangle \times \langle s_5 \rangle$, where the first factor is, tautologically, central in the full centraliser. Now we consider the -1 eigenspace for our representative, which is 3-dimensional. It is spanned by the root vectors r_1, r_5 (the -1 eigenspaces of s_1, s_5 respectively) together with the -1 eigenspace for $s_0s_6s_3$, which is spanned by $r_0 + r_3$. While $r_0 + r_3$ is not a root vector, the sum $r_0 + r_3 + r_1 + r_5$ is twice the root vector r_T , hence T belongs to the centraliser. Considering the root system $\{r_1, r_T, r_5\}$ see that the group $\langle s_1, T, s_5 \rangle \cong S_4$ and hence has trivial intersection with $\langle s_0s_6s_3s_1s_5 \rangle$. It follows that the centraliser is

$$Z_W(s_0s_6s_3s_1s_5) = \langle s_0s_6s_3s_1s_5 \rangle \times \langle s_1, T, s_5 \rangle \cong C_4 \times S_4.$$

- Case $A_5 \times A_1$, representative $s_0s_6s_3s_4s_5s_1$: The elementary part of the centraliser is the same as for the A_5 case, $\langle s_0s_6s_3s_4s_5 \rangle \times \langle s_1 \rangle \cong C_6 \times C_2$, and has index 3 in the full centraliser. The -1 eigenspace of the representative is 2 dimensional and is spanned by the -1 eigenvector $r_0 + r_3 + r_5$ of $s_0s_6s_3s_4s_5$ and the -1 eigenvector r_1 of s_1 . As in the previous case T is in the centraliser. We note that $\langle s_1, T \rangle$ is a copy of S_3 which has trivial intersection with $\langle s_0s_6s_3s_4s_5s_1 \rangle$ since $\langle s_1, T \rangle$ has trivial centre. Hence we obtain the centraliser subgroup as

$$Z_W(s_0s_6s_3s_4s_5s_1) = \langle s_0s_6s_3s_4s_5s_1 \rangle \times \langle s_1, T \rangle \cong C_6 \times S_3.$$

The remaining cases are $E_6[a_2]$, $D_4[a_1]$ and A_2^3 . These all correspond to Springer-regular elements, [12], and we can read off their structures from [12, Table 1] along

with the Shephard–Todd classification of complex reflection groups [11, Table VII] as the groups G_5, G_8, G_{25} respectively.

In order to identify the actions on the fixed sets in the maximal torus, we need to identify not just the structure of these groups but their elements which we do as follows.

- Case $D_4[a_1]$, representative $s_1 T s_5 s_0^T$: We choose this representative in the subgroup $\langle s_0, s_1, s_5, T \rangle$ of type D_4 , so that its square $(s_1 T s_5 s_0^T)^2 = s_0 s_1 s_5 s_3$ is the representative of the class of type A_1^4 , and hence its centraliser is a subgroup of index 12 in the A_1^4 centraliser $\langle s_0, s_1, s_5, T \rangle \rtimes \langle u_1, u_2 \rangle \cong W(D_4) \rtimes S_3$ (see above). We consider the standard 4-dimensional representation of $W(D_4)$ corresponding to the generators s_1, T, s_5, s_0 , with roots

$$\begin{aligned} \rho_1 &:= (1, -1, 0, 0), \quad \rho_T := (0, 1, -1, 0), \\ \rho_5 &:= (0, 0, 1, -1), \quad \rho_0 := (0, 0, 1, 1). \end{aligned}$$

The representative $s_1 T s_5 s_0^T$ corresponds to the block diagonal matrix $R_{\pi/2} \oplus R_{\pi/2}$ which has $\pi/2$ rotations in coordinates 1, 2 and 3, 4. We equip \mathbb{R}^4 with the structure of a complex vector space, declaring that the operator $s_1 T s_5 s_0^T$ is multiplication by i . Hence the elements of $W(D_4)$ centralising $s_1 T s_5 s_0^T$ are precisely those which are \mathbb{C} -linear in this space. Specifically the intersection of our centraliser with the subgroup $W(D_4)$ is the Pauli group P generated by the matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

These correspond to the elements $s_1^T s_5^T, T (s_1 s_5 s_0)^{T s_5}$ and $s_0 s_5$ respectively. The group $P = Z_W(s_1 T s_5 s_0^T) \cap W(D_4)$ has index 6 in the full centraliser, hence as the centraliser is contained in $W(D_4) \rtimes S_3$ it fits into an extension

$$1 \rightarrow P \rightarrow Z_W(s_1 T s_5 s_0^T) \rightarrow S_3 \rightarrow 1.$$

We now identify lifts of the generators of S_3 . The conjugation action of u_3 corresponds to interchanging ρ_5, ρ_0 and fixing ρ_1, ρ_T . In particular, the conjugation action of u_3 on $R_{\pi/2} \oplus R_{\pi/2}$ inverts the second rotation, as does the element s_5 , and therefore the element $s_5 u_3$ is in the centraliser of $s_1 T s_5 s_0^T$. It is represented as a complex 2×2 matrix it is the diagonal matrix with entries 1, i : we denote this complex matrix by α .

Now we lift u_1 , conjugation by which corresponds to exchanging ρ_1, ρ_5 while preserving ρ_T, ρ_0 . Conjugating $R_{\pi/2} \oplus R_{\pi/2}$ by this we obtain the matrix

$$\left(\begin{array}{c|c} 0 & R_{\pi/2} \\ \hline R_{\pi/2} & 0 \end{array} \right)$$

which can be conjugated back to $R_{\pi/2} \oplus R_{\pi/2}$ by the transposition (24), given by s_5^T . Hence $s_5^T u_1$ lies in the centraliser of $s_1 T s_5 s_0^T$. This is represented by the complex matrix

$$\beta := \frac{1}{2} \begin{pmatrix} 1 - i & 1 + i \\ 1 + i & 1 - i \end{pmatrix}.$$

To summarise, the centraliser of $s_1 T s_5 s_0^T$ is generated by the Pauli matrices $\sigma_1, \sigma_2, \sigma_3$ along with α, β .

Note that the lifts α and β each have order 4, indeed $\alpha^2 = \sigma_3$ and $\beta^2 = \sigma_1$. Moreover $\alpha \sigma_1 \alpha^{-1} = \sigma_2$, hence the generators α, β are sufficient. The centraliser is thus the group

$$\langle s_5 u_3, s_5^T u_1 \rangle$$

which is faithfully represented as the matrix group $\langle \alpha, \beta \rangle$. These complex reflections generate the symmetry group of the complex polygon 4(96)4 (see Sect. 1.6 of [10]), identifying $Z_W(s_1 T s_5 s_0^T)$ with the complex reflection group G_8 in the Shephard–Todd classification.

- Case A_2^3 , representative $s_0 s_6 s_1 s_2 s_5 s_4$: The elementary part of the centraliser is $\langle s_0 s_6 \rangle \times \langle s_1 s_2 \rangle \times \langle s_5 s_4 \rangle \cong C_3 \times C_3 \times C_3$. This has index 24 in the full centraliser, and it is evident that the missing elements include $\langle u_1, u_2 \rangle$, which permute the direct factors of the elementary part, giving a subgroup of index 4 in the centraliser isomorphic to $C_3 \wr_3 D_3$.

Applying Lemma 2.1 we see that $T s_3$ is in the centraliser. We will show that $\langle s_0 s_6, T s_3, s_5 s_4 \rangle$ is the complex reflection group G_{25} and hence is the full centraliser.

We start with the six dimensional real representation of the Weyl group $W(E_6)$ on the real vector space spanned by the E_6 root system. On this space define the operator

$$J := \frac{1}{\sqrt{3}} (2s_0 s_6 s_1 s_2 s_5 s_4 + I).$$

Note that

$$J^2 = \frac{1}{3} \left(4(s_0 s_6 s_1 s_2 s_5 s_4)^2 + 4s_0 s_6 s_1 s_2 s_5 s_4 + I \right) = -I$$

since $(s_0 s_6 s_1 s_2 s_5 s_4)^2 + s_0 s_6 s_1 s_2 s_5 s_4 + I = 0$. Hence we can define a complex scalar multiplication on the Lie algebra using J as multiplication by i . An element of the Weyl group $W(E_6)$ centralises $s_0 s_6 s_1 s_2 s_5 s_4$ if and only if it commutes with J , i.e. is \mathbb{C} -linear for this structure.

Moreover the elements $s_0 s_6, T s_3, s_5 s_4$ are complex reflections: taking r_0, r_1, r_5 as a basis of this complex space these elements are given by:

$$\alpha_1 = \begin{pmatrix} \zeta & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \alpha_2 = \frac{1}{\sqrt{3}} \begin{pmatrix} \eta & -\bar{\eta} & -\bar{\eta} \\ -\bar{\eta} & \eta & -\bar{\eta} \\ -\bar{\eta} & -\bar{\eta} & \eta \end{pmatrix},$$

$$\alpha_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \zeta \end{pmatrix}$$

where $\eta = e^{\pi i/6}$. These generating reflections satisfy the braid relations

$$\alpha_1\alpha_2\alpha_1 = \alpha_2\alpha_1\alpha_2, \alpha_2\alpha_3\alpha_2 = \alpha_3\alpha_2\alpha_3,$$

and also $\alpha_1\alpha_3 = \alpha_3\alpha_1$. Since the geometry of the fixed hyperplanes of the complex reflections is determined by the above relations, the reflection group is the Shephard–Todd group $G_{25} = \langle s_0s_6, Ts_3, s_5s_4 \rangle$.

- Case $E_6[a_2]$. The representative $s_6s_2s_0^T s_1^T s_4s_3$ squares to the representative $s_0s_6s_1s_2s_5s_4$ above, therefore its centraliser is a subgroup of G_{25} . In the above generators $\alpha_1, \alpha_2, \alpha_3$ we have

$$s_6s_2s_0^T s_1^T s_4s_3 = \alpha_1^2\alpha_2^2\alpha_3\alpha_2^2\alpha_1^2 = \begin{pmatrix} 0 & -\bar{\zeta} & 0 \\ -\bar{\zeta} & 0 & 0 \\ 0 & 0 & -\bar{\zeta} \end{pmatrix}.$$

This evidently commutes with $\langle \alpha_2, \alpha_3 \rangle$ which is which is the binary tetrahedral group $SL_2(3)$, that is G_4 in the Shephard–Todd classification. Its centre is a copy of C_2 generated by the element $(\alpha_1^2\alpha_2^2\alpha_3\alpha_2^2\alpha_1^2)^3 = (\alpha_3^2\alpha_2)^2$, hence $\langle (\alpha_1^2\alpha_2^2\alpha_3\alpha_2^2\alpha_1^2)^2 \rangle$ intersects trivially with $\langle \alpha_2, \alpha_3 \rangle$.

The group $C_3 \times SL_2(3)$ is the Shephard–Todd group G_5 , so we have the found the full centraliser

$$Z_W(s_6s_2s_0^T s_1^T s_4s_3) = \langle (\alpha_1^2\alpha_2^2\alpha_3\alpha_2^2\alpha_1^2)^2 \rangle \times \langle \alpha_2, \alpha_3 \rangle \cong C_3 \times SL_2(3).$$

3 Fixed sets and their quotients: the simply connected type

In this section we will determine the fixed sets of each conjugacy class representative in the standard maximal torus \mathcal{T} in the simply connected form of the Lie group of type E_6 . We will analyse in detail the actions of the centralisers on the fixed sets for each conjugacy class representative. In each case we will see that the quotient, which we refer to as a *sector*, has the homotopy type of a finite union of points or a finite union of circles. These results are summarised in Table 2.

Notation

- \mathbb{T}^m the m -torus
- Δ_k the k -simplex (not necessarily equilateral)
- $\Delta_k \tilde{\times} S^1$ a twisted bundle over the circle with fibre a k -simplex
- $SP^2(X)$ the 2-fold symmetric product of a space X
- C_m the cyclic group of order m
- \mathcal{D}_3 the dihedral group of order 6 generated by the special elements u_1, u_2
- $C_2 \wr \mathcal{D}_3$ is the permutation wreath product $C_2^3 \rtimes \mathcal{D}_3$
- \mathcal{D}_4 a dihedral group of order 8 of the form $\langle s_i \rangle \wr \langle u_j \rangle$
- S_m a symmetric group on m letters generated by a subset of the 36 root reflections
- V_4 a Klein 4 group.
- G_5, G_8, G_{25}, G_{28} exceptional complex reflection groups (cf. [11])
- rep is an abbreviation for the chosen representative of the conjugacy class.
- \mathcal{Z} the centre of the simply connected Lie group of type E_6 .

Table 2 Centralisers, fixed sets and quotients

Conjugacy class		Centraliser	Fixed set	Quotient
Type	Representative	Structure	Generators	
\emptyset	e	$W(E_6)$	$\langle s_1, \dots, s_6 \rangle$	\mathbb{T}^6
A_1	s_0	$C_2 \times W(A_5)$	$\langle s_0 \rangle \wr \langle s_1, \dots, s_5 \rangle$	\mathbb{T}^5
A_1^2	$s_0 s_1$	$D_4 \times S_4$	$\langle (s_0) \wr (u_2) \rangle \times \langle s_3, s_4, s_5 \rangle$	\mathbb{T}^4
A_2	$s_0 s_6$	$C_3 \times (S_3 \wr C_2)$	$\langle s_0 s_6 \rangle \times \langle (s_1, s_2) \wr (u_1) \rangle$	\mathbb{T}^4
A_1^3	$s_0 s_1 s_5$	$(C_2 \wr C_3) \times C_2$	$\langle (s_0) \wr (u_1, u_2) \rangle \times \langle s_3 \rangle$	\mathbb{T}^3
$A_2 \times A_1$	$s_0 s_6 s_1$	$C_3 \times C_2 \times S_3$	$\langle s_0 s_6, s_1 \rangle \times \langle s_4, s_5 \rangle$	\mathbb{T}^3
A_3	$s_0 s_6 s_3$	$C_4 \times D_4$	$\langle s_0 s_6 s_3 \rangle \times \langle (s_1) \wr (u_1) \rangle$	\mathbb{T}^3
A_1^4	$s_0 s_1 s_5 s_3$	G_{28}	$\langle s_0, s_1, s_5, T \rangle \rtimes \langle (u_1, u_2) \rangle$	$\mathbb{T}^2 \times V_4$
$A_2 \times A_1^2$	$s_0 s_6 s_1 s_5$	$C_3 \times D_4$	$\langle s_0 s_6 \rangle \times \langle (s_1) \wr (u_1) \rangle$	\mathbb{T}^2
A_2^2	$s_0 s_6 s_1 s_2$	$(C_3 \wr C_2) \times S_3$	$\langle (s_0 s_6) \wr (u_2) \rangle \times \langle s_4, s_5 \rangle$	$\mathbb{T}^2 \times \mathcal{Z}$
$A_3 \times A_1$	$s_0 s_6 s_3 s_1$	$C_4 \times V_4$	$\langle s_0 s_6 s_3 \rangle \times \langle s_1 \rangle \times \langle s_5 \rangle$	\mathbb{T}^2
A_4	$s_0 s_6 s_3 s_4$	$C_5 \times C_2$	$\langle s_0 s_6 s_3 s_4 \rangle \times \langle s_1 \rangle$	\mathbb{T}^2
D_4	$s_0 s_1 s_5 T$	$C_6 \times D_3$	$\langle s_0 s_1 s_5 T \rangle \times \langle (u_1, u_2) \rangle$	\mathbb{T}^2
$D_4[st_1]$	$s_1 T s_5 s_0 T$	G_8	$\langle s_5 u_3, s_5^T u_1 \rangle$	\mathbb{T}^2

Table 2 continued

Conjugacy class		Centraliser		Fixed set	Quotient
Type	Representative	Structure	Generators		
$A_2^2 \times A_1$	$s_0s_6s_5s_1s_2$	$(C_3 \wr C_2) \times C_2$	$(s_0s_6) \wr (u_2) \times (s_5)$	$\mathbb{T}^1 \times \mathcal{Z}$	$\Delta^1 \times \mathcal{Z}$
$A_3 \times A_1^2$	$s_0s_6s_3s_1s_5$	$C_4 \times S_4$	$(\text{rep}) \times (s_1, T, s_5)$	$\mathbb{T}^1 \times V_4$	$\mathbb{T}^1 \sqcup \mathbb{T}^1$
$A_4 \times A_1$	$s_0s_6s_3s_4s_1$	$C_5 \times C_2$	$(s_0s_6s_3s_4s_5s_1)$	\mathbb{T}^1	\mathbb{T}^1
A_5	$s_0s_6s_3s_4s_5$	$C_6 \times S_2$	$(s_0s_6s_3s_4s_5) \times (s_1)$	$\mathbb{T}^1 \times \mathcal{Z}$	$\Delta^1 \times \mathcal{Z}$
D_5	$s_0s_6s_3s_4s_5s_2s_4$	C_8	$(s_0s_6s_3s_4s_5s_2s_4)$	\mathbb{T}^1	\mathbb{T}^1
$D_5[a_1]$	$s_0s_6s_3s_4T$	C_{12}	$(s_0s_6s_3s_4T)$	\mathbb{T}^1	\mathbb{T}^1
A_3^2	$s_0s_6s_1s_2s_5s_4$	G_{25}	$(s_0s_6, T s_3, s_5s_4)$	$C_3 \times C_3 \times \mathcal{Z}$	Four points
$A_5 \times A_1$	$s_0s_6s_3s_4s_5s_1$	$C_6 \times S_3$	$(\text{rep}) \times (s_1, T)$	$V_4 \times \mathcal{Z}$	$\mathcal{Z} \sqcup \mathcal{Z}$
E_6	$s_1s_2s_3s_4s_5s_6$	C_{12}	$(s_1s_2s_3s_4s_5s_6)$	\mathcal{Z}	\mathcal{Z}
$E_6[a_1]$	$s_1s_2s_3s_4s_5s_6s_3$	C_9	$(s_1s_2s_3s_4s_5s_6s_3)$	\mathcal{Z}	\mathcal{Z}
$E_6[a_2]$	$s_6s_2s_0s_1s_4s_3$	G_5	$((\text{rep})^2, T s_3, s_5s_4)$	\mathcal{Z}	\mathcal{Z}

The fixed sets are identified as topological spaces using Theorem 4 combined with the following well known lemma. Recall that $\bar{l}(w)$ denotes the length of the element w with respect to the generating set of *all* root reflections.

Lemma 3.1 ([4, Lemma 2]) $\bar{l}(w)$ is the number of eigenvalues with multiplicity of w which are not equal to 1.

Using this lemma, for each $w \in W$ the torus T_1^w has dimension $6 - \bar{l}(w)$, where $\bar{l}(w)$ denotes the word length of w with respect to the generating set consisting of all root reflections in W .

Combining this with Theorem 5 determines the topology of the fixed sets in both the simply connected and adjoint forms:

Theorem 7 For each $w \in W$ the fixed sets T^w and $(T^\vee)^w$ are both homeomorphic to the disjoint union of g copies of a $6 - \bar{l}(w)$ -torus where g is the greatest common divisor of the $r \times r$ -minors of the matrix of $I - w$ and r is the rank of $I - w$.

We now turn to the actions of centralisers on the corresponding fixed sets.

Since the action of the Weyl group W on the maximal torus T is given by conjugation, all of the fixed sets contain the centre Z of the Lie group as a pointwise fixed subset. For example in the three elliptic cases of Carter-type $E_6, E_6[a_1], E_6[a_2]$, the fixed sets, and hence the quotients are identified with the centre Z .

In cases $A_4 \times A_1, D_5$ and $D_5[a_1]$ the fixed set consists of a single circle containing the (pointwise fixed) centre so any element of W preserving the circle must fix it pointwise. In particular the centralisers pointwise fix the circle identifying the quotient with the fixed set.

3.1 Alternative coordinate systems

To understand the actions of centralisers on the fixed sets it will be helpful to consider alternative coordinate systems on the torus.

- (1) \mathfrak{su}_6 coordinates: In many cases the fixed set will lie inside the 5-torus $(\alpha, \beta, \gamma, \delta, \epsilon, 1)$ which is the maximal torus corresponding to the A_5 root system given by $\{r_1, \dots, r_5\}$. When considering the A_n case it is often helpful to use coordinates where the roots are $(1, -1, 0, \dots, 0), (0, 1, -1, 0, \dots, 0)$ etc. and for which the maximal torus is correspondingly given by tuples with product 1 (as is the case for the standard maximal torus in SU_{n+1}). We obtain these coordinates on the above 5-torus by the change of coordinates:

$$(\alpha, \beta, \gamma, \delta, \epsilon, 1) \mapsto (\alpha, \beta\alpha^{-1}, \gamma\beta^{-1}, \delta\gamma^{-1}, \epsilon\delta^{-1}, \epsilon^{-1}).$$

Under this identification the A_5 Weyl group generated by s_1, \dots, s_5 acts by permuting the coordinates.

- (2) \mathfrak{su}_3^3 coordinates: Another alternative coordinate system is obtained by considering the roots $r_1, r_2, r_5, r_4, r_0, r_6$ as a basis for the Lie algebra. The advantage of this approach is that the dihedral group $\langle u_1, u_2 \rangle$ acts by signed permutations on the pairs $(r_1, r_2), (r_5, r_4)$ and (r_0, r_6) . Moreover the actions of s_1, s_5, s_0 will remain

relatively simple in our new coordinates. The omitted root vector r_3 can be obtained from

$$\begin{aligned} 3r_3 &= -r_0 - 2r_6 - r_1 - 2r_2 - r_5 - 2r_4 \\ &= -r_0 + r_6 - r_1 + r_2 - r_5 + r_4 - 3(r_6 + r_2 + r_4) \end{aligned}$$

hence the root lattice is the refinement of the lattice generated by $r_1, r_2, r_5, r_4, r_0, r_6$ by the addition of the vector $\frac{1}{3}(-r_0 + r_6 - r_1 + r_2 - r_5 + r_4)$.

Motivated by this we take the ordered basis $r_1, -r_2, r_5, -r_4, r_0, -r_6$ for the Lie algebra, for which coordinates the lattice is \mathbb{Z}^6 extended by the vector $(\frac{1}{3}, \dots, \frac{1}{3})$. The maximal torus is given by \mathbb{T}^6/C_3 where the C_3 acts by multiplication by (ω, \dots, ω) for ω a cube root of unity. We denote these new coordinates $(\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta')$.

The action of (u_1, u_2) on \mathbb{T}^6/C_3 is to permute and invert pairs of coordinates as follows.

$$\begin{aligned} u_1 : (\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta') \\ \mapsto ((\epsilon')^{-1} : (\delta')^{-1} : (\alpha')^{-1} : (\beta')^{-1} : (\eta')^{-1} : (\zeta')^{-1}) \\ u_2 : (\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta') \\ \mapsto ((\eta')^{-1} : (\zeta')^{-1} : (\epsilon')^{-1} : (\delta')^{-1} : (\alpha')^{-1} : (\beta')^{-1}) \\ u_3 : (\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta') \\ \mapsto ((\alpha')^{-1} : (\beta')^{-1} : (\eta')^{-1} : (\zeta')^{-1} : (\epsilon')^{-1} : (\delta')^{-1}) \end{aligned}$$

The conversion from these coordinates to the standard r_1, \dots, r_6 coordinates is given by the map

$$\begin{aligned} (\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta') \\ \mapsto (\eta'(\alpha')^{-1}, (\eta')^2\beta', (\eta')^3, (\eta')^2\delta', \eta'(\epsilon')^{-1}, (\eta')^2\zeta') \end{aligned}$$

which is well defined on C_3 -cosets.

Now applying s_1 changes the first of these coordinates to

$$(\eta'(\alpha')^{-1})^{-1}(\eta')^2\beta = \eta'\alpha'\beta'$$

while leaving the other coordinates unchanged. Thus in the new coordinates s_1 takes $(\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta')$ to $((\alpha'\beta')^{-1} : \beta' : \epsilon' : \delta' : \eta' : \zeta')$.

Similarly s_5 changes ϵ' to $(\epsilon'\delta')^{-1}$ leaving the other coordinates unchanged, and s_0 changes η' to $(\eta'\zeta')^{-1}$ leaving the other coordinates unchanged.

The action of T in the new coordinates is much simpler than in the original coordinates. In standard coordinates T is given by the matrix:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & -1 & 1 & -1 \\ 1 & -1 & 2 & -1 & 1 & -1 \\ 1 & -1 & 1 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & -1 & 1 & -1 & 1 & 0 \end{pmatrix}$$

Applying this to $(\eta'(\alpha')^{-1}, (\eta')^2\beta', (\eta')^3, (\eta')^3\delta', \eta'(\epsilon')^{-1}, (\eta')^2\zeta)$, and choosing ϕ to be a cube root of $(\alpha'\beta'\epsilon'\delta'\zeta'\eta')^{-1}$ we obtain

$$\begin{aligned} &(\eta'(\alpha')^{-1}, \eta'(\alpha'\epsilon'\delta'\zeta')^{-1}, (\eta')^2(\alpha'\beta'\epsilon'\delta'\zeta')^{-1}, \\ &\quad \eta'(\alpha'\beta'\epsilon'\zeta')^{-1}, \eta'(\epsilon')^{-1}, \eta'(\alpha'\beta'\epsilon'\delta')^{-1}) \\ &= (\eta'(\alpha')^{-1}, (\eta')^2\beta'\phi^3, (\eta')^3\phi^3, (\eta')^2\delta'\phi^3, \eta'(\epsilon')^{-1}, (\eta')^2\zeta'\phi^3) \\ &= ((\eta'\phi)(\alpha'\phi)^{-1}, (\eta'\phi)^2\beta'\phi, (\eta'\phi)^3, \\ &\quad (\eta'\phi)^2\delta'\phi, (\eta'\phi)(\epsilon'\phi)^{-1}, (\eta'\phi)^2\zeta'\phi) \end{aligned}$$

whence the action of T in the new coordinates simply multiplies each coordinate by ϕ . Since ϕ was well-defined up to a cube root of unity this makes the action well-defined on \mathbb{T}^6/C_3 .

3.2 The fixed sets and their quotients

We now proceed to compute the fixed sets and sectors.

- (1) Case \emptyset : The fixed set is \mathcal{T} and the centraliser is $W(E_6)$, which acts on \mathcal{T} with quotient a 6-simplex.
- (2) Case A_1 : The fixed variety is a single 5-torus $(\alpha, \beta, \gamma, \delta, \epsilon, 1)$. This is canonically identified with the maximal torus for the Lie group of type A_5 . The centraliser of s_0 is the direct product of $\langle s_0 \rangle$ acting trivially with the Weyl group $\langle s_1, s_2, s_3, s_4, s_5 \rangle$ of type A_5 acting on the A_5 maximal torus. The quotient of the fixed set by the centraliser is a 5-simplex.
- (3) Case A_1^2 : The fixed variety is a single 4-torus $(\alpha, \alpha^2, \gamma, \delta, \epsilon, 1)$ inside the 5-torus where $\zeta = 1$, hence we can use \mathfrak{su}_6 coordinates as above. The fixed set of our element is then identified with the subset of the SU_6 -torus

$$\begin{aligned} &\{(\alpha, \alpha, \gamma'', \delta'', \epsilon'', \zeta'') : \alpha, \gamma'', \delta'', \epsilon'', \zeta'' \in \mathbb{T}^1 \\ &\quad \text{and } \alpha^2\gamma''\delta''\epsilon''\zeta'' = 1\}. \end{aligned} \tag{1}$$

The centraliser is the direct product of $\langle s_0 \rangle \wr \langle u_2 \rangle$ with the symmetric group $S_4 = \langle s_3, s_4, s_5 \rangle$, where s_0 acts trivially and S_4 acts by permutation of the last four coordinates. First we factor out the action of the S_4 which we will

do in logarithmic coordinates $(x_1, x_1, x_3, \dots, x_6)$ subject to the constraint $x_1 + x_1 + x_3 + x_4 + x_5 + x_6 = 0$. We take x_1 to vary from $-\frac{1}{2}$ to $\frac{1}{2}$. In these coordinates the elements of the S_4 factor act by permuting the coordinates x_3, \dots, x_6 , and have fundamental domain D given by

$$D = \{(x_1, x_1, x_3, \dots, x_6) : x_1 \in [-\frac{1}{2}, \frac{1}{2}], 2x_1 + x_3 + x_4 + x_5 + x_6 = 0, x_3 \leq x_4 \leq x_5 \leq x_6 \leq x_3 + 1\}.$$

See [6, Eq. 11] for the details.

As noted in Example 7.2 *ibid* the quotient of the 4-torus (1) under the action of this S_4 is an orientable twisted 3-simplex bundle over the circle parameterised by α with monodromy defined by

$$\left(-\frac{1}{2}, -\frac{1}{2}, x_3, x_4, x_5, x_6\right) \mapsto \left(\frac{1}{2}, \frac{1}{2}, x_5 - 1, x_6 - 1, x_3, x_4\right).$$

Now we consider the action of the element u_2 . Since this commutes with the S_4 we can act directly on the quotient bundle described above. Recall that u_2 negates r_3, r_4, r_5 and exchanges r_0 with $-r_1$ and r_2 with $-r_6$, so the action of u_2 on the fixed set is:

$$(\alpha, \alpha^2, \gamma, \delta, \epsilon, 1) \mapsto (\alpha, \alpha^2, \alpha^3\gamma^{-1}, \alpha^2\delta^{-1}, \alpha\epsilon^{-1}, 1).$$

Changing coordinates as above this becomes

$$(\alpha, \alpha, \gamma'', \delta'', \epsilon'', \zeta'') \mapsto (\alpha, \alpha, (\alpha\gamma'')^{-1}, (\alpha\delta'')^{-1}, (\alpha\epsilon'')^{-1}, (\alpha\zeta'')^{-1}).$$

In logarithmic coordinates this is

$$\begin{aligned} &(x_1, x_1, x_3, x_4, x_5, x_6) \\ &\mapsto (x_1, x_1, -x_1 - x_3, -x_1 - x_4, -x_1 - x_5, -x_1 - x_6). \end{aligned}$$

While this does not preserve the fundamental domain D for the S_4 action, composing with the permutation that reverses the last 4 coordinates returns us to D , so we have the map

$$\begin{aligned} &(x_1, x_1, x_3, x_4, x_5, x_6) \\ &\mapsto (x_1, x_1, -x_1 - x_6, -x_1 - x_5, -x_1 - x_4, -x_1 - x_3). \end{aligned}$$

In the $x_1 = 0$ fibre the vertex set of the simplex is

$$\begin{aligned} v_0(0) &= (0, 0, 0, 0, 0, 0), \\ v_1(0) &= \left(0, 0, -\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4}, \frac{3}{4}\right), \end{aligned}$$

$$v_2(0) = \left(0, 0, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right),$$

$$v_3(0) = \left(0, 0, -\frac{3}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right)$$

and in a general fibre the vertex set $\{v_i(x_1)\}$ is given by translating each of these vectors by $(x_1, x_1, -\frac{x_1}{2}, -\frac{x_1}{2}, -\frac{x_1}{2}, -\frac{x_1}{2})$

In each simplex two of the vertices are fixed by the u_2 action, namely $v_0(x_1)$ and $v_2(x_1)$ while the other two are interchanged. We think of the fibres as given by the join of two 1-simplices corresponding to the edge $e_0 = [v_0(x_1), v_2(x_1)]$ and the edge $e_1 = [v_1(x_1), v_3(x_1)]$.

The action of u_2 on the simplex bundle preserves the fibres and moreover preserves the join structure on each, yielding again a simplex bundle with fibres given by the join of e_0 with the quotient of e_1 by inversion.

The quotient of the fixed set by the centraliser is therefore a simplex bundle over the circle, where the monodromy is given by the reflection of the 3-simplex which inverts the e_0 factor and fixes the e_1 factor. This yields a Seifert fibred 4-orbifold: the base space is the quotient of the join of e_0 and e_1 by the Klein 4 group inverting both of these edges, [9].

- (4) Case A_2 : The fixed variety is the 4-torus $(\alpha, \beta, 1, \delta, \epsilon, 1)$ inside the standard 5-torus from case 2. We again make use of \mathfrak{su}_6 coordinates. Under this identification the A_5 Weyl group $\langle s_1, \dots, s_5 \rangle$ acts by permuting the coordinates. The fixed set of our element is then identified with the subset $\{(\alpha, \beta\alpha^{-1}, \beta^{-1}, \delta, \epsilon\delta^{-1}, \epsilon^{-1}) : \alpha, \beta, \delta, \epsilon \in \mathbb{T}^1\}$. Since the first three coordinates and the last three coordinates each multiply to 1 this is naturally a product of two maximal tori of type A_2 .

The centraliser has the form $\langle s_0s_6 \rangle \times (\langle s_1, s_2 \rangle \wr \langle u_1 \rangle) \cong C_3 \times (S_3 \wr C_2)$ with the element s_0s_6 acting trivially. The subgroups $\langle s_1, s_2 \rangle$ and $\langle s_5, s_4 \rangle = \langle s_1, s_2 \rangle^{u_1}$ act as the Weyl groups on these 2-torus factors so the direct product $S_3 \times S_3$ acts in the natural way on $\mathbb{T}^2 \times \mathbb{T}^2$ with quotient a product of two equilateral triangles. The element u_1 acts to swap these, so the quotient is the symmetric product of two copies of an equilateral triangle.

- (5) Case A_1^3 : The fixed set is a single 3-torus $(\alpha, \alpha^2, \gamma, \epsilon^2, \epsilon, 1)$, and is the intersection of the fixed sets of each of the elements s_0, s_1, s_5 . The centraliser is the direct product of $\langle s_3 \rangle$ with the wreath product $\langle s_0 \rangle \wr \langle u_1, u_2 \rangle$, where the conjugates of $\langle s_0 \rangle$ are $\langle s_1 \rangle, \langle s_5 \rangle$. Since these act trivially the centraliser acts as $\langle u_1, u_2 \rangle \times \langle s_3 \rangle$.

Recall that u_1 exchanges r_1 with $-r_5, r_2$ with $-r_4$ and negates r_3, r_6 , so on the fixed set the action of u_1 exchanges the values of α and ϵ^{-1} , and inverts γ . On the other hand u_2 is given by

$$(\alpha, \alpha^2, \gamma, \epsilon^2, \epsilon, 1) \mapsto (\alpha, \alpha^2, \alpha^3\gamma^{-1}, \alpha^2\epsilon^{-2}, \alpha\epsilon^{-1}, 1),$$

see A_1^2 case. We decompose our 3-torus as the product

$$\mathbb{T}^2 \times \mathbb{T}^1 = \{(\alpha, \alpha^2, \alpha\epsilon, \epsilon^2, \epsilon, 1)\} \times \{(1, 1, \gamma, 1, 1, 1)\}.$$

and note that both factors are preserved by the actions of u_1, u_2 . Moreover s_3 pointwise fixes the first factor and inverts the second. Since u_1, u_2 also invert the second factor, the products u_1s_3, u_2s_3 (which also generate a dihedral group) pointwise fix the second factor, and act on the first as

$$\begin{aligned} u_1s_3 : (\alpha, \epsilon) &\mapsto (\epsilon^{-1}, \alpha^{-1}) \\ u_2s_3 : (\alpha, \epsilon) &\mapsto (\alpha, \alpha\epsilon^{-1}) \end{aligned}$$

which is the standard dihedral action with quotient an equilateral triangle. Hence the quotient of the fixed set of $s_0s_1s_5$ by the centraliser is

$$\mathbb{T}^2 / \langle u_1s_3, u_2s_3 \rangle \times \mathbb{T}^1 / \langle s_3 \rangle = \Delta^2 \times I.$$

- (6) Case $A_2 \times A_1$: The fixed set is the single 3-torus $(\alpha, \alpha^2, 1, \delta, \epsilon, 1)$ and the centraliser is $\langle s_0s_6 \rangle \times \langle s_1 \rangle \times \langle s_4, s_5 \rangle \cong C_3 \times C_2 \times S_3$. The first two factors act trivially so we are left with a residual action of $S_3 = \langle s_4, s_5 \rangle$. These generators act as the matrices $\begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix}$ on the free parameters (δ, ϵ) , fixing the parameter α . The non trivial part of the action is therefore the standard dihedral action on a 2-torus with quotient an equilateral triangle, crossed with the trivial action on the remaining circle. Hence the quotient is $\mathbb{T}^1 \times \Delta^2$.
- (7) Case A_3 : The fixed set is a single 3-torus $(\alpha, \beta, 1, \beta^{-1}, \epsilon, 1)$ and the centraliser of $s_0s_6s_3$ has the structure $\langle s_0s_6s_3 \rangle \times (\langle s_1 \rangle \wr \langle u_1 \rangle) \cong C_4 \times (C_2 \wr C_2)$, so we need to compute the quotient by the action of the wreath product $\langle s_1 \rangle \wr \langle u_1 \rangle$. We begin by identifying the quotient under the action of s_1 and $s_1^{u_1} = s_5$. The elements s_1 and s_5 both act trivially on the β factor of the 3-torus, with s_1 acting on the α, β subtorus by the matrix $\begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}$ and s_5 acting on the β, ϵ subtorus via $\begin{pmatrix} 1 & 0 \\ -1 & -1 \end{pmatrix}$.

We introduce logarithmic coordinates for β , i.e. set $\beta = e^{2\pi ix}$. The 3-torus is then

$$\{(\alpha, x, \epsilon) : \alpha, \epsilon \in \mathbb{T}^1, x \in [0, 1]\} / (\alpha, 1, \epsilon) \sim (\alpha, 0, \epsilon).$$

Now introduce coordinates: $\alpha' = \alpha e^{-\pi ix}$, and $\epsilon' = \epsilon e^{\pi ix}$. Under this change of coordinates $(\alpha, 1, \epsilon)$ goes to $(\alpha', 1, \epsilon') = (-\alpha, 1, -\epsilon)$ so in these coordinates the torus is:

$$\{(\alpha', x, \epsilon') : \alpha', \epsilon' \in \mathbb{T}^1, x \in [0, 1]\} / (\alpha', 1, \epsilon') \sim (-\alpha', 0, -\epsilon').$$

The action of s_1 fixes $\beta = e^{2\pi ix}$ and ϵ , and hence also fixes ϵ' , while

$$\alpha \mapsto \alpha^{-1}\beta = (\alpha' e^{\pi ix})^{-1} e^{2\pi ix} = (\alpha')^{-1} e^{\pi ix}.$$

Hence in primed coordinates, s_1 takes α' to $(\alpha')^{-1} e^{\pi ix} e^{-\pi ix} = (\alpha')^{-1}$.

Similarly s_5 fixes α' and x and takes ϵ to $\epsilon^{-1}\beta^{-1} = (\epsilon')^{-1}e^{-\pi ix}$. Hence $\epsilon' \mapsto (\epsilon')^{-1}$.

Setting $I = \{\phi \in \mathbb{T}^1 : \text{Im } \phi \geq 0\}$ which we identify in the natural way as the quotient of \mathbb{T}^1 by the map $\phi \mapsto \phi^{-1}$, the quotient of the torus by $\langle s_1, s_5 \rangle$ is given by

$$\{(\alpha', x, \epsilon') : \alpha', \epsilon' \in I, x \in [0, 1]\} / (\alpha', 1, \epsilon') \sim ((-\alpha')^{-1}, 0, (-\epsilon')^{-1}).$$

This is a square bundle over the circle, whose monodromy map is rotation by π .

Now the action of u_1 preserves β (and hence x), and exchanges α with ϵ^{-1} . In primed coordinates

$$u_1 : (\alpha', x, \epsilon') \mapsto ((\epsilon')^{-1}, x, (\alpha')^{-1})$$

so descending to the square bundle the action simply exchanges α' with ϵ' . The quotient of each fibre is a right-isosceles triangle, hence the quotient of the fixed set by the centraliser is a triangle bundle over the circle with monodromy map reflecting the triangle. This is a Seifert bundle over the $(4, 4, 2)$ triangle, [9].

- (8) Case A_1^4 : The fixed set is the product of a 2-torus indexed by α, ϵ with a Klein 4 group generated by independent square-roots of 1 denoted $\eta = \gamma(\zeta\alpha\epsilon)^{-1}$ and ζ . This gives coordinates $(\alpha, \zeta\alpha^2, \eta\zeta\alpha\epsilon, \zeta\epsilon^2, \epsilon, \zeta)$. The centraliser is

$$\langle s_0, s_1, s_5, T \rangle \rtimes \langle u_1, u_2 \rangle \cong W(D_4) \rtimes S_3.$$

We will now change to su_3^3 coordinates in which the actions of the generators of the centraliser are all relatively straightforward.

Recall that the tuple $(\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta')$ in new coordinates corresponds to $(\eta'(\alpha')^{-1}, (\eta')^2\beta', (\eta')^3, (\eta')^2\delta', \eta'(\epsilon')^{-1}, (\eta')^2\zeta')$ in the original coordinates so in the new coordinates a point of the fixed set must satisfy

$$\begin{aligned} (\eta')^2\beta' &= \zeta(\eta'(\alpha')^{-1})^2 \\ (\eta')^3 &= \eta\zeta(\eta'(\alpha')^{-1})(\eta'(\epsilon')^{-1}) \\ (\eta')^2\delta' &= \zeta(\eta'(\epsilon')^{-1})^2 \\ (\eta')^2\zeta' &= \zeta \end{aligned}$$

where η, ζ are ± 1 giving the element of the Klein 4 group and the other variables lie in the circle. This simplifies to

$$\begin{aligned} \beta' &= \zeta(\alpha')^{-2} \\ \alpha'\epsilon'\eta' &= \zeta\eta \\ \delta' &= \zeta(\epsilon')^{-2} \end{aligned}$$

$$\zeta' = \zeta(\eta')^{-2}$$

so for (η, ζ) in the Klein 4 group V_4 , the corresponding component of the fixed set is parametrised as

$$\{(\alpha', \epsilon', \zeta') : \alpha'\epsilon'\eta' = \zeta\eta\}/\langle(\omega, \omega, \omega)\rangle$$

where ω is a non-trivial cube root of 1.

In these coordinates the subgroup $\langle u_1, u_2 \rangle$ simply acts by the signed permutations:

$$\begin{aligned} u_1 : (\alpha' : \epsilon' : \zeta') &\mapsto ((\epsilon')^{-1} : (\alpha')^{-1} : (\zeta')^{-1}) \\ u_2 : (\alpha' : \epsilon' : \zeta') &\mapsto ((\zeta')^{-1} : (\epsilon')^{-1} : (\alpha')^{-1}) \end{aligned}$$

and preserve η, ζ . As noted in Sect. 2.1 this action on the PSU_3 torus is equivariantly isomorphic to the permutation action on the SU_3 torus, for which the quotient is an equilateral triangle.

The element T acts by multiplication by ϕ where ϕ is a cube root of

$$(\alpha'\zeta(\alpha')^{-2}\epsilon'\zeta(\epsilon')^{-2}\eta'\zeta(\eta')^{-2})^{-1} = \alpha'\epsilon'\eta'\zeta = \eta.$$

Since $\eta = \pm 1$ the action simply multiplies the projective coordinates $\alpha', \epsilon', \eta'$ by η and hence as $\alpha'\epsilon'\eta' = \zeta\eta$ the coordinate ζ is also multiplied by η .

We now turn to the actions of s_1, s_5 and s_0 . It suffices to consider the element s_1 as the group $\langle u_1, u_2 \rangle$ conjugates s_1, s_5 and s_0 . For a general point we have:

$$s_1 : (\alpha' : \beta' : \epsilon' : \delta' : \eta' : \zeta') \rightarrow ((\alpha'\beta')^{-1} : \beta' : \epsilon' : \delta' : \eta' : \zeta')$$

so applying this to an element $(\alpha' : \zeta(\alpha')^{-2} : \epsilon' : \zeta(\epsilon')^{-2} : \eta' : \zeta(\eta')^{-2})$ we obtain

$$(\zeta\alpha' : \zeta(\zeta\alpha')^{-2} : \epsilon' : \zeta(\epsilon')^{-2} : \eta' : \zeta(\eta')^{-2})$$

Since the product of the coordinates $\alpha'\epsilon'\zeta'$ changes by a factor of ζ , this means that η is changed by ζ .

In the identity component $\eta = \zeta = 1$, so T, s_1 (and hence also s_5, s_0) act trivially. Hence $\mathcal{T}_1^w/Z_W(w)$ is the quotient of the PSU_3 torus

$$\{(\alpha', \epsilon', \zeta') : \alpha'\epsilon'\zeta' = 1\}/C_3$$

by the signed permutation action of $\langle u_1, u_2 \rangle$. As noted above this can be identified with the quotient of the SU_3 torus by its Weyl group, giving an equilateral triangle.

Turning to the components indexed by the non-identity elements of the Klein 4 group we see that these are permuted transitively by $\langle s_1, T \rangle$, hence there is

just one more component in the quotient. We obtain this by taking the quotient of the $(\eta, \zeta) = (1, -1)$ component by its stabiliser in the centraliser, noting that this component is fixed pointwise by T . This component is parametrised by

$$\{(\alpha', \epsilon', \zeta') : \alpha'\epsilon'\zeta' = -1\}/C_3,$$

and we now determine its stabiliser.

As noted above the elements u_1, u_2 preserve all the components so we begin by considering the action of the subgroup $W(D_4)$ on the set of the three non-identity components. This action gives a map from $W(D_4) = \langle s_1, s_5, s_0, T \rangle$ to the permutation group S_3 in which s_1, s_5, s_0 all map to one transposition and T maps to another. Thus the component stabiliser in $W(D_4)$ is generated by T along with the kernel of this map. Clearly the elements $s_1s_5, s_0s_5, (s_1s_5)^T$ and $(s_0s_5)^T$ lie in the kernel. We note that $s_1s_5(s_0s_5)^T$ has order 4 (it corresponds to a pair of rotations by $\pi/2$ in the standard representation of $W(D_4)$) hence $\langle s_1s_5, (s_0s_5)^T \rangle$ is a dihedral group of order 8, and similarly for $\langle s_0s_5, (s_1s_5)^T \rangle$. All the other pairs from $s_1s_5, s_0s_5, (s_1s_5)^T$ and $(s_0s_5)^T$ have product of order 2 and hence pairwise commute. The intersection of the two dihedral groups is precisely the centre of each group (which has order 2) and hence together they generate a group of order 32 which is thus the whole of the kernel. Hence the component stabiliser in $W(D_4)$ is the semidirect product

$$\langle s_1s_5, s_0s_5, (s_1s_5)^T, (s_0s_5)^T \rangle \rtimes \langle T \rangle = \langle s_1s_5, s_0s_5, T \rangle.$$

As noted above the action of T on the $(\eta, \zeta) = (1, -1)$ component is trivial. On the other hand s_1s_5 acts by $(\alpha' : \epsilon' : \zeta') \mapsto (-\alpha' : -\epsilon' : \zeta')$ and similarly s_0s_5 acts by $(\alpha' : \epsilon' : \zeta') \mapsto (\alpha' : -\epsilon' : -\zeta')$. The quotient of the component under this action can be identified with

$$\{(\alpha'', \epsilon'', \zeta'') : \alpha''\epsilon''\zeta'' = 1\}/C_3,$$

via the map $(\alpha' : \epsilon' : \zeta') \mapsto ((\alpha')^2 : (\epsilon')^2 : (\zeta')^2)$.

It remains to take the quotient of this by the action of $\langle u_1, u_2 \rangle$ which acts by signed permutation of the coordinates $(\alpha')^2, (\epsilon')^2, (\zeta')^2$. The result therefore agrees with the quotient of the identity component (though geometrically one is 4 times the area of the other), giving the quotient $T^w/Z_W(w)$ as the union of two equilateral triangles.

- (9) Case $A_2 \times A_1^2$: The fixed set is the single 2-torus with coordinates $(\alpha, \alpha^2, 1, \epsilon^2, \epsilon, 1)$. The centraliser is $\langle s_0s_6 \rangle \times (\langle s_1 \rangle \wr \langle u_1 \rangle)$ and the elements s_0s_6, s_1 both act trivially on the fixed set in the Lie algebra, and hence on its fixed torus. This leaves the action of u_1 to consider, which exchanges and inverts α, ϵ . Replacing the variable ϵ by $\epsilon' = \epsilon^{-1}$, the quotient is therefore the symmetric product of two copies of \mathbb{T}^1 , which is a Möbius band.

- (10) Case A_2^2 : The fixed set is a family of three 2-tori indexed by the centre as follows:

$$\{(\beta^{-1}, \beta, 1, \beta^{-1}, \beta, 1) : \beta^3 = 1\} \times \{(1, 1, 1, \delta, \epsilon, 1) : \delta, \epsilon \in \mathbb{T}^1\}$$

Since the centre is pointwise fixed by the Weyl group, the quotient is 3 copies of the quotient of the identity component.

We write the centraliser as $(\langle s_1 s_2 \rangle \wr \langle u_2 \rangle) \times \langle s_4, s_5 \rangle$. Note that $s_1 s_2$ acts trivially on the fixed set, hence the action is just by the quotient group $\langle u_2 \rangle \times \langle s_4, s_5 \rangle = \langle s_4, s_5 u_2 \rangle \cong \mathcal{D}_6$.

The element $s_5 u_2$ takes $(1, 1, 1, \delta, \epsilon, 1)$ to $(1, 1, 1, \delta^{-1}, \delta^{-1} \epsilon, 1)$ while the element s_4 acts as

$$s_4 : (1, 1, 1, \delta, \epsilon, 1) \mapsto (1, 1, 1, \delta^{-1} \epsilon, \epsilon, 1)$$

Taking a hexagonal fundamental domain for the torus, (dual to the r_4, r_5 lattice) we have the standard action of the dihedral group \mathcal{D}_6 yielding a $(3, 4, 6)$ triangle.

Hence the quotient of the fixed set by the centraliser is three copies of the triangle.

- (11) Case $A_3 \times A_1$: The fixed set is a single 2-torus with coordinates $(\alpha, \alpha^2, 1, \alpha^{-2}, \epsilon, 1)$.

The centraliser is $\langle s_0 s_6 s_3 \rangle \times \langle s_1 \rangle \times \langle s_5 \rangle$, with $s_0 s_6 s_3, s_1$ acting trivially. The element s_5 takes $(\alpha, \alpha^2, 1, \alpha^{-2}, \epsilon, 1)$ to $(\alpha, \alpha^2, 1, \alpha^{-2}, \epsilon^{-1} \alpha^{-2}, 1)$. Changing coordinates to α, ϵ' where $\epsilon' = \alpha \epsilon$ we see that the action takes $\epsilon' = \alpha \epsilon$ to $\alpha(\epsilon^{-1} \alpha^{-2}) = (\epsilon')^{-1}$, while leaving α unchanged. Hence the quotient is a cylinder.

- (12) Case A_4 : The fixed set is a single 2-torus with coordinates $(\alpha, \beta, 1, \beta^{-1}, \beta^{-2}, 1)$.

The centraliser is $\langle s_0 s_6 s_3 s_4 \rangle \times \langle s_1 \rangle$ with the first factor acting trivially. The action of s_1 takes α to $\alpha^{-1} \beta$ and leaves β unchanged. Setting $\beta' = \alpha^{-1} \beta$ this action simply exchanges α, β' hence the quotient is the symmetric product of two copies of \mathbb{T}^1 (a Möbius band).

- (13) Case D_4 : The fixed set is a single 2-torus $(\alpha, \alpha^2, \alpha \epsilon, \epsilon^2, \epsilon, 1)$, which is the $\zeta = \eta = 1$ component of the fixed set of $s_0 s_1 s_5 s_3$ (case 8). Indeed our representative is chosen so that its cube equals $s_0 s_1 s_5 s_3$ so of course the fixed sets must be nested. The centraliser is $\langle s_0 s_1 s_5 T \rangle \times \langle u_1, u_2 \rangle$ with the first factor acting trivially. As in case 8 the action of $\langle u_1, u_2 \rangle$ can be equivariantly identified with the permutation action of \mathcal{D}_3 on the SU_3 torus yielding the quotient as an equilateral triangle.

- (14) Case $D_4[a_1]$: The fixed set is the same single 2-torus $(\alpha, \alpha^2, \alpha \epsilon, \epsilon^2, \epsilon, 1)$ as for the previous case of type D_4 , and here we note that the square of the representative yields the element $s_0 s_1 s_5 s_3$. The centraliser is the complex reflection group G_8 given by $\langle s_5 u_3, s_5^T u_1 \rangle$. Again we will use the calculations

from case 8, with the fixed set described as

$$\{(\alpha', \epsilon', \zeta') : \alpha' \epsilon' \zeta' = 1\} / C_3.$$

As noted in that case, the elements s_5, T act trivially on this set, hence we again reduce to the action of the $\langle u_3, u_1 \rangle = \langle u_1, u_2 \rangle$ giving an equilateral triangle.

- (15) Case $A_2^2 \times A_1$: The fixed set consists of 3 circles indexed by a cube root of unity β with coordinates $(\beta^2, \beta, 1, \epsilon^2, \epsilon, 1)$. Each of these circles contains one of the elements of the centre, indeed we can write elements of the fixed set in the form

$$(\beta^2, \beta, 1, \beta^2, \beta, 1)(1, 1, 1, (\epsilon')^2, \epsilon', 1).$$

The first factor is central, hence fixed by the whole Weyl group, and the second lies in the identity component of the fixed set, and hence is fixed by s_0, s_6, s_5, s_1, s_2 . The centraliser is $(\langle s_0 s_6 \rangle \wr \langle u_2 \rangle) \times \langle s_5 \rangle$ where only the element u_2 acts non-trivially. This acts on the second factor by inverting ϵ' . The quotient is therefore 3 intervals.

- (16) Case $A_3 \times A_1^2$: The fixed set is a family of four circles, indexed by ζ, η with $\zeta^2 = \eta^2 = 1$. We have coordinates $(\epsilon^{-1}\eta, \epsilon^{-2}\zeta, \zeta, \epsilon^2\zeta, \epsilon, \zeta)$. We factorise this as

$$(\epsilon^{-1}\eta, \epsilon^{-2}, \eta, \epsilon^2, \epsilon, 1)(1, \zeta, \eta\zeta, \zeta, 1, \zeta).$$

This factorisation is chosen so that for each choice of η, ζ we obtain a circle lying in the corresponding 2-torus appearing in case 8.

The centraliser is $\langle s_0 s_6 s_3 s_1 s_5 \rangle \times \langle s_1, T, s_5 \rangle$ with the first factor acting trivially. The second factor lies in the centraliser from case 8, which contains the group $W(D_4) = \langle s_1, T, s_5, s_0 \rangle$. The subgroup $\langle s_1, T, s_5 \rangle$ is sufficient to transitively permute the three non-identity components in case 8 and hence transitively permutes the circles in this case.

Since the centre lies in the circle $\zeta = \eta = 1$ it follows that this circle is fixed by the whole of $\langle s_0 s_6 s_3 s_1 s_5 \rangle \times \langle s_1, T, s_5 \rangle$.

For the non-identity components, as in case 8 consider the component $(\eta, \zeta) = (1, -1)$ which is pointwise fixed by T . This component is stabilised by $\langle T, s_1 s_5 \rangle \cong D_4$ and this must be the whole stabiliser as it has index 3 in $\langle s_1, T, s_5 \rangle \cong S_4$. As the element T acts trivially it remains to take the quotient by the action of $s_1 s_5$. This fixes $(\epsilon^{-1}\eta, \epsilon^{-2}, \eta, \epsilon^2, \epsilon, 1)$ and takes $(1, \zeta, \eta\zeta, \zeta, 1, \zeta)$ to

$$(\zeta, 1, 1, 1, \zeta, 1)(1, \zeta, \eta\zeta, \zeta, 1, \zeta)$$

hence it has the effect of negating ϵ . Thus the quotient of the non-identity components yields a circle.

The quotient of the fixed set by the centraliser is therefore two circles.

- (17) Case $A_4 \times A_1$: The fixed set is a single circle with coordinates $(\alpha, \alpha^{-2}, 1, \alpha^{-2}, \alpha^{-4}, 1)$. As noted after Theorem 7 this circle contains the centre \mathcal{Z} of the Lie group, hence the action of the centraliser must be trivial.
- (18) Case A_5 : The fixed set is a family of three circles indexed by a choice ϵ of a cube root of unity and with coordinates $(\alpha, \epsilon, 1, \epsilon^2, \epsilon, 1)$. As in case 15 this can be written as a product of a circle with the centre so that points of the fixed set take the form

$$(\epsilon^2, \epsilon, 1, \epsilon^2, \epsilon, 1)(\alpha\epsilon, 1, 1, 1, 1).$$

The centraliser is $\langle s_0s_6s_3s_4s_5 \rangle \times \langle s_1 \rangle$ with the first factor acting trivially. The second factor inverts $\alpha\epsilon$ hence gives a reflection on each of the three circles. Hence the quotient is three copies of the interval.

- (19,20) Cases $D_5, D_5[a_1]$: In both cases the fixed set is a single circle with coordinates $(\delta^{-2}, \delta^{-1}, 1, \delta, \delta^2, 1)$. As in case 17, this must contain the three elements of the centre so is fixed pointwise, yielding a circle as the quotient.
- (21) Case A_2^3 : The fixed set is the finite group $C_3 \times C_3 \times C_3$ indexed by three choices of cube roots of 1: β, ϵ, ζ . In coordinates it is given by: $(\beta^2\zeta, \beta, 1, \epsilon^2\zeta, \epsilon, \zeta)$. We can split the central points $(\epsilon^2, \epsilon, 1, \epsilon^2, \epsilon, 1)$ off from this and writing $\beta' = \beta\epsilon^2\zeta^2$ yields a factorisation as a product as follows:

$$(\epsilon^2, \epsilon, 1, \epsilon^2, \epsilon, 1)(1, \zeta, 1, \zeta, 1, \zeta)((\beta')^2, \beta', 1, 1, 1, 1).$$

The first factor is the centre so fixed by everything. We identify the fixed set as an \mathbb{F}_3 vector space V with basis

$$e_1 = (\omega^2, \omega, 1, \omega^2, \omega, 1), e_2 = (1, \omega, 1, \omega, 1, \omega), e_3 = (\omega^2, \omega, 1, 1, 1, 1)$$

where $\omega = e^{2\pi i/3}$. We take the dual basis $\{e_1^*, e_2^*, e_3^*\}$ for V^* .

As the first basis vector e_1 is fixed by the centraliser we obtain an affine action of the centraliser on the affine plane $\mathbb{A} = \{v^* \in V^* : \langle v^*, e_1 \rangle = 1\}$.

We now consider the action of the three generators s_0s_6, Ts_3, s_5s_4 on V . Starting with s_0s_6 , this fixes e_1 and e_3 , and takes e_2 to $e_1 + e_2 + e_3$. Similarly s_5s_4 fixes e_1 and e_3 , and takes e_2 to $2e_1 + e_2 + e_3$.

The element Ts_3 fixes e_1, e_2 , and takes e_3 to $e_3 + 2e_2$.

In matrix form the generators s_0s_6, Ts_3, s_5s_4 act on V as:

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}$$

Dually, the action on V^* is given by:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix}$$

It is not hard to see that these generate the Hessian group, i.e. the group of all orientation preserving affine maps on \mathbb{A} .

Dualising the action back to V is therefore all orientation preserving linear maps which fix the vector e_1 . The quotient is therefore 4 points, corresponding to the three points on the line spanned by e_1 (the centre of the Lie group) and a single orbit for all the remaining vectors in V .

(22) Case $A_5 \times A_1$: In coordinates the fixed set is given by

$$(\eta\epsilon^2, \zeta\epsilon, \zeta, \zeta\epsilon^2, \epsilon, \zeta) = (\eta\epsilon^2, \epsilon, \eta, \epsilon^2, \epsilon, 1)(1, \zeta, \zeta\eta, \zeta, 1, \zeta)$$

where ϵ is a cube root of unity and $\zeta^2 = \eta^2 = 1$. We choose this factorisation to the copy of the Klein group V_4 matches with that appearing in cases 8 and 16.

The centraliser is $\langle s_0s_6s_3s_4s_5s_1 \rangle \times \langle s_1, T \rangle$ with the first factor acting trivially. The fixed set lies in the fixed set from case 16, and the group $\langle s_1, T \rangle$ is a subgroup of the centraliser from that case.

The $\eta = \zeta = 1$ part is the centre of the Lie group and is thus fixed by the centraliser. As noted in case 16, the three other possible values for (η, ζ) are transitively permuted by $\langle s_1, T \rangle$. The triple with $(\eta, \zeta) = (1, -1)$ is pointwise fixed by T and its stabiliser in the group $\langle s_1, T \rangle$ has index 3 and is thus $\langle T \rangle$. Hence the 6 points given by $(\eta, \zeta) = (1, 1)$ and $(\eta, \zeta) = (1, -1)$ form a strict fundamental domain for the action of the centraliser.

The quotient is therefore 6 points.

(23–25) Cases $E_6, E_6[a_1], E_6[a_2]$: In each case the fixed set is the centre of the Lie group $(\epsilon^2, \epsilon, 1, \epsilon^2, \epsilon, 1)$ where ϵ ranges over the cube roots of 1. Each of these three cases provides three points in the quotient.

4 Fixed sets and their quotients: the group of adjoint type

In this section we consider the case of the E_6 Lie group of adjoint type. The maximal torus in this case is the torus $T^\vee = T/\mathcal{Z}$ where \mathcal{Z} denotes the centre of the Lie group of simply connected type. We will determine the quotients $(T^\vee)^w/Z_W(w)$ and thus establish the following:

Theorem 6 *Let W be the Weyl group of type E_6 and let T and T^\vee denote the corresponding (real) maximal tori. For each $w \in W$ there is a homotopy equivalence*

$$T^w/Z_W(w) \sim (T^\vee)^w/Z_W(w).$$

As discussed in Sect. 1 there are two possibilities for the fixed sets $(T^\vee)^w$: either the centre lies in the identity component T_1^w , in which case $(T^\vee)^w = T^w/\mathcal{Z}$, or \mathcal{Z} injects into the component group F_w , in which case $(T^\vee)_1^w = T_1^w$. The first case we will refer to as having ramified torus while the latter case has ramified component group.

The component group is ramified in cases $A_2^2, A_2^2 \times A^1, A_5$ as well as the elliptic cases $A_2^3, A_5 \times A_1, E_6, E_6[a_1], E_6[a_2]$. In all the other cases the torus is ramified.

4.1 Ramified torus

In this case the fixed set $(T^\vee)^w$ is given by T^w/\mathcal{Z} where $T^w \cong T_1^w \times F_w$. Hence $(T^\vee)^w \cong T_1^w/\mathcal{Z} \times F_w$ (with $F_w = F_w^\vee$).

Since the action of the Weyl group arises from the conjugation action in the Lie group, this commutes with the action of the centre \mathcal{Z} . Hence the quotient $(T^\vee)^w/Z_W(w)$ is the quotient of T^w by the action of the product $Z_W(w) \times \mathcal{Z}$ and indeed we can therefore compute this as the quotient $(T^w/Z_W(w))/\mathcal{Z}$. The quotients $T^w/Z_W(w)$ were evaluated in Sect. 3.

Note that these quotients carry a natural metric, inherited from the Lie algebra \mathfrak{t} . The action of \mathcal{Z} on the quotients is induced from a translation action on \mathfrak{t} and is thus isometric. In many cases this is sufficient to determine the quotient up to homotopy.

- Case \emptyset : The quotient $(T^\vee)^w/Z_W(w)$ is Δ^6/\mathcal{Z} where the contraction of Δ^6 to its barycentre is equivariant, hence Δ^6/\mathcal{Z} is contractible.
- Case A_1 : The same argument shows that the quotient Δ^5/\mathcal{Z} is also contractible.
- Case A_1^2 : The quotient $T^w/Z_W(w)$ in this case is a Δ^3 bundle over \mathbb{T}^1 . Writing Δ^3 as the join of two edges e_0 and e_1 , the monodromy is given by the map inverting the edge e_0 and fixing e_1 .

In logarithmic coordinates the Δ^3 fibres are described by the sum of the base fibre with a point $(x_1, x_1, -\frac{x_1}{2}, -\frac{x_1}{2}, -\frac{x_1}{2}, -\frac{x_1}{2})$. The centre acts by translation by multiples of $(\frac{2}{3}, \frac{2}{3}, -\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3})$ which tautologically preserves the coordinates of the fibre. A fundamental domain is given by $x_1 \in [0, \frac{1}{3}]$. The monodromy in the identification of the fibres at $0, \frac{1}{3}$ is the composition of the trivial identification of the $0, \frac{4}{3}$ fibres with the (inverse of the) original monodromy in $T^w/Z_W(w)$. Hence the monodromy maps agree and the quotient $(T^\vee)^w/Z_W(w)$ is identified with $T^w/Z_W(w)$.

- Case A_2 : The quotient $T^w/Z_W(w)$ is $SP^2(\Delta^2) = (\Delta^2 \times \Delta^2)/C_2$, hence the quotient $(T^\vee)^w/Z_W(w)$ is given by $(\Delta^2 \times \Delta^2)/(C_2 \times \mathcal{Z})$. The action of $C_2 \times \mathcal{Z}$ is isometric and hence preserves the product of the barycentres, giving a contractible quotient.
- Case A_1^3 : The quotient is $(\Delta^2 \times \Delta^1)/\mathcal{Z}$. Since the action is isometric it preserves the Cartesian product structure. Since \mathcal{Z} has order 3 it preserves the Δ^1 direction and the contraction to the barycentre of Δ^2 times the interval Δ^1 is equivariant. Hence $(\Delta^2 \times \Delta^1)/\mathcal{Z}$ is contractible.
- Case $A_2 \times A_1$: The quotient $T^w/Z_W(w)$ is $\mathbb{T}^1 \times \mathbb{T}^2/S_3$ giving a product $\mathbb{T}^1 \times \Delta^2$. The action of \mathcal{Z} acts diagonally by rotations on both factors. This action is fixed point free and yields a solid torus which carries a Seifert fibration over the cone Δ^2/\mathcal{Z} , [9].
- Case A_3 : The quotient $T^w/Z_W(w)$ in this case is a triangle bundle over \mathbb{T}^1 , with the triangle obtained as the quotient of a square by a diagonal reflection. The monodromy is obtained from the rotation of the square by π , which yields a diagonal flip on the triangle. The action of \mathcal{Z} is to translate by $1/3$ along the circle while again giving a rotation by π on the square. Note that the cube of this element agrees with the monodromy of the original bundle as required. Hence the quotient $(T^\vee)^w/Z_W(w)$ is identified with $T^w/Z_W(w)$.

- Case A_1^4 : The action of \mathcal{Z} on $\Delta^2 \sqcup \Delta^2$ must preserve the components fixing the barycentre of each, and the contraction to the barycentres is equivariant.
- Case $A_2 \times A_1^2$: The action of \mathcal{Z} on the symmetric product $\text{SP}^2(\mathbb{T}^1)$ is given by the diagonal action of \mathcal{Z} on the two copies of the circle \mathbb{T}^1 . Since this is a fixed point free action on the Möbius band, the quotient is again a Möbius band.
- Case $A_3 \times A_1$: The quotient $\mathcal{T}^w/Z_W(w)$ is $\mathbb{T}^2/C_2 = \Delta^1 \times \mathbb{T}^1$. The action of \mathcal{Z} rotates the \mathbb{T}^1 factor, again yielding a copy of $\Delta^1 \times \mathbb{T}^1$.
- Case A_4 : As in case $A_2 \times A_1^2$ the centre \mathcal{Z} acts diagonally on $\text{SP}^2(\mathbb{T}^1)$ giving quotient a Möbius band.
- Cases D_4 and $D_4[a_1]$: In these cases the quotient $\mathcal{T}^w/Z_W(w)$ are equal, giving a copy of Δ^2 . The group \mathcal{Z} acts by rotation on this giving a cone Δ^2/\mathcal{Z} .
- Case $A_3 \times A_1^2$: The action of \mathcal{Z} must preserve the two components $\mathbb{T}^1 \sqcup \mathbb{T}^1$, acting isometrically and fixed point freely on each circle, hence the quotient is a disjoint union $\mathbb{T}^1 \sqcup \mathbb{T}^1$.
- Cases $A_4 \times A_1, D_5, D_5[a_1]$: In each of these cases the centre acts fixed-point freely on \mathbb{T}_1 and the quotient is again a circle.

4.2 Ramified component groups

We now derive the quotients in the ramified component group cases. The results are summarised in Table 3. We denote elements of the centre \mathcal{Z} by $z_\omega = (\omega^2, \omega, 1, \omega^2, \omega, 1)$ where ω is a cube root of 1.

- Case A_2^2 :
Recall that $w = s_0s_6s_1s_2$ and $\mathcal{T}^w = T_1^w \times \mathcal{Z}$ where (in standard coordinates on \mathbb{T}^6) the identity component is

$$T_1^w = \{(1, 1, 1, \delta, \epsilon, 1) : \delta, \epsilon \in \mathbb{T}^1\}.$$

For ω a cube root of 1, the action of w takes $p_\omega = (1, \omega^2, 1, 1, 1, \omega)$ to

$$(\omega^2, 1, 1, \omega^2, \omega, \omega) = z_\omega p_\omega$$

so p_ω is fixed by w up to the action of \mathcal{Z} .

When $\omega \neq 1$ the point p_ω does not lie in \mathcal{T}^w , hence the image of p_ω in the quotient by \mathcal{Z} lies in a non-identity component of $(\mathcal{T}^\vee)^w$. We can therefore index the components of $(\mathcal{T}^\vee)^w$ by ω .

We then have

$$(\mathcal{T}^\vee)^w = (\{p_\omega : \omega^3 = 1\} \times \mathcal{T}^w)/\mathcal{Z},$$

and taking a fundamental domain for the action of \mathcal{Z} , we can identify $(\mathcal{T}^\vee)^w$ with

$$\{p_\omega : \omega^3 = 1\} \times \{(1, 1, 1, \delta, \epsilon, 1) : \delta, \epsilon \in \mathbb{T}^1\}.$$

The centraliser of the element w is $\langle s_1s_2 \rangle \wr \langle u_2 \rangle \times \langle s_4, s_5 \rangle$ and we now consider its action on $(\mathcal{T}^\vee)^w$.

We first consider the action of the normal subgroup $\langle s_1s_2, s_0s_6 \rangle$ in the centraliser. (Recall that $(s_1s_2)^{u_2} = s_0s_6$.) Since $s_1s_2s_0s_6$ acts trivially it suffices to consider the action of s_1s_2 . This does not preserve the fundamental domain for the action of \mathcal{Z} , so we act by s_1s_2 and then by the central element z_ω . This takes p_ω to $(1, \omega^2, 1, \omega^2, \omega, \omega) = p_\omega(1, 1, 1, \omega^2, \omega, 1)$.

Hence the action of s_1s_2 preserves components of $(T^\vee)^w$ and acts on the component indexed by ω by multiplication by $(1, 1, 1, \omega^2, \omega, 1) \in (T^\vee)_1^w = T_1^w$.

Recall that the action of the \mathcal{D}_6 group $\langle u_2, s_4, s_5 \rangle = \langle s_4, u_2s_5 \rangle$ on T_1^w is given by the standard dihedral action by the group of symmetries of the hexagonal fundamental domain for the torus, which we will denote \diamond . The element s_5 acts trivially while u_2 acts to invert the ϵ coordinate. For the identity component $(T^\vee)_1$, indexed by $\omega = 1$, the element s_1s_2 acts trivially, and the quotient by the centraliser is a $(2, 4, 6)$ triangle as in the case of T^w .

When ω is a nontrivial cube root of the element 1, our \mathcal{D}_6 group still preserves the component, but s_1s_2 no longer acts trivially so we first factor out its action, which we recall is given by multiplication by $(1, 1, 1, \omega^2, \omega, 1)$. This yields the torus with dual hexagon as fundamental domain and we denote this fundamental domain by \diamond . So the quotient of this component by the centraliser is given by taking the dual action of \mathcal{D}_6 on the dual hexagon \diamond . But these actions are equivalent so we again obtain as quotient a $(2, 4, 6)$ triangle, here with $1/3$ the area of the $\omega = 1$ case, and the quotient $(T^\vee)^w$ by the centraliser of w is three $(2, 4, 6)$ triangles.

- Case $A_2^2 \times A_1$: Here the representative is $w = s_0s_6s_5s_1s_2$. Since s_5 acts trivially on p_ω , we have $w \cdot p_\omega = z_\omega p_\omega$ using the previous case. We thus have

$$(T^\vee)^w = (\{p_\omega : \omega^3 = 1\} \times T^w) / \mathcal{Z}.$$

where $T^w = T_1^w \times \mathcal{Z}$ and $T_1^w = \{(1, 1, 1, \epsilon^2, \epsilon, 1) : \epsilon \in \mathbb{T}^1\}$.

Taking a fundamental domain for the action of \mathcal{Z} , we can identify $(T^\vee)^w$ with

$$\{p_\omega : \omega^3 = 1\} \times \{(1, 1, 1, \epsilon^2, \epsilon, 1) : \epsilon \in \mathbb{T}^1\}.$$

The centraliser is $\langle s_1s_2 \rangle \wr \langle u_2 \rangle \times \langle s_5 \rangle$.

As noted in the A_2^2 case, for ω a cube root of 1, the action of s_1s_2 followed by z_ω takes p_ω to $(1, \omega^2, 1, \omega^2, \omega, \omega)$, so in $(T^\vee)^w$ the element s_1s_2 has the effect of preserving the three components and on each it multiplies the coordinate ϵ by ω . It is easy to see that the generator s_5 acts trivially on the above fundamental domain. The generator u_2 also fixes p_ω because it interchanges r_2 and $-r_6$. As in the case of T^w the element u_2 acts on the second factor by inverting ϵ .

Hence the quotient of each component is an interval: the element s_1s_2 can act either trivially or by rotation on the circle. For $\omega = 1$ we have the former, just leaving the action of u_2 . The latter occurs when $\omega \neq 1$, and we have the dihedral action of the group $(\langle s_1s_2 \rangle \wr \langle u_2 \rangle) / \langle s_1s_2(s_1s_2)^{u_2} \rangle \cong \mathcal{D}_3$.

- Case A_5 : Consider the action of the representative $w = s_0s_6s_3s_4s_5$ on the point $q_\omega = (1, 1, 1, 1, \omega^2, \omega^2) \in \mathcal{T}$, where ω is a cube root of unity. The action of w translates this by the element z_w in the centre, so it represents a fixed point in the

dual torus T^\vee . Hence the fixed set under the dual action of w is given by

$$(T^\vee)^w = (\{q_\omega : \omega^3 = 1\} \times T^w) / \mathcal{Z}.$$

where $T^w = T_1^w \times \mathcal{Z}$ and $T_1^w = \{(\alpha, 1, 1, 1, 1, 1) : \alpha \in \mathbb{T}^1\}$. As in the previous case, by taking a fundamental domain for the action of \mathcal{Z} , we can identify the fixed set with the set

$$\{q_\omega : \omega^3 = 1\} \times \{(\alpha, 1, 1, 1, 1, 1) : \alpha \in \mathbb{T}^1\}.$$

The centraliser is $\langle s_0 s_6 s_3 s_4 s_5 \rangle \times \langle s_1 \rangle$ with the first factor acting trivially. The element s_1 preserves the three circles and in each case acts by a reflection taking α to α^{-1} . The quotient is therefore, again, three copies of the interval.

- Case A_3^3 : The dual case is considered in detail in the derivation of the action of the centraliser on T^w , so we will not repeat it here. In summary, the dual fixed set can be identified with a 3-dimensional vector space over \mathbb{F}_3 , equipped with a quotient to \mathbb{F}_3 . The cosets of the kernel give two affine planes and a linear plane, and the action of the centraliser is given by the action of the Hessian group by orientation preserving affine maps on the affine planes, with its linear part on the linear plane. The quotient is therefore a set of four points.

For the sake of completeness we note that $w = s_0 s_6 s_1 s_2 s_5 s_4$ takes p_ω to $(\omega^2, 1, 1, \omega^2, \omega, \omega) = z_\omega p_\omega$. The fixed set T^w is

$$\{(1, \zeta, 1, \zeta, 1, \zeta)(\beta^2, \beta, 1, 1, 1, 1) : \zeta^3 = \beta^3 = 1\} \times \mathcal{Z}.$$

The fixed set $(T^\vee)_1^w$ is therefore identified with a fundamental domain

$$\{(1, \omega^2, 1, 1, 1, \omega)(1, \zeta, 1, \zeta, 1, \zeta)(\beta^2, \beta, 1, 1, 1, 1) : \omega^3 = \zeta^3 = \beta^3 = 1\}.$$

- Case $A_5 \times A_1$: Consider the action of the representative $w = s_0 s_6 s_3 s_4 s_5 s_1$ on the point $(1, 1, 1, 1, \omega^2, \omega^2) \in T$, where ω is a cube root of unity. As noted in the A_5 case, this element is fixed by s_1 and is fixed up to the action of \mathcal{Z} by $s_0 s_6 s_3 s_4 s_5$. Hence

$$(T^\vee)^w = (\{q_\omega : \omega^3 = 1\} \times T^w) / \mathcal{Z}.$$

Here elements of T^w have the form $(\eta\epsilon^2, \zeta\epsilon, \zeta, \zeta\epsilon^2, \epsilon, \zeta)$ where $\epsilon^3 = \zeta^2 = \eta^2 = 1$. A fundamental domain for the action of \mathcal{Z} is given by

$$\{q_\omega : \omega^3 = 1\} \times \{(\eta, \zeta, \zeta, \zeta, 1, \zeta) : \zeta^2 = \eta^2 = 1\}.$$

The centraliser is $\langle w, s_1, T \rangle$ where w acts trivially (on the quotient by \mathcal{Z}). Clearly the element s_1 fixes the first factor, and T also fixes this because T fixes the vector $r_5 + r_6$. We note that the second factor is obtained by applying the exponential map to $\frac{1}{2}\mathbb{Z}r_1 + \frac{1}{2}\mathbb{Z}r_T$ and $r_1, -r_T$ have inner product -1 (giving the group $\langle s_1, T \rangle$ of type A_2).

Table 3 Fixed sets in \mathcal{T}^\vee for ramified component group cases

Conjugacy class	Fixed sets in \mathcal{T}^\vee lifted to \mathcal{T}
A_2^2	$\{(1, 1, 1, \delta, \epsilon, 1) : \delta, \epsilon \in \mathbb{T}^1\} \times \{(1, \omega^2, 1, 1, 1, \omega) : \omega^3 = 1\}$
$A_2^2 \times A_1$	$\{(1, 1, 1, \epsilon^2, \epsilon, 1) : \epsilon \in \mathbb{T}^1\} \times \{(1, \omega^2, 1, 1, 1, \omega) : \omega^3 = 1\}$
A_5	$\{(\alpha, 1, 1, 1, 1, 1) : \alpha \in \mathbb{T}^1\} \times \{(1, 1, 1, 1, \omega^2, \omega^2) : \omega^3 = 1\}$
A_3^2	$\{(\beta^2, \beta\zeta, 1, \zeta, 1, \zeta) : \zeta^3 = \beta^3 = 1\} \times \{(1, \omega^2, 1, 1, 1, \omega) : \omega^3 = 1\}$
$A_5 \times A_1$	$\{(\eta, \zeta, \zeta, \zeta, 1, \zeta) : \zeta^2 = \eta^2 = 1\} \times \{(1, 1, 1, 1, \omega^2, \omega^2) : \omega^3 = 1\}$
E_6	$\{(1, \omega, 1, 1, \omega, 1) : \omega^3 = 1\}$
$E_6[a_1]$	$\{(1, \omega, 1, \omega, 1, \omega^2) : \omega^3 = 1\}$
$E_6[a_2]$	$\{(1, \omega, 1, 1, 1, \omega^2) : \omega^3 = 1\}$

The action of $\langle s_1, T \rangle$ on pairs (η, ζ) is thus the standard action $s_1 : (\eta, \zeta) \rightarrow (\eta^{-1}\zeta, \zeta) = (\eta\zeta, \zeta)$ and $T : (\eta, \zeta) \rightarrow (\zeta, \zeta) = (\eta, \eta\zeta)$.

Hence the quotient is 6 points: a fundamental domain is given by $\zeta = 1$.

- Cases $E_6, E_6[a_1], E_6[a_2]$. In each of these cases the fixed set in \mathcal{T} consists entirely of the centre of the Lie group, and is fixed pointwise. It follows from the discussion in Sect. 1, Remark 1.8, that the fixed set in \mathcal{T}^\vee can be identified with the Pontryagin dual of the centre, which is again cyclic of order 3, and that the centraliser acts on this by the (dual of) the trivial action. All that remains, for the sake of completeness is to identify the three fixed point groups in these cases.

In the E_6 case the element $w = s_1s_2s_3s_4s_5s_6$ takes $(\omega, 1, 1, \omega, 1, 1)$ to $(1, \omega, 1, 1, \omega, 1) = z_\omega(\omega, 1, 1, \omega, 1, 1)$ for ω a cube root of unity, so the fixed set is identified with $\{(1, \omega, 1, 1, \omega, 1) : \omega^3 = 1\}$.

For $E_6[a_1]$, the element $w = s_1s_2s_3s_4s_5s_6^{s_3}$ takes $(1, \omega, 1, \omega, 1, \omega^2)$ to $z_\omega(1, \omega, 1, \omega, 1, \omega^2)$, giving fixed set $\{(1, \omega, 1, \omega, 1, \omega^2) : \omega^3 = 1\}$.

Finally for $E_6[a_2]$, the element $w = s_6s_2s_0^T s_1^T s_4s_3$ takes $(1, \omega, 1, 1, 1, \omega^2)$ to $z_\omega(1, \omega, 1, 1, 1, \omega^2)$, giving fixed set $\{(1, \omega, 1, 1, 1, \omega^2) : \omega^3 = 1\}$.

5 Applications

There are two major applications for the results in this article.

5.1 Extended affine Weyl groups of type E_6

Our first application is the computation of the rational K -theory of the group C^* -algebras for extended affine Weyl groups. From [5] we know that Langlands duality induces an isomorphism in K -theory, and the computations given here make it possible to exhibit explicit classes and to visualise this isomorphism.

Theorem 8 *Let W'_a be an extended affine Weyl group of one of the two Lie groups of type E_6 (simply connected or adjoint type). Up to torsion, $K_*(C_r^* W'_a)$ is*

$$\begin{aligned} \mathbb{Z}^{47} & \text{ in dimension } 0 \\ \mathbb{Z}^{11} & \text{ in dimension } 1. \end{aligned}$$

Proof Let G be a compact connected simply-connected semisimple Lie group of type E_6 , with centre \mathcal{Z} . Let \mathcal{T} be a maximal torus in G , and let Γ be the kernel of the exponential map $\exp : \mathfrak{t} \rightarrow T$, which is simply the root lattice since G is simply connected.

The Langlands dual $G^\vee = G/\mathcal{Z}$ is the adjoint form and its extended affine Weyl group is

$$W'_a(G^\vee) := \Gamma^\vee \rtimes W$$

where Γ^\vee is the dual lattice. Its group C^* -algebra $C^*(W'_a(G^\vee))$ is isomorphic to $C(\mathcal{T}) \rtimes W$ and by the Green–Julg theorem its K -theory is isomorphic to $K^*_W(\mathcal{T})$

Now we apply the equivariant Chern character [2] for the discrete group W :

$$K^*_W(\mathcal{T}) \otimes_{\mathbb{Z}} \mathbb{C} \simeq H^*(\mathcal{T} // W; \mathbb{C})$$

where the geometric extended quotient $\mathcal{T} // W$ is the quotient of the inertia space

$$\tilde{\mathcal{T}} = \{(w, x) \in W \times \mathcal{T} : wx = x\}$$

by the action $g(w, x) = (w^g, gx)$. The quotient is given by the formula

$$\mathcal{T} // W := \bigsqcup \mathcal{T}^w / Z_W(w)$$

with one w chosen in each W -conjugacy class. Putting all this together, we obtain

$$K_* C^*(W'_a(G^\vee)) \otimes_{\mathbb{Z}} \mathbb{C} \simeq \bigoplus H^*(\mathcal{T}^w / Z_W(w); \mathbb{C})$$

The 25 sectors are listed in the right-hand column of Table 2, and the K -theory is readily computed.

For the extended affine Weyl group $W'_a(G)$ the K -theory is obtained by replacing the sectors $\mathcal{T}^w / Z_W(w)$ by $(\mathcal{T}^\vee)^w / Z_W(w)$ each of which is homotopic to the corresponding sector for the action of W on \mathcal{T} by Theorem 6. □

5.2 p -adic groups of type E_6

The Baum–Connes conjecture does not help with computing the K -theory of p -adic groups since the left hand side is not itself tractable. The ABPS-framework developed in [1] is designed to fix this, providing a more effective approach to p -adic groups. Indeed, Conjecture 5, *ibid* provides a much finer and more precise formula in K -theory than Baum–Connes alone provides.

Let \mathcal{G} denote a split group of type E_6 over a p -adic field; the group may be of adjoint type or simply connected. Let $C_r^*(\mathcal{G})$ denote the reduced C^* -algebra of \mathcal{G} . The reduced C^* -algebra admits the Bernstein decomposition

$$C_r^*(\mathcal{G}) = \bigoplus_{\mathfrak{s} \in \mathfrak{B}(\mathcal{G})} C_r^*(\mathcal{G})^{\mathfrak{s}}$$

where $\mathfrak{B}(\mathcal{G})$ is the Bernstein spectrum of \mathcal{G} . Each point \mathfrak{s} in the Bernstein spectrum is an equivalence class of cuspidal pairs (M, σ) where M is a Levi subgroup of \mathcal{G} and σ is an irreducible cuspidal representation of M . In particular, we have the *Iwahori point* \mathfrak{i} defined by the pair $(\mathcal{T}_p, 1)$ where \mathcal{T}_p is a maximal torus of (the p -adic group) \mathcal{G} and 1 is the trivial representation of \mathcal{T}_p . We will write

$$\mathfrak{A} = C_r^*(\mathcal{G})^{\mathfrak{i}}.$$

This is called the reduced Iwahori-spherical C^* -algebra. The spectrum of the C^* -algebra \mathfrak{A} comprises all irreducible tempered representations of \mathcal{G} which admit a nonzero Iwahori-fixed vector.

Let \mathcal{G}^\vee denote the (complex) Langlands dual of \mathcal{G} , let $\mathcal{T}_{\mathbb{C}}$ denote a maximal torus in \mathcal{G}^\vee , and let \mathcal{T} denote the maximal compact subgroup of $\mathcal{T}_{\mathbb{C}}$.

According to [1, Eq. (4.9)], we have

$$K_j(\mathfrak{A}) \otimes_{\mathbb{Z}} \mathbb{Q} \simeq K_W^j(\mathcal{T}) \otimes_{\mathbb{Z}} \mathbb{Q}$$

The results in Sect. 5.1 now lead immediately to the following answer for the K -theory of \mathfrak{A} :

$$K_0(\mathfrak{A}) \otimes_{\mathbb{Z}} \mathbb{C} = \mathbb{C}^{47}, \quad K_1(\mathfrak{A}) \otimes_{\mathbb{Z}} \mathbb{C} = \mathbb{C}^{11}.$$

From the point of view of noncommutative geometry, the C^* -algebra \mathfrak{A} behaves, at the level of K -theory (after tensoring by \mathbb{C}), as if its spectrum was equal to the extended quotient $\mathcal{T} // W$.

Acknowledgements The authors would like to thank the referee for their comments and the community of developers and mathematicians who write and maintain the GAP computer algebra system. While none of the arguments given here rely on computer calculations, and the conjugacy class representatives were built carefully by hand to simplify the construction of centralisers and their actions, we found the Lie Algebra data package provided by Willem Adriaan de Graaf and Thomas Breuer very helpful in our initial investigations.

Data Availability The research in this paper is self-contained and does not rely on any dataset.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Aubert, A.-M., Baum, P.F., Plymen, R., Solleveld, M.: Conjectures about p -adic groups. In: Brumley, F., Aparicio, M.P., Mínguez, A. (eds.) *Around Langlands Correspondences*, pp. 15–51. *Contemporary Mathematics*, vol. 691. American Mathematical Society, USA (2017)
2. Baum, P., Connes, A.: Chern character for discrete groups. In: Matsumoto, Y., Mizutani, T., Morita, S. (eds.) *A fête of Topology*, pp. 163–232. Academic Press, Cambridge (1988)
3. Brewer, J.P.: Complex reflection subgroups of real reflection groups. *J. Algebra* **232**, 310–330 (2000)
4. Carter, R.W.: Conjugacy classes in the Weyl group. *Compos. Math.* **25**(1), 1–59 (1972)
5. Niblo, G.A., Plymen, R., Wright, N.: Poincaré duality and Langlands duality for extended affine Weyl groups. *Ann. K-Theory* **3**(3), 491–522 (2018)
6. Niblo, Graham A., Plymen, Roger, Wright, Nick: Stratified Langlands duality in the A_n tower. *Journal of Noncommutative Geometry* **13**(1), 193–225 (2019)
7. Reeder, M.: Elliptic centralizers in Weyl groups and their coinvariant representations. *Represent. Theory* **15**, 63–111 (2011)
8. Reeder, M., Levy, P., Yu, J.-K., Benedict, H.: Gross gradings of positive rank on simple Lie algebras. *Transform. Groups* **17**(4), 1123–1190 (2012)
9. Scott, P.: The geometries of 3-manifolds. *Bull. Lond. Math. Soc.* **15**(5), 401–487 (1983)
10. Shephard, G.C.: Regular complex polytopes. *Proc. Lond. Math. Soc.* **3**(1), 82–97 (1952)
11. Shephard, G.C., Todd, J.A.: Finite unitary reflection groups. *Can. J. Math.* **6**, 274–304 (1954)
12. Springer, T.A.: Regular elements of finite reflection groups. *Invent. Math.* **25**, 159–198 (1974)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.