LETTER

A Response to: Letter to the Editor Regarding Combination of Tocilizumab and Steroids to Improve Mortality in Patients with Severe COVID-19 Infection: A Spanish, Multicenter, Cohort

Belén Ruiz-Antorán · Aránzazu Sancho-López · Ferrán Torres ·
Ana Fernández-Cruz on behalf of the TOCICOV study group

Received: March 25, 2021 / Accepted: March 26, 2021 / Published online: April 29, 2021
© The Author(s) 2021

Keywords: Tocilizumab; Steroids; COVID-19; SARS-CoV-2; Mortality

DIGITAL FEATURES

To view digital features for this article go to
<https://doi.org/10.6084/m9.figshare.14308061>.

Dear Editor,

We appreciate the interest in our manuscript [1], and thank Dr. Janapala et al. [2] for the

letter, and the Editor for giving us the opportunity to respond.

In response to Dr. Janapala, we would like to note that we are fully aware of the limitations of retrospective observational studies and the risk of bias derived from diverse factors, but mainly from differences in baseline characteristics between groups. Nevertheless, there are different strategies that can be implemented to minimize these limitations and increase the robustness of the conclusions reached. In our study [1], specifically to avoid bias derived from differences between groups and account for potential confounders, we have used a well-established and scientifically sound methodology, the inverse probability of the treatment weights (IPTW) technique. The details, and some key scientific references on which these methods rely on, are provided in our manuscript. The IPTW technique generates a pseudo-population in which the two groups are balanced across baseline covariates. Any reader can therefore check the baseline comparability in Table 1 of our manuscript, and confirm that all variables achieved a $< |11\%$ standardized difference (STD), except for the interleukin-6 (IL-6). The latter remained unbalanced since the amount of missingness did not allow us to compensate for that observed. The IPTW analysis for assessment of the study outcomes has been based on this well-balanced population.

B. Ruiz-Antorán (✉) · A. Sancho-López
Clinical Pharmacology Department, Instituto de Investigación Sanitaria Puerta de Hierro-Segovia de Arana, Hospital Universitario Puerta de Hierro-Majadahonda, Madrid, Spain
e-mail: bruizantorán@gmail.com

F. Torres
Medical Statistics Core Facility, August Pi i Sunyer Biomedical Research Institute, Hospital Clinic, Barcelona, Spain

F. Torres
Biostatistics Unit, Faculty of Medicine, Autonomous University of Barcelona, Barcelona, Spain

A. Fernández-Cruz
Infectious Diseases Unit, Internal Medicine Department, Instituto de Investigación Sanitaria Puerta de Hierro-Segovia de Arana, Hospital Universitario Puerta de Hierro-Majadahonda, Madrid, Spain

Dr. Janapala et al. might have misunderstood the methodology, focusing on the raw data instead of the IPTW analysis for assessing baseline comparability. Also, by no means have we substantiated the baseline comparability on a hypothetical lack of statistical significance but on small STD. Please note that, in fact, no *p* values have been provided for the IPTW baseline comparability.

With regard to the time from the onset of symptoms of COVID-19 to the inclusion of patients in the study (day 0 for the control group), it is noted that, as a matter of fact, the mean (SD) results are also provided in Table 1, and for the IPTW analysis were 10.41 (4.71) and 10.12 (5.42) for the tocilizumab and the control group, respectively. The median [25th–75th percentiles] were not reported in the manuscript but they were also similar: 10 [7–13] and 9 [6–13] for tocilizumab and the control group, respectively. Notably, the STD were -0.2% , which may be considered very small on the STD scale, thus ruling out any potential impact of this variable on the study outcomes.

Finally, concerning the use of medicinal drugs other than the study treatment of interest (tocilizumab), we would like to point out that all these treatments were used in the context of clinical practice, and all of them were used concomitantly with tocilizumab (and not instead of). At the time of the analysis of our study, continuous evidence became available regarding the lack of efficacy of hydroxy-chloroquine, azithromycin, and lopinavir/ritonavir in the treatment of COVID-19, so it was not considered appropriate to carry out a subgroup analysis to account for the potential influence of these drugs, as they had already proved to be ineffective.

Additionally, the hypothesis of our study was to evaluate the efficacy of tocilizumab as anti-inflammatory and immunomodulatory therapy in patients with COVID-19 who are purportedly in the inflammatory phase of the disease, where steroids would have a similar and potentially synergic role. This, together with the positive results that had been reported for steroids, led us to consider it appropriate to analyze the effect of combined use of steroids plus tocilizumab therapy.

In conclusion, in spite of the fact that our study is an observational one and, as such, has the inherent limitations thereof, our findings raised the need to conduct appropriate randomized clinical studies to confirm results. We are pleased to see that consistent benefits have been shown in several published randomized controlled studies, e.g., the RECOVERY trial [3], which has recently demonstrated that tocilizumab is an effective treatment for hospitalized patients with COVID-19 who have hypoxia and evidence of inflammation, where the benefits of tocilizumab were clearly seen among those also receiving treatment with a systemic corticosteroid.

ACKNOWLEDGEMENTS

We thank the members of the TOCICOV study group for their contribution to the work.

Funding. No funding or sponsorship was received for this study or publication of this article.

Authorship. All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship for this article, take responsibility for the integrity of the work as a whole, and have given their approval for this version to be published.

Authors' Contributions. Conceptualization and study design: A.S.L., B.R.A., A.F.C. Methodology: B.R.A., A.F.C., A.S.L., F-T. Data collection: All authors. Data interpretation: A.S.L., B.R.A., A.F.C. F.T. Writing first draft: A.S.L., B.R.A., A.S.L., F-T, A.F.C. Critical revision for important intellectual content: All authors. Final approval: All authors. All authors agree to be accountable for all aspects of the work by ensuring that questions related to the accuracy or integrity of any part of the work will be appropriately investigated and resolved. A.F.C. and B.R.A. had full access to all the data in this study and take complete responsibility for the integrity of the data and the accuracy of the data analysis.

Disclosures. Belén Ruiz-Antorán, Aránzazu Sancho López, Ferrán Torres and Ana Fernández-Cruz have nothing to disclose.

Compliance with Ethics Guidelines. This article is based on previously conducted studies and does not contain any new studies with human participants or animals performed by any of the authors.

Open Access. This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Ruiz-Antorán B, Sancho-López A, Torres F, Moreno-Torres V, et al. Combination of tocilizumab and steroids to improve mortality in patients with severe COVID-19 infection: a Spanish, multicenter, cohort study. *Infect Dis Ther.* 2020. <https://doi.org/10.1007/s40121-020-00373-8>.
2. Janapala RN, Patel J, Belfaqeeh O, Alhashmi A, Pourmand A. Tocilizumab and steroids, mortality benefits in severe COVID-19 patients. *Infect Dis Ther.* 2021
3. Horby PW, Pessoa-Amorim G, Peto L, et al. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): preliminary results of a randomised, controlled, open-label, platform trial. *medRxiv.* 2021. <https://doi.org/10.1101/2021.02.11.21249258>.