Erratum to: Characterization of Five Kinds of Wood Vinegar Obtained from Agricultural and Forestry Wastes and Identification of Major Antioxidants in Wood Vinegar

LI Zhanchao¹, ZHANG Zhiquan¹, WU Lijie², ZHANG Hanqi¹ and WANG Ziming^{1*}

1. College of Chemistry, Jilin University, Changchun 130012, P. R. China;

2. College of Chinese Materia Medica, Tianjin University of Traditional Chinese Medicine, Tianjin 300193, P. R. China

Erratum to: Chem. Res. Chinese Universities

© Jilin University, The Editorial Department of Chemical Research in Chinese Universities and Springer-Verlag GmbH 2019

doi: 10.1007/s40242-019-8207-5

After publication of our article^[1], it came to our attention that there were errors in some of the figure legends.

The legend to Fig.2 was incorrect. The corrected legend to Fig.2 is given below.

Fig.2 VCEACs of wood vinegar and fractions from wood vinegar

WV1: oak wood vinegar; WV2: walnut shell wood vinegar; WV3: peanutshell woodvinegar; WV4: ricehuskwood vinegar; WV5: corn cob wood vinegar. The concentration of WV: 10 mL/L.

The presentation of r values in Fig.3(D), Fig.4(D), Fig.4(E) and Fig.4(H) were incorrect. The corrected r values are given below. Fig.3(D): $r=-0.6800$, Fig.4(D): $r=-0.9837$, Fig.4(E): $r=-0.1714$, Fig.4(H): $r=-0.1005$.

The legend to Fig.3 was uncomplete. The corrected legend to Fig.3 is given below.

Fig.3 Regression equations and Pearson's correlation coefficients between the contents of components in phenolic extractives and VCEACs of phenolic extractives ($n=3$, $R=\text{adjusted } r^2$)

(A) Total phenolic compounds; (B) 2-methoxyphenol; (C) 2,6-dimethoxyphenol; (D) phenol; (E) 1-(4-hydroxy-3-methoxyphenyl)-2-propanone; (F) 3-methyl-1,2-cyclopentanedione; (G) 2-methoxy-4-methyl-phenol; (H) 4-ethyl-2-methoxy-phenol; (I) 4-methylphenol.

The legend to Fig.4 was uncomplete. The corrected legend to Fig.4 is given below.

Fig.4 Regression equations and Pearson's correlation coefficients between the contents of components in five kinds of wood vinegar and VCEACs of five kinds of wood vinegar($n=3$, $R=\text{adjusted } r^2$)

(A) Total phenolic compounds; (B) 2-methoxyphenol; (C) 2-methoxy-4-methyl-phenol; (D) phenol; (E) 4-methylphenol; (F) 4-ethyl-2-methoxyphenol; (G) 2,6-dimethoxyphenol; (H) 1-(4-hydroxy-3-methoxyphenyl)-2-propanone; (I) 3-methyl-1,2-cyclopentanedione; (J) 2-methylphenol.

References

- [1] Li Z., Zhang Z., Wu L., Zhang H., Wang Z., *Chem. Res. Chinese Universities*, **2019**, 35(1), 12(<https://doi.org/10.1007/s40242-019-8207-5>)

*Corresponding author. Email: wangziming@jlu.edu.cn

Received February 19, 2019; accepted May 29, 2019.

© Jilin University, The Editorial Department of Chemical Research in Chinese Universities and Springer-Verlag GmbH