Mechanisms and Management of Chimeric Antigen Receptor T-Cell Therapy-Related Toxicities

Bhagirathbhai R. Dholaria¹ · Christina A. Bachmeier² · Frederick Locke¹

Published online: 17 December 2018
© Springer Nature Switzerland AG 2018

Abstract

Chimeric antigen receptor T-cell (CAR-T) therapy has proven to be a very effective cancer immunotherapy. Axicabtagene cilomelimumab and tisagenlecleumab are the first-in-class anti-CD19 CAR-T currently available for relapsed/refractory adult large B-cell lymphoma. Tisagenlecleumab is also available for pediatric and young adult (up to age 25 years) patients with relapsed/refractory B-acute lymphoblastic leukemia. Cytokine release syndrome (CRS) and CAR-T-associated encephalopathy syndrome (neurotoxicity) are the most common adverse effects associated with CAR-T therapy. They can lead to significant morbidity and preclude widespread use of this treatment modality. Treatment-related deaths from severe CRS and cerebral edema have been reported. There is a significant heterogeneity in the side-effect profile of different CAR-T products under investigation and there is a need to develop standardized guidelines for toxicity grading and management. Here, we summarize the current literature on pathogenesis, clinical presentation, and management of CRS and neurotoxicity. The different grading systems of CRS and management protocols used in different trials have made it difficult to compare the outcomes of different CAR-T therapies. Several prevention strategies such as predictive biomarkers of CRS and neurotoxicity and modified CAR-T with ‘built-in’ safety mechanisms are being studied, with the potential to greatly expand the safety and applicability of CAR-T treatment across various malignancies.

Key Points

Cytokine release syndrome (CRS) and neurotoxicity are two important chimeric antigen receptor T-cell (CAR-T) therapy-associated toxicities.

Incidence and severity of CRS and neurotoxicity varies by CAR-T product, underlying malignancy, and patient characteristics.

Tocilizumab (anti-IL-6 receptor antibody) and steroids are the current mainstays for management of CAR-T therapy-associated toxicities.

Further study of the pathophysiology behind CAR-T therapy-associated toxicities will provide new insights into prevention and management strategies.

1 Introduction

Chimeric antigen receptor (CAR) T-cell therapy is an immunotherapeutic approach that utilizes gene transfer to reprogram T cells to recognize and eliminate cancerous cells by targeting and interacting with cell surface antigens specific to the tumor. The CAR consists of an extracellular domain that can bind to a target molecule expressed on the surface of tumor cells, a transmembrane domain, and an intracellular domain that provides an activation signal to T cells when the extracellular domain is engaged with its target. The extracellular domain usually comprises the antigen-recognition regions of an antibody, in the form of a single-chain variable fragment (scFv) or ligands of cell-surface receptors. The intracellular domain usually incorporates a region of the T-cell receptor (TCR) CD3ζ chain to provide an activation signal. First-generation CARs had very limited efficacy in pre-clinical models. Early clinical trials limited to solid organ malignancies without routine use of conditioning lymphoablative chemotherapy (i.e., fludarabine, cyclophosphamide) failed to show meaningful clinical responses and had minimal systemic toxicities [1, 2]. Second-generation CARs have an additional domain from a co-stimulatory receptor, such as CD28,

✉ Frederick Locke
frederick.Locke@moffitt.org

¹ Department of Blood and Marrow Transplantation and Cellular Immunotherapy, Moffitt Cancer Center, 12902 Magnolia Drive, FOB-3, Tampa, FL 33612, USA

² Department of Pharmacy, Moffitt Cancer Center, Tampa, FL, USA

OX40 (CD134), and/or 4-1BB (CD137) to provide a second activation signal: axicabtagene ciloleucel (axi-cel) has a CD28 domain and tisagenlecleucel (tisa-cel) has a 4-1BB domain. An elegant clinical trial demonstrated the increased expansion capacity of second-generation CD19 CAR T cells (CAR-T) as compared with first-generation for lymphoma; however, this was conducted without conditioning chemotherapy [3]. The addition of these co-stimulatory receptors and routine use of conditioning lymphoablative chemotherapy prior to CAR-T infusion have greatly enhanced the efficacy of CAR-T therapy, yet it also led to increased systemic toxicities.

The production of CAR-T is a multistep process that includes collection of white blood cells (including T cells) via leukapheresis, introduction of CAR construct typically via a viral vector (replication-defective lentivirus or gammaretrovirus) followed by cell expansion and cryopreservation [4]. Multiple clinical trials are ongoing in both hematological and solid organ malignancies targeting various cell surface tumor antigens with autologous or allogenic CAR-T [5]. ZUMA-1, a phase II trial of axi-cel in refractory large B-cell lymphoma, showed a complete response rate (CR) of 54% with overall survival (OS) of 52% at 18 months median follow-up [6]. Another CD19 CAR-T product from the University of Pennsylvania (CTL019) showed similar responses in relapsed/refractory diffuse large B-cell lymphoma (DLBCL) or follicular lymphoma with CR in 64% of the patients [7]. In the pivotal JULIET trial, tisa-cel showed an overall remission rate of 81% after a single infusion in relapsed or refractory B-cell acute lymphoblastic lymphoma (B-ALL) [8]. The recent approvals of tisa-cel for refractory pediatric and adolescent/young adult B-ALL and both tisa-cel and axi-cel for adult relapsed large B cell lymphoma by the United States FDA and European Commission (EC) are major advancements in the field of cancer immunotherapy [9–11]. These therapies offer a new hope to the patients who are refractory to conventional treatments with no effective salvage options.

CAR-T therapy is associated with unique toxicities related to immune-system activation. Cytokine release syndrome (CRS) and CAR-T-related encephalopathy syndrome (neurotoxicity) are two major complications that can lead to significant morbidity and mortality [12, 13]. Hemophagocytic lymphohistiocytosis/macrophage activation syndrome (HLH/MAS) is a rare and potentially fatal complication of CAR-T therapy [6]. These toxicities limit the application of CAR-T therapy to patients with good performance status and adequate organ function. Along with clinical development of CAR-T therapy, there have been significant efforts to understand the pathophysiology and improve prevention and management strategies of

CAR-T-related toxicities. Prompt diagnosis and treatment is imperative for the management of CRS and neurotoxicity to prevent adverse outcomes. In this article, we provide a review of the clinical presentation, pathophysiology, and management of CRS and neurotoxicity. We also consider potential future developments in CAR-T to prevent and effectively treat these complications.

We performed a thorough review using PubMed and meeting abstract databases from the American Society of Clinical Oncology (ASCO), the American Association of Cancer Research (AARC) and the American Society of Clinical Oncology (ASH) updated through June 30, 2018. We narrowed our search with the following keywords and MeSH terms: chimeric antigen receptor T-cell therapy, cellular immunotherapy, high-grade B-cell lymphoma, diffuse large B-cell lymphoma, multiple myeloma, acute lymphoblastic leukemia, CRS, neurotoxicity. Studies reviewed here must have had at least preliminary results released before the date of the search. Two authors (BD and CB) reviewed the full-text articles and meeting abstracts and summarized the clinical data. Senior author (FL) acted as a curator. Some observations regarding clinical presentations of CRS and neurotoxicity are based on our institutional experience. At the Moffitt Cancer Center, we had infused 50 patients with commercial CAR-T products and are approaching 100 additional patients with investigational products on clinical trials, as of November 1, 2018.

2 Cytokine Release Syndrome (CRS)

The most common toxicity associated with CAR-T is CRS (Table 1). The precise pathophysiology behind CAR-T-associated CRS remains to be defined. CRS is a constellation of inflammatory symptoms caused by the activation of T cells and subsequent release of cytokines, as well as the recruitment and activation of other immune cells [14]. These cytokines include interleukin (IL)-6, interferon- γ , IL-10, and IL-2 and may be produced by the CAR-T directly or by other cells such as monocytes/macrophages in response to cytokines produced by the CAR-T. Recent reports demonstrated that host-derived monocyte/macrophage and CAR-T interactions play an important role in CRS pathophysiology. In a murine model, CAR-T promoted recruitment and proliferation of monocytes by direct cell contact between CD40 (dendritic cell/monocyte/macrophage) and CD40 ligand (T cell), which in turn produced IL-1, IL-6, and nitric oxide (NO) [15]. It was also demonstrated that depletion of macrophages before CAR-T infusion leads to elimination of CRS in xenograft human leukemia mouse models treated with CD19 CAR-T [15, 16]. Additional investigation as to the interactions

between CAR-T and other cells, whether it be cytokine driven or cell contact-mediated effects, is warranted.

2.1 Clinical and Laboratory Manifestations of CRS

CRS usually manifests with constitutional symptoms with the hallmark being fever; however, symptoms vary greatly and can affect any organ system, including cardiovascular, gastrointestinal, hepatic, renal, respiratory, hematological, and nervous systems. Hay et al. examined 133 adult patients with relapsed/refractory B-cell malignancies who received CD19 CAR-T with a 4-1BB costimulatory domain. Patients with grade 4 CRS developed a fever earlier (median days after CAR-T infusion 0.4, vs 3.9 for grades 1-3) and the fever peaked sooner (median days after CAR-T infusion 2.8, vs 5.7 for grades 1-3), with a higher maximum temperature. Temperature ≥ 38.9 °C within 36 h of CAR-T infusion has 100% sensitivity and 84% specificity for grade ≥ 4 CRS [17]. Thus, the onset and peak time for fever after CAR-T infusion may guide CRS prevention strategies.

The onset of CRS varies from hours to several days after CAR-T infusion with differences noted between CAR-T products, disease state, and severity of CRS [12–14, 17]. Patients treated with anti-CD19-CD28-CD3 ζ CARs may experience CRS earlier than those treated with anti-CD19-4-1BB-CD3 ζ CARs [12]. This disparity may be due to differences in pharmacokinetic profiles as CD28 CAR constructs exhibit a greater peak expansion in vivo, while 4-1BB CAR T cells may exhibit greater longevity [18]. In ZUMA-1, the onset of CRS was a median of 2 days following infusion (range 1–12) [6]. In contrast, the median time to onset of CRS was 3 days (range 1–22) in the ELIANA trial with tisa-cel in pediatric and young adult acute B-ALL [19].

Commensurate with the immune-mediated signs and symptoms of CRS, patients with severe CRS typically have higher levels of IL-6, interferon- γ , C-reactive protein (CRP), and ferritin than patients who did not experience severe CRS, although their ability to be predictive of impending toxicity is unverified [20]. In the previously mentioned report of 133 recipients of CD19 CAR-T biomarkers were evaluated to determine those associated with severe CRS. Patients who had grade ≥ 4 CRS demonstrated higher ferritin and CRP compared with those who experienced grade ≤ 3 CRS. Furthermore, the patients with grade ≥ 4 CRS exhibited higher concentration of interferon- γ , IL-6, IL-8, IL-10, IL-15, monocyte chemoattractant protein-1 (MCP-1), tumor necrosis factor receptor p55 (TNFRp55), and macrophage inflammatory protein-1 β (MIP-1 β) within 36 h of CAR-T infusion compared with the degree of elevation of these cytokines in the patients with grade ≤ 3 CRS.

Higher tumor burden and CAR-T dose were also associated with severe CRS [17]. This is consistent with previously reported CD19 CAR-T data [20–22]. In the pivotal ZUMA-1 trial, several biomarkers were significantly associated with grade 3 or higher CRS, including but not limited to IL-15, IL-6, interferon- γ , IL-10, IL-8, and granzyme B. Of note, it was also demonstrated that CAR-T peak expansion and area under the curve were significantly associated with grade 3 or higher neurologic events, but not grade 3 or higher CRS [6].

Unfortunately, real-time rapid cytokine level measurement is not currently feasible or widely available, necessitating the use of available surrogate indicators for CRS. CRP has demonstrated a correlation with CRS progression, increasing with the onset of CRS and returning to baseline with CRS resolution [22]. CRP is a useful daily monitoring tool for CRS in the CAR-T therapy patients; however, this correlation is not detected in all patients and intervention for CRS should not be based on CRP alone [12]. Our experience with CAR-T-treated patients on clinical trials and commercial Yescarta[®] indicates that CRP elevation does not always adequately precede CRS to be a reliable marker to guide an intervention, as the elevation of CRP appears to occur concurrently with the clinical signs and symptoms of CRS. However, anecdotally we have seen its decrease herald resolution of CRS signs and symptoms and help guide the tapering of systemic therapy. The verification of predictive biomarkers for CRS remains incomplete and additional confirmatory research is necessary. The number one goal of CRS management is to prevent life-threatening toxicities. However, any intervention to block cytokine release could in theory block CAR-T activity and compromise anti-tumor activity. Current evidence suggests that anti-tumor activity can be preserved even in patients treated with tocilizumab and corticosteroids [6].

2.2 CRS Grading

The Common Terminology Criteria for Adverse Events version 4.03 (CTCAE v4.03) [23] was used for grading of organ toxicities throughout many CAR-T trials. Typically, the organ toxicity associated with CRS was graded by this scale, but CRS requires grading of the entire syndrome. The CTCAE v4 contains grading criteria for CRS (Table 2); however, this was not created for cellular therapy and is directed towards management of infusion reactions with immunotherapies. This grading system is insufficient for CRS symptoms that occur days after the CAR-T infusion. Single-center and pivotal trials of CAR-T therapy have utilized differing criteria, making cross-comparison of toxicity severities difficult. Lee et al. created the National Cancer Institute (NCI) consensus criteria to define mild, moderate, severe, and life-threatening CRS associated with high-risk immunotherapies and to guide treatment recommendations based on this

Table 1 Incidence and treatment of CAR-T-related cytokine release syndrome and neurotoxicity in published clinical trials

Study	CAR-T construct	Diseases	Number of patients	Rate of CRS	Grade III-IV CRS	Death from CRS	Rate of neuro-toxicity	Grade III-IV neurotoxicity	Death from neurotoxicity	Treatment (CRS + neuro-toxicity)
Borchmann et al. (2018) (JULIET) [8]	CD19-4-1BB (tisagenlecleucel; CTL019)	DLBCL	111	57 (58%)	23 (23%)	0	23 (21%)	12 (12%)	0	Tocilizumab 15%, Glucocorticoids 11%
Maude et al. (2018) (ELI-ANA) [19]	CD19-4-1BB	B-ALL	75	58 (77%)	35 (46%)	1 (1%)	30 (40%)	10 (13%)	0	
Neelapu et al. (2017) (ZUMA-1) [6]	CD19-28- ζ (axicabtagene ciloleucel; axi-cel, KTE-C19)	DLBCL	101	94 (93%)	13 (13%)	2 (2%)	65 (64%)	28 (28%)	0	Glucocorticoids 27%, Tocilizumab 43%
Schuster et al. (2017) (Novartis/U. Penn) [7]	CD19-4-1BB	DLBCL	28	16 (57%)	5 (18%)	0	11 (39%)	3 (11%)	1 (4%)	Glucocorticoids 0%, Tocilizumab 4%
Abramson et al. (2018) (TRAN-SCEND NHL 001) [45]	CD19-4-1BB (lisocabtagene maraleucel; liso-cell; JCAR017)	DLBCL	102	38 (37%)	1 (1%)	0	23 (23%)	13 (13%)	0	Glucocorticoids 21%, Tocilizumab 17%
Park et al. (2018) (MSKCC) [43]	CD19-28- ζ	B-ALL	53	45 (85%)	14 (26%)	1 (2%)	23 (43%)	22 (42%)	0	Glucocorticoids 7.5%, Tocilizumab 11%, Glucocorticoids + Tocilizumab 25%
Maude et al. (2014) (U. Penn) [20]	CD19-4-1BB	B-ALL	30	30 (100%)	8 (27%)	0	13 (43%)	NA	0	Glucocorticoids 20%, Tocilizumab 30%
Ghorashian et al. (2017) (NHS) [64]	CD19-4-1BB (low affinity)	B-ALL	8	8 (100%)	0	0	3 (38%)	0	0	Tocilizumab 0%
Qasim et al. (2017) (NHS) [75]	CD19-4-1BB with RQR8 safety switch (allogeneic CAR-T)	B-ALL	5	5 (100%)	1 (20%)	0	2 (40%)	0	0	Tocilizumab 20%

Table 1 (continued)

Study	CAR-T construct	Diseases	Number of patients	Rate of CRS	Grade III-IV CRS	Death from CRS	Rate of neuro-toxicity	Grade III-IV neurotoxicity	Death from neurotoxicity	Treatment (CRS + neuro-toxicity)
Gust et al. (2017) (Juno) [39, 51]	CD19-4-1BB	B-ALL, NHL, CLL	133	93 (70%)	16 (12%)	5 (4%)*	53 (40%)	28 (21%)	4 (3%)	Glucocorticoids 11%, Tocilizumab 2%, Glucocorticoids + Tocilizumab 38% NA
DeAngelo et al. (2017) (ROCKET) [40]	CD19-28-ζ (JCAR015)	B-ALL	38	NA	8 (21%)	0	NA	20 (52%)	5 (13%)	NA
Cohen et al. (2017) (U. Penn) [58]	BCMA-4-1BB	MM	21	18 (86%)	6 (29%)	0	3 (14%)	1 (5%)	0	Tocilizumab 19%
Raje et al. (2018) [55]	BCMA-4-1BB	MM	43	27 (63%)	2 (5%)	0	14 (33%)	1 (2%)	0	Glucocorticoids 9%, Tocilizumab 21%

*Combined deaths from CRS and neurotoxicity

B-ALL B-cell acute lymphoblastic lymphoma, CAR-T chimeric antigen receptor T cell, CLL chronic lymphocytic leukemia, CRS cytokine release syndrome, DLBCL diffuse large B-cell lymphoma, MM multiple myeloma, MSKCC Memorial Sloan Kettering Cancer Center, NA not available, NHL non-Hodgkin lymphoma, NHS National Health Service

grade. The Lee et al. grading criteria is included in Table 2 and is based on constitutional symptoms, oxygen requirement, indication for vasopressors, and grade of organ toxicity [14]. The Lee grading system was used in the ZUMA-1 trial [6]. Of note, CTCAE version 5.0, released in November 2017, has modified the grading of CRS to correlate closely with the original Lee criteria (Table 3) [24].

A modification of the Lee grading system was proposed by Neelapu and colleagues (CARTOX group). These recommendations expand upon the Lee et al. grading system to develop a more consistent approach to monitoring, grading, and management of CAR-T-associated toxicities in adult patients. CRS grade was concordant with the Lee et al. CRS grading, considering temperature, systolic blood pressure, oxygen requirement, and organ toxicity with only slight modification in relation to transaminase elevation. The CARTOX grading was then combined with additional CARTOX consensus management and intervention strategies [12, 25].

An alternative grading system based on clinical parameters created by Porter et al. was used in the ELIANA and JULIET trials [19]. This grading scale was originally created and used for CRS classification in chronic lymphocytic leukemia (CLL) patients receiving CAR-T at the University of Pennsylvania. The PENN grading criteria is included in Table 2 and, as with the Lee et al. scale, is based on clinical features and organ toxicity as well as need for vasopressors and oxygen. Need for hospitalization is also included, although many patients have CAR-T infused in the inpatient setting [26, 27].

When interpreting rates of reported CRS, it is important to be aware of the grading system utilized. The PENN grading scale created by Porter et al. considers all patients with hypoxemia requiring oxygen grade 3 CRS, whereas the Lee et al. scale considers patients with an oxygen requirement < 40% a grade 2. Additionally, the Lee et al. scale classifies hypotension responding to intravenous (IV) fluids or a low-dose vasopressor as grade 2; hypotension requiring intervention is grade 3 CRS under the PENN scale. Subsequently, hypotension requiring high-dose or multiple vasopressors is a grade 4 under the PENN scale, but a grade 3 when using the Lee et al. scale. Given these differences, the PENN CRS grading scale likely leads to a higher number of grade 3 and 4 CRS ratings than if the Lee et al. system is used for the same subset of patients and graded in an otherwise consistent manner. Additional observational studies comparing these grading scales would be helpful to interpret pivotal trial data for different products. With a goal to unify the CRS grading system, The American Society of Blood and Marrow Transplantation (ASBMT) is in the process of developing both immune effector cell CRS and neurotoxicity grading systems for CAR-T and related therapies [28].

2.3 Management of CRS

The clinical manifestations and severity of CRS varies greatly from mild constitutional symptoms to life-threatening severe toxicities. Adverse outcomes likely can be avoided with accurate and prompt evaluation and patient management. The management of these patients is complicated by overlapping conditions. The hallmark of CRS, fever, as well as other symptoms of CRS mirrors the presentation of infection. CRS symptoms typically resolve within 2 weeks of CAR-T infusion. This toxicity can be self-limiting, requiring only symptomatic care, or may require treatment with an IL-6 antagonist and/or glucocorticoids. The goal of treatment of CRS is to avoid harmful toxicities while maximizing the anti-tumor effect of the cellular therapy.

2.3.1 Location of Care

There is no consensus on need for hospitalization of patients receiving CAR-T therapy. In the ZUMA-1 study of adult patients receiving axicabtagene ciloleucel, all patients were hospitalized for the CAR-T infusion and for a minimum of 7 days afterwards [6]. In the pivotal JULIET trial that led to approval of tisa-cel for adult patients with DLBCL, 26% received CAR-T infusion in the outpatient setting and 77% of these patients remained as outpatients for ≥ 3 days following infusion [29]. Pediatric and young adult patients receiving tisa-cel on the ELIANA trial were also able to receive product infusions in the outpatient setting [6, 19]. Long-term follow-up data on the patients treated in outpatient settings is needed. Although this demonstrates that CAR-T therapy can be administered in a well controlled outpatient setting, it is important to note that the centers doing this had significant experience with CAR-T therapy and already well established robust outpatient hematopoietic stem-cell transplant programs. Further development of predictive biomarkers is needed to identify patients at risk of severe early CRS requiring intensive monitoring and an inpatient admission before severe symptoms develop [30].

2.3.2 Supportive Care

Supportive care measures begin following cell infusion and continue throughout all stages of CRS. Daily monitoring typically includes a complete blood count with differential and complete metabolic panel. At our center we perform daily CRP and ferritin as markers of inflammation associated with CRS. These can be particularly helpful in high-risk patients or those experiencing severe CRS to trend the inflammatory state. Intravenous fluids are used to maintain hydration; however, fluid balance (including

daily bodyweight) must be monitored closely due to the risk of volume overload and pulmonary edema. Due to the risk of cardiac arrhythmias, telemetry monitoring should be considered from the time of CAR-T cell infusion until resolution of CRS, especially in patients with additional cardiac risk factors [12].

Acetaminophen may be administered for management of fever in patients with normal hepatic function; cooling blankets may also be used. Non-steroidal anti-inflammatory drugs (NSAIDs) may be used as an alternative; however, caution must be taken in the setting of thrombocytopenia. Additionally, NSAIDs may contribute to hemorrhage, gastritis, and renal insufficiency [31]. As many of these patients are neutropenic, and all receive lymphodepletion, it becomes imperative to monitor for infection. Febrile patients should be assessed for infection including blood and urine cultures and chest radiography. Broad-spectrum antibiotics should also be initiated.

2.3.3 Anti IL-6 Therapy

Tocilizumab is a humanized monoclonal antibody (mAb) against the IL-6 receptor (IL-6R), recently FDA approved for CAR-T-induced severe or life-threatening CRS [32]. Rapid resolution of CRS has been demonstrated following administration of tocilizumab in most patients. In addition, published reports suggest tocilizumab does not negatively impact CAR-T expansion or persistence [6, 22, 33]. Tocilizumab is infused over 60 min at a dose of 12 mg/kg for patients with a body weight < 30 kg and 8 mg/kg for patients weighing ≥ 30 kg. The FDA-approved dosing strategy includes the ability to administer up to three additional doses if there is no clinical improvement in the signs and symptoms of CRS, with a minimum of 8 h between consecutive doses. Additional infusions may not be warranted if a patient has responded to the therapy and CRS symptoms are not recurring.

Although the approved indication is for severe or life-threatening CRS, there are no clear recommendations on the optimal timing for administration of tocilizumab. Lee and colleagues provide a treatment algorithm based on their CRS grading assessment; tocilizumab should be administered to patients experiencing CRS of grade 3 or greater and to patients with grade 2 CRS with comorbidities. On the ZUMA-1 trial, 43% of patients received tocilizumab with no difference in response rate compared with patients that did not receive tocilizumab [6]. There was a decrease in the incidence of CRS and neurologic events of grade 3 or higher in ZUMA-1 over the course of the study (18% grade ≥ 3 CRS at the interim analysis of $n = 62$ patients to 13% grade ≥ 3 CRS at the final analysis of $n = 101$ patients [6]). This decrease might be attributed to a protocol amendment allowing for earlier intervention

Table 2 Summary of cytokine release syndrome grading systems

CRS grading	CTCAE v4 [23]	CTCAE v5 [24]	Lee et al. [14]	Porter et al. [26]
Grade 1	Mild reaction; infusion interruption not indicated; intervention not indicated	Fever with or without constitutional symptoms	Symptoms are not life threatening and require symptomatic treatment only (e.g., fever, nausea, fatigue, headache, myalgias, malaise)	Mild reaction: treated with supportive care such as anti-pyretics, anti-emetics
Grade 2	Therapy or infusion interruption indicated but responds promptly to symptomatic treatment (e.g., antihistamines, NSAIDS, narcotics, IV fluids); prophylactic medications indicated for ≤ 24 h	Hypotension responding to fluids; hypoxia responding to $< 40\%$ oxygen	Symptoms require and respond to moderate intervention Oxygen requirement $< 40\%$ or hypotension responsive to fluids or low dose of one vasopressor or Grade 2 organ toxicity	Moderate: requiring IV therapies or parenteral nutrition; some signs of organ dysfunction (i.e., grade 2 Cr or grade 3 LFTs) related to CRS and not attributable to any other condition. Hospitalization for management of CRS-related symptoms including fevers with associated neutropenia
Grade 3	Prolonged (e.g., not rapidly responsive to symptomatic medication and/or brief interruption of infusion); recurrence of symptoms following initial improvement; hospitalization indicated for clinical sequelae (e.g., renal impairment, pulmonary infiltrates)	Hypotension managed with one pressor; hypoxia requiring $\geq 40\%$ oxygen	Symptoms require and respond to aggressive intervention Oxygen requirement $\geq 40\%$ or hypotension requiring high dose or multiple vasopressors or Grade 3 organ toxicity or grade 4 transaminitis	More severe reaction: hospitalization required for management of symptoms related to organ dysfunction including grade 4 LFTs or grade 3 Cr related to CRS and not attributable to any other conditions; excludes management of fever or myalgias. Includes hypotension treated with intravenous fluids or low-dose pressors, coagulopathy requiring FFP or cryoprecipitate, and hypoxia requiring supplemental oxygen (nasal cannula oxygen, high flow oxygen, CPAP or BiPAP). Patients admitted for management of suspected infection due to fevers and/or neutropenia may have grade 2 CRS
Grade 4	Life-threatening consequences; pressor or ventilatory support indicated	Life-threatening consequences; urgent intervention indicated	Life-threatening symptoms Requirement for ventilator support or Grade 4 organ toxicity (excluding transaminitis)	Life-threatening complications such as hypotension requiring 'high-dose pressors', hypoxia requiring mechanical ventilation

BiPAP bilevel positive airway pressure, *CPAP* continuous positive airway pressure, *Cr* creatinine, *CRS* cytokine release syndrome, *CTCAE* Common Terminology Criteria for Adverse Events, *FFP* fresh frozen plasma, *IV* intravenous, *LFTs* liver function tests, *NSAIDS* non-steroidal anti-inflammatory drugs

Table 3 Comparison of different grading systems for cytokine release syndrome

Clinical parameter	CTCAE v5 [24]	Lee et al. [14]	Porter et al. [26]
Constitutional symptoms	Grade 1	Grade 1	Grade 1
Hypotension responsive to fluids	Grade 2	Grade 2	Grade 3
< 40% FiO ₂	Grade 2	Grade 2	Grade 3
High dose vasopressor requirement	Grade 3	Grade 3	Grade 4
Ventilator support	Grade 4	Grade 4	Grade 4

CTCAE Common Terminology Criteria for Adverse Events, FiO₂ fraction of inspired oxygen

for the treatment of CRS and neurologic toxicity partway through the trial. Essentially this amendment changed treatment guidelines from the Lee et al. guidelines to the CARTOX guidelines, allowing for administration of tocilizumab for grade 2 CRS possibly leading to fewer patients progressing to grade 3 CRS [6, 25].

The grading system from the University of Pennsylvania was used in conjunction with management recommendations in the ELIANA trial of tisa-cel in pediatric and young adult ALL. Tocilizumab was administered as second-line management for high fevers, hypoxia, and hypotension in response to any of the following: hemodynamic instability despite IV fluids and vasopressor support, worsening respiratory distress or rapid clinical deterioration. Based on this algorithm, tocilizumab is given later in CRS progression than when utilizing the Lee et al. algorithm or CARTOX guidelines for management of CRS. Forty percent of patients received tocilizumab for the management of CRS in this study [19]. The JULIET study of tisa-cel in adult DLBCL patients also employed the PENN CRS grading scale and a similar algorithm with 15% of patients receiving tocilizumab [29].

The CARTOX group recommended earlier administration of tocilizumab, as compared with the Lee et al. guidelines, with consideration in both grade 1 and grade 2. In grade 1 this is specifically for patients with a refractory fever lasting over 3 days and in grade 2 CRS tocilizumab is recommended for hypotension that is refractory to fluid boluses. Repeat doses of tocilizumab can also be considered in grade 2 or higher. Anti-IL-6 therapy is also recommended for patients categorized as a grade 2 CRS due to persistent hypoxia at a fraction of inspired oxygen (FiO₂) < 40% and other grade 2 organ toxicities. Both tocilizumab and glucocorticoids are recommended for management of grades 3 and 4 CRS [12]. We suggest that additional doses of tocilizumab should be considered if CRS does not improve with initial dosing. Further studies are needed to determine optimal timing between doses and efficacy with repeat doses.

Siltuximab (anti-IL-6 chimeric mAb) is another drug being used off-label for CRS, especially in tocilizumab and steroid refractory cases. It can rapidly reverse CRS symptoms when used in different clinical studies [12, 13, 20, 34].

This drug is currently approved by the US FDA for Multicentric Castleman's disease [35]. It binds directly to IL-6 with a high affinity, so there is a theoretical advantage of more complete blockage of IL-6 activity over tocilizumab, which blocks membrane-bound and soluble IL-6R. There have been reports of transient increase in IL-6 levels after the first dose of tocilizumab, possibly due to decreased IL-6 clearance in peripheral tissue via IL-6R [33, 36]. Prospective randomized studies might be considered to compare efficacy of tocilizumab and siltuximab as therapeutics for CRS patients.

2.3.4 Glucocorticoids

Glucocorticoids have also demonstrated efficacy in ameliorating CRS due to the ability to suppress inflammatory responses. Published evidence suggests glucocorticoids dampen CAR-T expansion and anti-tumor effect in ALL patients after CD-19-directed CAR-T infusion [37], [22]. In contrast, data from the ZUMA-1 study suggests glucocorticoids used for treatment of CAR-T-related toxicities do not impact objective response rates. The objective response rate for the 27 patients who received steroids was no different from that for patients that did not receive steroids [6]. Of note, this conflicting data is in patients with different disease states (ALL versus DLBCL). Due to concerns of CAR-T suppression, steroids remain a second-line treatment for CRS refractory to tocilizumab, except in extremely rapid onset cases of severe CRS, and should not be used for other non-life-threatening indications.

The trigger to initiate steroids as well as the optimal steroid and dose is not clearly defined. Lee and colleagues recommend if the patient's condition does not improve or stabilize within 24 h of the initial tocilizumab dose, a second dose of tocilizumab or glucocorticoid should be considered. This applies to patients with grade 3 and higher CRS, and elderly patients or those with significant comorbidities with grade 2 CRS. The CARTOX group includes glucocorticoids as an option of consideration for grade 2 CRS, specifically for patients at high risk of severe CRS or if hypotension persists after one to two doses of anti-IL-6 therapy. Tocilizumab

and steroids, specifically dexamethasone 10 mg IV every 6 h, is recommended in grade 3. Grade 4 CRS includes a recommendation for methylprednisolone 1000 mg IV per day followed by a rapid taper. Teachey et al. recommended restricting the starting dose of prednisone to 1 mg/kg based on their pediatric experience using CD19-4-1-BB CAR-T product [38].

In summary, optimal timing and dosages of tocilizumab and glucocorticoids in the management of CRS need to be studied systemically across different CAR-T products. Prospective intervention studies with a standardized CRS grading scale are urgently needed at this point.

3 CAR-T-Associated Neurotoxicity

3.1 Clinical Manifestations and Incidence of Neurotoxicity

Neurotoxicity is the second most common toxicity related to CAR-T therapy. Typical manifestations of neurotoxicity range from minor headache and diminished attention to seizure, severe encephalopathy, and death. The most characteristic manifestation is encephalopathy, typified by confusion progressing to expressive aphasia and at the extreme, obtundation. Early signs are language and handwriting impairment followed by confusion, agitation, hallucinations, tremors, and headaches. Seizures, motor weakness, incontinence, mental obtundation, increased intracranial pressure, papilledema, and cerebral edema can be seen in severe cases of neurotoxicity (grade > 2) [12]. The manifestations may be bi-phasic with early confusion coinciding with high fevers and CRS, and later encephalopathy often following the resolution of CRS. The incidence and severity of neurotoxicity varies by different CAR constructs. Neurotoxicity may be more frequent in the patient with pre-existing neurological conditions, younger patients, and heavily pretreated patients [39, 40].

Electroencephalography (EEG) findings in the patients with neurotoxicity-induced encephalopathy are very non-specific, with diffuse generalized slowing with or without triphasic waves [12]. The main utility of EEG is to rule out seizure activity, which can occur in a minority of these patients. Brain imaging is typically described as normal, even in the patients with severe neurotoxicity. Occasional T2/fluid attenuated inversion recovery (FLAIR) MRI hyperintensity involving the thalami, mid-brain, and cerebral edema have been reported [39, 41]. Elevated opening pressure (≥ 20 mmHg) during lumbar puncture is common in moderate to severe neurotoxicity. Elevated cerebrospinal fluid (CSF) protein and leukocyte count secondary to increased blood–brain barrier (BBB) permeability is seen in neurotoxicity patients; however, the cell count within the

CSF does not seem to be correlated with the severity of neurotoxicity [39].

Gust et al. studied 133 adults with B-ALL, non-Hodgkin lymphoma (NHL), or CLL who received CD19 CAR-T containing a 4-1BB costimulatory domain. A total of 53 of 133 patients (40%) had one or more grade ≥ 1 neurological adverse event; 7 (5%) developed grade ≥ 4 neurotoxicity and 4 patients (3%) died due to neurotoxicity within the first 28 days of CAR-T infusion. The median time for presentation of neurotoxicity was 4 days after CAR-T infusion and the majority of these patients had preceding CRS (91%) before the onset of neurotoxicity [39]. In the phase II ZUMA-1 trial, CD19 CAR-T with CD28 co-stimulatory domain were given to 101 patients with relapsed refractory large B-cell lymphoma. Neurologic events occurred in 65 patients (64%); 28% were grade 3 or higher, and no deaths were attributed to neurotoxicity. The median time of the onset of neurologic events was on day 5 (range 1–17), with median resolution on day 17 after infusion. Median duration of neurotoxicity was around 5 days (range 1–21) and typically lasted 2–4 days [6]. Prolonged memory impairment and tremor beyond 3–4 weeks after CAR-T infusion have been reported [39, 42, 43]. In the JULIET trial testing tisa-cel (CD19-4-1BB) in 111 relapsed/refractory DLBCL, incidence of overall and grade 3–4 neurotoxicity was reported at 21% and 12%, respectively; however, the FDA-approved label reported ($N=106$) 58% and 18%, suggesting that different observers may attribute neurotoxicity to CAR-T or other factors, thereby skewing summary neurotoxicity data [8, 44]. There seems to be a general trend of lower rate of neurotoxicity with 4-1BB compared with CD28 costimulatory CD19 CAR-T construct; however, comparative studies controlling for confounding factors are needed.

Table 1 describes published clinical experience with different CARs tested in multi-center trials and different rates of neurotoxicity. Due to differences in CAR construct, dose of infused CAR-T, lymphodepleting chemotherapy, and target diseases, it is difficult to cross-compare rates of neurotoxicity among different clinical trials. Most of the currently available information about neurotoxicity is based on CD19 CAR-T products, and CNS toxicities associated with non-CD19 CAR-T therapies are yet to be characterized. For DLBCL, the TRANSCEND NHL 001 study (CD19-4-1BB, Juno therapeutics) reported a low rate of neurotoxicity [45]. Alternatively, five deaths in the ROCKET trial (JCAR015, CD19-28- ζ) from cerebral edema raised an alarm, which led to termination of the trial [40]. Again, comparing neurotoxicity rates across pivotal trials is fraught with complexities based upon attribution and definition of what exact CTCAE toxicities should be considered part of the spectrum of CAR-T therapy-related encephalopathy and neurotoxicity. Improved consensus neurotoxicity definitions and grading

scales are needed and must be prospectively validated so they can be utilized universally in CAR-T clinical trials.

3.2 Pathophysiology of CAR-T-Associated Neurotoxicity

Precise underlying mechanisms behind neurotoxicity are not fully understood. Since most neurotoxicity is preceded by CRS, one hypothesis is that neurotoxicity is a manifestation of passive diffusion of cytokines into the brain in the presence of a permeable BBB. Alternatively, a recent report by Santomaso et al. demonstrated that there were disproportionately higher levels of IL-6, IL-8, MCP-1, and interferon- γ -induced protein 10 (IP10) in CSF compared with serum in the patients with severe neurotoxicity. There were also increased levels of endogenous excitatory neurotransmitters (glutamate, quinolinic acid) in CSF in those patients [46]. The data showing higher CSF cytokine levels compared with peripheral blood, and the fact that CAR-T can be found in the CSF of patients with or without severe neurotoxicity, suggests that active cytokine release from, or induced by, local CAR-T within the CNS may be a driver [33, 46, 47].

In a study involving CD19 CARs, patients with severe neurotoxicity had evidence of endothelial activation and increased BBB permeability. The concentrations of IFN γ , TNF α , IL-6, and TNFR p55 had increased significantly and were comparable between serum and CSF during the acute phase of neurotoxicity [39]. Autopsies on two patients who died from neurotoxicity followed by CD19 CAR-T showed widespread vascular lesions, cerebral edema, and necrosis with perivascular CD8 + T-cell infiltration, suggesting pervasive endothelial dysfunction and destruction [39]. Another study of B-ALL patients treated with CD19 CAR-T showed elevated levels of IL-1 α , IL-2, IL-3, IL-5, IL-6, IL-10, IL-15, IP10, INF γ , granulocyte-colony stimulating factor (GCSF), granulocyte-macrophage colony-stimulating factor (GMCSF), and MCP-1 by day 3 in patients with severe neurotoxicity [46]. Pre-existing endothelial activation before CAR-T infusion can also increase the risk for CRS and neurotoxicity. Angiopoietin-1 (ANG1) is an endothelial stabilizing cytokine and angiopoietin-2 (ANG2) promotes endothelial activation via the ANG-TIE2 axis [48, 49]. A higher ANG2: ANG1 ratio was associated with severe neurotoxicity in a study by Santomaso et al. [46]. High-intensity lymphodepletion with fludarabine before CAR-T infusion can lead to increased levels of IL-15, which is associated with greater peak CAR-T expansion and resultant neurotoxicity [50, 51]. High rates of cerebral edema in JCAR015 (phase II ROCKET trial) were attributed to rapid CAR-T expansion and elevated IL-15 levels before cell infusion [40].

It is unlikely that CNS invasion by CAR-T leads directly to neuronal cytotoxicity. Most patients exhibit a complete neurological recovery, which would be impossible

with widespread neuronal destruction. CAR-T have been detected in the CSF of patients with neurotoxicity without CNS malignancy [52, 53]. However, there is no evidence of CD19 expression in neurological tissue and neurotoxicity has been reported in the patients who received CD22 CAR-T for B-ALL and B-cell maturation antigen (BCMA) CAR-T for multiple myeloma [54, 55]. It is unknown at this point how the incidence and severity of neurotoxicity differ between CD19 and non-CD19 CAR-T therapies due to the paucity of data.

Most of the T cells in the brain parenchyma (93%) and CSF (95%) were CAR-T. A higher fraction of the CD4 + CAR-T subset in the CSF compared with blood suggests there might be a difference in the migration pattern of CD4 + and CD8 + CAR T across the BBB and their role in neurotoxicity [33, 39]. Similarly, an association with higher CD4 + CAR-T in the CSF was seen in patients that later developed severe neurotoxicity as compared with those that did not, although the ratio of CD4: CD8 T cells in the infused product did not predict the rates of CRS and neurotoxicity [6].

Only JCAR017 (Juno Therapeutics) has a 1:1 ratio of CD4 + : CD8 + CAR-T in the final product, whereas other CAR-T therapies are not separately manufacturing and combining CD4 and CD8 T cells [21]. More studies are needed with other CAR-T products in different diseases with exploratory immunophenotyping of blood, CSF, and brain tissue to validate the findings by Gust et al. [39]. Animal models of neurotoxicity have been developed, including a murine and a non-human primate model of CD20 CAR-T-mediated CRS and neurotoxicity [16, 47]. These models will allow the study of pathogenesis and test different therapeutic interventions for neurotoxicity. As with any animal model, species barrier will be a limiting factor when applying these pre-clinical findings in designing clinical trials.

3.3 Management of Neurotoxicity

CAR-T therapy-associated neurotoxicity management is guided by toxicity grading and may be informed by the severity of concurrent CRS. The CARTOX group coined the term CAR-T-associated encephalopathy syndrome (CRES) which encompasses some of the neurotoxicity symptoms and signs, although it does not address nonspecific neuropsychiatric symptoms that may or may not have been attributed to CAR-T in the pivotal trials. Table 4 shows CRES grading based on the CARTOX group's experience with CD19 CAR-T therapy in adult patients with high-grade B-cell lymphoma [12]. The CAR-T therapy-associated toxicity 10-point neurological assessment (CARTOX-10) is an easy-to-use clinical tool for bedside assessment of patients at risk for neurotoxicity and it has been incorporated into the neurotoxicity grading score [42]. It can be easily performed multiple

times per day by providers or nursing staff. It is important to consider that the CARTOX-10 and therefore the CRES neurotoxicity grading system have not been prospectively validated and that it is based primarily on axi-cel usage in adults with lymphoma. An additional shortcoming of the CRES grading scale is that it relies upon fundoscopic exam and CSF opening pressures, tests that may not be universally available or performed correctly. Mini-mental state examination (MMSE) [56] and Glasgow Coma Scale (GCS) [57] are other tools for clinical evaluation of neurotoxicity. Additional updated grading systems will be helpful to further delineate the CAR-related neurotoxicity from other likely unrelated symptoms like headache.

Neurotoxicity management involves frequent neurological evaluation and early involvement of neurology and critical care experts. The treatment is mainly supportive in grade 1/2 neurotoxicity without significant CRS with close monitoring, aspiration precautions, EEG, and CNS imaging. Lumbar puncture is ideal for opening pressure measurement but not always feasible in delirious patients with coagulopathy from CAR-T-related disseminated intravascular coagulation.

3.3.1 Anti-IL-6 Therapy and Glucocorticoids

Neurotoxicity may occur early, concurrent with CRS (approximately day 1–7), and/or later independently of CRS [12]. Patients experiencing grade ≥ 1 neurotoxicity with concurrent grade ≥ 2 CRS may benefit from anti-IL-6 therapy [10]. In contrast, neurotoxicity occurring independently of CRS should not be managed with tocilizumab as anti-IL-6 therapy does not cross the BBB and has not been associated with resolution of CAR-T-related encephalopathy [12].

Tocilizumab can cause a transient increase in IL-6 levels after the initial administration, whether its use for CRS

can initiate or exacerbate the neurotoxicity is unknown [36]. In the report by Gust et al., the peak grade of neurotoxicity occurred after the first dose of tocilizumab in 67% of patients, and in eight of those patients, the first presentation of neurotoxicity occurred after tocilizumab had been administered for CRS [39]. It is unknown if siltuximab has any advantage over tocilizumab in patients with neurotoxicity with CRS, and a prospective study is warranted.

In patients with grade ≥ 2 neurotoxicity without CRS, initial treatment with a steroid should be considered over anti-IL-6 therapy. Optimal dose, duration, and choice of steroid agent remain to be standardized. Our preferred regimens for grade 2–3 neurotoxicity are dexamethasone 10 mg IV every 6 h or methylprednisolone 1 mg/kg IV every 12 h with rapid taper over 7–10 days depending on clinical improvement. In refractory cases with grade 4 neurotoxicity, high-dose steroid therapy with methylprednisolone 1 g/day IV for 3 days may be considered followed by a rapid taper. Close monitoring for recurrence of neurotoxicity is needed during steroid taper. Anti-IL-6 therapy is added if a patient develops concurrent CRS as described above [42]. The median time to resolution of neurotoxicity was 4 days (range 1–64) in CD19 CAR-T-treated adult patients with relapsed B-cell malignancies [39]. Neurotoxicity treatment response is generally slower than response to CRS symptoms. The published clinical experience with different CD19 CAR-T therapies so far suggests that there may not be a co-relation between the use of anti-IL-6 therapy and steroids and the efficacy of CAR-T therapy [6, 22, 43, 51].

Of note, in the clinical trials evaluating the use of tisa-cel in both pediatric and adult B-ALL (ELIANA trial) and adult DLBCL (JULIET trial), glucocorticoids were not mandated for the treatment of neurotoxicity, and management may have consisted of supportive care only [8, 19].

Table 4 Grading of CAR-T-related neurotoxicity per CARTOX group^a

Symptom or sign	Grade 1	Grade 2	Grade 3	Grade 4
Neurological assessment score (by CARTOX-10)	7–9 (mild impairment)	3–6 (moderate impairment)	0–2 (severe impairment)	Patient in critical condition, and/or obtunded and cannot perform assessment of tasks
Raised intracranial pressure	NA	NA	Stage 1–2 papilledema ^b , or CSF opening pressure < 20 mmHg	Stage 3–5 papilledema ^b , or CSF opening pressure ≥ 20 mmHg, or cerebral edema
Seizures or motor weakness	NA	NA	Partial seizure, or non-convulsive seizures on EEG with response to benzodiazepine	Generalized seizures, or convulsive or non-convulsive status epilepticus, or new motor weakness

^aReprinted with permission from Neelapu et al., Nat Rev Clin Oncol; 2017 [12]

^bPapilledema grading is performed according to the modified Frisén scale

CAR-T chimeric antigen receptor T cell, CARTOX-10 CAR-T therapy-associated toxicity 10-point neurological assessment, CSF cerebrospinal fluid, EEG electroencephalogram, NA not applicable

3.3.2 Anti-Epileptics and Supportive Care

Seizure prophylaxis with levetiracetam 750 mg orally or intravenously every 12 h for the first 30 days is commonly used in patients receiving CAR-T therapy with known risk of neurotoxicity/CRS, especially with CD19 CAR-T with CD28 co-stimulatory construct and BCMA CAR-T therapy [12, 58]. CD19 CAR-T with a 4-1-BB construct may have a lower rate of neurotoxicity in pediatric patients and routine seizure prophylaxis may not be required [38]. Ideally, neurotoxicity grade ≥ 3 should be monitored in an intensive care unit (ICU) as many of these patients require mechanical ventilation for airway protection and permissive hypercapnia for cerebral edema. Non-convulsive and convulsive status epilepticus should be managed with benzodiazepines and additional antiepileptics as needed. Levetiracetam and phenobarbital are commonly used antiepileptics, in refractory cases endotracheal intubation and anesthesia is required.

4 Prediction and Prevention of CAR-T Therapy-Associated Toxicities

Based on the risk factors defined above, different groups have started studying various preventative strategies to reduce the incidence and severity of CRS and neurotoxicity from CAR-T therapy. For lymphoma, the amount of disease burden corresponds to severe neurotoxicity rates: more disease leads to higher risk [59]. Debulking chemotherapy before CAR-T infusion may reduce antigenic exposure and could be pursued. We have used bridging chemotherapy or pulse steroids between lymphocyte apheresis and CAR-T infusion with the goal to reduce disease burden and improve patient's functional status, and it appears safe in our experience. Bridging chemotherapy was allowed and was used in almost all patients treated on several of the pivotal trials for lymphoma. However, the effect of bridging therapy on CAR-T therapy efficacy is unknown at this point. Other approaches under investigation are early biomarker-driven treatment with anti-IL-6 and/or steroids and self-inactivating CAR-T, which can be turned off in the patients with life-threatening CRS or neurotoxicity.

4.1 Predictive Biomarkers of CRS and Neurotoxicity

Given the significant morbidity and mortality attributed to CRS and neurotoxicity, the development of predictive biomarkers in the serum and CSF is an active area of investigation. Elevated cytokines in serum before and during CRS and neurotoxicity provides the logical basis for the approach, as they likely play a central role in pathogenesis to CAR-T therapy-related toxicities [6, 39, 48, 51, 60]. The goal is to develop a validated cytokine profile to predict the

occurrence and severity of these complications with a goal of prevention and reduction in the severity of these complete dictations by early cytokine-directed therapy. Gardner et al. have developed an early intervention protocol with tocilizumab $\pm$ dexamethasone, which appear to reduce the severity of CRS without affecting the efficacy of CD19 CAR-T therapy [61]. Locke et al. presented their interim analysis findings of a ZUMA-1 safety expansion cohort of patients who received a prophylactic dose of tocilizumab on day 2 post axi-cel infusion. Only one patient (3%) experienced ≥ 3 (grade 4) CRS; however, incidence of ≥ 3 neurotoxicity was 41% compared with 28% in the original ZUMA-1 cohort [6, 33]. In one reported death from cerebral edema, the patient had very high levels of serum IL-15, IL-8, TNF- β , CVAM 1, TNF- γ , IL-1RA, CCL17, and IP-10, consistent with the presence of activated myeloid and lymphoid cells on the day of axi-cel infusion [33]. Other studies are ongoing to further explore the role of prophylactic anti-IL-6 therapy to prevent or reduce the severity of CAR-T therapy-associated toxicities (ClinicalTrials.com identifiers NCT02906371, NCT02926833). Preclinical models showed a central role of host monocyte-derived IL-1, IL-6, and NO in CRS and neurotoxicity. Treatment with anakinra (anti-IL-1 mAb) successfully prevented both CRS and neurotoxicity in human leukemia xenograft models treated with CD19 CAR-T [16]. A simple correlation between elevated cytokine and CRS/neurotoxicity will not be adequate if it fails to predict the occurrence ahead of severe symptoms. CRP and ferritin are associated with CRS, but both biomarkers fail to predict development of severe CRS [34]. A group from the University of Pennsylvania has developed a cytokine profile by measuring IFN- γ , IL-13, and MIP1 α concentrations within 72 h after CD19 CAR-T infusion in pediatric patients with B-ALL, which has a sensitivity of 100% and a specificity of 96% in predicting CRS [34]. Hay et al. showed that elevated serum MCP-1 in patients with fever ≥ 38.9 °C within 36 h of CAR-T infusion has better sensitivity and specificity to predict grade ≥ 4 CRS than CRP and ferritin (sensitivity 100%; specificity 95%). Adding IL-6 levels ≥ 16 pg/mL in the first 36 h after CAR-T infusion in this algorithm can predict grade ≥ 4 neurotoxicity with sensitivity of 100% and specificity of 94% [39, 48]. Gust et al. showed an earlier peak IL-6 serum concentration was associated with a higher risk of grade ≥ 4 neurotoxicity [39]. It is important to note here that tocilizumab and siltuximab can interfere with the measurement of serum IL-6 and soluble IL-6R [62]. In ZUMA-1, patients developing grade ≥ 3 neurotoxicity had elevated IL-15 ($p=0.0006$) and decreased perforin ($p=0.001$) on day 0 before CAR-T infusion and increased IL-15, MCP-1, and IL-6 on day 1 after CAR-T infusion [40]. Severe neurotoxicity correlated with higher peak concentration of C-reactive protein, ferritin, and multiple cytokines including IL-6, IFN- γ , and TNF- α [39]. In the ZUMA-1

trial, elevated IL-2, GM-CSF, and ferritin were associated with neurological events without CRS [6]. In a Memorial Sloan Kettering Cancer Center (MSKCC) cohort of B-ALL patients treated with CD19 CAR, elevated IL-2 and IL-5 at day 3 were unique to neurotoxicity [63]. Measurement of endothelial activation biomarkers such as von Willebrand factor (VWF) and Ang-2 before and after CAR-T infusion can potentially be explored to predict permeability of the BBB and neurotoxicity [16]. The MSKCC group has shown baseline clinical characteristics associated with severe neurotoxicity. Baseline platelet count < 60 or mean corpuscular hemoglobin concentration > 33.2% and morphologic disease (> 5% blasts) predicted severe neurotoxicity with 95% sensitivity and 70% specificity [63]. Consumptive coagulopathy with prolonged prothrombin time (PT), activated partial thromboplastin time (aPTT), elevated D-dimer, and hyperfibrinogenemia was associated with grade ≥ 4 CRS [48].

4.2 Designing Safer CAR-T Cells

The balance between anti-tumor effect and toxicity reduction may be hard to achieve as most approaches to prevent CRS/neurotoxicity can potentially reduce CAR-T proliferation and persistence, both of which are key to successful tumor elimination [64]. Pre-clinical studies have shown that optimizing the binding affinity of scFv in the CAR can improve anti-tumor activity and reduce associated toxicities. Excessive target affinity can lead to early exhaustion, poor persistence of CAR-T, and increased toxicity [65–68]. Park et al. showed that ICAM-1 avid CAR-T with micromolar affinity have improved efficacy and safety compared with ones with nanomolar affinity [66]. Rapid advances in gene editing technology such as Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) have made it possible to generate new CAR constructs with the goal to increase the efficacy and safety of the CAR-T product. The MSKCC group has developed a CD19-specific CAR inserted next to the T-cell receptor α constant (TRAC) locus using the CRISPR platform, which resulted in uniform CAR expression and enhanced potency of CAR-T [69]. This may allow a reduced dose of CAR-T with a goal to reduce the toxicity. Although these findings are intriguing, gene editing is associated with a number of safety concerns, mainly insertional mutagenesis and acquired defects in DNA repair function [70].

Other modalities currently under investigation include the insertion of genetic constructs within CAR to selectively turn off CAR-T when serious toxicity develops [71–75]. Tasian et al. from University of Pennsylvania developed anti-CD123-4-1BB-CD3 ζ T cells with co-expression of CD20. In pre-clinical models, it allowed successful eradication of CD123 acute myeloid leukemia cells with subsequent elimination of CAR-T with rituximab [74]. Similarly,

Minagawa et al. showed that insertion of caspase9 allowed selective apoptosis of CAR-T with the administration of a non-therapeutic dimerizer, which activated the suicide gene [73]. Eliminating CAR-T to control short-term toxicities will invariably compromise tumor control when long-term persistence of CAR-T is important. This issue might be circumvented by insertion of an inducible gene regulatory system that enables controlled expression of CARs upon drug administration. Sakemura et al. developed CD19 CAR with a tetracycline regulation system (Tet-on), which allowed controlled activation of CAR-T only in the presence of doxycycline. This approach provided the ability to turn the CAR-T ‘on’ and ‘off’ by doxycycline administration [76]. Ma et al. showed that CAR-T activity can be controlled by soluble intermediary ‘switch’ molecules. Modified CAR-T are dependent on these switch molecules to form a ternary complex between the CAR-T, switch, and tumor-associated antigen. The theoretical advantage of this approach is that the activity of CAR-T could be titrated by adjusting the concentration of the switch molecule instead of completely turning it ‘on’ or ‘off’. These CAR-T could also be re-directed towards different tumor-associated antigens based on specificity of the switch molecules, which can ultimately help to treat the disease relapse due to ‘antigen-escape’ [77]. At the same time, immunogenicity of these switch molecules and slower CAR-T inactivation compared with CAR-T with a suicide gene limits the usefulness of this approach in the setting of acute toxicity. Self-activating CAR-T have been generated by fusing the oxygen sensing domain of hypoxia-inducing factor-1 α (HIF-1 α), which is only active in a hypoxic environment, commonly found in neoplastic tissue. These CAR-T became inactive in normal tissue in the presence of normoxia and avoided off-target toxicities [78]. Roybal et al. constructed ‘AND-gate’ CAR-T using a synNotch receptor that required two antigen engagements to get activated. In vivo, T cells engineered with dual-receptor circuits recognizing combinations of antigens can efficiently kill target tumor cells, while sparing bystander cells [79]. Giavridis et al. recently showed that the CAR-T engineered to produce endogenous IL-1 receptor antagonist successfully prevented CRS without anti-leukemia efficacy in a human leukemia mouse model [15]. All these approaches are still in their infancy and will require robust clinical validation.

5 Conclusions

CAR-T therapy is the latest advance in cancer immunotherapy with promising results in various hematological malignancies. CAR-T therapy-related toxicities, mainly CRS and neurotoxicity, remain major hurdles that will need to be overcome before its widespread use. Efforts are underway to

understand host-tumor-CAR-T interactions, which will lead to a better understanding of the pathophysiologies behind CRS and neurotoxicity. Timely diagnosis and multi-disciplinary management are the cornerstones for optimal outcomes. Development of toxicity grading scales and protocol-based management are important advances; however, newer strategies are urgently needed to predict CRS/neurotoxicity and salvage those patients who are refractory to anti-IL-6 therapy and steroids. There is significant heterogeneity between the different CAR-T therapies (lymphodepleting chemotherapy, dose of CAR-T, co-stimulatory molecules, tumor types etc.), grading scales, and attributions of toxicities, which prevents generalization of CRS and neurotoxicity management protocols across disease types. Multi-institutional collaborations and standardized diagnosis and grading criteria are needed. Predictive biomarkers and next-generation CAR-T will guide the future treatment strategies of CAR-T-associated toxicities. In the future, we may have the luxury to select a particular CAR-T construct based on the patient and disease characteristics to maximize efficacy and safety.

Compliance with Ethical Standards

Funding This work was supported by an NCI K23 award (PI- Locke, F; CA201594, NIH/NCI) and an NCI Cancer Center support grant P30 (CA076292-18S4, NIH/NCI). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest Frederick Locke declares being a scientific advisor to Kite (a Gilead subsidiary) and Novartis; and a consultant for Cellular BioMedicine Group. Bhagirathbhai R. Dholaria and Christina Bachmeier declare that they have no conflicts of interest that might be relevant to the contents of this manuscript.

References

- Lamers CH, Klaver Y, Gratama JW, Sleijfer S, Debets R. Treatment of metastatic renal cell carcinoma (mRCC) with CAIX CAR-engineered T-cells—a completed study overview. *Biochem Soc Trans*. 2016;44(3):951–9.
- Brocker T, Karjalainen K. Signals through T cell receptor-zeta chain alone are insufficient to prime resting T lymphocytes. *J Exp Med*. 1995;181(5):1653–9.
- Savoldo B, Ramos CA, Liu E, Mims MP, Keating MJ, Carrum G, et al. CD28 costimulation improves expansion and persistence of chimeric antigen receptor-modified T cells in lymphoma patients. *J Clin Invest*. 2011;121(5):1822–6.
- Wang X, Riviere I. Clinical manufacturing of CAR T cells: foundation of a promising therapy. *Mol Ther Oncol*. 2016;3:16015.
- June CH, Riddell SR, Schumacher TN. Adoptive cellular therapy: a race to the finish line. *Sci Transl Med*. 2015;7(280):280ps7.
- Neelapu SS, Locke FL, Bartlett NL, Lekakis LJ, Miklos DB, Jacobson CA, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med*. 2017;377(26):2531–44.
- Schuster SJ, Svoboda J, Chong EA, Nasta SD, Mato AR, Anak O, et al. Chimeric antigen receptor T cells in refractory B-cell lymphomas. *N Engl J Med*. 2017;377(26):2545–54.
- Borchmann P, Tam C, Jäger U, McGuirk J, Holte H, Waller E, et al. An updated analysis of JULIET, a global pivotal phase 2 trial of tisagenlecleucel in adult patients with relapsed or refractory diffuse large B-cell lymphoma. *European Hematology Association*, abstract S799; June 2018.
- FDA approves second CAR T-cell therapy. *Cancer Discov*. 2018;8(1):5–6. <http://cancerdiscovery.aacrjournals.org/content/8/1/5.full>
- Heymach J, Krilov L, Alberg A, Baxter N, Chang SM, Corcoran R, et al. Clinical Cancer advances 2018: annual report on progress against cancer from the american society of clinical oncology. *J Clin Oncol*. 2018;36(10):1020–44.
- First two CAR-T cell medicines recommended for approval in the European Union 2018. <https://www.ema.europa.eu/en/news/first-two-car-t-cell-medicines-recommended-approval-european-union>. Accessed 19 Oct 2018.
- Neelapu SS, Tummala S, Kebriaei P, Wierda W, Gutierrez C, Locke FL, et al. Chimeric antigen receptor T-cell therapy—assessment and management of toxicities. *Nat Rev Clin Oncol*. 2017;Sep:19.
- Brudno JN, Kochenderfer JN. Toxicities of chimeric antigen receptor T cells: recognition and management. *Blood*. 2016;127(26):3321–30.
- Lee DW, Gardner R, Porter DL, Louis CU, Ahmed N, Jensen M, et al. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood*. 2014;124(2):188–95.
- Giavridis T, van der Stegen SJC, Eyquem J, Hamieh M, Piersigilli A, Sadelain M. CAR T cell-induced cytokine release syndrome is mediated by macrophages and abated by IL-1 blockade. *Nat Med*. 2018;24(6):731–8.
- Norelli M, Camisa B, Barbiera G, Falcone L, Purevdorj A, Genua M, et al. Monocyte-derived IL-1 and IL-6 are differentially required for cytokine-release syndrome and neurotoxicity due to CAR T cells. *Nat Med*. 2018;24(6):739–48.
- Hay KA, Hanafi LA, Li D, Gust J, Liles WC, Wurfel MM, et al. Kinetics and biomarkers of severe cytokine release syndrome after CD19 chimeric antigen receptor-modified T cell therapy. *Blood*. 2017;Sep:18.
- van der Stegen SJ, Hamieh M, Sadelain M. The pharmacology of second-generation chimeric antigen receptors. *Nat Rev Drug Discov*. 2015;14(7):499–509.
- Maude SL, Laetsch TW, Buechner J, Rives S, Boyer M, Bittencourt H, et al. Tisagenlecleucel in children and young adults with B-cell lymphoblastic leukemia. *N Engl J Med*. 2018;378(5):439–48.
- Maude SL, Frey N, Shaw PA, Aplenc R, Barrett DM, Bunin NJ, et al. Chimeric antigen receptor T cells for sustained remissions in leukemia. *N Engl J Med*. 2014;371(16):1507–17.
- Turtle CJ, Hanafi LA, Berger C, Gooley TA, Cherian S, Hudecek M, et al. CD19 CAR-T cells of defined CD4+ :CD8+ composition in adult B cell ALL patients. *J Clin Invest*. 2016;126(6):2123–38.
- Davila ML, Riviere I, Wang X, Bartido S, Park J, Curran K, et al. Efficacy and toxicity management of 19-28z CAR T cell therapy in B cell acute lymphoblastic leukemia. *Sci Transl Med*. 2014;19;6(224):224ra25.
- NCI Common Terminology Criteria for Adverse Events (CTCAE) v.4 data file. 2009; 4.0:[NCI]. http://evs.nci.nih.gov/ftp1/CTCAE/CTCAE_4.03_2010-06-14_QuickReference_5x7.pdf. Accessed 13 Sept 2014.
- Common Terminology Criteria for Adverse Events (CTCAE) Version 5.0. 2017; <https://ctep.cancer.gov/protocoldevelopment/elect>

- [ronic_applications/docs/CTCAE_v5_Quick_Reference_5x7.pdf](#). Accessed 22 Apr 2018.
25. Neelapu SS, Tummala S, Kebriaei P, Wierda W, Locke FL, Lin Y, et al. Toxicity management after chimeric antigen receptor T cell therapy: one size does not fit 'ALL'. *Nat Rev Clin Oncol*. 2018;15(4):218.
 26. Porter DL, Hwang W-T, Frey NV, Lacey SF, Shaw PA, Loren AW, et al. Chimeric antigen receptor T cells persist and induce sustained remissions in relapsed refractory chronic lymphocytic leukemia. *Sci Transl Med*. 2015;7(303):303ra139–303ra139.
 27. Porter D, Frey N, Wood PA, Weng Y, Grupp SA. Grading of cytokine release syndrome associated with the CAR T cell therapy tisagenlecleucel. *J Hematol Oncol*. 2018;11(1):35.
 28. CAR-T toxicity rating system to be finalized by ASBMT by year-end—Drug Development Technology. <https://www.drugdevelopment-technology.com/comment/car-t-toxicity-rating-system-finalized-asbmt-year-end/>. Accessed 11 Sept 2018.
 29. Schuster SJB, Tam CS. Primary analysis of JULIET: a global, pivotal, phase 2 trial of CTL019 in adult patients with relapsed or refractory diffuse large B-cell lymphoma. *Blood*. 2017;130(suppl 1).
 30. Davila ML, Sadelain M. Biology and clinical application of CAR T cells for B cell malignancies. *Int J Hematol*. 2016;104(1):6–17.
 31. Brudno JN, Kochenderfer JN. Chimeric antigen receptor T-cell therapies for lymphoma. *Nat Rev Clin Oncol*. 2017;Aug:31.
 32. Center for Drug Evaluation and Research. Approved Drugs—FDA approves tisagenlecleucel for B-cell ALL and tocilizumab for cytokine release syndrome. <https://www.fda.gov/drugs/informationondrugs/approveddrugs/ucm574154.htm>. Accessed 21 Nov 2018.
 33. Locke FL, Neelapu SS, Bartlett NL, Lekakis LJ, Jacobson CA, Braunschweig I, et al. Preliminary results of prophylactic tocilizumab after axicabtagenechicleucel (axi-cel; KTE-C19) treatment for patients with refractory, aggressive non-hodgkin lymphoma (NHL). *Blood*. 2017;130(Suppl 1):1547.
 34. Teachey DT, Lacey SF, Shaw PA, Melenhorst JJ, Maude SL, Frey N, et al. Identification of predictive biomarkers for cytokine release syndrome after chimeric antigen receptor T-cell therapy for acute lymphoblastic leukemia. *Cancer Discov*. 2016;6(6):664–79.
 35. van Rhee F, Wong RS, Munshi N, Rossi JF, Ke XY, Fossa A, et al. Siltuximab for multicentric Castleman's disease: a randomised, double-blind, placebo-controlled trial. *Lancet Oncol*. 2014;15(9):966–74.
 36. Nishimoto N, Terao K, Mima T, Nakahara H, Takagi N, Kakehi T. Mechanisms and pathologic significances in increase in serum interleukin-6 (IL-6) and soluble IL-6 receptor after administration of an anti-IL-6 receptor antibody, tocilizumab, in patients with rheumatoid arthritis and Castleman disease. *Blood*. 2008;112(10):3959–64.
 37. Brentjens RJ, Davila ML, Riviere I, Park J, Wang X, Cowell LG, et al. CD19-targeted T cells rapidly induce molecular remissions in adults with chemotherapy-refractory acute lymphoblastic leukemia. *Sci Transl Med*. 2013;20(177):177ra38.
 38. Teachey DT, Bishop MR, Maloney DG, Grupp SA. Toxicity management after chimeric antigen receptor T cell therapy: one size does not fit 'ALL'. *Nat Rev Clin Oncol*. 2018;15(4):218.
 39. Gust J, Hay KA, Hanafi LA, Li D, Myerson D, Gonzalez-Cuyar LF, et al. Endothelial activation and blood-brain barrier disruption in neurotoxicity after adoptive immunotherapy with CD19 CAR-T cells. *Cancer Discov*. 2017;7(12):1404–19.
 40. DeAngelo D, Ghobadi A, Park J, Dinner S, Mannis G, Lunning M, et al. Clinical outcomes for the phase 2, single-arm, multicenter trial of JCAR015 in adult B-ALL (ROCKET Study). 32nd annual meeting and pre-conference programs of the Society for Immunotherapy of Cancer (SITC 2017): part one. *J Immunother Cancer*. 2017;5(2):86.
 41. Santomaso B, Park JH, Riviere I, Mead E, Halton E, Diamonte C, et al. Biomarkers associated with neurotoxicity in adult patients with relapsed or refractory B-ALL (R/R B-ALL) treated with CD19 CAR T cells. *J Clin Oncol*. 2017;35(15_suppl):3019.
 42. Neelapu SS, Tummala S, Kebriaei P, Wierda W, Gutierrez C, Locke FL, et al. Chimeric antigen receptor T-cell therapy—assessment and management of toxicities. *Nat Rev Clin Oncol*. 2018;15(1):47–62.
 43. Park JH, Riviere I, Gonen M, Wang X, Senchal B, Curran KJ, et al. Long-term follow-up of CD19 CAR therapy in acute lymphoblastic leukemia. *N Engl J Med*. 2018;378(5):449–59.
 44. Package insert—KYMRIAHA. 2018. <https://www.fda.gov/downloads/BiologicsBloodVaccines/CellularGeneTherapyProducts/ApprovedProducts/UCM573941.pdf>. Accessed 1 May 2018.
 45. Abramson JS, Gordon LI, Palomba ML, Lunning MA, Arnason JE, Forero-Torres A, et al. Updated safety and long term clinical outcomes in TRANSCEND NHL 001, pivotal trial of lisocabtagene maraleucel (JCAR017) in R/R aggressive NHL. *J Clin Oncol*. 2018;36(15_suppl):7505.
 46. Santomaso BD, Park JH, Salloum D, Riviere I, Flynn J, Mead E, et al. Clinical and biologic correlates of neurotoxicity associated with CAR T cell therapy in patients with B-cell Acute lymphoblastic leukemia (B-ALL). *Cancer Discov*. 2018;8(8):958–71.
 47. Taraseviciute A, Tkachev V, Ponce R, Turtle CJ, Snyder JM, Liggett HD, et al. Chimeric antigen receptor T cell-mediated neurotoxicity in nonhuman primates. *Cancer Discov*. 2018;8(6):750–63.
 48. Hay KA, Hanafi LA, Li D, Gust J, Liles WC, Wurfel MM, et al. Kinetics and biomarkers of severe cytokine release syndrome after CD19 chimeric antigen receptor-modified T-cell therapy. *Blood*. 2017;130(21):2295–306.
 49. Higgins SJ, Purcell LA, Silver KL, Tran V, Crowley V, Hawkes M, et al. Dysregulation of angiopoietin-1 plays a mechanistic role in the pathogenesis of cerebral malaria. *Sci Transl Med*. 2016;28(358):358ra128.
 50. Kochenderfer JN, Somerville RPT, Lu T, Shi V, Bot A, Rossi J, et al. Lymphoma remissions caused by anti-CD19 chimeric antigen receptor T cells are associated with high serum interleukin-15 levels. *J Clin Oncol*. 2017;35(16):1803–13.
 51. Turtle CJ, Hay KA, Juliane G, Hanafi L-A, Li D, Chaney C, et al. Biomarkers of Cytokine Release Syndrome and Neurotoxicity after CD19 CAR-T cells and mitigation of toxicity by cell dose. *Blood*. 2016;128(22):1852.
 52. Lee DW, Kochenderfer JN, Stetler-Stevenson M, Cui YK, Delbrook C, Feldman SA, et al. T cells expressing CD19 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young adults: a phase 1 dose-escalation trial. *Lancet (London, England)*. 2015;385(9967):517–28.
 53. Hu Y, Sun J, Wu Z, Yu J, Cui Q, Pu C, et al. Predominant cerebral cytokine release syndrome in CD19-directed chimeric antigen receptor-modified T cell therapy. *J Hematol Oncol Engl*. 2016;9(1):70.
 54. Prudent V, Breitbart WS. Chimeric antigen receptor T-cell neuropsychiatric toxicity in acute lymphoblastic leukemia. *Palliat Support Care*. 2017;15(4):499–503.
 55. Raje NS, Berdeja JG, Lin Y, Munshi N, Siegel BA, Liedtke M, et al. bb2121 anti-BCMA CAR T-cell therapy in patients with relapsed/refractory multiple myeloma: Updated results from a multicenter phase I study. *J Clin Oncol*. 2018;36(suppl; abstr 8007).
 56. Pangman VC, Sloan J, Guse L. An examination of psychometric properties of the mini-mental state examination and the standardized mini-mental state examination: implications for clinical practice. *ANR*. 2000;13(4):209–13.
 57. Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet (London, England)*. 1974;2(7872):81–4.

58. Cohen AD, Garfall AL, Stadtmauer EA, Lacey SF, Lancaster E, Vogl DT, et al. Safety and efficacy of B-cell maturation antigen (BCMA)-specific chimeric antigen receptor T cells (CART-BCMA) with cyclophosphamide conditioning for refractory multiple myeloma (MM). *Blood*. 2017;130(Suppl 1):505.
59. Locke F, Ghobadi A, Lekakis LJ, Miklos DB, Jacobson CA, Jacobsen ED, et al. Outcomes by prior lines of therapy (LoT) in ZUMA-1, the pivotal phase 2 study of axicabtagene ciloleucel (Axi-Cel) in patients (Pts) with refractory large B cell lymphoma. *J Clin Oncol*. 2018;36(no. 15_suppl):3039.
60. Locke FL, Neelapu SS, Bartlett NL, Siddiqi T, Chavez JC, Hosing CM, et al. Phase 1 results of ZUMA-1: a multicenter study of KTE-C19 anti-CD19 CAR T cell therapy in refractory aggressive lymphoma. *Mol Ther*. 2017;25(1):285–95.
61. Gardner R, Leger KJ, Annesley CE, Summers C, Rivers J, Gust J, et al. Decreased rates of severe CRS seen with early intervention strategies for CD19 CAR-T cell toxicity management. *Blood*. 2016;128(22):586.
62. Chen F, Teachey DT, Pequignot E, Frey N, Porter D, Maude SL, et al. Measuring IL-6 and sIL-6R in serum from patients treated with tocilizumab and/or siltuximab following CAR T cell therapy. *J Immunol Methods*. 2016;434:1–8.
63. Park JH, Santomasso B, Riviere I, Senecal B, Wang X, Purdon T, et al. Baseline and early post-treatment clinical and laboratory factors associated with severe neurotoxicity following 19-28z CAR T cells in adult patients with relapsed B-ALL. *J Clin Oncol*. 2017;35(15_suppl):7024.
64. Ghorashian S, Kramer AM, Albon SJ, Wright G, Castro F, Popova B, et al. A Novel Low affinity CD19CAR results in durable disease remissions and prolonged CAR T cell persistence without severe CRS or neurotoxicity in patients with paediatric ALL. *Blood*. 2017;130(Suppl 1):806.
65. Watanabe K, Terakura S, Uchiyama S, Martens AC, Meerten TV, Kiyoi H, et al. Excessively high-affinity single-chain fragment variable region in a chimeric antigen receptor can counteract T-cell proliferation. *Blood*. 2014;124(21):4799.
66. Park S, Shevlin E, Vedvyas Y, Zaman M, Hsu YS, Min IM, et al. Micromolar affinity CAR T cells to ICAM-1 achieves rapid tumor elimination while avoiding systemic toxicity. *Sci Rep*. 2017;7(1):14366.
67. Schmid DA, Irving MB, Posevitz V, Hebeisen M, Posevitz-Fejfar A, Sarria JC, et al. Evidence for a TCR affinity threshold delimiting maximal CD8 T cell function. *J Immunol*. 2010;184(9):4936–46.
68. Caruso HG, Hurton LV, Najjar A, Rushworth D, Ang S, Olivares S, et al. Tuning sensitivity of CAR to EGFR density limits recognition of normal tissue while maintaining potent anti-tumor activity. *Cancer Res*. 2015;75(17):3505–18.
69. Eyquem J, Mansilla-Soto J, Giavridis T, van der Stegen SJ, Hamieh M, Cunanan KM, et al. Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances tumour rejection. *Nature*. 2017;543(7643):113–7.
70. Ihry RJ, Worringer KA, Salick MR, Frias E, Ho D, Theriault K, et al. p53 inhibits CRISPR-Cas9 engineering in human pluripotent stem cells. *Nat Med*. 2018;24(7):939–46.
71. Zhou X, Brenner MK. Improving the safety of T-Cell therapies using an inducible caspase-9 gene. *Exp Hematol*. 2016;44(11):1013–9.
72. Casucci M, Falcone L, Camisa B, Norelli M, Porcellini S, Stornaiuolo A, et al. Extracellular NGFR spacers allow efficient tracking and enrichment of fully functional CAR-T cells co-expressing a suicide gene. *Front Immunol*. 2018;9:507.
73. Minagawa K, Jamil MO, Al-Obaidi M, Pereboeva L, Salzman D, Erba HP, et al. In vitro pre-clinical validation of suicide gene modified anti-CD33 redirected chimeric antigen receptor T-cells for acute myeloid leukemia. *PLoS One*. 2016;11(12):e0166891.
74. Tasian SK, Kenderian SS, Shen F, Ruella M, Shestova O, Kozlowski M, et al. Optimized depletion of chimeric antigen receptor T cells in murine xenograft models of human acute myeloid leukemia. *Blood*. 2017;129(17):2395–407.
75. Qasim W, Ciocarlie O, Adams S, Inglott S, Murphy C, Rivat C, et al. Preliminary results of UCART19, an allogeneic anti-CD19 CAR T-cell product in a first-in-human trial (PALL) in pediatric patients with CD19+relapsed/refractory B-cell acute lymphoblastic leukemia. *Blood*. 2017;130(Suppl 1):1271.
76. Sakemura R, Terakura S, Watanabe K, Julamanee J, Takagi E, Miyao K, et al. A tet-on inducible system for controlling CD19-chimeric antigen receptor expression upon drug administration. *Cancer Immunol Res*. 2016;4(8):658–68.
77. Ma JS, Kim JY, Kazane SA, Choi SH, Yun HY, Kim MS, et al. Versatile strategy for controlling the specificity and activity of engineered T cells. *Proc Natl Acad Sci USA*. 2016;113(4):E450–8.
78. Juillerat A, Marechal A, Filhol JM, Valogne Y, Valton J, Duclert A, et al. An oxygen sensitive self-decision making engineered CAR T-cell. *Sci Rep*. 2017;20(7):39833.
79. Roybal KT, Rupp LJ, Morsut L, Walker WJ, McNally KA, Park JS, et al. Precision tumor recognition by T cells with combinatorial antigen-sensing circuits. *Cell*. 2016;164(4):770–9.