



Correction to: Meta-analysis of Placebo Response in Randomized Clinical Trials of Antipsychotic Drugs Using PANSS Focusing on Different Approaches to the Handling of Missing Data

Akiko Matsusaki¹ · Masayuki Kaneko¹ · Mamoru Narukawa¹

Published online: 19 June 2018
© Springer International Publishing AG, part of Springer Nature 2018

Correction to: Clin Drug Investig
<https://doi.org/10.1007/s40261-018-0661-1>

In the original publication, the Estimate column with values under Multivariate meta-regression (LOCF) in Table 2 was missed. The corrected Table 2 is shown below:

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s40261-018-0661-1>.

✉ Akiko Matsusaki
matsusakiakiko_kitasato@yahoo.co.jp

¹ Department of Clinical Medicine (Pharmaceutical Medicine), Graduate School of Pharmaceutical Sciences, Kitasato University, 5-9-1 Shirokane, Minato-ku, Tokyo 108-8641, Japan

Table 2 Findings by meta-regression analyses in placebo-controlled randomized clinical trials by MMRM and LOCF

Variable	Univariate meta-regression (MMRM)			Univariate meta-regression (LOCF)		
	Estimate	SE	p value	Estimate	SE	p value
Patient factors						
Diagnosis				-9.304	3.107	0.003
Severity	-0.127	0.104	0.225	-0.101	0.164	0.539
Age	0.747	0.480	0.120	0.707	0.483	0.144
% Male	0.149	0.094	0.111	0.104	0.112	0.352
Investigator factor						
Rater training	-4.321	2.032	0.033	-2.384	2.279	0.295
Study design factors						
Study duration	-2.130	0.740	0.004	2.267	0.757	0.003
Placebo lead-in	-1.039	1.878	0.580	5.618	2.057	0.006
Number of treatment arms	-1.241	1.205	0.303	0.774	1.123	0.491
Placebo randomization rate	-0.014	0.236	0.953	0.160	0.162	0.324
Regimen	2.466	1.658	0.137	-1.191	2.309	0.606
Active comparator	-2.324	1.818	0.201	4.407	2.573	0.087
Treatment setting	-3.463	1.425	0.015	-1.329	2.737	0.627
Operational factors						
Number of study sites	-0.096	0.048	0.047	-0.023	0.049	0.640
Number of countries	-0.653	0.297	0.028	-0.328	0.339	0.333
Study period	-0.042	0.152	0.781	0.126	0.144	0.381
Enrollment Speed	5.106	1.674	0.002	-0.782	3.293	0.812
Variable	Multivariate meta-regression (MMRM)			Multivariate meta-regression (LOCF)		
	Estimate	SE	p value	Estimate	SE	p value
Patient factors						
Diagnosis				-4.490	3.297	0.173
Investigator factor						
Rater training	-3.607	2.990	0.228			
Study design factors						
Study duration	0.866	1.231	0.482	1.567	0.747	0.036
Placebo lead-in				2.994	2.119	0.158
Active comparator				1.914	2.384	0.422
Treatment setting	-4.217	1.971	0.032			
Operational factors						
Number of study sites	0.149	0.092	0.106			
Number of countries	-0.864	0.399	0.030			
Enrollment speed	3.876	2.943	0.188			

PANSS positive and negative syndrome scale, LOCF last observation carried forward, MMRM mixed-effect models for repeated measures