



Targeting Sigma Receptors for the Treatment of Neurodegenerative and Neurodevelopmental Disorders

Dicson S. Malar¹ · Premrutai Thitilertdecha² · Kanokphorn S. Ruckvongacheep³ · Sirikalaya Brimson³ ·
Tewin Tencomnao¹ · James M. Brimson^{1,4}

Accepted: 18 April 2023 / Published online: 11 May 2023
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2023

Abstract

The sigma-1 receptor is a 223 amino acid-long protein with a recently identified structure. The sigma-2 receptor is a genetically unrelated protein with a similarly shaped binding pocket and acts to influence cellular activities similar to the sigma-1 receptor. Both proteins are highly expressed in neuronal tissues. As such, they have become targets for treating neurological diseases, including Alzheimer's disease (AD), Huntington's disease (HD), Parkinson's disease (PD), multiple sclerosis (MS), Rett syndrome (RS), developmental and epileptic encephalopathies (DEE), and motor neuron disease/amyotrophic lateral sclerosis (MND/ALS). In recent years, there have been many pre-clinical and clinical studies of sigma receptor (1 and 2) ligands for treating neurological disease. Drugs such as blarcamesine, dextromethorphan and pridopidine, which have sigma-1 receptor activity as part of their pharmacological profile, are effective in treating multiple aspects of several neurological diseases. Furthermore, several sigma-2 receptor ligands are under investigation, including CT1812, rivastigmine and SAS0132. This review aims to provide a current and up-to-date analysis of the current clinical and pre-clinical data of drugs with sigma receptor activities for treating neurological disease.

Key Points

The sigma receptors represent an interesting target for the treatment of neurodegenerative and neurodevelopmental disorders.

Investigational sigma-1/2 drugs including blarcamesine (ANAVEX2-73), ANAVEX3-71 and CT1812 have shown promising results in clinical trials, and old drugs with sigma-1/2 affinities such as amantadine, dextromethorphan, fenfluramine, naluzotan, phenytoin, pridopidine and rivastigmine are finding new purpose.

Dicson S. Malar and Premrutai Thitilertdecha contributed equally to this manuscript.

✉ James M. Brimson
jamesmichael.b@chula.ac.th

- ¹ Natural Products for Neuroprotection and Anti-ageing Research Unit, Chulalongkorn University, Bangkok, Thailand
- ² Siriraj Research Group in Immunobiology and Therapeutic Sciences, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand
- ³ Department of Clinical Microscopy, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok, Thailand
- ⁴ Research, Innovation and International Affairs, Faculty of Allied Health Sciences, Chulalongkorn University, Room 409, ChulaPat-1 Building, 154 Rama 1 Road, Bangkok 10330, Thailand

1 Introduction

1.1 The Sigma Receptors

The sigma-1 receptor ($\sigma 1R$) is a 223 amino acid chaperone protein that resides at the endoplasmic reticulum (ER); it is also found at the mitochondria-associated ER membrane

(MAM), and it can translocate to the plasma membrane (Fig. 1). The σ 1R is coded by the *SIGMAR1* gene (found on chromosome 9 in humans and chromosome 4 in mice), which comprises four exons that code for the 223 amino acid protein. The σ 1R has no homology with any other mammalian protein; however, it does share some homology with the yeast *ERG2*, a fungal gene encoding a C8–C7 sterol isomerase from *Magnaporthe grisea*, *Saccharomyces cerevisiae* and *Ustilago maydis* [1], although the σ 1R has no isomerase activity.

The σ 1R is concentrated in lipid-enriched microdomains at the ER and mitochondria. In NG108 neuroblastoma cells, the σ 1R was shown to be important in the compartmentalization of ER-synthesized lipids. At the ER, the σ 1R colocalizes with caveolin-2, a cholesterol-binding protein [2]. The σ 1R also interacts with and modulates the actions of various other proteins, including the IP3R [3], ion channels [4–6] and the small molecule G-protein (Rac1) [7].

The σ 1R has no confirmed endogenous ligand, although there are several candidates. The hallucinogen, *N,N*-dimethyltryptamine (DMT) [8]; neuro-active steroids such as progesterone [but also testosterone, pregnenolone sulphate and dehydroepiandrosterone sulphate (DHEA-S) bind to the σ 1R] [9]; and *N*-alkyl amines, such as the endogenous sphingolipids [10], are all candidates as endogenous σ 1R ligands.

The σ 1R ligands can be divided up into several categories: the traditional agonists and antagonists that, when used at physiologically relevant doses, provide a physiological response, and the σ 1R modulatory ligands, which do not provide a physiological response alone, rather they exhibit positive or negative modulatory effects when


certain physiological processes are activated (such as during cellular stress or neuron firing).

The σ 1R (allosteric) modulatory ligands may also act synergistically with subtherapeutic concentrations of σ 1R agonists or antagonists [11, 12]. Defining σ 1R ligands into these groups can be difficult since there was confusion about the identity of the sigma binding site due to a lack of specific ligands [13]. Furthermore, many early studies saw behavioural effects using what are now called σ 1R agonists. In contrast, molecular pharmacological studies have shown that what are now termed σ 1R antagonists produce cellular responses in calcium flux [6, 14, 15]. In general terms, σ 1R agonists are ligands that have neuroprotective, mood modifying, analgesic or cognitive effects, and the antagonists mostly (there are several exceptions) have cytotoxic effects.

More specifically, laboratory tests have been suggested to define σ 1R ligands into agonist or antagonist categories. One proposed, generally accepted test is to check for the dissociation of binding immunoglobulin protein (BiP), with agonists causing the dissociation whereas antagonists do not [16]. The ability to potentiate neurite outgrowth in PC12 cells indicates agonistic activity at the σ 1R. Another proposed way to define σ 1R ligands is the Hill slope of their binding isotherms, with a standard slope factor (slope \approx 1) indicating an agonist and a shallow slope factor (slope \approx 0.5) indicating an antagonist [5, 6].

The σ 1R is found throughout much of the tissues in the body. However, it appears more concentrated in the central nervous system (CNS), heart, liver and eyes [17, 18]. Many cancers also overexpress the σ 1R, suggesting that the σ 1R

Fig. 1 Structure of the *Xenopus* sigma-1 receptor homotrimer (left) bound to (+)-pentazocine (ligand removed from image) and the Bovine sigma-2 receptor homodimer (right) bound to PB28, (ligand removed from image) at the membrane of the endoplasmic reticulum.



plays an important role in cell survival under stressful conditions [19].

Another receptor with a similar ligand-binding profile is the sigma-2 receptor (σ 2R), although it is structurally and genetically unrelated to the σ 1R [20]. The σ 2R has a molecular weight of 18–21.5 kDa, is located on chromosome 17 (location: 17q11.2) and was previously thought to be part of the progesterone receptor membrane component 1 (PGRMC1) [21]. A more recent study has identified the σ 2R to be the transmembrane protein 97 (TMEM97), also known as meningioma-associated protein 30 (MAC30) [22]. The σ 2R is an endoplasmic reticulum (ER) resident membrane protein [23, 24] and is also located in the lipid raft containing cholesterol and sphingolipids [25, 26] of the plasma membrane [27]. The structure of the σ 2R has been identified using x-ray crystallography showing four transmembrane helices (Fig. 1) [28]. Although genetically unrelated to the σ 1R, the binding pockets of the two proteins are remarkably similar. There are functionally comparable amino acids in similar spatial positions, which likely explains the convergent evolution of these two proteins and reveals why these two proteins bind such a similar portfolio of compounds [28]. σ 2Rs are found primarily in the CNS [29], breast, colon, stomach, oesophagus and lung. The σ 2R is highly expressed in a large number of cancers [30], such as colorectal cancer, oral/oropharyngeal squamous cell carcinoma, breast cancer cell lines (MCF7, MDA-MB-231), neuroblastoma SK-N-SH cells [31], ovarian cancer cells [32] and prostate cancer [33].

This review aims to cover the current understanding of the sigma receptors' role in neurodegenerative and neurodevelopmental disorder development, discuss the relevant approved or investigational drugs and critically review the available clinical data.

Literature searches were carried out (with the last search carried out in April 2023) using PubMed, Scopus and Google Scholar using search terms related to Alzheimer's disease, Parkinson's disease, motor neuron disease, multiple sclerosis, Rett syndrome, and developmental and epileptic encephalopathies in combination with the terms related to the σ -1/2 receptors. Clinical trials were identified via www.clinicaltrials.gov and by refining searches on PubMed only to include clinical trials. No specific time period was chosen for article publication, although more emphasis was given to recent and up-to-date articles.

1.2 Overview of Neurological Disease and the Role of the Sigma Receptors

1.2.1 Alzheimer's Disease

Alzheimer's disease (AD) is a neurodegenerative disorder with a progressive loss of cognitive functions, commonly associated with age and neuropathology. The noticeable

symptoms, such as memory loss and language problems, begin when damage occurs to neurons responsible for these cognitive functions. Other symptoms involving basic bodily functions, such as walking and swallowing, subsequently occur along with the disease progression [34]. The key components of the disease have been reported to be the accumulation of amyloid precursor protein (APP) fragment amyloid-beta ($A\beta$) from its overproduction or reduced clearance [35], and the downstream process in hyperphosphorylation of the protein tau, leading to cell death, neurotransmitter deficits [36], reduced synaptic strength and neurodegeneration [37], as well as induction of inflammation related to dysregulated glucose metabolism [38, 39].

The σ 1R is expressed highly throughout many organs in the body; however, it is particularly concentrated in the CNS. Much research into the σ 1R has focused on pain and neurotransmission [40–44]. In studies of normal healthy ageing, in rats or monkeys, the levels of the σ 1R in the brain are maintained and possibly increased [45–47]. Receptor increases with healthy ageing have only been reported for σ 1R and muscarinic M2 receptors. Furthermore, this increase in σ 1R has been seen in human ageing studies.

Using ^{11}C -labelled donepezil, it was shown that cardiac levels of the σ 1R were elevated in aged healthy individuals [48]. This apparent increase of σ 1Rs in the body appears to coincide with the age-related loss of other receptors such as the M_1 and M_4 muscarinic acetylcholine receptors [49], serotonin 5HT_{2A} receptors [50] and D_1/D_2 dopamine (DA) receptors [51]. It has been hypothesized that the increase in σ 1R expression is a compensatory mechanism to offset the loss of the other receptors [52, 53]. In the early stages of AD, there is a significant loss of σ 1R expression in the brain [54, 55].

Genetic studies have identified that gene mutations in apolipoprotein E (*ApoE*) are related to AD risk. The *ApoE2* allele is associated with a reduced risk of AD, and the *ApoE3* allele is the most frequently found and is considered a neutral allele regarding AD risk. Carriers of the *ApoE4* allele have an earlier onset of AD, and the amyloid plaques are more diffuse.

The σ 1R gene is found on chromosome 9p13 and is approximately 7 kbp long, consisting of four exons coding for a 223 bp amino acid. There have been two reported variants of the σ 1R receptor that may affect AD development. The G–241T/C–240T (*rs. 1799729*) in the proximal promoter region and A61C (*rs. 1800866*) resulted in an amino acid substitution Q2P) in the first exon [56]. The haplotype TT241-240P2 (TT-P) has been shown to express σ 1R receptor protein at a level 43% lower than the allele GC-241-240Q2 [56].

Several studies have investigated the link between polymorphisms in the σ 1R and the risk of AD [56–59]. A Japanese study has suggested that the TT-P haplotype is

protective against AD [56]. However, this would appear to go against what is already known about the σ 1R and has been contradicted by several other studies. Studies from Hungarian, Polish, Australian and Chinese populations have shown a link between the σ 1R *TT-P* haplotype and increased susceptibility to AD [57–59].

Furthermore, there is evidence that there may be an interaction between the effects of the *ApoE4* allele and the *σ 1R* allele. In the Hungarian study [59], those who inherited both the *TT-P* haplotype and the *E4* allele had a considerably higher risk for AD than those who carried only one. In the Australian/Chinese cohorts [58], the study found the *Q2P* polymorphism to be associated with a less severe form of AD in non-*E4* allele carriers (i.e. *ApoE2* or *ApoE3*); however, expression of the *Q2P* polymorphism in combination with *ApoE4* had the potential to create a more severe cognitive dysfunction.

Reduction or loss of expression of the σ 1R may make neuronal cells vulnerable to stress due to $A\beta$ -mediated toxicity and damage to intracellular lipid metabolism, cell skeleton network and immune responses. These lead to neuronal cell death and contribute to the pathology of AD [60]. σ 1R knockdown has been shown multiple times in primary hippocampal neurons to reduce dendritic spine formation [61–63].

The σ 1R also plays a role in controlling tau phosphorylation. The σ 1R is essential in proper tau phosphorylation and axon extension by promoting p35 turnover through the interaction of the receptor with myristic acid. Furthermore, *σ 1R* knockout (KO) in neurons results in the accumulation of p35 due to its slower degradation. Conversely, overexpression of the σ 1R results in a decrease of p35. In the σ 1R-KO, neurons show shorter axons with lower densities. Interestingly the σ 1R agonist myristic acid mitigates p35 accumulation, diminishes tau phosphorylation and restores axon elongation [64].

Autophagy is a cellular function that involves destroying and recycling cellular organelles and proteins, either when damaged or when the cell needs building blocks to survive. The σ 1R regulates autophagy, which may be an important target for treating neurodegenerative disease (reviewed in [53]). σ 1R agonists can activate autophagy, reversing some of the effects of APP overexpression [65]. The dysregulation of the autophagic process is thought to be a possible part of AD pathology [66–68]. This, and the fact that the σ 1R is lost in the early stages of AD, could be a potential cause of, at least part of, the pathology of AD.

Calcium homeostasis is significant in AD disease development (reviewed in [69]). Vaccines/antibodies that have been trialled, which remove $A\beta$ plaque build-up, have not stopped dementia progression despite their efficacy at removing $A\beta$ [70, 71], suggesting that non-plaque forming $A\beta$ -oligomers are responsible for neuron cell death [72].

Oligomers of soluble $A\beta$ can form ion channels and dysregulate calcium influx into neurons [73]. Fibrillar $A\beta$ aggregates may cause cell death via the induction of oxidative stress, depolarization of synaptic membranes, excessive calcium influx and mitochondrial stress [74].

The σ 1R appears to play an essential role in regulating calcium homeostasis. Disrupting the σ 1R in cancer cells with an antagonist such as IPAG (1-(4-iodophenyl)-3-(2-adamantyl)guanidine) or rimcazole results in a large influx of $[Ca^{2+}]$ into the cytosol, that can be reversed or prevented by σ 1R knockdown or a σ 1R agonists such as pentazocine, SKF10047 or PRE084 [6, 14, 75]. During stress, the σ 1R prevents cytosolic $[Ca^{2+}]$ dysregulation in neuron cells [76]. The σ 1R potentially helps buffer cytosolic $[Ca^{2+}]$ by shuttling calcium from the ER via the MAM to the mitochondria, stimulating ATP production and improving cellular health [16, 77, 78].

1.2.2 Developmental and Epileptic Encephalopathies

Developmental and epileptic encephalopathy (DEE) has recently been classified as gene-associated developmental encephalopathy without epilepsy which continues progressing even when seizures are controlled. It is then difficult to judge whether developmental or epileptic components contribute more to clinical presentation in DEE. On the other hand, developmental encephalopathy (DE) generally has only developmental impairment (e.g. intellectual disability and autistic spectrum disorder), without epileptic activity associated with regression or further slowing of development.

In contrast, epileptic encephalopathy (EE) is the epileptic activity itself contributing to severe cognitive and behavioural impairments with no pre-existing developmental delay [79]. Since most patients with DEE have aetiologies due to genetic variants, and patients with EE also have a molecular genetic basis, genetic testing to distinguish DEE becomes essential. The most frequent and studied DEE-associated genes include *CDKL5*, *SCN1A*, *KCNQ2/KCNQ3*, *PCDH19*, *SCN2A/SCN8A* and *TSC1/TSC2*. Other promising gene variants associated with an increased risk of expressing DEE comprise *KCNT1*, *STXBP1*, *CACNA1A*, *SYNGAP1*, *GABRB3* and *GRIN2A/GRIN2D* [80]. Besides genetic variants, the causation of DEE also includes structural disorder [e.g. tuberous sclerosis complex (TSC), hypothalamic hamartomas and hemimegalencephaly], metabolic disorder (e.g. pyridoxine or biotinidase deficiency and GLUT1 deficiency) and immune disorder (e.g. Rasmussen syndrome) [81–83].

The current DEE treatments are anti-seizure medications (ASMs). Other emerging pharmacologic agents to potentially treat DEE patients include serotonin modulators [e.g. EPX-100 (clemizole hydrochloride) [84, 85], EPX-200,

fenfluramine [86] and lorcaserin [87], TAK935 (soticlestat, a cholesterol 24-hydroxylase inhibitor), huperzine analogue, stiripentol (an anti-convulsant with an unknown mechanism of action) [86], ataluren (a suppressant of premature stop codons) [87], selective sodium channel modulators (e.g. verapamil) [87] and antisense oligonucleotide therapy (e.g. STK-001) [87] are in the developmental stage [88]. The first precision gene-targeted therapy specific to DEE remains challenging.

The role of the σ 1R in the development of DEE is thoroughly reviewed by Martin et al. (2021) [89]. The σ 1R, in the brain, acts on neuronal networks at the cellular level to fine tune neuronal signals, thus maintaining a balance between excitatory and inhibitory signals [90]. The σ 1R achieves this by interacting with multiple signalling pathways and proteins and becoming activated by physiological stressors, including the excitatory/inhibitory neuronal imbalances that lead to seizure activity. In σ 1R KO mice, there is a higher susceptibility to convulsive seizures caused by GABA antagonists pentylenetetrazol and (+)-bicuculline [90]. An imbalance in excitatory and inhibitory neuron signalling results in seizures. Inadequate control of glutamate signalling at the NMDAR by GABA receptors results in increased seizure activity and σ 1R activation. Since there are many psychoactive σ 1R ligands, there is the potential for treating DEEs [22].

1.2.3 Huntington's Disease

Huntington's disease (HD) is a hereditary and progressive neurological disorder caused by a Huntingtin (*HTT*) gene mutation. The mutation results in an abnormal form of the protein that accumulates in brain cells, leading to impairment of several neuronal signalling [91] and protein degradation [92] pathways and mitochondrial function [93], resulting in brain cell degeneration and eventual death.

The symptoms of HD usually appear in adulthood but can sometimes start in childhood or adolescence. They include uncontrolled movements, such as chorea (involuntary jerking), and cognitive and psychiatric symptoms, including impaired judgment, difficulty concentrating, depression and personality changes [94].

HD is inherited in an autosomal dominant manner, meaning that if a person has the mutated *HTT* gene, they have a 50% chance of passing it on to their children. There is currently no cure for Huntington's disease, and treatment focuses on managing symptoms, such as through medication and therapy. The disease can have a significant impact on an individual's quality of life, as well as on their family and caregivers [95].

The σ 1R has been shown to play a role in HD. The exact mechanism by which the σ 1R contributes to the development

and progression of the disease is still being studied. There are several possible ways in which it may be involved.

One way the σ 1R is thought to be involved in HD is through its interaction with the mutant huntingtin protein. In cell models of HD that express the mutated HTT protein, the σ 1Rs are expressed in the nuclear inclusions and help to control the degradation of malformed proteins. Knockdown of the σ 1R using small interfering RNA (siRNA) increased the volume of aggregates in the nucleus and cytoplasm and caused nuclear inclusions. Moreover, overexpression of the σ 1R reduced the nuclear inclusions [96]. Studies have shown that the σ 1R can bind to mutant huntingtin and may help to promote its toxic effects on brain cells. In addition, the σ 1R may also be involved in regulating cellular processes disrupted in Huntington's disease, such as protein folding and calcium signalling.

Studies have shown that expression of the σ 1R is increased in the striatum of mouse models and patients with severe HD [97]. This contradicts the previous experiments; however, it may be explained by the fact that the huntingtin protein and the σ 1R are involved in transporting vesicles and proteins. Therefore, overexpression of one may lead to the downregulation of the other. In the case of HD, neither protein is overexpressed, but the mutated huntingtin protein causes loss of function. As a result, the σ 1R is upregulated to compensate for this loss of function.

Evidence suggests that the σ 1R may be a potential target for treating Huntington's disease. Studies have shown that drugs that activate the σ 1R can help to reduce symptoms and slow the progression of the disease in animal models [98, 99]. However, more research is needed to determine whether these drugs will be effective in humans.

1.2.4 Motor Neuron Disease/Amyotrophic Lateral Sclerosis

Motor neuron disease (MND) is a progressive neurological disorder with particular damage to motor neurons responsible for controlling skeletal muscle activities involving limbs, speech, swallowing and respiratory wall muscles.

MND encompasses a spectrum of different motor neuron syndromes, including amyotrophic lateral sclerosis (ALS), progressive muscular atrophy (PMA), primary lateral sclerosis (PLS), and upper and lower motor neuron dominance [100, 101]. Among them, the most common form of MND is ALS which affects both upper and lower motor neurons and begins with limb weakness (i.e. arms, hands and legs) before progressive paralysis and death within 2–5 years after the symptom onset. The onset of MND is reported to be sporadic and varied among patients, and its pathogenesis is not yet fully understood.

A complex interaction between genetic and environmental factors is reported to trigger pathologic development and expansions. This involves glutaminergic pathways leading

to glutamate-induced excitotoxicity and mutations of genes such as *c9orf72*, *TDP-43* and fused in sarcoma (*FUS*), resulting in abnormalities of translation and formation of intracellular neuronal aggregates. In contrast, mutations in the superoxide dismutase-1 (*SOD-1*) gene increases oxidative stress [102, 103].

For treatment, palliative care is primarily advised. No effective medication is available apart from riluzole, a glutamatergic neurotransmission inhibitor, and sodium phenylbutyrate/ursodexicoltaurine, an inhibitor of histone deacetylases (HDACs), the only drugs approved by the US Food and Drug Administration (FDA) that can delay ALS progression and improve survival by 2–3 months [104, 105].

Another drug recently approved by the US FDA, but limited to use only in Japan and the USA, is edaravone, a free radical scavenger that halts progression in early stages [106]. Although edaravone treatment has beneficial effects, its precise mechanism and cellular/molecular targets are uncertain. Masitinib, a tyrosine kinase inhibitor, has successfully passed a randomized controlled phase II/III trial in ALS patients as an additional drug for riluzole by showing a significant improvement in ALS progression, quality of life, respiratory function and time delay to death [107]. However, further trials were halted in 2021 when observational studies indicated an increased risk of ischaemic heart disease [108].

The σ 1R is highly expressed in motor neurons, particularly in the ER at the postsynaptic subsurface cisternae of cholinergic C-terminals. It is hypothesized to help transport ions across voltage-gated potassium and small conductance calcium-activated potassium channels (SK) channels, thereby reducing motor neuron excitability [109, 110]. It is also speculated that higher conductance of potassium channels can also be indirectly mediated by the interaction of σ 1R with the IP3R3 receptor and the subsequent amplification of calcium release [16, 111].

Consistently, reports imply that genetic mutation in the σ 1R gene and the lack of functioning of the protein can result in MND [112]. The σ 1R^{E102Q} mutation (familial ALS) causes the cytoplasmic aggregation of σ 1R, making them unable to bind with IP3R with loss of function to control Ca²⁺ flux, activated calpain and reduced ATP synthesis alongside mitochondrial-associated membrane collapse [113]. Also, studies from transgenic *Drosophila* expressing σ 1R^{E102Q} confirmed the same, and showed that overexpression of IP3R or glucose transporter GluT-3 could counteract σ 1R^{E102Q} toxicity [114]. Upon σ 1R knockdown, the lipid raft markers, including Flotillin-1, Erlin-2 and Caveolin-1, were observed to be localized in non-raft regions, indicating the disruption of lipid rafts and associated intracellular calcium dyshomeostasis [115].

Ectopic expression of σ 1R^{E102Q} causes structural alteration in the mitochondria and ER, and induces defective autophagy, vesicular transport and mislocalization of

RNA-binding proteins, leading to protein quality control modulation [116]. In ALS model mice, *SOD1**G93A, knockout of σ 1R exhibited deregulated motor performance, ultimately resulting in a faster death [117]. Overexpression of σ 1R^{E102Q} missense mutation in motor neuron cellular models showed high susceptibility to apoptosis triggered by endoplasmic reticulum stress [118].

Furthermore, the cells overexpressing σ 1R^{E102Q} exhibited abnormal mislocalization of the TAR DNA-binding protein (TDP-43), resulting in a toxic cytoplasmic gain of function [119, 120]. Impaired TDP-43 regulation results in unfolded protein accumulation, defective RNA metabolism and oxidative damages [121]. Misfolding or mutation in TDP-43 increases its aggregation propensity, and the intensity of motor neuron degeneration is directly correlated with TDP-43 aggregation [122]. In the cellular ALS model, σ 1R stabilises nuclear pore proteins (Nups) by co-localizing with Ran-activating protein (RanGAP) and rescues nucleocytoplasmic transport deficit to reverse the ALS pathology, further extending the function and confirming the importance of the receptor [123].

1.2.5 Multiple Sclerosis

Multiple sclerosis (MS) is a neurodegenerative disease of the CNS, predominantly mediated by the autoimmune system. The myelin sheaths of neurons are permanently destroyed by immune cells causing acute inflammation and axonal transection lesions. After that, the chronic phase enters with remyelination and an unrepaired inflammation [124]. T- and B-cell infiltrates are correlated with demyelination [124]. MS is commonly found in young adults, especially women, aged between 20 and 30 years [125]. MS symptoms and the degree of severity are various, depending on the number, type and location of damaged nerves, and lead to a non-traumatic neurological disability affecting physical function, cognition, quality of life and employment. Fatigue has been reported to be a common symptom in 75% of individuals with MS [126]. Fatigue could be due to the involvement of the immune system, axonal loss, sleep disorders and circadian rhythm abnormalities during MS [127].

The treatment of MS is considered according to the disease status, whether relapsing–remitting MS (RRMS, defined as relapses at the onset with stable neurologic disability between episodes), secondary progressive MS with activity (SPMS, defined as steadily increasing neurologic disability following a relapsing course with evidence of ongoing inflammatory activity) or primary progressive MS (PPMS, defined as steadily progression from onset). Currently, there is no cure for MS, and disease-modifying therapies (DMTs) are primarily prescribed as they reduce annual relapse rates and short-term disability via modulation in immune systems, including sequestration of lymphocytes,

Th1/Th2 shift, DNA synthesis in lymphocytes and cytokine production [125]. Interferon (IFN)- β -1b was the first DMT approved by the US FDA to replace the former immunosuppressants such as azathioprine, methotrexate and corticosteroids [128].

Several monoclonal antibodies (mAbs) with well-clarified mechanisms of action, including ocrelizumab, natalizumab, ofatumumab and alemtuzumab, are also approved [129]. Ocrelizumab is selective to B cells expressing CD20 and causes cell destruction and reduced inflammation. This is the only drug approved to treat PPMS. Natalizumab is specific to the α 4 integrin subunit on lymphocytes and prevents lymphocytes from entering the CNS, whereas alemtuzumab selectively binds CD52 antigen on lymphocytes, leading to their depletion [129].

Besides those mAbs, sphingosine 1-phosphate receptor (S1P) modulators (e.g. fingolimod, siponimod, ponesimod and ozanimod) have been shown to bind a subtype of S1P receptors, resulting in sequestration of lymphocytes in lymph nodes [130–132]. Glatiramer acetate is another relatively safe DMT, even though its action is unclear [133]. Other DMTs, including fumarates (e.g. dimethyl fumarate and diroximel fumarate), teriflunomide (a pyrimidine synthesis inhibitor with anti-inflammatory and immunomodulatory activities) and cladribine (an immunomodulatory agent selectively targeting lymphocyte subtypes) are also available for treatment [134, 135].

In CNS, oligodendrocytes form myelin sheaths, including proliferation, migration and differentiation of oligodendrocyte precursor cells (OPCs). Studies have reported the co-localization of σ 1R in progenitor and matured oligodendrocytes, implying a role in myelination [136]. In line with this, a study has indicated that σ 1R regulated oligodendrocyte differentiation and myelin sheath generation, while σ 1R siRNA showed the opposite effect [137]. Also, σ 1R could interact with cholesterol, and dysfunction of the protein causes perturbations in cholesterol levels, which in turn can cause myelin degeneration [2, 138, 139]. The σ 1R agonist donepezil could restore mitochondrial functions in the induced circadian rhythm disruption cellular model, suggesting the role of the receptor in maintaining a healthy circadian rhythm and possibly reducing fatigue [140]. Further, targeting σ 1R reduces neuroinflammation and demyelination in the experimental autoimmune encephalomyelitis MS model, indicating the potential of the receptor to counteract the pathological effects [141].

1.2.6 Parkinson's Disease

Parkinson's disease (PD) is a common neurodegenerative disease with impairment in the CNS and motor symptoms. The early signs are usually tremors in one hand causing stiffness (rigid muscles) or slow movement (bradykinesia).

Other symptoms can be poor balance; loss of automatic movements such as blinking, smiling and swinging arms while walking; and limitations in speech and writing.

Several biomarkers have been introduced to identify PD at the early stage because most of the cardinal symptoms occur late and dopaminergic neurons are already destroyed [142]. However, only a few have been implemented in clinical practice [143]. Neuroimaging and biochemical measurements of biofluids are the current focus due to a better prediction of PD [144].

There is currently no therapy slowing or arresting the PD progression available, so symptomatic treatment using levodopa (or in combination with carbidopa) as the first-line drug together with personalized non-pharmacological management (e.g. deep brain stimulation, strength training, gait and balance training and hydrotherapy) from a multidisciplinary team approach are primarily suggested [145, 146]. Although levodopa has been proven to improve disability and discomfort in PD patients through the conversion of L-3,4-dihydroxyphenylalanine (L-DOPA) into DA in striatum, the complete mechanism of actions and causes of its adverse events, such as dyskinesias and levodopa resistance from long-term use, are not fully understood [147, 148]. α -Synuclein deposition in the brain and the formation of Lewy bodies are thought to be a hallmark of PD-related dementia characterized by changes in thinking, movement, behaviour and mood [149]. PD-related dementia and dementia with Lewy bodies are the second most common causes of dementia behind AD, accounting for 15–20% of the incidences of dementia globally [150]. Drugs targeting the build-up of Lewy bodies could potentially help prevent PD-related dementia.

There is a reduction in σ 1R expression in PD, much like in AD [151], plus there is an age-related loss of dopaminergic neurons in σ 1R knockout mice [152]. The σ 1R may also alleviate DA toxicity in PD. High availability of DA may act as a neurotoxin, and ischaemia, hypoxia and local exposure to neurotoxins, such as high concentrations of excitatory amino acid and methamphetamine, all result in increased concentrations of DA. The DA neurotoxicity mechanism is linked to an imbalance in REDOX management. DA can spontaneously oxidize in vitro or in an enzyme-catalysed reaction in vivo to form reactive oxygen species (ROS) and DA-quinones. DA-quinones can bind to cysteine or cysteinyl residues on proteins, resulting in an imbalance of glutathione homeostasis (a key cellular antioxidant) [153]. ROS is increased in wild-type CHO and σ 1R knockdown CHO cells treated with physiologically relevant DA concentrations (below 10 μ M). However, apoptosis only increases in the σ 1R knockdown cells [154]. Furthermore, DA-treated Neuro2A cells expressing APP^{SWE} undergo ROS-mediated apoptosis [155], and this can be prevented using this σ 1R agonist dipentylamine

(this protective effect itself was reversed by σ -1 antagonist BD1047) [156].

Mutations in genes associated with PD, such as *α -synuclein*, *DJ-1*, *LRRK-2* (which encodes leucine-rich repeat kinase 2) and *GBA* (which encodes glucocerebrosidase) have been identified, which cause abnormal α -synuclein proteostasis leading to mitochondrial dysfunction, oxidative stress, neuroinflammation and motor circuit pathophysiology [157].

Oxidative damage to the substantia nigra has been demonstrated in patients with PD [158]. Oxidative stress accelerates α -synuclein aggregation and cell death [159, 160]. Multiple studies have identified σ 1R ligands to reduce oxidative stress through multiple in vivo and in vitro models via various signalling pathways involving protecting the ER and mitochondria [161–165].

Similarly to A β in AD, α -synuclein oligomers bind to PGRMC1 at synaptosomes in the brain [166]. Furthermore, in chaperone-controlled autophagy, α -synuclein binds to lysosome-associated membrane protein 2A (LAMP-2A) [167, 168] (LAMP-2A is the receptor for chaperone-mediated autophagy). α -Synuclein disrupts cholesterol-rich lipid membranes where the σ 2R is located and disrupts downstream signalling events, including autophagy [169–171]. Without efficient autophagy, neurons gather ubiquitinated protein aggregates (proteins tagged with ubiquitin for degradation). Thus, allosteric antagonists of the σ 2R may prevent the effects of α -synuclein oligomers via various mechanisms (possibly similar to AD and A β accumulation at the synaptosomes).

A high-throughput screening assay making use of the National Institutes of Health (NIH) small molecule repository (725 compounds that have been tested in phase I/III clinical trials) and the CogRx library of high-affinity σ 2R allosteric antagonists, investigated the effects of these compounds on lipid vesicle trafficking deficits caused by α -synuclein [172]. The study found that the most effective compounds were the σ 2R allosteric antagonists [172]. Furthermore, an α -synuclein oligomer-induced increase in LAMP-2A expression was blocked by the σ 2R allosteric antagonists.

1.2.7 Rett Syndrome

Rett syndrome (RS) is a rare genetic neurodevelopmental disorder affecting brain development and consequently developing into physical, cognitive and intellectual disabilities. RS is mostly found in females [173]. The most precise cause of RS is demonstrated to be the loss-of-function mutation of methyl CpG binding protein 2 (*MECP2*) gene located at the X chromosome Xq28 [174].

Dysfunction of MeCP2, an epigenetic modulator of gene expression acting as both a transcriptional activator

and repressor, induces either positive or negative changes in many other gene expressions at transcriptional and post-transcriptional levels [175, 176]. MeCP2 deficiency also impacts successive stages of brain development and contributes to mitochondria dysfunction [177], impaired protein synthesis [178] and reduced synaptogenesis [179], leading to pathology of RS. The important symptoms for differential diagnoses are cerebral palsy, autism, Angelman syndrome and non-specific developmental delay [173]. Spectrums of these symptoms, together with progression and severity, are also broad and individualized [180].

There is no cure available [181], so medical care and management aim to provide symptomatic relief through pharmacological and non-pharmacological approaches. Medication is needed for breathing and motor difficulties. Since over half of RS patients have seizures, anti-convulsant/anti-epileptic drugs such as valproate, lamotrigine, levetiracetam and carbamazepine are usually prescribed [182]. Serotonin-based therapy, such as serotonin 1A receptor agonists and serotonin reuptake inhibitors (SSRIs) are also reported to help reduce symptoms of sleep apnoea [183].

A wide variety of neurologic impairments characterizes RS. Unlike some neurodegenerative diseases (PD and AD in particular), there appears to be little or no causal link established between the σ 1R and the progression of RS. However, due to the promiscuous relationship of the σ 1R with multiple proteins and signalling pathways in the brain and its relative success as a target for treating other neurodegenerative diseases, the σ 1R is a target for treating the neurological damage seen in RS.

Most patients carry a mutation in the *MeCP2* gene [184]. MeCP2 is found at high levels in neurons and is associated with the maturation and synaptic plasticity of the CNS [185, 186]. MeCP2 is largely thought to negatively regulate gene expression (although there is some evidence that it activates some genes).

However, when mutated in Rett syndrome, MeCP2 target genes that would normally be activated remain inactive, and genes that would be inactivated remain active, thus producing unneeded proteins. Brain-derived neurotrophic factor (BDNF) is one protein that MeCP2 is thought to negatively regulate [187], along with the mTOR/AKT signalling pathways [178]. The σ 1R is known to have a role in regulating BDNF expression in multiple cellular systems, including the brain [188], eyes [189] and heart [190]. Furthermore, many studies have identified that σ 1R agonist treatment of various cells results in AKT phosphorylation [191]. Also, σ 1R activity has a role in autophagy regulation, becoming a novel target for treating neurological disease [53, 65, 192].

1.2.8 Wolfram Syndrome

Wolfram syndrome (WS), also known as diabetes insipidus, diabetes mellitus, optic atrophy and deafness (DIDMOAD), is a rare genetic disorder affecting many body parts. It is caused by mutations in the *WFS1* gene, which provides instructions for making a protein called wolframin [193].

The most common features of the disorder include diabetes mellitus (a condition that affects how the body uses and stores glucose), diabetes insipidus (a disorder that causes excessive thirst and urination), optic atrophy (damage to the optic nerve that can cause vision loss) and deafness. WS may also include neurological problems such as ataxia (difficulty with coordination and balance), seizures, behavioural problems, urinary tract problems and psychiatric disorders.

There is no cure for WS, and treatment is based on the individual's symptoms. Management may involve regularly monitoring blood sugar levels, hormone replacement therapy and other supportive measures such as hearing aids and visual aids [194].

Wolframin protein is expressed at the ER, interacts with the MAM and is involved in calcium transfer between the ER and the mitochondria. The mutated form of wolframin disrupts calcium transfer, leading to mitochondrial dysfunction and cell death [195]. The σ 1R is heavily involved in chaperoning proteins, such as the IP3R, in calcium shuttling between the ER and mitochondria. σ 1R activation, either during cellular stress or with an agonist, results in calcium transfer from the ER to the mitochondria reducing ER stress, protecting the mitochondria and improving cellular ATP levels and the general health of the cell [196]. In mouse and

zebrafish models of WS, overexpression of the σ 1R alleviated the symptoms and restored calcium homeostasis in vivo and in vitro [197].

2 Sigma Receptor Ligands for Neurological Disease


2.1 Amantadine

Amantadine (Fig. 2) is a weak NMDA antagonist that increases DA release and blocks DA reuptake [198, 199]. Amantadine also has a reported affinity for the σ 1R of 7 μ M ($[^3\text{H}]$ -(+)-SKF-10,047 in rat cerebellum or forebrain homogenates [200]) or 740 μ M ($[^3\text{H}]$ -(+)-pentazocine in permeabilized MDA-MB-468 breast cancer cells [2, 3]), and also has an affinity for the σ 2R of 9 μ M ($[^3\text{H}]$ -DTG in membranes prepared from MCF7 breast cancer cells [5, 201]). Amantadine was previously used to treat influenza-A in the early 1960s and 1970s, although it is no longer recommended due to widespread resistance [202]. Amantadine has a mild side effect profile. The common neurological side effects include drowsiness, lightheadedness, dizziness and confusion [203].

2.1.1 Parkinson's Disease

Amantadine is a drug long used in treating PD [204] and has been used for treating movement disorders in PD for over 50 years. Despite its long use in treating PD, the pharmacological activity of amantadine is still not fully understood. Its

Fig. 2 Chemical structures of drugs with sigma receptor activity discussed within this manuscript.



use in PD has declined since the introduction of L-DOPA treatment. However, it is still used in 10% of patients today, alone or in combination with L-DOPA [205].

The blocking of [^3H]-(+)-SKF-10,047 binding by amantadine in vivo in various mouse brain structures correlated with the data from the rat cerebellum or forebrain homogenates [200], and in post-mortem studies of human brains [206], suggesting a physiologically relevant interaction of amantadine with the σ1R .

The use of amantadine in treating PD and L-DOPA-induced dyskinesia has been debated, with one systematic review suggesting there is no evidence for its safety and benefit in treating PD [207]. This may be due to the dose used and the pharmacokinetics of amantadine. As plasma concentrations of amantadine increase, there is a greater risk for toxicity, particularly in patients with renal diseases (the kidneys remove most amantadine) [208]. New formulations of amantadine that release over a longer period have been trialed and approved for PD in the USA [209–212] (Table 1). The extended-release formulation is typically used to help manage PD symptoms throughout the day rather than providing short-term relief. Slowly releasing the medication over several hours can help control motor symptoms such as tremors, stiffness and difficulty with movement during both dyskinesia and OFF episodes. Immediate-release amantadine is typically taken two to four times a day, and its effects can wear off relatively quickly, which may result in fluctuations in symptom control [209]. On the other hand, extended-release amantadine is designed to slowly release the medication over several hours, providing a more stable blood level of the medication and, therefore, more stable symptom control.

Extended-release amantadine may also have a lower incidence of certain side effects, such as nausea and dizziness, compared with immediate-release amantadine. This is because extended-release amantadine releases the medication more slowly, which can reduce the peak concentration of the drug in the blood and potentially minimize side effects [213–215].

2.1.2 Multiple Sclerosis

Amantadine treatment improved fatigability associated with MS in several randomized controlled studies [216–219]. Amantadine treatment has been earlier reported to induce sleep spindles in PD individuals, implying the role in rectifying sleep disorders and associated fatigue during MS [220]. Amantadine has been trialed to relieve fatigue in MS patients (NCT03185065). Using a randomized cross-over study design, 100 mg of amantadine (increasing to 200 mg if tolerated) was given for 6 weeks per treatment period, followed by a 2-week washout. Fatigue (the primary endpoint) was measured using the modified fatigue impact

scale (MFIS) between days 26 and 35. However, the study showed no significant improvement over the placebo and increased adverse events [221]. There have been three other clinical trials for MS with extended-release formulations of amantadine concerning drug safety and walking impairment (NCT02471222, NCT03567057 and NCT03436199). Amantadine extended release at 137 mg, (165 patients) and 274 mg (132 patients) improved walking speed in the Timed 25 Foot Walk compared with the placebo (174 patients) (NCT03436199). Furthermore, these improvements were also seen in the open-label follow-up study (NCT03567057). The data from NCT02471222 have not been made available.

2.2 ANAVEX2-73 (Blarcamesine)

Blarcamesine (ANAVEX2-73; Fig. 2) is a σ1R receptor agonist with an affinity of 860 nM (IC_{50}) [222] developed for treating AD. Blarcamesine also has agonist activity at the muscarinic M1 receptor, and the NMDA receptor with affinities of 5 and 8 μM , respectively [222].

2.2.1 Alzheimer's Disease

Pre-clinical studies have shown that blarcamesine can reverse scopolamine's long-term amnesic effects in mice [222]. Furthermore, blarcamesine can reverse the effect of learning deficits in mice injected with $\text{A}\beta_{25-35}$ [222–224]. Blarcamesine given to mice injected with $\text{A}\beta_{25-35}$ restored respiration rates in hippocampal mitochondria and reduced lipid peroxidation levels. Blarcamesine could also prevent tau-hyperphosphorylation and amyloid- β (1–42) generation in mouse models of AD [223]. Furthermore, indicators of increased toxicity, including Bax/Bcl-2 ratio and cytochrome C release into the cytosol, were reduced [224]. Blarcamesine has protective effects in *Caenorhabditis elegans* and cell culture models of AD by enhancing autophagic flux and increasing proteostasis [65].

A phase 1 clinical study in healthy male volunteers showed that doses up to 50 mg were well tolerated, with no serious adverse events reported in a poster at the CNS summit in 2014 [225]. At 60 mg, three out of four subjects presented dose-limiting CNS symptoms (dizziness and headache). The study concluded with pharmacokinetic analysis showing extensive biotransformation of blarcamesine to AV19-144 (the bioactive form of the drug) with assumed linear pharmacokinetics after single oral dosing of 1 to 60 mg blarcamesine [225].

A biomarker analysis, phase 2a study (NCT02244541), that include some exploratory study into the benefits of blarcamesine, where blarcamesine was administered over 57 weeks, showed a significant reduction in cognitive decline [226]. When the study was extended to 265 weeks (NCT02756858), there was confirmation of this prevention

Table 1 Clinical trials of extended-release amantadine

Study title	Conditions studied	Primary outcome measures	Study results	Trial Phase	Trial number
E-R amantadine safety and efficacy study in LID (EASED Study)	Dyskinesia LID PD	Change in UDysRS total score from baseline to week 8	E-R amantadine appeared relatively safe with minimal adverse effects at doses of 260 and 340 mg. There was also a statistically significant reduction in LID (measured using the change in UDysRS) total score. Amantadine reduced LID compared with placebo	2/3	NCT01397422
Efficacy and safety study of ADS-5102 in PD patients with LID (EASE LID 3)	Dyskinesia LID PD	Change in UDysRS total score from baseline to week 12	A statistically significant improvement was measured using UDysRS total score for 340 mg amantadine compared with placebo [210–212, 214].	3	NCT02274766
ADS-5102 for the treatment of LID (EASE LID Study)	Dyskinesia LID PD	Change in UDysRS total score from baseline to week 12	A statistically significant change in UDysRS of -15.9 (1.6) for ADS-5102. A statistically significant change in OFF time, which decreased by a mean (SE) of 0.6 (0.3) hours for ADS-5102 and increased by 0.3 (0.3) hours for placebo [214].	3	NCT02136914
Open-Label safety study of ADS-5102 in PD patients with LID	Dyskinesia LID PD	Number of participants with reported AEs and safety-related study drug discontinuations	Patients switching from IR to E-R amantadine experienced an improvement in motor complications, without exacerbating adverse events. Plus a statistically significant reduction in their MDS-UPDRS Part IV scores at the first post-baseline visit (week 8) plus many patients were able to reduce their levodopa use [212].	3	NCT02202551

AEs adverse events, E-R extended release, IR immediate release LID levodopa-induced dyskinesia, PD Parkinson's disease, SE standard error, UDysRS Unified Dyskinesia Rating Scale

of decline at 148 weeks with analysis of blood concentrations of blarcamesine and its metabolite AV19-144, showing that those with high concentrations of blarcamesine had improved therapeutic responses of 78% and 88% in adjusted MMSE and adjusted AD cooperative study–activities of daily living (ADCS–ADL), respectively, relative to the low/medium concentrations of blarcamesine.

Furthermore, genomic analysis also indicated the significant effects of biomarkers on clinical outcomes of blarcamesine. Two patients showed an exceptional therapeutic response at 148 weeks. Both subjects had $\sigma 1R$ wild type and a high mean concentration of blarcamesine in plasma, with a baseline MMSE > 20 [226].

The study of biomarkers that predict the potential therapeutic outcome of a drug is especially useful in a disease with such heterogeneity as AD [226]. The ApoE $\epsilon 4$ genotype alone lacks predictive power since only approximately 25% of people carry this genotype. The study discussed above identified that patients with a baseline MMSE greater than 20 who had the $\sigma 1R$ wild type with a high mean blarcamesine serum concentration and *ApoE3* alleles had significantly better outcomes in the ADCS–ADL. The *SIGMAR1* Q2P variant, having a baseline MMSE score lower than 20 and a low mean serum concentration of blarcamesine were predictive of poor outcomes in the ADCS–ADL [226].

Blarcamesine is undergoing further phase 2b/3 clinical trials (ongoing: NCT04314934; completed with data yet to be published: NCT03790709). The results of these clinical trials will be interesting, given the promise blarcamesine has shown in the pre-clinical studies.

2.2.2 Multiple Sclerosis

Blarcamesine increased the proliferation of oligodendrocyte progenitor cells to myelin sheath-producing oligodendrocytes in a $\sigma 1R$ -dependent manner and protected them from excitotoxic insults [227]. There are no currently listed clinical trials for blarcamesine in MS on www.clinicaltrials.gov.

2.2.3 Rett Syndrome

Long-term administration of blarcamesine to mice deficient in *MeCP2* restores physiological and neurological abnormalities that mimic the human Rett syndrome, including ameliorating motor and acoustic startle deficits and reversing expiratory apnoea [228]. Several $\sigma 1R$ -related mechanisms could benefit the neurological and motor symptoms seen in this and other models of Rett syndrome. Enhancement of calcium homeostasis via $\sigma 1R$ shuttling of calcium to the mitochondria leads to improved mitochondrial function and the activation of autophagy via $\sigma 1R$ activation, resulting in a reduction of misfolded proteins and fewer damaged organelles or the restoring/upregulation of BDNF [65, 228, 229].

This leads to potential improvements in synaptic function in multiple brain regions.

Blarcamesine is currently in phase 2/3 clinical trials (NCT03758924, NCT03941444 and NCT04304482). Blarcamesine has shown positive results in a phase 2 clinical trial (NCT03758924) in female patients diagnosed with Rett syndrome (positive *MeCP2* gene mutation), meeting both primary and secondary endpoints with no reported serious adverse events (press release) [230]. In another phase 3 (NCT03941444) clinical trial the primary and secondary endpoints were met with significant improvements versus placebo in the clinical global-impression of improvement scale (CGI-I); the anxiety, depression and mood scale (ADAMS); and drug exposure-dependent response of the Rett Syndrome Behaviour Questionnaire (RSBQ), all with a low incidence of adverse events (press release) [230]. It is important to note that the data available from these studies has not been peer reviewed and formally published. There are further ongoing phase 3 studies organized by Anavex Life Sciences (NCT04304482). However, while the data look promising, it will be important to see some independent studies with peer-reviewed results to fully assess the efficacy and safety of blarcamesine for Rett syndrome treatment.

2.3 ANAVEX3-71

ANAVEX3-71 (AF710B) (Fig. 2) is an agonist at the muscarinic M_1 receptor and $\sigma 1R$, and the second drug developed by for AD treatment. ANAVEX3-71 has an affinity for the $\sigma 1R$ of 1.3 nM and an affinity for the $\sigma 2R$ of 10 μM [62]. The affinity for the M_1 receptor is approximately 0.05 nM. ANAVEX3-71 also has affinities for the μ opioid receptor and serotonin receptor (approximately 10 μM for each) [62].

2.3.1 Alzheimer's Disease

Given the complex pathology of AD, it is of interest that a drug such as AF710B may interact with multiple target sites to have a beneficial effect in AD [52].

Pre-clinical studies have shown ANAVEX3-71 to be worthy of further study. AF710B has been shown to modify the disease-defining hallmarks of AD in transgenic (3 \times Tg-AD) mice, such as cognitive deficits, amyloid and tau pathologies, and beneficial effects on mitochondrial dysfunction and neuroinflammation [62, 231].

Phase 1 clinical trials (NCT04442945) in healthy individuals have been reported on Anavex Life Sciences' website, suggesting that in healthy volunteers, 5–200 mg daily had no serious adverse events. The pharmacokinetics of ANAVEX3-71 (serum concentration of AF710B) was also proportional to the dose for doses up to 160 mg. Furthermore, there were no clinically significant electrocardiogram

(ECG) parameters throughout the study (press release) [232].

There are no further registered (www.clinicaltrials.gov) ongoing clinical trials with ANAVEX3-71. However, Anavex states on its website that they plan further phase 2 clinical trials. More independent studies are required. However, pre-clinical data on ANAVEX3-71 suggest it could be useful in treating dementia and AD.

2.4 CogTX Compounds

CT1812 (Fig. 2) is one of several σ 2R antagonist compounds developed by Cognitive Therapeutics Inc. (CogTx) for treating AD [233]. Other similar σ 2R compounds from this company that have a strong affinity for the σ 2R are CT1978 (Kd = 9.2 nM), CT1970 (Kd = 1.8 nM), CT2168 (Kd = 2 nM), CT1681 (Kd = 12 nM), CT1696 (Kd = 11 nM) and CT1925 (Kd = 18 nM), and act as allosteric σ 2R antagonists [172].

CT1812 has become the company's lead compound, with higher selectivity of the σ 2R and improved metabolic stability. Compounds CT01346 and CT01344 had a low polar surface area and off-target effects at the Ether-a-go-go-related gene (*ERG*) and at other sites. CT1812 has an affinity of 8.5 nM for the σ 2R [233].

2.4.1 Alzheimer's Disease

CT1812 has been demonstrated in rats to prevent A β -oligomer binding to neurons and to displace bound A β -oligomers from neuronal receptors [234, 235]. CT1812 does not block oligomer formation, suggesting that its ability to stop A β -oligomer binding and trafficking deficits are a result of σ 2R binding. A β oligomer concentrations were measured in the brain interstitial fluid (ISF) of 12-month-old APP^{Swe}/PS1dE9 transgenic mice using electrodes coated with the oligomer-specific antibody placed in the hippocampus. CT1812 resulted in a rapid and significant increase in A β -oligomer levels (not seen with the vehicle control) but did not affect A β monomer levels.

Furthermore, post-mortem tissue samples observed a similar displacement of A β -oligomers after treatment with CT1812 [175]. CT1812 is the first drug for AD to work by displacing A β -oligomer from synaptic receptors and being the first σ 2R antagonist. Interestingly, CT1812 mimics the protective effects of the Icelandic A673T mutation in the *APP* gene, which may be predictive of clinical success [236].

CT1812 has been studied in phase 1 clinical trials and is relatively safe (NCT03716427 and NCT02907567) at doses up to 560 mg, resulting in minimal adverse events in healthy volunteers (reported in an abstract from the Alzheimer's Association) [237, 238]. In a placebo-controlled biomarker study (NCT02907567) in mild AD patients treated with CT1812, there was a significant increase in

CSF A β -oligomer concentrations and decreased synaptic protein fragments neurogranin and synaptotagmin relative to baseline compared with placebo-treated patients [235]. There are several phase II clinical trials, either ongoing (SEQUEL study, NCT04735536, SHINE study, NCT03507790, NCT05225415 and NCT05531656) or completed, without published results or results made available (SNAP study: NCT03522129, NCT05225389, NCT05248672 and NCT02570997). The SPARC study (NCT03493282) aimed to assess the effects of CT1812 on synaptic density and suggests an increase in synaptic density based on the available data found at www.clinicaltrials.gov, along with imaging results and statistical analysis, which have yet to be formally published. In the same study, results posted online appear to show a reduction in CSF A β ₄₀, CSF A β ₄₂, CSF tau, CSF phospho-tau, CSF neurogranin (NRGN), CSF(SNAP25) and CSF, and increases in CSF synaptotagmin and neurofilament light (NFL). However, there appeared to be no obvious improvements in the cognition scores, MMSE or the AD Assessment Scale–Cognitive Subscale (ADAS–COG), although without complete data analysis and statistical study, it is difficult to comment on these study aspects.

2.4.2 Parkinson's Disease

α -Synuclein binds to LAMP-2A, and mutated forms bind to LAMP-2A tighter, preventing degradation of other substrates and inhibiting the autophagic process. The CogTx compounds were screened in primary neuronal cells against α -synuclein oligomers. Each compound inhibited α -synuclein oligomer-induced lipid vesicle trafficking deficits in a dose-dependent manner [CT1978 (EC50 = 1.48 μ M), CT1970 (EC50 = 6.50 μ M), CT2168 (EC50 = 1.5 μ M), CT1681 (EC50 = 9.44 μ M), CT1696 (EC50 = 15.53 μ M) and CT1925 (EC50 = 8.95 μ M)]. Two more potent compounds, CT1978 and CT2168, were also tested against LAMP-2A expression, and both prevented α -synuclein-induced upregulation of LAMP-2A [172]. Taken together, the data from this study suggest that σ 2R ligands (antagonists) may potentially regulate chaperone-mediated (LAMP-2A) autophagy and have a potential role in PD therapy.

The CogTx σ 2R compound CT1812 discussed above regarding AD is also in phase 2 clinical trials to assess the efficacy of CT1812 in patients with Lewy body dementia (NCT05225415).

No other studies/trials are registered at www.clinicaltrials.gov regarding the above compounds or any other σ 2R ligand. However, these compounds appear to have potential in treating PD, and their activities through the σ 2R warrant studying other σ 2R antagonists in relation to PD.

2.5 Dextromethorphan

Dextromethorphan (DXM) (Fig. 2) is a drug used in cough-suppressing medicines and targets multiple receptors, including inhibiting serotonin reuptake [239] and acting as an agonist at the σ 1R [240]. DXM has an affinity for the σ 1R of between 142 and 652 nM, and between 2 and 9 μ M for the NMDA receptor [241]. DXM is metabolized into its active metabolite dextrorphan (DXO) by the enzyme cytochrome P450 (CYP)2D6, a much more potent NMDA receptor antagonist (400–900 nM) but a much less potent serotonin reuptake inhibitor than (400–800 nM) DXM (20–40 nM) [240]. Dextromethorphan/quinidine (AVP-923; Nuedexta) is developed and marketed by Avanir Pharmaceuticals. Adding quinidine to the DXM reduces its metabolism to DXO by inhibiting CYP2D6. Deuteration of DXM (AVP-786, also produced by Avanir Pharmaceuticals) further hinders its metabolism by CYP2D6, allowing for lower doses of quinidine in the combination. AXS-05 combines DXM and the atypical antidepressant drug bupropion (developed and marketed by Axsome Therapeutics). In this combination, bupropion also acts as a potent CYP2D6 inhibitor, preventing the bioactivation of DXM.

2.5.1 Alzheimer's Disease

Pseudobulbar affect (PBA) is a condition linked to various neurological disorders and manifests as uncontrollable laughing or crying. There is evidence to suggest that σ 1R agonists act to reduce PBA in patients with neurological diseases. The mechanism by which DXM is thought to treat PBA is related to NMDA and σ 1R binding; it is also noted that dextromethorphan most prominently binds in the brainstem and cerebellum, areas particularly rich in σ 1R and is implicated in PBA [242]. DXM, with the addition of quinidine inhibiting its metabolism, has been shown to positively affect the symptoms of PBA [243]. Dextromethorphan/quinidine is the first FDA-approved drug for the treatment of PBA.

Agitation is another symptom of AD and is one of the most distressing for caregivers, resulting in institutionalization, further isolation and reduced quality of life for the AD patient. There have been multiple studies investigating DXM in combination with other drugs (AVP-923 and AXS-05) or the deuterated form of DXM in combination with quinidine (AVP-786) [244–246]. The registered trials with DXM and DXM-related drug combinations are listed in Table 2.

DXM has an affinity of 205 nM for the σ 1R, whereas its metabolites DN and dimemorfan (DF) have affinities of 151 and 144 nM, respectively. All three compounds have a low affinity for the σ 2R (4–11 μ M). DN has a higher affinity for the PCP site of the NDMAR (900 nM), whereas DXM and DF have affinities of 7 and 17 μ M, respectively [247].

2.5.2 Parkinson's Disease

Intraperitoneal injections (IP) of DXM (20 mg/kg) to 6-OHDA-treated rats significantly attenuated 6-OHDA-induced DAT and SERT loss, correlating to rotational behaviours [4]. DXM also has anti-dyskinetic activity [248, 249]. For a long time, it was assumed that these effects were mediated via the NMDA receptor since DXM has moderate to low affinity at this receptor and acts as an antagonist. However, it has been confirmed that the σ 1R is responsible (at least in part) for the anti-dyskinetic effects since other NMDA antagonists, MK-801 and HA-966, failed to suppress dyskinesias in the 6-OHDA rat model, whereas DXM did and the σ 1R antagonist BMY-14802 could reverse this [250].

Analysis of patient data between January 2005 and December 2013 showed that low cumulative doses of DXM may potentially achieve a protective effect for PD; however, high cumulative doses seem to be a risk for PD [251].

DXM/quinidine 45 mg/10 mg twice daily has been clinically trialled (NCT01767129) to treat levodopa-induced dyskinesia in PD. The small study provided evidence of the clinical benefit of the drug 10 mg twice daily (BID) for treating L-DOPA-induced dyskinesia in patients with PD [252]. These results aligned with two older studies showing similar anti-dyskinetic effects of DXM alone [249, 253]. However, at the time of writing, DXM (and its formulations) have only been approved for the treatment of PBA and uses in other conditions have been discontinued.

In a case study of a patient with Lewy body dementia who exhibited agitation and aggression (common behavioural and psychological symptoms of Lewy body dementia), the patient was prescribed an acetylcholinesterase inhibitor (rivastigmine) in combination with different anti-psychotics, antidepressants with anti-histaminic effects, anti-convulsants and benzodiazepines, without effect on the outbursts of aggression (paradoxically, the agitation and aggression worsened). The caregiving physician then tried DXM in combination with the CYP2D6 enzyme-inhibiting drug paroxetine. Within 1 day, the patient's behaviour improved with the cessation of the aggression. The patient remained on the DXM/paroxetine combination plus the anti-psychotic drugs. However, 6 weeks post-discharge from the hospital, the patient committed suicide. It is unclear whether any prescribed drugs had a causal relationship with the patient taking their life [254]. However, there is reported evidence that DMX abuse can lead to assault, suicide or homicide [255].

2.5.3 Motor Neuron Disease/Amyotrophic Lateral Sclerosis

As the first FDA-approved drug for the treatment of PBA, dextromethorphan/quinidine enhanced the bulbar function (speech, swallowing and salivation) in ALS individuals in

Table 2 Clinical trials of dextromethorphan, and drug formulations of dextromethorphan for AD and AD-related symptoms.

Study title	Conditions studied	Primary outcome measures	Study results	Trial phase	Trial number
Efficacy, safety and tolerability study of AVP-923 (dextromethorphan/quinidine) for treatment of symptoms of agitation in participants with Alzheimer's disease	Agitation in AD	Change in the NIP Agitation/Aggression Domain Score from day 1 to day 36 and from day 36 to day 70	A significantly reduced NPI Agitation/Aggression scores for dextromethorphan-quinidine versus placebo [244]	2	NCT01584440
Nuedexta in the treatment of pseudobulbar affect in patients with Alzheimer's disease	PBA in AD	Reduction of PBA frequency	No results posted (last update 2019)	4	NCT01852350
Open-label safety study of AXS-05 in subjects with Alzheimer's disease agitation	Agitation in AD	Incidence of treatment-emergent adverse events following dosing with AXS-05	Study ongoing (last update 2023)	3	NCT04947553
A study to assess the efficacy and safety of AXS-05 in subjects with Alzheimer's disease agitation	Agitation in AD	Cohen-Mansfield Agitation Inventory (CMAI)	AXS-05 significantly relieved agitation, a mean percentage reduction from baseline of 48% for AXS-05 versus 38% for placebo [246]	3	NCT05557409
Assessing clinical outcomes in Alzheimer's disease agitation (AXS-05)	Agitation in AD	Time from randomization to relapse of agitation symptoms	Study ongoing (last update 2022)	3	NCT04797715
Long-term extension study of the safety and efficacy of AVP-786 for the treatment of agitation in patients with dementia of the Alzheimer's type	Agitation in AD	Number of participants with any type of adverse event. Sheehan Suicidality Tracking Scale, Mini-Mental State Examination and Epworth Sleepiness Scale	Study ongoing (last update 2023)	3	NCT02446132
Efficacy, safety and tolerability of AVP-786 for the treatment of agitation in patients with dementia of the Alzheimer's type	Agitation in AD	Change from baseline in the CMAI Composite Score to week 6 and week 12	AVP-786 showed some statistically significant reductions in agitation measured by CMAI and mADCS-CGIC-Agitation score	3	NCT02442765
Efficacy, safety and tolerability of AVP-786 for the treatment of agitation in participants with dementia of the Alzheimer's type	Agitation in AD	Change from baseline to week 12 in the CMAI Composite Score	AVP-786 showed some statistically significant improvement versus placebo in the secondary outcome in change from baseline to week 12 in the NPI Agitation/Aggression Domain score	3	NCT02442778
Comparative effectiveness of MCI and dementia treatments in a community-based dementia practice	Mild cognitive impairment	Cognitive testing – Memory Orientation Screening Test (MOST)	No results published (last update 2016)	3	NCT02860338
Safety study of AVP-923 in the treatment of IEED also known as PBA	PBA in AD	Emotional control	No results published (last update 2016)	3	NCT00056524

AD Alzheimer's disease, CMAI Cohen-Mansfield Agitation Inventory, mADCS-CGIC Mean Modified Alzheimer's Disease Cooperative Study-Clinical Global Impression of Change, IEED Involuntary Emotional Expression Disorder, NPI Neuropsychiatric Inventory, PBA pseudobulbar affect.

a randomized clinical trial [256, 257], and several other trials have been completed (NCT01806857, NCT03883581, NCT01806857 and NCT00573443). However, in the long term, the drug does not affect the deterioration of bulbar function or avert non-invasive respiratory management in ALS individuals [258].

2.5.4 Multiple Sclerosis

Long-term treatment with DXM in experimental animal models of MS showed a reduction in neutrophil infiltration through the inhibition of NOX-2 and attenuated axonal demyelination through the upregulation of σ 1R [259]. In addition, DXM stimulated the proliferation and maturation of oligodendroglia progenitor cells to oligodendrocytes and protected against excitotoxicity, indicating neuroprotective effects [260].

Two placebo-controlled trials with dextromethorphan/quinidine have shown an effective therapy for PBA in MS patients [261, 262], and treatment with fluoxetine and DXM also produced a similar effect in women with MS [263].

A common symptom of MS is chronic pain [264]. The σ 1R has long been associated with pain since its identification and misidentification as a novel opioid receptor [265]. Panitch et al. [266] showed evidence supporting the off-label use of dextromethorphan/quinidine for pain reduction in MS. However, pain measurement was not a primary outcome measure for those particular trials. DXM has been trialled for treating central neuropathic pain in multiple sclerosis (NCT01324232). There have been at least five studies into the use of DXM in pain treatment from various conditions other than MS, with the results pointing towards a possible pain-relieving effect of DXM (unrelated to MS) [266–271]. Concerning the σ 1R, pain-relieving drugs studied tend to be antagonists [40], and DXM is a σ 1R agonist. Thus, any pain-relieving effect may not be due to DXM's activation of the σ 1R.

2.5.5 Rett Syndrome

DXM has also been trialled in Rett syndrome patients (NCT00593957, NCT01520363, NCT00069550 and NCT00004807). Two trials have published results at www.clinicaltrials.gov (NCT01520363 and NCT00593957) [272]. In the two groups (PCTDMRTT–NCT01520363), 24 patients in DXM (5 mg/kg per day) and 26 in placebo, were evaluated in the Mullen Scales of Early Learning (Mullen) at the end of the three-month treatment period. The open-label study had three groups, each receiving either 0.25, 2.5 or 5 mg/kg/day. At six months, the improvement in receptive language was measured by the Mullen Scale and the Screen for Social Interaction (SSI) score (mean and difference from baseline) were recorded. The data from these

studies do not look promising. While the safety of the drug was not an issue, with only a few minor adverse events, the study showed little efficacy in the primary and secondary outcomes in both studies.

2.6 Dipentylammonium

Dipentylammonium (DPA) (Fig. 2) is an *N*-alkyl-amine similar in structure to that of (2*R*-trans)-2-butyl-5-hetylpyrrolidine (σ 1R K_i 2nM [273]), tridemorph (σ 1R K_i 0.3–30 nM [274]), stearylamine (σ 1R K_i 8 μ M [275]), dodecylamine (σ 1R K_i 183 nM [274]) and endogenous sphingolipids [276]. DPA has an affinity for the σ 1R (K_i 17 nM), and it binds to the same binding site as (+)-pentazocine in a competitive fashion [2, 5, 156].

2.6.1 Alzheimer's Disease

DPA has been shown to prevent oxidative stress and ER stress in HT-22 cells exposed to 5 mM glutamate. Reducing cell death and ROS production in a σ 1R antagonist sensitive manner. Furthermore, DPA increased neurite outgrowth in Neuro2A cells and prevented DA toxicity in APP^(Swe) over-expressing Neuro2A cells [156].

There are no clinical safety or efficacy data for DPA, and its simple structure will likely hinder potential clinical studies due to concerns about its specificity. DPA, however, is particularly selective for the σ 1R, over the σ 2R, monoamine transporters, DA receptors and muscarinic AchR [2].

2.7 Donepezil

Donepezil (Aricept) (Fig. 2) is one of the current standard acetylcholinesterase inhibitors (AChEI) that are prescribed for mild-to-moderate AD, along with galantamine (Reminyl), rivastigmine (Exelon) and memantine (Ebixa), according to the online guidance from the National Institute for Health and Care Excellence (NICE) in the UK [276]. Donepezil appears to provide improved cognition in mild-to-moderate AD, although it does not halt the progression of the disease. Donepezil has an affinity for the σ 1R (K_i 14.6 nM) [277], and studies using rats have shown that there is 90% occupancy of σ 1Rs in the brain when dosed with 10 mg/kg [278].

2.7.1 Alzheimer's Disease

Donepezil has been associated with reduced cardiovascular mortality in patients with AD [279, 280], likely due to cardioprotective effects of σ 1R ligands [281]. Studies investigating donepezil's σ 1R targeting activities have shown that it can potentiate nerve growth factor (NGF)-induced neurite outgrowth in PC12 cells [282] (a common test for σ 1R

agonist activity). Furthermore, positron emission tomography (PET) scans in mice have shown that donepezil can displace ^{11}C -SA4503 from the brain dose dependently at physiologically relevant doses [278].

Donepezil has been thoroughly tested clinically in moderate-to-severe AD, with many studies (discussed in multiple meta-analyses and systematic reviews [283–285]) indicating a statistically significant improvement in the standardized Mini-Mental State Examination (SMMSE, on which scores range from 0 to 30, with higher scores indicating better cognitive function) and the Bristol Activities of Daily Living Scale (BADLS, on which scores range from 0 to 60, with higher scores indicating greater impairment) compared with baseline [286]. In most clinical studies of donepezil, patients treated with donepezil maintained or improved cognition and overall dementia severity in contrast to placebo-treated patients who showed a decline in cognition [287].

The accumulation of A β in the brain is part of the pathology of AD, and in animal studies, donepezil reduces brain A β accumulation [288, 289]. However, using ^{11}C -Pittsburgh compound B, a clinical study of AD patients receiving donepezil (5 or 10 mg) for at least 1 year showed that there was no reduction in A β accumulation at either dose [290]. Another symptom of AD, other than memory loss, is agitation [245]. Despite appearing to be effective in the cognition portions of clinical trials, donepezil does not appear to be effective at managing agitation in AD patients [291].

2.8 Eliprodil

Eliprodil (Fig. 2) is an NMDA antagonist that selectively inhibits the NR2B (GLUN2B) and is a σ 1R agonist. Eliprodil failed a phase III clinical trial to treat acute ischaemic stroke [292], and has also been trialled for schizophrenia [293]. Eliprodil has a moderate affinity for the σ 1R ($K_i = 630$ nM) [294].

2.8.1 Parkinson's Disease

In a pre-clinical study of eliprodil, a low dose of eliprodil stimulated locomotion in mouse PD models, whereas a high dose had more sedative action. In the same study of reserpine-treated mice, eliprodil reversed akinesia, although this was not statistically significant. Furthermore, eliprodil failed to improve motor recovery from the DA D1 agonist SKF38393, or the DA D2 agonist RU24213, and inhibited the motor stimulation induced by L-DOPA [295]. Eliprodil has been clinically trialled for movement disorders, including PD (NCT00001929), although the data were never published. Based on this limited evidence, eliprodil is unlikely to be an effective PD treatment.

2.8.2 Multiple Sclerosis

Eliprodil has been reported to exhibit a neuroprotectant effect against toxic insults. A study by Demerens et al. [296] has indicated the ability of eliprodil to induce myelination in cellular models, suggesting that the molecule can be of therapeutic interest for the treatment of MS.

2.9 Fabomotizole

Fabomotizole (Afobazole) (Fig. 2) was launched in Russia in the early 2000s as an anxiolytic drug. As well as acting on the melatonin receptor 1 and 3 (MT1 and MT3), Afobazole has GABAergic [297], NGF and BDNF-releasing properties [298]. Afobazole is a mixed σ 1/2R ligand [299].

2.9.1 Alzheimer's Disease

Afobazole has been shown to inhibit A β_{25-35} -induced calcium increase and nitric oxide production in both rat cortical neurons and microglial cultures [300, 301]. Afobazole shows some promise as a σ 1/2R ligand; however, its use in AD is still in the pre-clinical stage, and there appear to be no current or planned clinical trials.

2.10 Fenfluramine

Fenfluramine (Fintepla) (Fig. 2) is a positive modulator of the σ 1R [89], with an affinity of 266 nM (K_i guinea pig brain) [302]. It also acts to cause the release of serotonin by disrupting the vesicular storage of the neurotransmitter and reversing serotonin transporter function.

2.10.1 Developmental and Epileptic Encephalopathies

The serotonin activity of fenfluramine fails to explain its efficacy in Dravet syndrome (a genetic epilepsy characterized by temperature-sensitive/febrile seizures). Furthermore, fenfluramine could potentiate the BiP dissociating effects of σ 1R agonist PRE084 (although it did not affect BiP dissociation by itself), indicating that fenfluramine is a positive modulator of the σ 1R rather than a σ 1R agonist. However, while the positive modulator SOMCL-668 reduced epileptic activity in zebrafish, the σ 1R agonist PRE084 inhibited fenfluramine-mediated anti-epileptic activity in zebrafish models [303]. This may be explained by the biphasic nature of many σ 1R ligands in vivo [304].

In a clinical trial of fenfluramine (NCT03355209) in patients with Lennox–Gastaut syndrome (another developmental and epileptic encephalopathy), fenfluramine, compared with placebo, decreased drop seizures and may be particularly good for patients who experience generalized tonic–clonic seizures [304]. In another pair of fenfluramine

studies in Dravet syndrome, where the two identical studies were merged (NCT02682927 and NCT02826863), fenfluramine provided a significantly greater reduction in convulsive seizure frequency compared with placebo [305]. In total, 119 patients, with a minimum number of convulsive seizures per 4-week period for 12 weeks before screening, were randomized to receive placebo, fenfluramine 0.2 mg/kg/day or fenfluramine 0.7 mg/kg/day. The study met its primary efficacy endpoint, with 0.7 mg/kg/day showing a 62% greater reduction in mean monthly convulsive seizure

frequency than placebo. Furthermore, the low dosage of fenfluramine (0.2 mg/kg/day) showed a 32% reduction in mean MCSF compared with placebo [305].

Another study of fenfluramine provided a clinically meaningful reduction in convulsive seizure frequency in most patients in a real-world study (71.1% of patients had a greater than 50% reduction of convulsive seizures, 53.3% had a more than 75% reduction and 11.1% were seizure free) [306]. Several other studies are ongoing, and studies with data yet to be published are detailed in Table 3.

Table 3 Ongoing clinical trials of fenfluramine

Study title	Conditions	Primary outcome measures	Study results	Phase	Trial number
Fenfluramine for adult Dravet patients	Dravet syndrome Dravet syndrome, intractable	Seizure reduction Comparing the proportion of participants with >50% reduction in the monthly countable convulsive seizure frequency between 4 week baseline period (observational phase) and after 12 week treatment phase using seizure diary	No results posted (last update 2022)	3	NCT05560282
Treatment of Dravet syndrome with fenfluramine (expanded access protocol)	Dravet syndrome	Evaluate the safety of fenfluramine	No results posted (last update 2020)	Unknown	NCT04437004
A study to investigate the long-term safety of ZX008 (fenfluramine hydrochloride) oral solution in children and adults with epileptic encephalopathy including Dravet syndrome and Lennox–Gastaut syndrome	Dravet syndrome Lennox–Gastaut syndrome Epileptic encephalopathy	Evaluate the safety of fenfluramine Change in convulsive seizure response Change in cognitive CGI by parent/caregiver and investigator Change in behavioral CGI by parent/caregiver and investigator Change in motor CGI by parent/caregiver and investigator Change in clinical global impression by parent/caregiver and investigator	No results posted (last update 2023)	3	NCT03936777
Fenfluramine for the treatment of different types of developmental and epileptic encephalopathies: a pilot trial exploring epileptic and non-epileptic outcomes (FENDEEP)	Refractory epilepsy SYNGAP1 encephalopathy STXBP1 encephalopathy with epilepsy Inv Dup(15) Encephalopathy Multifocal or bilateral malformations of cortical development Continuous spike and waves during slow sleep	Seizure frequency, measured using a seizure diary	No results posted (last update 2022)	4	NCT05232630

CGI clinical global impression

2.11 Fluoxetine

Fluoxetine (Fig. 2) is an SSRI, broadly administered as an antidepressant. Besides its use as an antidepressant, fluoxetine has been suggested for many off-label uses, such as cancer treatment [307] and COVID-19 [308]. Fluoxetine can regulate the inflammatory response by reducing tumour necrosis factor (TNF)- α ; interleukin (IL)-1 β , -2 and -6; and interferon (INF)- γ , while increasing IL-2, -4 and -10 [309]. Fluoxetine has an affinity for the σ 1R of 214–240 nM [310].

2.11.1 Multiple Sclerosis

In a pilot study with relapsing MS patients, fluoxetine reduced the formation of new-enhancing inflammatory lesions in the brain [311]. Fluoxetine attenuated microglial activation and protected oligodendrocytes from apoptosis, preventing myelin loss through the inhibition of c-Jun [312]. Oral administration of fluoxetine (NCT01910259) in MS patients was found to reduce axonal degeneration as measured by proton magnetic resonance spectroscopy [309]. However, in secondary progressive MS individuals, fluoxetine treatment did not exhibit any significant changes in brain atrophy [313–317].

2.12 Fluvoxamine

The SSRI fluvoxamine is a strong σ 1R agonist and has been proven to have strong efficacy against depression. Recent studies indicate it may help treat COVID-19 [318]. It could potentiate neurite outgrowth in PC12 cells in a σ 1R-dependent mechanism, indicating its neuroprotective potential [319]. In an open-label study, the administration of fluvoxamine exhibited antidepressant activity in MS individuals [320].

2.12.1 Multiple Sclerosis

Fluvoxamine increased the proliferation of neural stem cells and facilitated their differentiation into oligodendrocytes through the Notch signalling pathway. Further, in an experimental autoimmune encephalitis animal model, fluvoxamine treatment reduced the severity of the condition by inhibiting IFN- γ , and reducing inflammatory infiltration and demyelination, suggesting therapeutic potential against MS [321].

2.13 Ketamine

Ketamine (Fig. 2) is an NMDA receptor antagonist acting to block the open ion channel similarly to memantine. Ketamine also has a moderate affinity for the σ 1R (K_i = 139.60 μ M). It can potentiate PC12 cell neurite outgrowth in response to NGF (a common assay of σ 1R agonism) in

a bell-shaped dose-response curve (also a common feature among σ 1R agonists), with the optimal concentration being 1 nM [322]. Furthermore, NE100 (a potent σ 1R antagonist) could inhibit the neurite outgrowth potentiation. As we have discussed above, developing new neuron connections in the hippocampus is important in the progression and treatment of AD. Treatment of mice with low-dose ketamine (8 mg/kg) reduced behavioural abnormalities and improved cognitive function in mouse models of Rett syndrome (a neurodegenerative disease discussed further below). Furthermore, 10 mg/kg ketamine improved cognitive flexibility in rats subjected to repeated restraint stress [323].

2.13.1 Alzheimer's Disease

Ketamine has been shown to prevent lipopolysaccharide (LPS)-induced TNF- α production in glial cells, astrocyte cultures and microglial cultures [324], suggesting that ketamine can prevent neuroinflammation.

A clinical trial (NCT02433041) investigating ketamine to prevent post-operative delirium found, as part of the secondary outcomes, a significant decrease in inflammatory markers with the use of ketamine post-operatively when compared with the placebo [325]. This suggests ketamine has neuroprotective action on neurons, glial cells and astrocytes.

Several studies have shown that low doses of ketamine helped improve visual memory, simple working memory and complex working memory in patients with treatment-resistant depression [326, 327]. Depression is a common symptom of AD; furthermore, depression is linked to AD development [328]. Ketamine relieves depressive symptoms within hours of treatment, especially in cases where other antidepressants have failed [329]. In a case report of a 76-year-old female AD patient suffering from depression who had not responded to conventional antidepressant therapy, the patient was reported to have recovered from severe treatment-resistant depressive decline after only two doses of ketamine [330].

Recently, a meta-analysis review of 25 randomized controlled trials that included 14 commonly prescribed antidepressants failed to show efficacy in AD patients [331]. Thus, with ketamine's potential for neuroprotection and its efficacy in treating depression, it is possible that ketamine could be a useful intervention in treating depression in AD. However, studies of prolonged treatment (3 months) with ketamine in mice and monkeys show an increase in tau hyperphosphorylation, a key part of AD pathology, before the formation of neurofibril tangles (NFTs). The study did not show A β plaque formation, another pathological hallmark of AD [332].

There are several safety issues concerning ketamine use over long-term treatment (discussed below regarding PD, although many concerns also apply to AD). Therefore,

further study of the efficacy and safety of ketamine at various doses and over short- and long-term treatment is required before ketamine can be seriously considered as a treatment for AD.

2.13.2 Parkinson's Disease

Ketamine in combination with xylazine (α_2 adrenergic agonist) has been studied pre-clinically in rat models of PD (MPTP and 6-OHDA). The ketamine/xylazine (85/3 mg/kg) combination given before the MPTP or 6-OHDA significantly reduced the loss of tyrosine hydroxylase immunostained cells. Furthermore, ketamine/xylazine treatment reversed the amnesic effects of the two neurotoxins, with ketamine/xylazine-treated rats performing the same as control rats in the hidden platform test (MPTP- or 6-OHDA-treated rats showed a 10-second latency in finding the hidden platform) [333]. Another study of rats subjected to 6-OHDA showed reduced mobility time in the forced swim test (suggesting depressive-like behaviour). Rats treated with ketamine for 28 days (starting 21 days after the brain lesions were created with 6-OHDA) increased mobility times in the water at all doses of ketamine. Furthermore, ketamine also reduced symptoms of anhedonia (caused by the 6-OHDA lesions) measured using the sucrose preference test. The same study also investigated short-term memory in the rats using the social recognition test, and ketamine significantly (at all doses 5–15 mg/kg) reversed the amnesic effects of 6-OHDA treatment [334]. In a mouse model of PD using MPTP to induce PD-like symptoms in the mice, ketamine (8 mg/kg) improved the coordinating movement, learning and memory ability of PD model mice.

Furthermore, ketamine also prevented tyrosine hydroxylase-positive neurons from being attacked by MPTP, therefore, protecting nigra dopaminergic neurons. MPTP treatment of mice significantly increased mTOR expression and lowered the expression of several autophagy markers (LC3-II, Beclin1, Parkin and PINK). Ketamine restored expression levels of LC3-II, Beclin1, Parkin and PINK1 to levels of the untreated controls, and reduced the expression of mTOR. This indicates that the subanaesthetic dose of ketamine can restore autophagy in the PD mouse model [335].

In a case report of an 88-year-old female patient with PD exhibiting severe bilateral dyskinesia who required surgery, ketamine [20 mg intravenous (IV)] followed a dose of midazolam (1 mg IV) (to prevent the possible psychotomimetic effects of ketamine). It resolved her severe tremor both at rest and during intentional movement. Following the ketamine, the patient remained responsive, communicative and able to follow commands. Several days later, during post-operative care, she remembered receiving the pre-operative ketamine with no adverse consequences [336]. Two previous cases have used ketamine to permit surgery in cases of PD,

both of which were completed with this drug combination resulting in no major side effects [337]. Ketamine has now entered phase 2 clinical trials for treating depression in PD (NCT04944017) and treating levodopa-induced dyskinesia (NCT04912115). Both studies are still ongoing.

There are several possible drawbacks to the use of ketamine in the treatment of depression and levodopa-induced dyskinesia in PD. In treating depression (in general), ketamine IV injection relieves depression symptoms for 1–2 weeks, with some patients reporting only a few days. One study of eight patients where ketamine (0.5 mg/kg IV) was administered up to every two weeks showed promising results [338]. Furthermore, ketamine dosing requires injections, as it has poor bioavailability and a high first-pass hepatic metabolism, thus requiring hospital/clinical visits. Trips to the hospital/clinic every 3–7 days for an IV infusion of ketamine are not practical. Ketamine also has psychotomimetic effects and dissociative symptoms (although these tend to dissipate within 4 hours of treatment) [339] and has the potential for addiction [340]. There are also concerns about the long-term safety of ketamine treatment as transient tachycardia, hypertension, neurotoxicity, bladder toxicity and tolerance are reported with repeated ketamine infusion [341].

2.13.3 Multiple Sclerosis

In the experimental autoimmune encephalitis mice model, (R)-ketamine reduced microglial activation, preserved blood–brain barrier (BBB) integrity and ameliorated the demyelination [342]. Further, under cuprizone-induced demyelination in mice, (R)-ketamine facilitated remyelination through activating TrkB and altered the abnormal amount of *Eisenbergiella massiliensis* in the gut microbiota, which is positively correlated with demyelination [343]. Oral administration of ketamine in a 60-year-old woman with MS reduced neuropathic pain (allodynia) associated with the disease [344]. In a patient with treatment-resistant depression associated with MS, IV ketamine administration significantly reduced the depression index score with no worsening of her condition over a two-year observation period [345]. Likewise, ketamine also improved fatigue, which is associated with MS, in an active placebo-controlled double-blind trial (NCT03500289), suggesting its beneficial effects. Many of the drawbacks mentioned above regarding ketamine and PD also apply here.

2.13.4 Rett Syndrome

In a pre-clinical study, daily treatment with ketamine reduced Rett syndrome-like symptoms and extended the life span of MeCP2-null mice without adverse side effects. Furthermore, cortical processing and connectivity were

fully restored to wild-type levels [346]. Functional deficits in forebrain circuits in mutant MeCP2 mice can be restored by ketamine treatment [347].

Two clinical trials (phase 1 and phase 2) are registered at www.clinicaltrials.gov (NCT02562820 and NCT03633058). The phase 1 safety study was terminated due to funding withdrawal, and the phase 2 study is listed as complete; however, there are no listed results or peer-reviewed publications yet. As well as the drawbacks of ketamine described above, ketamine may also provoke epileptic discharges in some patients with a seizure disorder [348].

2.14 Memantine

Memantine (Ebixa, Axura, Namenda, Akatinol) (Fig. 2) is a low–moderate affinity NMDA receptor antagonist often prescribed for AD patients who do not tolerate AChEI drugs such as donepezil [349]. Memantine, in combination with donepezil, is approved as a treatment for AD in many countries [350]. Memantine is prescribed for moderate-to-severe AD. It acts as an uncompetitive channel blocker of the NMDA receptor, i.e. memantine does not interact with the binding sites of the NMDA agonists, rather it fits into the channel, competing with magnesium ions, blocking ion flow through the channel into and out of the cell. It is somewhat surprising that an antagonist of the glutamate receptor can benefit AD symptoms since AD sees a reduction in glutamate and glutamate receptors [351].

2.14.1 Alzheimer's Disease

It has been hypothesized that several mechanisms mediate the beneficial effects of memantine in AD, including decreasing synaptic 'noise' from excessive NMDAR activation and reducing the excitotoxicity [351, 352] or inhibiting A β production or toxicity [353]. It is hypothesized that memantine's fast on/off pharmacokinetics allow for the physiological activation of NMDARs, while at the same time inhibiting pathological overactivation because of its mechanism of NMDAR inhibition.

The σ 1R has been shown to play a role in regulating the NMDA receptor and has been well documented [41, 354, 355]. For example, σ 1R KO impairs neurogenesis in the hippocampus of adult mice via the downregulation of the NMDA receptor [356]. Furthermore, activation of the σ 1R results in the upregulation and transport to the plasma membrane of NMDA receptors [357].

Memantine, like other NMDA ion-channel blockers (amantadine [5] and ketamine [358], among others), has some affinity for the σ 1R. However, the affinity of memantine is rather low (2.5 μ M) [200], and thus the σ 1R is unlikely to play a full role in memantine's therapeutic effects. However, since activating the σ 1R receptor

modulates NMDA function and upregulates NMDA expression and trafficking to the plasma membrane, it is possible that the combination of drugs such as memantine and a σ 1R agonist could be beneficial in AD. Monotherapy with memantine has been clinically trialled in multiple studies and is generally safe and well tolerated, plus it has beneficial effects compared with a placebo. A meta-analysis of six placebo-controlled phase 3 trials supports the clinically relevant efficacy of memantine in patients with moderate-to-severe AD [359]. The combination therapy of memantine and donepezil has been trialled in several studies with mixed results (NCT00866060, NCT02580305, NCT03316898 and NCT02162251). Multiple studies have identified significant improvements in preventing cognitive decline when memantine is added to stabilize the treatment of patients with stable donepezil treatment for moderate-to-severe AD [360–364]. However, other studies have been less convincing, with the data suggesting only a mild to no benefit to the combination therapy over monotherapy with either donepezil or memantine [365].

2.14.2 Parkinson's Disease

Memantine is not used in treating the motor symptoms of PD. However, it has been trialled for treating PD-related dementia [366, 369]. Memantine is relatively safe and well tolerated in PD-related dementia treatment, and there appear to be some studies with positive outcomes [367].

However, a meta-analysis study of ten trials indicates that memantine does not improve scores on the MMSE, whereas cholinesterase inhibitors did [370]. The study did find that there was a reported improvement in global impression but not cognitive function. The clinical trials are listed in Table 4.

2.15 Naluzotan

Naluzotan (Fig. 2) acts as a selective and potent 5-HT1A receptor partial agonist, and it has been claimed that it binds to and activates the σ 1R, although the evidence for this is weak. Naluzotan has a claimed affinity of 100 nM for the σ 1R; however, few references to the σ 1R of naluzotan can be found, with no binding data to back up the claims [371]. It has been trialled for depression and anxiety, with the study suggesting that naluzotan is relatively safe and higher doses of naluzotan may be effective in anxiety and depression [372].

2.15.1 Developmental and Epileptic Encephalopathies

Naluzotan has been tested for its effects in localization-related epilepsy (NCT01281956) [373], with the data suggesting that there is no significant difference between

Table 4 Clinical trials of memantine for Parkinson's disease and dementia-related symptoms

Study title	Primary outcome measures	Study results	Phase	Trial Number
Namenda (memantine) for non-motor symptoms in Parkinson's disease	UPDRS Analyses will be computed for the categorical DV: Global Tremor Assessment by examiner	Memantine was well tolerated in Parkinson's disease with few adverse events. However, specific measures of sleepiness, fatigue, depression and attention did not significantly improve [367]	4	NCT00646204
Parkinson's disease evaluated by PET and the effect of memantine	Decreases in cerebral blood flow and oxygen metabolism in these areas will be the consequence and this can be detected by PET	No results posted (last update 2006)	N/A	NCT00375778
Study of memantine for gait disorders and attention deficit in Parkinson's disease	Stride length by gait analysis with an optoelectronic system	In the memantine treated group, statistically significantly better results (versus placebo) were observed for the overall UPDRS score [368]	4	NCT01108029
Memantine versus placebo in Parkinson's disease dementia or DLB	Assess the global benefits of memantine compared with placebo in outpatients with a diagnosis of PDD or DLB (mild to moderate severity) over a 6 month period	Patients with DLB treated with memantine showed greater improvement according to Alzheimer's disease cooperative study (ADCS)-clinical global impression of change scores than those who received placebo. No significant differences were noted between the two treatments in patients with PDD [366]	4	NCT00855686
Memantine for treatment of cognitive impairment in patients with Parkinson's disease and dementia	DRS Memory Subscore CIBIC-Plus Score	DRS scores appeared to improve with a score of 1.6 (95% CI -0.13 to 3.33) in memantine treated compared with -1.4 (95% CI -3.19 to 0.39) in placebo treated. There was also an apparent improvement in the CIBIC-Plus Score in the memantine treated group ^e	N/A	NCT00294554
Efficacy and safety of memantine for PDD and dementia with Lewy bodies (DLB) (MEMPDD)	Clinical Global Impression of Change	No results posted (last update 2015)	2	NCT00630500
Monitoring anti-dementia drugs by serum levels (MONANTI)	Change of MMSE Change of ACE Change of NPI-Q Change of DAD Change of clinical serum level of memantine Change in GDS	No results posted (last update 2020)	4	NCT04117178

ACE Adenbrooke's Cognitive Exam, DV dependent variable, DAD Disability Assessment for Dementia, DRS Dementia Rating Scale, DLB dementia with Lewy bodies, GDS Geriatric Depression Scale, MMSE Mini Mental State Examination, NPI-Q Neuropsychiatric Inventory Questionnaire, PDD Parkinson's disease dementia, PET positron emission tomography, UPDRS Unified Parkinson Disease Rating Scale

placebo and naluzotan; furthermore, the number of seizures in the naluzotan group appeared to be higher than that of the placebo and the trial was terminated.

Given the lack of evidence for the role of the σ 1R in the pharmacology of naluzotan, and that it had no effect of seizure frequency when trialled, naluzotan is unlikely to benefit DEE patients.

2.16 Phenytoin

Phenytoin (Dilantin, amongst others) (Fig. 2) has long been used clinically for seizures [374]. The primary mechanism for phenytoin is thought to be the inhibition of voltage-gated sodium channels [375]. Phenytoin is a non-selective allosteric modulator of the σ 1R. There is much evidence that the σ 1R interacts with and modulates the function of various ion channels, including calcium channels, with many studies indicating that the σ 1R acts to negatively regulate calcium movement [376–379].

Phenytoin is a positive modulator of the σ 1R as it increases σ 1R agonist binding to the receptor when co-administered with a σ 1R agonist. Furthermore, it can be used to distinguish between σ 1R agonists and antagonists as phenytoin does not increase antagonist binding [380].

2.16.1 Developmental and Epileptic Encephalopathies

Many clinical studies have investigated phenytoin (a total of 123 listed on www.clinicaltrials.gov, of which 36 involve seizures). However, phenytoin is considered a last resort for treating encephalopathy since the severe side effects of phenytoin maintenance therapy, especially cognitive impairment, are undesirable in these already cognitively impaired patients [381]. There has been one case report that suggests phenytoin treatment in a two-year-old male with epileptic encephalopathy that did not improve with phenobarbital, pyridoxine, pyridoxal phosphate and levetiracetam showed dramatic response to an IV loading dose of phenytoin, followed by a transfer over to carbamazepine to prevent seizure relapses because of the difficulty in maintaining appropriate blood levels of phenytoin [382].

2.17 PRE-084

The selective σ 1R agonist, 2-(4-morpholino ethyl)1 phenylcyclohexanecarboxylate (PRE-084) (Fig. 2) contains the three pharmacophoric moieties for a σ 1R ligand: a hydrophobic cluster, an amine group and an intermediate chain [383], and has an affinity of 2.2 nM. PRE-084 has been a very useful pharmacological tool in the discovery of other σ 1R ligands [384]. Although thought to be

specific to the σ 1R, a study has shown that it can have some off-target effects. PRE-084 could elicit an enhanced barrier function of human umbilical vein endothelial cell (HUVEC) monolayers in a dose-dependent fashion, and even with significant siRNA knockdown of the σ 1R, PRE-084 maintained the barrier function effect, meaning that either only a low occupancy of σ 1R is required for the barrier function effect, or that another receptor is involved [385].

2.17.1 Alzheimer's Disease

PRE-084 has many characteristics that could make it a potential treatment for AD. Pre-clinical data have shown that PRE-084 can attenuate the blood–brain barrier (BBB) disruption caused by A β . The study showed that PRE-084 protected the integrity of the BBB by preventing the deposition of A β (insoluble) A β _{1–40} and A β _{1–42} levels in the brains of mice, and increasing vascular endothelial growth factor (VEGF) and ApoE receptor [386]. PRE-084 also increases the expression of glial cell line-derived neurotrophic factor (GDNF) [387] and BDNF [388].

The hippocampal region of the brain is responsible for learning and memory and is one of the first regions to show damage in AD [389]. Neurogenesis occurs in the adult hippocampus, although with age, this begins to slow [390]. A decreased rate of neurogenesis, increased destruction of mature neurons, and increased inflammation could contribute to the cognitive symptoms seen in AD. It is thought that this damage in the hippocampus can occur up to 10 years before cognitive symptoms appear in the patient. Studies using A β _{1–42} [391] or A β _{25–35} [392] injections into mouse brains have shown reduced proliferation and neurite growth, increased death of newly formed cells and an increase in the number of immature cells in mice brains. PRE-084 treatment promotes neurogenesis in mice treated with A β _{1–42} or A β _{25–35} [391, 392]. Furthermore, the density of hyper-reactive astrocytes was reduced by the administration of PRE-084 [391], suggesting PRE-084 may play an anti-inflammatory role. PRE-084 could also increase mushroom dendritic spine formation in the hippocampus of transgenic rats expressing presenilin-1 (PS1) and human APP^(Swe) [393].

There has been no clinical study registered or completed with PRE-084 for AD or any other neurodegenerative disease, and there are no data on the safety or pharmacokinetic profile for PRE-084 in human trials. PRE-084 is a well-known selective σ 1R agonist [380] and has been used over the past 30 years as an important prototypical drug to uncover potential therapeutic benefits of the σ 1R. However, it has yet to be investigated clinically, possibly

due to its potential off-target effects, which are thought to increase with dose [384].

2.17.2 Amyotrophic Lateral Sclerosis/Motor Neuron Disease

PRE-084 protected spinal cord organotypic culture explants from glutamate toxicity and induced the elongation of neurites in motor neurons through the activation of PKC [394]. Administration of PRE-084 in adult rats prevented the loss of axotomized motoneurons upon spinal root avulsion injury with decreased astroglial immunoreactivity, alongside increased BiP and GDNF expression in motor neurons and astrocytes, respectively [387]. In SOD1^{G93A} mice, PRE-084 treatment improved locomotor performance, preserved motor neurons, extended survival of animals and modulated NMDA receptor calcium influx through the PKC pathway [395–397]. In wobbler mice, chronic treatment of PRE-084 increased BDNF expression, reduced reactive astrocytes and induced the survival of motor neurons indicating σ 1R and their agonists can also be used for treating non-SOD1-linked ALS [398]. Nrf-2 co-localization with σ 1R has been reported in photoreceptor cells, implying Nrf-2 could be directly targeted by σ 1R [399]. However, possibly due to a difference in the study design, there are also data suggesting that NRF-2 does not interact with the Nrf-2 [400]. PRE-084 treatment rescued motor neuron defects in TDP43G348C zebrafish by enhancing ATP generation, boosting the unfolded protein response (UPR) and activating the Nrf-2 pathway [401].

2.17.3 Huntington's Disease

PRE-084 has been tested pre-clinically as a therapeutic agent for HD. A study in 2013 [402] showed that PRE-084 promotes cell survival and counteracts the detrimental effects of mutant N-terminal huntingtin protein in neuronal PC6.3 cells. PRE-084 increased cellular antioxidants through the NF- κ B pathway (a pathway normally compromised by the mutant huntingtin proteins), supporting the future use of σ 1R agonists as therapeutics for HD.

Treatment of transgenic cortical cultures from mice expressing 128 CAG repeat expansion in the *Htt* gene (YAC128 transgenic mice) with PRE-084 (100 nM) restored synaptic connections between the cortical and striatal neurons but did not affect their number in the cultures when compared with wild-type mice; furthermore, this was reversed with the σ 1R antagonist NE-100 [403].

2.17.4 Wolfram Syndrome

PRE-084 has been shown in a pre-clinical study to attenuate learning deficits in male *Wfs1* ^{Δ Exon8} mice (a transgenic

WS model) and reduce anxiety in female *Wfs1* ^{Δ Exon8} mice [197]. In the same study, *wfs1ab*^{KO} (zebrafish larvae with the *Wfs1* gene knocked out) showed a clear locomotor alteration when their response to light variation was tested. PRE-084 treatment (0.1–10 μ M) in the fish water resulted in a dose-dependent decrease in the hyperlocomotion response, back down to the level of the wild-type larvae (*Wfs1ab*^{WT}) [197]. Furthermore, these effects were reversed by the σ 1R antagonist NE-100, suggesting that the σ 1R is responsible for these beneficial effects.

2.18 Pridopidine

Pridopidine (Fig. 2), developed by Prilenia Therapeutics, is a drug currently in the late stages of clinical trials for PD, ALS and HD [52]. Pridopidine is a selective σ 1R ligand with low affinity for DA D2/D3 receptors, the adrenergic α 2C receptor and the σ 2R [404], and has the potential as an AD therapy. At clinically relevant doses, positron emission tomography (PET) imaging studies in rats and humans show high occupancy of the σ 1R, with no or negligible occupancy of the D₂/D₃ receptors [405, 406].

2.18.1 Alzheimer's Disease

AD cognitive impairment results from damage to synaptic connections and neuronal atrophy. The plasticity of dendritic spine structures in developing neurons underlies memory formation, as thin spines convert into mushroom spines during information storage in response to long-term potentiation (LTPs) [407–409]; mushroom spine deficiency may reflect learning and memory defects in models of AD. Pre-clinical studies have shown that pridopidine stabilizes mushroom spines in mouse models of AD. In PS1 knock-in (PS1-KI) mouse hippocampal cultures and wild-type cultures treated with A β _{1–42}, the percentage of mushroom spines was significantly reduced. This reduction could be prevented by 100 nM pridopidine. Furthermore, this was repeated in vivo in PS1-KI mice [63].

Pridopidine protects against NMDA-induced oxidative stress in primary hippocampal cultures and can increase mushroom dendritic spine formation in the hippocampus of transgenic rats expressing presenilin-1 (PS1) and human APP^(Swe), as well as reduce deficits in spatial learning [393]. The study also showed that pridopidine induced ERK and AKT phosphorylation. Thus, the involvement of the mitogen-activated protein kinase/extracellular signal-regulated kinases (MAPK/ERK) and phosphoinositide 3-kinase/protein kinase B/mammalian target of rapamycin (PI3K/Akt/mTOR) signalling pathways, through which the σ 1R may control the neuroprotective properties and synaptogenesis [393].

Pridopidine has yet to be studied clinically for AD; however, given the success of pridoipidine in treating other neurodegenerative diseases (discussed further below) and its pre-clinical data in AD, it is possible that pridoipidine could show some benefit in patients with AD.

2.18.2 Amyotrophic Lateral Sclerosis

Pridopidine targets $\sigma 1R$ in SOD1^{G93A} ALS models (both in vitro and in vivo) and ameliorates the pathology by maintaining neuromuscular junction integrity and enhancing axonal transport through the activation of ERK pathway [410]. Also, cachexia and motor neuron deficits were prevented in SOD1G93A mice upon pridoipidine administration [411]. One of the important steps during autophagy is the transport of transcription factor EB (TFEB) into the nucleus through POM121/KPNB1/importin β 1. The occurrence of hexanucleotide RNA expansion (HRE) because of defective *C9orf72* gene interferes with the $\sigma 1R$ and POM121 interaction, causing reduced TFEB transport and autophagy. In NSC34-expressing (G4C2)₃₁ RNA repeats potentiated $\sigma 1R$ chaperone activity, and enhanced POM121 and autophagy [412]. Under the HEALEY ALS Platform Trial umbrella, two ongoing phase 2/3 clinical trials are active for ALS patients to study pridoipidine (as well as other drugs, such as zilucoplan, verdiperstat, CNM-Au8, SLS-005 Trehalose and ABBV-CLS-7262) on disease progression, respiratory function, muscle strength, bulbar function and survival (NCT04615923 and NCT04297683).

2.18.3 Huntington's Disease

Treatment of YAC128 HD mice with pridoipidine reduces anxiety and depression-like symptoms [413], prevents striatal medium spiny neuron (MSN) loss and facilitates Ca²⁺ homeostasis and signalling at the ER via activation of the $\sigma 1R$ [97]. In both animal and human studies of HD, the BDNF pathway has been shown to be affected [414]. Under non-disease conditions, the Htt protein is involved in the transport of BDNF. However, the mutated version of Htt protein reduces BDNF release [415]. Furthermore, BDNF overexpression can rescue HD phenotypes in YAC128 mice [416]. This is particularly relevant to the $\sigma 1R$, as $\sigma 1R$ activation has been shown in multiple studies to induce BDNF expression and activity [188, 416–418].

Pridopidine has undergone randomized, double-blind clinical trials for HD treatment [419, 420]. Pridopidine treatment was ineffective at 20 mg/day and 45 mg/day. However, the higher 90 mg/day improved movement ability measured by the total motor scores (TMS) [419].

Higher doses of pridoipidine or longer treatment periods could show further improvements. However, another larger

study failed to replicate the improvements in TMS. However, some improvements were seen at 26 and 52 weeks in an exploratory assessment of rater-independent Q-Motor and total functional capacity (TFC) using a dose of 45 mg BID (90 mg/day) [421].

Furthermore, an open label study of pridoipidine covering 60 months showed that 45 mg BID (90 mg/day) can maintain stable TFC and TMS, when using historical placebo controls as a comparison [422]. Pridopidine, although not a cure-all drug, does show some promising improvements in some aspects of HD and is tolerable orally at doses up to 112.5 mg BID (225 mg/day), therefore, further investigation is warranted into its benefits for HD treatment.

2.18.4 Parkinson's Disease

Pridopidine has been investigated pre-clinically for its use in PD. In a mouse model of PD where 6-OHDA was used to create lesions that resulted in several PD-like symptoms in the mice, pridoipidine significantly improved deficits in forelimb use, offered significant protection of nigral DA cell bodies and increased dopaminergic fibre density in the striatum. In $\sigma 1R$ KO mice, pridoipidine did not protect against 6-OHDA-induced lesions. The study also showed the upregulation of GDNF, BDNF and phosphorylated ERK1/2 in the striatal region of the brain [226]. In a similar rat model, pridoipidine decreased L-DOPA-induced sensitized locomotor activity while permitting a beneficial motor-stimulant effect of the L-DOPA [423]. In primates [macaques treated with the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)], Pridopidine produced a dose-dependent reduction in dyskinesia caused by L-DOPA without modulating the anti-parkinsonian benefit of L-DOPA [302].

Dyskinesia is involuntary hyperkinetic movements, and the risk of developing dyskinesia is approximately 25–40% after 4–6 years of levodopa therapy [424]. It is thought that L-DOPA does not cause the development of dyskinesia in PD. Rather it sensitizes the nigrostriatal system, since multiple clinical studies have shown that delaying L-DOPA treatment does not prolong the time before dyskinesia onset [425]. In a clinical trial of pridoipidine in PD patients with dyskinesia (NCT03922711), pridoipidine appeared to reduce Unified Dyskinesia Rating Scale (UDysRS) scores compared with the placebo after 5 weeks (the trial was terminated due to the COVID-19 outbreak). At time of writing, this is the only trial of pridoipidine concerning PD, and, unfortunately, this trial was stopped early due to COVID, as the results looked promising.

2.19 Rivastigmine

Rivastigmine (Fig. 2) is a therapeutic agent that acts as an AChEI and is used in treating mild-to-moderate AD [426]. Rivastigmine is also thought to be active at the σ 2R [29].

2.19.1 Alzheimer's Disease

Rivastigmine enhances NGF-induced neurite outgrowth in PC12 cells via σ 1R and σ 2R since antagonists of each were both required to block the NGF-induced neurite outgrowth fully. Furthermore, σ 1R and σ 2R knockdown also abolished NGF-induced neurite outgrowth in PC12 cells [427].

Rivastigmine provides meaningful management of symptoms, maintains cognitive ability and daily function, and prevents some behavioural problems in AD [428–430].

2.20 SAS0132

SAS0132 (Fig. 2) is a novel potent σ 2R ligand with a nine-fold selectivity over the σ 1R. SAS0132 has an affinity of 90 nM for the σ 2R, acting as a σ 2R antagonist reversing intracellular calcium influx in human SK-N-SH neuroblastoma cells in response to σ 2R agonist DKR-1051 [431].

2.20.1 Alzheimer's Disease

In transgenic *C. elegans* overexpressing the APP gene, SAS0132 reduced the age-dependent neurodegeneration compared with controls. Furthermore, in the Thy-1 hAPP^{Lond/Swe+} transgenic mouse model of AD, SAS0132 improves cognitive performance [431]. SAS0132 is a promising σ 2R antagonist with interesting results in pre-clinical studies; however, it has yet to enter clinical trials.

2.21 T-817 MA

T-817MA (edonergic maleate) (Fig. 2) is a σ 1R ligand with an affinity (K_i) of 16 nM. It has been shown to induce the outgrowth of neurites in cultured rat cortical neurons and protect against sodium nitroprusside-induced ROS production and cell death, which could be reversed by the σ -1 R antagonist BD1047 [432].

2.21.1 Alzheimer's Disease

In animal studies of AD, edonergic maleate ameliorated the deficits in adult neurogenesis and spatial memory in rats infused intra-cerebroventricularly (ICV) with A β peptide and showed neuroprotective effects on learning deficits induced by continuous infusion of A β peptide (1–40) [433, 434]. The doses required to see a therapeutic effect in these studies are higher than the reported K_i of 16 nM, with doses

of 0.5–1 μ M required to see positive effects. This possibly rules out the σ -1 R as the drug target of edonergic maleate. However, this phenomenon is seen with other σ 1R ligands, including IPAG and dipentylammonium [6, 156]. It is possibly explained by a non-sigmoidal dose-response curve, such as a bell-shaped or a multiple-binding sites curve [6].

In phase 1 clinical trials, edonergic maleate was well tolerated at a single dose of up to 896 mg and once-daily doses of 672 mg for up to two weeks [435]. An initial phase 2 proof of concept study (NCT00663936) demonstrated the safety and tolerability of edonergic maleate at a 224 mg/day dose for one year. Although it did not meet its primary and secondary outcome targets, post hoc analysis of a positive cognitive effect was seen in patients with moderate impairment (MMSE scores < 20) (reported in a poster at the Alzheimer's Association international conference in 2013 [436]). When a larger phase 2 study of edonergic maleate (NCT02079909) was conducted, it was shown to be generally safe and tolerable at the two doses studied (224 and 448 mg/day). However, in the primary measures of AD [AD Assessment Scale–Cognitive Subscale (ADAS–COG) and AD Cooperative Study–Clinical Global Impression of Change (ADCS–CGIC)], no significant difference was observed compared with the placebo. Furthermore, there were no significant improvements in secondary outcome, the AD Cooperative Study–Activities of Daily Living (ADCS–ADL) scale [435].

It is possible that since edonergic maleate was tested at too high a dose, it fell on the far side of the bell-shaped dose-response curve, and therefore the positive effects were not seen. However, two doses were tested, one low and one high (224 and 448 mg/day), all of which put the expected serum concentrations well above that needed to occupy the σ 1R. It is also possible that, like with blarcamesine, the heterogeneity of AD means that only a certain population of AD patients may benefit from edonergic maleate treatment. However, further clinical study, including patient biomarker analysis, is required. Therefore, with the available (albeit limited) clinical evidence, it is likely that edonergic maleate will not significantly help treat AD.

3 Conclusions

The σ 1R interacts with multiple other protein complexes to modulate multiple signalling pathways and influence cellular health and survival (Fig. 3). This makes the σ 1R an interesting target for the development of drugs for neurological diseases such as AD, HD, PD, MS and MND where neuron cell loss is a key part of the pathology. The fact that the expression of σ 1R diminishes early on in AD and PD, and that there are genotypes for the σ 1R that increase the risk of AD, PD and MS also add to the weight of evidence that




Fig. 3 Some of the documented roles of the sigma-1 receptor in preventing neurological disease. (1) Activation of the sigma-1 receptor by an agonist result in the disassociation from binding immunoglobulin protein (BiP) and chaperoning of calcium channels at the mitochondria-associated ER membrane (MAM), resulting in calcium shuttling into the mitochondria and increased ATP production. (2) NMDA receptors are upregulated on sigma-1 activation and trafficked to the plasma membrane, and the sigma-1 receptor modulates NMDA receptor synaptic transmission and plasticity via SK channels.

(3) Sigma-1 activation down-regulates calcium currents in L-type Ca^{2+} channels in retinal ganglion cells and upregulates calcium currents through L-type Ca^{2+} channels in the hippocampus. (4) Sigma-1 activation down-regulates Na^{+} currents through voltage-gated Na^{+} channels in intracardiac ganglion neurons. (5) Sigma-1 activation upregulates potassium currents through A-type K^{+} channels in neurohypophysial terminals and *Xenopus* oocytes. Parasympathetic intracardiac neurons CA1 field of hippocampus. Sigma-1 activation can upregulate human ether-a-go go K^{+} currents in *Xenopus* oocytes.

the $\sigma 1\text{R}$ is important to these diseases and that $\sigma 1\text{R}$ ligands may be of benefit.

There have been vast strides taken in recent years for the early detection of diseases such as AD [437–440] and PD [441, 442], which means that new drug treatments targeting the $\sigma 1\text{R}$ may benefit more if they can prevent damage to the brain before the disease can progress. Drugs that target multiple targets, including the $\sigma 1\text{R}$, such as blarcamesine or pridopidine in combination with $\text{A}\beta$ antibodies, may prove to be effective in preventing neurological decline in AD. This possibly could result in the repurposing of older drugs, less specific to the $\sigma 1\text{R}$ in a multi-targeting and repurposing approach to drug development [52].

With the ever-evolving understanding of genetic variations and other biomarkers in the populations of people with these neurological diseases, there is a great opportunity to select medicines that are more likely to work in a specific population. For example, in AD, patients with the wild-type $\sigma 1\text{R}$ and higher scores on the MMSE fared better with blarcamesine than those with lower scores and a mutant $\sigma 1\text{R}$.

The $\sigma 2\text{R}$ is often investigated concerning cancer treatment and imaging. However, in recent years, there has been a growing interest in both the $\sigma 2\text{R}$ and $\sigma 1\text{R}$ for treatments in neurological diseases. Despite their lack of homology, their similar binding profile and activity of the proteins mean both receptors need to be considered when designing novel drugs, since the $\sigma 1\text{R}$ agonists and $\sigma 2\text{R}$ antagonists are neuroprotective and $\sigma 1\text{R}$ antagonists and $\sigma 2\text{R}$ agonists are used in cancer study. In brief, with the current research and planned clinical studies, it is expected that there will be an advancement of new $\sigma 1\text{R}$ drugs for the treatment of the neurological disease including AD, HD and MS.

Acknowledgements D.S.M. was supported by The Second Century Fund (C2F) for Postdoctoral Fellowship, Chulalongkorn University. The Chulalongkorn University Graduate School Thesis Grant supported K.S.R. All figures were created with www.biorender.com.

Declarations

Funding No funding was received for this manuscript.

Conflict of interest The authors have no conflicts of interest to declare.

Availability of data and material Data sharing does not apply to this article as no datasets were generated or analysed during the current study.

Ethics approval Not applicable to this article.

Consent to participate: Not applicable to this article.

Consent for publication Not applicable to this article.

Code availability Not applicable to this article.

Author contributions Drafting manuscript: JMB, DSM, PT, KSR, SB, TT. Editing Manuscript: JMB, SB. Preparation of Figures: TT, JMB. Literature searching: JMB, PT, KSR, SB, DSM. Manuscript revision: JMB, PT, DSM. All authors have read and approved the final version of the manuscript and agree to be accountable for the work.

References

- Hanner M, Moebius FF, Flandorfer A, Knaus HG, Striessnig J, Kempner E, et al. Purification, molecular cloning, and expression of the mammalian sigma-1-binding site. *Proc Natl Acad Sci USA*. 1996;93(15):8072–7. <https://doi.org/10.1073/pnas.93.15.8072>.
- Hayashi T, Su TP. σ -1 Receptors (σ 1 binding sites) form raft-like microdomains and target lipid droplets on the endoplasmic reticulum: roles in endoplasmic reticulum lipid compartmentalization and export. *J Pharmacol Exp Ther*. 2003;306(2):718–25. <https://doi.org/10.1124/jpet.103.051284>.
- Hayashi T, Su TP. Regulating ankyrin dynamics: roles of sigma-1 receptors. *Proc Natl Acad Sci USA*. 2001;98(2):491–6. <https://doi.org/10.1073/pnas.021413698>.
- Kinoshita M, Matsuoka Y, Suzuki T, Mirrieles J, Yang J. Sigma-1 receptor alters the kinetics of Kv1.3 voltage gated potassium channels but not the sensitivity to receptor ligands. *Brain Res*. 2012;1452:1–9. <https://doi.org/10.1016/j.brainres.2012.02.070>.
- Brimson JM, Akula KK, Abbas H, Ferry DR, Kulkarni SK, Russell ST, et al. Simple ammonium salts acting on sigma-1 receptors yield potential treatments for cancer and depression. *Sci Rep*. 2020;10(1):9251. <https://doi.org/10.1038/s41598-020-65849-6>.
- Brimson JM, Brown CA, Safrany ST. Antagonists show GTP-sensitive high-affinity binding to the sigma-1 receptor. *Br J Pharmacol*. 2011;164(2B):772–80. <https://doi.org/10.1111/j.1476-5381.2011.01417.x>.
- Natsvlshvili N, Gogvadze N, Zhuravliova E, Mikeladze D. Sigma-1 receptor directly interacts with Rac1-GTPase in the brain mitochondria. *BMC Biochem*. 2015;16(1):1–7. <https://doi.org/10.1186/s12858-015-0040-y>.
- Fontanilla D, Johannessen M, Hajipour AR, Cozzi NV, Jackson MB, Ruoho AE. The hallucinogen N, N-dimethyltryptamine (DMT) is an endogenous sigma-1 receptor regulator. *Science*. 2009;323(5916):934–7. <https://doi.org/10.1126/science.1166127>.
- Hayashi T, Su TP. Sigma-1 receptor ligands: potential in the treatment of neuropsychiatric disorders. *CNS Drugs*. 2004;18(5):269–84. <https://doi.org/10.2165/00023210-200418050-00001>.
- Ruoho AE, Chu UB, Ramachandran S, Fontanilla D, Mavlyutov T, Hajipour AR. The ligand binding region of the sigma-1 receptor: studies utilizing photoaffinity probes, sphingosine and N-alkylamines. *Curr Pharm Des*. 2012;18(7):920–9. <https://doi.org/10.2174/138161212799436584>.
- Vavers E, Zvejniece L, Maurice T, Dambrova M. Allosteric modulators of sigma-1 receptor: a review. *Front Pharmacol*. 2019;10:223. <https://doi.org/10.3389/fphar.2019.00223>.
- Hayashi T, Su TP. The sigma receptor: evolution of the concept in neuropsychopharmacology. *Curr Neuropharmacol*. 2005;3(4):267–80. <https://doi.org/10.2174/157015905774322516>.
- Quirion R, Chicheportiche R, Contreras PC, Johnson KM, Lodge D, Tam SW, et al. Classification and nomenclature of phencyclidine and sigma receptor sites. *Trends Neurosci*. 1987;10(11):444–6. [https://doi.org/10.1016/0166-2236\(87\)90094-4](https://doi.org/10.1016/0166-2236(87)90094-4).
- Spruce BA, Campbell LA, McTavish N, Cooper MA, Appleyard MVL, O'Neill M, et al. Small molecule antagonists of the sigma-1 receptor cause selective release of the death program in tumor and self-reliant cells and inhibit tumor growth in vitro and in vivo. *Cancer Res*. 2004;64(14):4875–86. <https://doi.org/10.1158/0008-5472.CAN-03-3180>.
- Wang L, Prescott AR, Spruce BA, Sanderson J, Duncan G. Sigma receptor antagonists inhibit human lens cell growth and induce pigmentation. *Invest Ophthalmol Vis Sci*. 2005;46(4):1403–8. <https://doi.org/10.1167/iovs.04-1209>.
- Hayashi T, Su TP. Sigma-1 receptor chaperones at the ER-mitochondrion interface regulate Ca²⁺ signaling and cell survival. *Cell*. 2007;131(3):596–610. <https://doi.org/10.1016/j.cell.2007.08.036>.
- Hellewell SB, Bruce A, Feinstein G, Orringer J, Williams W, Bowen WD. Rat liver and kidney contain high densities of σ 1 and σ 2 receptors: characterization by ligand binding and photoaffinity labeling. *Eur J Pharmacol*. 1994;268(1):9–18. [https://doi.org/10.1016/0922-4106\(94\)90115-5](https://doi.org/10.1016/0922-4106(94)90115-5).
- Wolfe SA Jr, Culp SG, De Souza EB. σ -Receptors in endocrine organs: identification, characterization, and autoradiographic localization in rat pituitary, adrenal, testis, and ovary. *Endocrinology*. 1989;124(3):1160–72. <https://doi.org/10.1210/endo-124-3-1160>.
- Borde P, Cosgrove N, Charmsaz S, Safrany ST, Young L. An investigation of Sigma-1 receptor expression and ligand-induced endoplasmic reticulum stress in breast cancer. *Cancer Gene Ther*. 2023;30(2):368–74. <https://doi.org/10.1038/s41417-022-00552-4>.
- Huang YS, Lu HL, Zhang LJ, Wu Z. Sigma-2 receptor ligands and their perspectives in cancer diagnosis and therapy. *Med Res Rev*. 2014;34(3):532–66. <https://doi.org/10.1002/med.21297>.
- Nicholson H, Mesangeau C, McCurdy CR, Bowen WD. Sigma-2 receptors play a role in cellular metabolism: stimulation of glycolytic hallmarks by CM764 in human SK-N-SH neuroblastoma. *J Pharmacol Exp Ther*. 2016;356(2):232–43. <https://doi.org/10.1124/jpet.115.228387>.
- Rousseaux CG, Greene SF. Sigma receptors [σ Rs]: biology in normal and diseased states. *J Recept Signal Transduct Res*. 2016;36(4):327–88. <https://doi.org/10.3109/10799893.2015.1015737>.
- Liu C, Yu CF, Wang SC, Li HY, Lin CM, Wang HH, et al. Sigma-2 receptor/TMEM97 agonist PB221 as an alternative drug for brain tumor. *BMC Cancer*. 2019;19(1):473. <https://doi.org/10.1186/s12885-019-5700-7>.
- Yang K, Wang C, Sun T. The roles of intracellular chaperone proteins, sigma receptors, in Parkinson's disease (PD) and major depressive disorder (MDD). *Front Pharmacol*. 2019;10:528. <https://doi.org/10.3389/fphar.2019.00528>.
- Georgiadis MO, Karoutzou O, Foscolos AS, Papanastasiou I. Sigma receptor (σ R) ligands with antiproliferative and anticancer

- activity. *Molecules*. 2017;22:9. <https://doi.org/10.3390/molecules22091408>.
26. Gebreselassie D, Bowen WD. Sigma-2 receptors are specifically localized to lipid rafts in rat liver membranes. *Eur J Pharmacol*. 2004;493(1–3):19–28. <https://doi.org/10.1016/j.ejphar.2004.04.005>.
 27. Munro S. Lipid rafts: elusive or illusive? *Cell*. 2003;115(4):377–88. [https://doi.org/10.1016/S0092-8674\(03\)00882-1](https://doi.org/10.1016/S0092-8674(03)00882-1).
 28. Alon A, Lyu J, Braz JM, Tummino TA, Craik V, O'Meara MJ, et al. Structures of the σ_2 receptor enable docking for bioactive ligand discovery. *Nature*. 2021;600(7890):759–64. <https://doi.org/10.1038/s41586-021-04175-x>.
 29. Terada K, Migita K, Matsushima Y, Kamei C. Sigma-2 receptor as a potential therapeutic target for treating central nervous system disorders. *Neural Regen Res*. 2019;14(11):1893–4. <https://doi.org/10.4103/1673-5374.259609>.
 30. Abate C, Niso M, Berardi F. Sigma-2 receptor: past, present and perspectives on multiple therapeutic exploitations. *Future Med Chem*. 2018;10(16):1997–2018. <https://doi.org/10.4155/fmc-2018-0072>.
 31. Cantonero C, Camello PJ, Abate C, Berardi F, Salido GM, Rosado JA, et al. NO1, a new Sigma 2 receptor/TMEM97 fluorescent ligand, downregulates SOCE and promotes apoptosis in the triple negative breast cancer cell lines. *Cancers (Basel)*. 2020;12(2):257. <https://doi.org/10.3390/cancers12020257>.
 32. Zeng C, Vangveravong S, McDunn JE, Hawkins WG, Mach RH. Sigma-2 receptor ligand as a novel method for delivering a SMAC mimetic drug for treating ovarian cancer. *Br J Cancer*. 2013;109(9):2368–77. <https://doi.org/10.1038/bjc.2013.593>.
 33. Zeng C, Riad A, Mach RH. The biological function of Sigma-2 receptor/TMEM97 and its utility in PET imaging studies in cancer. *Cancers (Basel)*. 2020;12(7):1877. <https://doi.org/10.3390/cancers12071877>.
 34. 2021 Alzheimer's disease facts and figures. *Alzheimers Dement*. 2021;17(3):327–406. <https://doi.org/10.1002/alz.12328>.
 35. Querfurth HW, LaFerla FM. Alzheimer's disease. *N Engl J Med*. 2010;362(4):329–44. <https://doi.org/10.1056/NEJMra0909142>.
 36. Hyman BT. Amyloid-dependent and amyloid-independent stages of Alzheimer disease. *Arch Neurol*. 2011;68(8):1062–4. <https://doi.org/10.1001/archneurol.2011.70>.
 37. Soria Lopez JA, González HM, Léger GC. Alzheimer's disease. *Handb Clin Neurol*. 2019;167:231–55. <https://doi.org/10.1016/b978-0-12-804766-8.00013-3>.
 38. Hampel H, Caraci F, Cuello AC, Caruso G, Nisticò R, Corbo M, et al. A path toward precision medicine for neuroinflammatory mechanisms in Alzheimer's disease. *Front Immunol*. 2020;11:456. <https://doi.org/10.3389/fimmu.2020.00456>.
 39. Johnson ECB, Dammer EB, Duong DM, Ping L, Zhou M, Yin L, et al. Large-scale proteomic analysis of Alzheimer's disease brain and cerebrospinal fluid reveals early changes in energy metabolism associated with microglia and astrocyte activation. *Nat Med*. 2020;26(5):769–80. <https://doi.org/10.1038/s41591-020-0815-6>.
 40. Sánchez-Fernández C, Entrena JM, Baeyens JM, Cobos EJ. Sigma-1 receptor antagonists: a new class of neuromodulatory analgesics. *Adv Exp Med Biol*. 2017;964:109–32. https://doi.org/10.1007/978-3-319-50174-1_9.
 41. Su TP, Hayashi T. Understanding the molecular mechanism of sigma-1 receptors: towards a hypothesis that sigma-1 receptors are intracellular amplifiers for signal transduction. *Curr Med Chem*. 2003;10(20):2073–80. <https://doi.org/10.2174/0929867033456783>.
 42. De B, Nadal X, Portillo-salido E, Sánchez-arroyos R, Ovalle S, Palacios G, et al. Sigma-1 receptors regulate activity-induced spinal sensitization and neuropathic pain after peripheral nerve injury. *Pain*. 2009;145(3):294–303. <https://doi.org/10.1016/j.pain.2009.05.013>.
 43. Nieto FR, Cendán CM, Sánchez-Fernández C, Cobos EJ, Entrena JM, Tejada MA, et al. Role of sigma-1 receptors in paclitaxel-induced neuropathic pain in mice. *J Pain*. 2012;13(11):1107–21. <https://doi.org/10.1016/j.jpain.2012.08.006>.
 44. Zamanillo D, Romero L, Merlos M, Vela JM. Sigma 1 receptor: a new therapeutic target for pain. *Eur J Pharmacol*. 2013;716(1–3):78–93. <https://doi.org/10.1016/j.ejphar.2013.01.068>.
 45. Kawamura K, Kimura Y, Tsukada H, Kobayashi T, Nishiyama S, Kakiuchi T, et al. An increase of sigma1 receptors in the aged monkey brain. *Neurobiol Aging*. 2003;24(5):745–52. [https://doi.org/10.1016/S0197-4580\(02\)00152-5](https://doi.org/10.1016/S0197-4580(02)00152-5).
 46. Wallace DR, Mactutus CF, Booze RM. Sigma binding sites identified by [³H] DTG are elevated in aged Fischer-344x Brown Norway (F1) rats. *Synapse*. 2000;35(4):311–3. [https://doi.org/10.1002/\(SICI\)1098-2396\(20000315\)35:4%3c311::AID-SYN9%3e3.0.CO;2-5](https://doi.org/10.1002/(SICI)1098-2396(20000315)35:4%3c311::AID-SYN9%3e3.0.CO;2-5).
 47. Ishiwata K, Kobayashi T, Kawamura K, Matsuno K. Age-related changes of the binding of [³H] SA4503 to sigma1 receptors in the rat brain. *Ann Nucl Med*. 2003;17(1):73–7. <https://doi.org/10.1007/BF02988264>.
 48. Horsager J, Fedorova TD, Berge NV, Klinge MW, Knudsen K, Hansen AK, et al. Cardiac 11C-donepezil binding increases with age in healthy humans: potentially signifying Sigma-1 receptor upregulation. *J Cardiovasc Pharmacol Ther*. 2019;24(4):365–70. <https://doi.org/10.1177/1074248419838509>.
 49. Norbury R, Travis MJ, Erlandsson K, Waddington W, Owens J, Pimlott S, et al. In vivo imaging of muscarinic receptors in the aging female brain with (R, R)[¹²³I]-QNB and single photon emission tomography. *Exp Gerontol*. 2005;40(3):137–45. <https://doi.org/10.1016/j.exger.2004.10.002>.
 50. Sheline YI, Mintun MA, Moerlein SM, Snyder AZ. Greater loss of 5-HT_{2A} receptors in midlife than in late life. *Am J Psychiatry*. 2002;159(3):430–5. <https://doi.org/10.1176/appi.ajp.159.3.430>.
 51. Inoue M, Sahara T, Sudo Y, Okubo Y, Yasuno F, Kishimoto T, et al. Age-related reduction of extrastriatal dopamine D₂ receptor measured by PET. *Life Sci*. 2001;69(9):1079–84. [https://doi.org/10.1016/s0024-3205\(01\)01205-x](https://doi.org/10.1016/s0024-3205(01)01205-x).
 52. Brimson JM, Brimson S, Chomchoei C, Tencomnao T. Using Sigma-ligands as part of a multi-receptor approach to target diseases of the brain. *Expert Opin Ther Targets*. 2020;24(10):1009–28. <https://doi.org/10.1080/14728222.2020.1805435>.
 53. Prasanth MI, Malar D, Tencomnao T, Brimson J. The emerging role of the sigma-1 receptor in autophagy: hand-in-hand targets for the treatment of Alzheimer's Disease. *Expert Opin Ther Targets*. 2021;25(5):401–14. <https://doi.org/10.1080/14728222.2021.1939681>.
 54. Mishina M, Ohyama M, Ishii K, Kitamura S, Kimura Y, Oda K, et al. Low density of sigma 1 receptors in early Alzheimer's disease. *Ann Nucl Med*. 2008;22(3):151–6. <https://doi.org/10.1007/s12149-007-0094-z>.
 55. Minoshima S, Giordani B, Berent S, Frey KA, Foster NL, Kuhl DE. Metabolic reduction in the posterior cingulate cortex in very early Alzheimer's disease. *Ann Neurol*. 1997;42(1):85–94. <https://doi.org/10.1002/ana.410420114>.
 56. Uchida N, Ujike H, Tanaka Y, Sakai A, Yamamoto M, Fujisawa Y, et al. A variant of the sigma receptor type-1 gene is a protective factor for Alzheimer disease. *Am J Geriatr Psychiatry*. 2005;13(12):1062–6. <https://doi.org/10.1176/appi.ajgp.13.12.1062>.
 57. Maruszak A, Safranow K, Gacia M, Gabryelewicz T, Słowik A, Styczńska M, et al. Sigma receptor type 1 gene variation in a group of polish patients with Alzheimer's disease and mild cognitive impairment. *Dement Geriatr Cogn Disord*. 2007;23(6):432–8. <https://doi.org/10.1159/000101990>.

58. Huang Y, Zheng L, Halliday G, Dobson-Stone C, Wang Y, Tang H-D, et al. Genetic polymorphisms in sigma-1 receptor and apolipoprotein E interact to influence the severity of Alzheimer's disease. *Curr Alzheimer Res.* 2011;8(7):765–70. <https://doi.org/10.2174/156720511797633232>.
59. Feher A, Juhasz A, Laszlo A, Kalman J Jr, Pakaski M, Kalman J, et al. Association between a variant of the sigma-1 receptor gene and Alzheimer's disease. *Neurosci Lett.* 2012;517(2):136–9. <https://doi.org/10.1016/j.neulet.2012.04.046>.
60. Jin JL, Fang M, Zhao YX, Liu XY. Roles of sigma-1 receptors in Alzheimer's disease. *Int J Clin Exp Med.* 2015;8(4):4808–20.
61. Tsai SY, Hayashi T, Harvey BK, Wang Y, Wu WW, Shen RF, et al. Sigma-1 receptors regulate hippocampal dendritic spine formation via a free radical-sensitive mechanism involving Rac1•GTP pathway. *Proc Natl Acad Sci USA.* 2009;106(52):22468–73. <https://doi.org/10.1073/pnas.0909089106>.
62. Fisher A, Bezprozvanny I, Wu L, Ryskamp DA, Bar-Ner N, Natan N, et al. AF710B, a novel M1/σ1 agonist with therapeutic efficacy in animal models of Alzheimer's disease. *Neurodegener Dis.* 2016;16(1–2):95–110. <https://doi.org/10.1159/000440864>.
63. Ryskamp D, Wu L, Wu J, Kim D, Rammes G, Geva M, et al. Pridopidine stabilizes mushroom spines in mouse models of Alzheimer's disease by acting on the sigma-1 receptor. *Neurobiol Dis.* 2019;124:489–504. <https://doi.org/10.1016/j.nbd.2018.12.022>.
64. Tsai SYA, Pokrass MJ, Klauer NR, Nohara H, Su TP. Sigma-1 receptor regulates Tau phosphorylation and axon extension by shaping p35 turnover via myristic acid. *Proc Natl Acad Sci USA.* 2015;112(21):6742–7. <https://doi.org/10.1073/pnas.1422001112>.
65. Christ MG, Huesmann H, Nagel H, Kern A, Behl C. Sigma-1 receptor activation induces autophagy and increases proteostasis capacity in vitro and in vivo. *Cells.* 2019;8(3):211. <https://doi.org/10.3390/cells8030211>.
66. Romeo MA, Faggioni A, Cirone M. Could autophagy dysregulation link neurotropic viruses to Alzheimer's disease? *Neural Regen Res.* 2019;14(9):1503–6. <https://doi.org/10.4103/1673-5374.253508>.
67. McBrayer M, Nixon RA. Lysosome and calcium dysregulation in Alzheimer's disease: partners in crime. *Biochem Soc Trans.* 2013;41(6):1495–502. <https://doi.org/10.1042/BST20130201>.
68. Uddin MS, Stachowiak A, Mamun AA, Tzvetkov NT, Takeda S, Atanasov AG, et al. Autophagy and Alzheimer's disease: from molecular mechanisms to therapeutic implications. *Front Aging Neurosci.* 2018;10:04. <https://doi.org/10.3389/fnagi.2018.00004>.
69. Callens M, Loncke J, Bultynck G. Dysregulated Ca²⁺ homeostasis as a central theme in neurodegeneration: lessons from Alzheimer's disease and Wolfram syndrome. *Cells.* 2022;11(12):1963. <https://doi.org/10.3390/cells11121963>.
70. Holmes C, Boche D, Wilkinson D, Yadegarfar G, Hopkins V, Bayer A, et al. Long-term effects of Aβ42 immunisation in Alzheimer's disease: follow-up of a randomised, placebo-controlled phase I trial. *Lancet.* 2008;372(9634):216–23. [https://doi.org/10.1016/S0140-6736\(08\)61075-2](https://doi.org/10.1016/S0140-6736(08)61075-2).
71. van Dyck CH, Swanson CJ, Aisen P, Bateman RJ, Chen C, Gee M, et al. Lecanemab in early Alzheimer's disease. *N Engl J Med.* 2023;388(1):9–21. <https://doi.org/10.1056/NEJMoa2212948>.
72. Lacor PN, Buniel MC, Furlow PW, Clemente AS, Velasco PT, Wood M, et al. Aβ oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci.* 2007;27(4):796–807. <https://doi.org/10.1523/JNEUROSCI.3501-06.2007>.
73. Pannuzzo M. On the physiological/pathological link between Aβ peptide, cholesterol, calcium ions and membrane deformation: a molecular dynamics study. *Biochim Biophys Acta.* 2016;1858(6):1380–9. <https://doi.org/10.1016/j.bbamem.2016.03.018>.
74. Games D, Adams D, Alessandrini R, Barbour R, Borthellette P, Blackwell C, et al. Alzheimer-type neuropathology in transgenic mice overexpressing V717F β-amyloid precursor protein. *Nature.* 1995;373(6514):523–7. <https://doi.org/10.1038/373523a0>.
75. Brimson JM. The Pharmacology of the Sigma-1 Receptor [Doctoral Thesis]. In: The University of Bath online repository. 2010. <http://opus.bath.ac.uk/19712/>. Accessed 1 Dec 2022.
76. Katnik C, Guerrero WR, Pennypacker KR, Herrera Y, Cuevas J. Sigma-1 receptor activation prevents intracellular calcium dysregulation in cortical neurons during in vitro ischemia. *J Pharmacol Exp Ther.* 2006;319(3):1355–65. <https://doi.org/10.1124/jpet.106.107557>.
77. Delprat B, Crouzier L, Su TP, Maurice T. At the crossing of ER stress and MAMs: a key role of sigma-1 receptor? *Adv Exp Med Biol.* 2020;1131:699–718. https://doi.org/10.1007/978-3-030-12457-1_2878.
78. Hayashi T, Rizzuto R, Hajnoczky G, Su TP. MAM: more than just a housekeeper. *Trends Cell Biol.* 2009;19(2):81–8. <https://doi.org/10.1016/j.tcb.2008.12.00279>.
79. Scheffer IE, Berkovic S, Capovilla G, Connolly MB, French J, Guilhoto L, et al. ILAE classification of the epilepsies: position paper of the ILAE Commission for Classification and Terminology. *Epilepsia.* 2017;58(4):512–21. <https://doi.org/10.1111/epi.13709>.
80. Specchio N, Curatolo P. Developmental and epileptic encephalopathies: what we do and do not know. *Brain.* 2021;144(1):32–43. <https://doi.org/10.1093/brain/awaa371>.
81. Raga S, Specchio N, Rheims S, Wilmshurst JM. Developmental and epileptic encephalopathies: recognition and approaches to care. *Epileptic Disord.* 2021;23(1):40–52. <https://doi.org/10.1684/epd.2021.1244>.
82. Scheffer IE, Liao J. Deciphering the concepts behind “Epileptic encephalopathy” and “Developmental and epileptic encephalopathy.” *Eur J Paediatr Neurol.* 2020;24:11–4. <https://doi.org/10.1016/j.ejpn.2019.12.023>.
83. Trivisano M, Specchio N. What are the epileptic encephalopathies? *Curr Opin Neurol.* 2020;33(2):179–84. <https://doi.org/10.1097/wco.0000000000000793>.
84. EPX-100 (Clemizole Hydrochloride) as add-on therapy to control convulsive seizures in patients with Dravet syndrome (ARGUS). In: NIH U.S. National Library of Medicine. 2020. <https://clinicaltrials.gov/ct2/show/NCT04462770>. Accessed 1 Dec 2022.
85. Griffin A, Hamling KR, Knupp K, Hong S, Lee LP, Baraban SC. Clemizole and modulators of serotonin signalling suppress seizures in Dravet syndrome. *Brain.* 2017;140(3):669–83. <https://doi.org/10.1093/brain/aww342>.
86. Vasquez A, Buraniqi E, Wirrell EC. New and emerging pharmacologic treatments for developmental and epileptic encephalopathies. *Curr Opin Neurol.* 2022;35(2):145–54. <https://doi.org/10.1097/wco.0000000000001029>.
87. Miziak B, Czuczwar S. Advances in the design and discovery of novel small molecule drugs for the treatment of Dravet Syndrome. *Expert Opin Drug Discov.* 2021;16(5):579–93. <https://doi.org/10.1080/17460441.2021.1857722>.
88. Samanta D. Changing landscape of Dravet syndrome management: an overview. *Neuropediatrics.* 2020;51(2):135–45. <https://doi.org/10.1055/s-0040-1701694>.
89. Martin P, Reeder T, Sourbron J, de Witte PAM, Gammaitoni AR, Galer BS. An emerging role for Sigma-1 receptors in the treatment of developmental and epileptic encephalopathies. *Int J Mol Sci.* 2021;22(16):8416. <https://doi.org/10.3390/ijms22168416>.
90. Vavers E, Zvejniece B, Stelfa G, Svalbe B, Vilks K, Kupats E, et al. Genetic inactivation of the sigma-1 chaperone protein

- results in decreased expression of the R2 subunit of the GABA-B receptor and increased susceptibility to seizures. *Neurobiol Dis.* 2021;150:105244. <https://doi.org/10.1016/j.nbd.2020.105244>.
91. Schaffar G, Breuer P, Boteva R, Behrends C, Tzvetkov N, Strippe N, et al. Cellular toxicity of polyglutamine expansion proteins: mechanism of transcription factor deactivation. *Mol Cell.* 2004;15(1):95–105. <https://doi.org/10.1016/j.molcel.2004.06.029>.
 92. Martinez-Vicente M, Talloczy Z, Wong E, Tang G, Koga H, Kaushik S, et al. Cargo recognition failure is responsible for inefficient autophagy in Huntington's disease. *Nat Neurosci.* 2010;13(5):567–76. <https://doi.org/10.1038/nn.2528>.
 93. Mochel F, Haller RG. Energy deficit in Huntington disease: why it matters. *J Clin Invest.* 2011;121(2):493–9. <https://doi.org/10.1172/JCI45691>.
 94. Caron NS, Wright GEB, Hayden MR. Huntington Disease. In: Adam MP, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, et al., editors. *GeneReviews*, University of Washington, Seattle. 1993. <https://www.ncbi.nlm.nih.gov/books/NBK1305/>.
 95. Frank S. Treatment of Huntington's disease. *Neurotherapeutics.* 2014;11(1):153–60. <https://doi.org/10.1007/s13311-013-0244-z>.
 96. Miki Y, Tanji K, Mori F, Wakabayashi K. Sigma-1 receptor is involved in degradation of intranuclear inclusions in a cellular model of Huntington's disease. *Neurobiol Dis.* 2015;74:25–31. <https://doi.org/10.1016/j.nbd.2014.11.005>.
 97. Ryskamp D, Wu J, Geva M, Kusko R, Grossman I, Hayden M, et al. The sigma-1 receptor mediates the beneficial effects of pridopidine in a mouse model of Huntington disease. *Neurobiol Dis.* 2017;97:46–59. <https://doi.org/10.1016/j.nbd.2016.10.006>.
 98. Jabłońska M, Grzelakowska K, Wiśniewski B, Mazur E, Leis K, Gałązka P. Pridopidine in the treatment of Huntington's disease. *Rev Neurosci.* 2020;31(4):441–51. <https://doi.org/10.1515/revneuro-2019-0085>.
 99. Nguyen L, Lucke-Wold BP, Mookerjee SA, Cavendish JZ, Robson MJ, Scandinaro AL, et al. Role of sigma-1 receptors in neurodegenerative diseases. *J Pharmacol Sci.* 2015;127(1):17–29. <https://doi.org/10.1016/j.jphs.2014.12.005>.
 100. Swinnen B, Robberecht W. The phenotypic variability of amyotrophic lateral sclerosis. *Nat Rev Neurol.* 2014;10(11):661–70. <https://doi.org/10.1038/nrneuro.2014.184>.
 101. Machts J, Keute M, Kaufmann J, Schreiber S, Kasper E, Petri S, et al. Longitudinal clinical and neuroanatomical correlates of memory impairment in motor neuron disease. *Neuroimage Clin.* 2021;29:102545. <https://doi.org/10.1016/j.nicl.2020.102545>.
 102. Roggenbuck J, Quick A, Kolb SJ. Genetic testing and genetic counseling for amyotrophic lateral sclerosis: an update for clinicians. *Genet Med.* 2017;19(3):267–74. <https://doi.org/10.1038/gim.2016.107>.
 103. van den Bos MAJ, Geevasinga N, Higashihara M, Menon P, Vucic S. Pathophysiology and diagnosis of ALS: insights from advances in neurophysiological techniques. *Int J Mol Sci.* 2019;20(11):2818. <https://doi.org/10.3390/ijms20112818>.
 104. Bensimon G, Lacomblez L, Meininger V. A controlled trial of riluzole in amyotrophic lateral sclerosis. *ALS/Riluzole Study Group.* *N Engl J Med.* 1994;330(9):585–91. <https://doi.org/10.1056/nejm199403033300901>.
 105. Heo YA. Sodium phenylbutyrate and ursodoxicoltaurine: first approval. *CNS Drugs.* 2022;36(9):1007–13. <https://doi.org/10.1007/s40263-022-00945-x>.
 106. Jaiswal MK. Riluzole and edaravone: a tale of two amyotrophic lateral sclerosis drugs. *Med Res Rev.* 2019;39(2):733–48. <https://doi.org/10.1002/med.21528>.
 107. Mora JS, Genge A, Chio A, Estol CJ, Chaverri D, Hernández M, et al. Masitinib as an add-on therapy to riluzole in patients with amyotrophic lateral sclerosis: a randomized clinical trial. *Amyotroph Lateral Scler Frontotemporal Degener.* 2020;21(1–2):5–14. <https://doi.org/10.1080/21678421.2019.1632346>.
 108. Latham BD, Oskin DS, Crouch RD, Vergne MJ, Jackson KD. Cytochromes P450 2C8 and 3A catalyze the metabolic activation of the tyrosine kinase inhibitor masitinib. *Chem Res Toxicol.* 2022;35(9):1467–81. <https://doi.org/10.1021/acs.chemrestox.2c00057>.
 109. Casanovas A, Salvany S, Lahoz V, Tarabal O, Piedrafita L, Sabater R, et al. Neuregulin 1-ErbB module in C-bouton synapses on somatic motor neurons: molecular compartmentation and response to peripheral nerve injury. *Sci Rep.* 2017;7:40155. <https://doi.org/10.1038/srep40155>.
 110. Herrando-Grabulosa M, Gaja-Capdevila N, Vela JM, Navarro X. Sigma 1 receptor as a therapeutic target for amyotrophic lateral sclerosis. *Br J Pharmacol.* 2021;178(6):1336–52. <https://doi.org/10.1111/bph.15224>.
 111. Mavlyutov TA, Epstein ML, Liu P, Verbny YI, Ziskind-Conhaim L, Ruoho AE. Development of the sigma-1 receptor in C-terminals of motoneurons and colocalization with the N, N'-dimethyltryptamine forming enzyme, indole-N-methyltransferase. *Neuroscience.* 2012;206:60–8. <https://doi.org/10.1016/j.neuroscience.2011.12.040>.
 112. Mavlyutov TA, Epstein ML, Andersen KA, Ziskind-Conhaim L, Ruoho AE. The sigma-1 receptor is enriched in postsynaptic sites of C-terminals in mouse motoneurons. An anatomical and behavioral study. *Neuroscience.* 2010;167(2):247–55. <https://doi.org/10.1016/j.neuroscience.2010.02.022>.
 113. Watanabe S, Ilieva H, Tamada H, Nomura H, Komine O, Endo F, et al. Mitochondria-associated membrane collapse is a common pathomechanism in SIGMAR 1-and SOD 1-linked ALS. *EMBO Mol Med.* 2016;8(12):1421–37. <https://doi.org/10.15252/emmm.201606403>.
 114. Couly S, Khalil B, Viguier V, Roussel J, Maurice T, Liévens J-C. Sigma-1 receptor is a key genetic modulator in amyotrophic lateral sclerosis. *Hum Mol Genet.* 2020;29(4):529–40. <https://doi.org/10.1093/hmg/ddz267>.
 115. Vollrath J, Sechi A, Dreser A, Katona I, Wiemuth D, Vervoorts J, et al. Loss of function of the ALS protein SigR1 leads to ER pathology associated with defective autophagy and lipid raft disturbances. *Cell Death Dis.* 2014;5(6):e1290. <https://doi.org/10.1038/cddis.2014.243>.
 116. Dreser A, Vollrath JT, Sechi A, Johann S, Roos A, Yamoah A, et al. The ALS-linked E102Q mutation in Sigma receptor-1 leads to ER stress-mediated defects in protein homeostasis and dysregulation of RNA-binding proteins. *Cell Death Differ.* 2017;24(10):1655–71. <https://doi.org/10.1038/cdd.2017.88>.
 117. Mavlyutov TA, Epstein ML, Verbny YI, Huerta MS, Zaitoun I, Ziskind-Conhaim L, et al. Lack of sigma-1 receptor exacerbates ALS progression in mice. *Neuroscience.* 2013;240:129–34. <https://doi.org/10.1016/j.neuroscience.2013.02.035>.
 118. Al-Saif A, Al-Mohanna F, Bohlega S. A mutation in sigma-1 receptor causes juvenile amyotrophic lateral sclerosis. *Ann Neurol.* 2011;70(6):913–9. <https://doi.org/10.1002/ana.22534>.
 119. Fukunaga K, Shinoda Y, Tagashira H. The role of SIGMAR1 gene mutation and mitochondrial dysfunction in amyotrophic lateral sclerosis. *J Pharmacol Sci.* 2015;127(1):36–41. <https://doi.org/10.1016/j.jphs.2014.12.012>.
 120. Barmada SJ, Skibinski G, Korb E, Rao EJ, Wu JY, Finkbeiner S. Cytoplasmic mislocalization of TDP-43 is toxic to neurons and enhanced by a mutation associated with familial amyotrophic lateral sclerosis. *J Neurosci.* 2010;30(2):639–49. <https://doi.org/10.1523/JNEUROSCI.4988-09.2010>.
 121. Prasad A, Bharathi V, Sivalingam V, Girdhar A, Patel BK. Molecular mechanisms of TDP-43 misfolding and pathology in

- amyotrophic lateral sclerosis. *Front Mol Neurosci.* 2019;12:25. <https://doi.org/10.3389/fnmol.2019.00025>.
122. Brettschneider J, Del Tredici K, Toledo JB, Robinson JL, Irwin DJ, Grossman M, et al. Stages of pTDP-43 pathology in amyotrophic lateral sclerosis. *Ann Neurol.* 2013;74(1):20–38. <https://doi.org/10.1002/ana.23937>.
 123. Lee P-T, Liévens J-C, Wang S-M, Chuang J-Y, Khalil B, Wu H-E, et al. Sigma-1 receptor chaperones rescue nucleocytoplasmic transport deficit seen in cellular and *Drosophila* ALS/FTD models. *Nat Commun.* 2020;11(1):5580. <https://doi.org/10.1038/s41467-020-19396-3>.
 124. Frischer JM, Bramow S, Dal-Bianco A, Lucchinetti CF, Rauschka H, Schmidbauer M, et al. The relation between inflammation and neurodegeneration in multiple sclerosis brains. *Brain.* 2009;132(Pt 5):1175–89. <https://doi.org/10.1093/brain/awp070>.
 125. McGinley MP, Goldschmidt CH, Rae-Grant AD. Diagnosis and treatment of multiple sclerosis: a review. *JAMA.* 2021;325(8):765–79. <https://doi.org/10.1001/jama.2020.26858>.
 126. Krupp L. Fatigue is intrinsic to multiple sclerosis (MS) and is the most commonly reported symptom of the disease. *Mult Scler.* 2006;12(4):367–8. <https://doi.org/10.1191/13524850ms1373ed>.
 127. Braley TJ, Chervin RD. Fatigue in multiple sclerosis: mechanisms, evaluation, and treatment. *Sleep.* 2010;33(8):1061–7. <https://doi.org/10.1093/sleep/33.8.1061>.
 128. Jacobs LD, Beck RW, Simon JH, Kinkel RP, Brownschield CM, Murray TJ, et al. Intramuscular interferon beta-1a therapy initiated during a first demyelinating event in multiple sclerosis. CHAMPS Study Group. *N Engl J Med.* 2000;343(13):898–904. <https://doi.org/10.1056/nejm200009283431301>.
 129. Lycke J. Monoclonal antibody therapies for the treatment of relapsing-remitting multiple sclerosis: differentiating mechanisms and clinical outcomes. *Ther Adv Neurol Disord.* 2015;8(6):274–93. <https://doi.org/10.1177/1756285615605429>.
 130. Kappos L, Radue EW, O'Connor P, Polman C, Hohlfeld R, Calabresi P, et al. A placebo-controlled trial of oral fingolimod in relapsing multiple sclerosis. *N Engl J Med.* 2010;362(5):387–401. <https://doi.org/10.1056/NEJMoa0909494>.
 131. Kappos L, Bar-Or A, Cree BAC, Fox RJ, Giovannoni G, Gold R, et al. Siponimod versus placebo in secondary progressive multiple sclerosis (EXPAND): a double-blind, randomised, phase 3 study. *Lancet.* 2018;391(10127):1263–73. [https://doi.org/10.1016/s0140-6736\(18\)30475-6](https://doi.org/10.1016/s0140-6736(18)30475-6).
 132. Comi G, Kappos L, Selmaj KW, Bar-Or A, Arnold DL, Steinman L, et al. Safety and efficacy of ozanimod versus interferon beta-1a in relapsing multiple sclerosis (SUNBEAM): a multicentre, randomised, minimum 12-month, phase 3 trial. *Lancet Neurol.* 2019;18(11):1009–20. [https://doi.org/10.1016/s1474-4422\(19\)30239-x](https://doi.org/10.1016/s1474-4422(19)30239-x).
 133. Johnson KP, Brooks BR, Cohen JA, Ford CC, Goldstein J, Lisak RP, et al. Copolymer 1 reduces relapse rate and improves disability in relapsing-remitting multiple sclerosis: results of a phase III multicenter, double-blind placebo-controlled trial. The Copolymer 1 Multiple Sclerosis Study Group. *Neurology.* 1995;45(7):1268–76. <https://doi.org/10.1212/wnl.45.7.1268>.
 134. Confavreux C, O'Connor P, Comi G, Freedman MS, Miller AE, Olsson TP, et al. Oral teriflunomide for patients with relapsing multiple sclerosis (TOWER): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Neurol.* 2014;13(3):247–56. [https://doi.org/10.1016/s1474-4422\(13\)70308-9](https://doi.org/10.1016/s1474-4422(13)70308-9).
 135. Giovannoni G, Comi G, Cook S, Rammohan K, Rieckmann P, Soelberg Sørensen P, et al. A placebo-controlled trial of oral cladribine for relapsing multiple sclerosis. *N Engl J Med.* 2010;362(5):416–26. <https://doi.org/10.1056/NEJMoa0902533>.
 136. Palacios G, Muro A, Verdú E, Pumarola M, Vela JM. Immunohistochemical localization of the sigma1 receptor in Schwann cells of rat sciatic nerve. *Brain Res.* 2004;1007(1–2):65–70. <https://doi.org/10.1016/j.brainres.2004.02.013>.
 137. Hayashi T, Su T-P. Sigma-1 receptors at galactosylceramide-enriched lipid microdomains regulate oligodendrocyte differentiation. *Proc Natl Acad Sci USA.* 2004;101(41):14949–54. <https://doi.org/10.1073/pnas.0402890101>.
 138. Zhemkov V, Geva M, Hayden MR, Bezprozvanny I. Sigma-1 receptor (S1R) interaction with cholesterol: mechanisms of S1R activation and its role in neurodegenerative diseases. *Int J Mol Sci.* 2021;22(8):4082. <https://doi.org/10.3390/ijms22084082>.
 139. Berghoff SA, Spieth L, Saher G. Local cholesterol metabolism orchestrates remyelination. *Trends Neurosci.* 2022;45(4):272–83. <https://doi.org/10.1016/j.tins.2022.01.001>.
 140. Kenche H, Singh M, Smith J, Shen K. Neuronal mitochondrial dysfunction in a cellular model of circadian rhythm disruption is rescued by donepezil. *Biochem Biophys Res Commun.* 2021;567:56–62. <https://doi.org/10.1016/j.bbrc.2021.06.029>.
 141. Oxombre B, Lee-Chang C, Duhamel A, Toussaint M, Giroux M, Donnier-Maréchal M, et al. High-affinity σ 1 protein agonist reduces clinical and pathological signs of experimental autoimmune encephalomyelitis. *Br J Pharmacol.* 2015;172(7):1769–82. <https://doi.org/10.1111/bph.13037>.
 142. Lotankar S, Prabhavalkar KS, Bhatt LK. Biomarkers for Parkinson's disease: recent advancement. *Neurosci Bull.* 2017;33(5):585–97. <https://doi.org/10.1007/s12264-017-0183-5>.
 143. Chen-Plotkin AS, Albin R, Alcalay R, Babcock D, Bajaj V, Bowman D, et al. Finding useful biomarkers for Parkinson's disease. *Sci Transl Med.* 2018;10(454):eaam6003. <https://doi.org/10.1126/scitranslmed.aam6003>.
 144. Li T, Le W. Biomarkers for Parkinson's disease: how good are they? *Neurosci Bull.* 2020;36(2):183–94. <https://doi.org/10.1007/s12264-019-00433-1>.
 145. Bloem BR, Okun MS, Klein C. Parkinson's disease. *Lancet.* 2021;397(10291):2284–303. [https://doi.org/10.1016/s0140-6736\(21\)00218-x](https://doi.org/10.1016/s0140-6736(21)00218-x).
 146. Parkinson disease. In: World Health Organization website. 2022. <https://www.who.int/news-room/fact-sheets/detail/parkinson-disease>. Accessed 1 Dec 2022.
 147. LeWitt PA. Levodopa therapy for Parkinson's disease: pharmacokinetics and pharmacodynamics. *Mov Disord.* 2015;30(1):64–72. <https://doi.org/10.1002/mds.26082>.
 148. Nonnekes J, Timmer MH, de Vries NM, Rascol O, Helmich RC, Bloem BR. Unmasking levodopa resistance in Parkinson's disease. *Mov Disord.* 2016;31(11):1602–9. <https://doi.org/10.1002/mds.26712>.
 149. Walker Z, Possin KL, Boeve BF, Aarsland D. Lewy body dementias. *Lancet.* 2015;386(10004):1683–97. [https://doi.org/10.1016/S0140-6736\(15\)00462-6](https://doi.org/10.1016/S0140-6736(15)00462-6).
 150. Aarsland D, Andersen K, Larsen J, Lolk A, Nielsen H, Kragh-Sørensen P. Risk of dementia in Parkinson's disease: a community-based, prospective study. *Neurology.* 2001;56(6):730–6. <https://doi.org/10.1212/wnl.56.6.730>.
 151. Mishina M, Ishiwata K, Ishii K, Kitamura S, Kimura Y, Kawamura K, et al. Function of sigma 1 receptors in Parkinson's disease. *Acta Neurol Scand.* 2005;112(2):103–7. <https://doi.org/10.1111/j.1600-0404.2005.00432.x>.
 152. Hong J, Wang L, Zhang T, Zhang B, Chen L. Sigma-1 receptor knockout increases α -synuclein aggregation and phosphorylation with loss of dopaminergic neurons in substantia nigra. *Neurobiol Aging.* 2017;59:171–83. <https://doi.org/10.1016/j.neurobiolaging.2017.08.007>.
 153. Luo Y, Roth GS. The roles of dopamine oxidative stress and dopamine receptor signaling in aging and age-related neurodegeneration. *Antioxid Redox Signal.* 2000;2(3):449–60. <https://doi.org/10.1089/15230860050192224>.

154. Mori T, Hayashi T, Su TP. Compromising σ -1 receptors at the endoplasmic reticulum render cytotoxicity to physiologically relevant concentrations of dopamine in a nuclear factor- κ B/Bcl-2-dependent mechanism: Potential relevance to Parkinson's disease. *J Pharmacol Exp Ther*. 2012;341(3):663–71. <https://doi.org/10.1124/jpet.111.190868>.
155. Cagnin M, Ozzano M, Bellio N, Fiorentino I, Follo C, Isidoro C. Dopamine induces apoptosis in APPswe-expressing Neuro2A cells following Pepstatin-sensitive proteolysis of APP in acid compartments. *Brain Res*. 2012;1471:102–17. <https://doi.org/10.1016/j.brainres.2012.06.025>.
156. Brimson JM, Safrany ST, Qassam H, Tencomnao T. Dipentylammonium binds to the Sigma-1 receptor and protects against glutamate toxicity, attenuates dopamine toxicity and potentiates neurite outgrowth in various cultured cell lines. *Neurotox Res*. 2018;34(2):263–72. <https://doi.org/10.1007/s12640-018-9883-5>.
157. Poewe W, Seppi K, Tanner CM, Halliday GM, Brundin P, Volkman J, et al. Parkinson disease. *Nat Rev Dis Primers*. 2017;3:17013. <https://doi.org/10.1038/nrdp.2017.13>.
158. Jenner P, Olanow CW. Oxidative stress and the pathogenesis of Parkinson's disease. *Neurology*. 1996;47(6 Suppl 3):S161–70. https://doi.org/10.1212/wnl.47.6_suppl_3.161s.
159. Cristóvão AC, Guhathakurta S, Bok E, Je G, Yoo SD, Choi D-H, et al. NADPH oxidase 1 mediates α -synucleinopathy in Parkinson's disease. *J Neurosci*. 2012;32(42):14465–77. <https://doi.org/10.1523/JNEUROSCI.2246-12.2012>.
160. Guo JD, Zhao X, Li Y, Li GR, Liu XL. Damage to dopaminergic neurons by oxidative stress in Parkinson's disease. *Int J Mol Med*. 2018;41(4):1817–25. <https://doi.org/10.3892/ijmm.2018.3406>.
161. Pal A, Fontanilla D, Gopalakrishnan A, Chae Y-K, Markley JL, Ruoho AE. The sigma-1 receptor protects against cellular oxidative stress and activates antioxidant response elements. *Eur J Pharmacol*. 2012;682(1–3):12–20. <https://doi.org/10.1016/j.ejphar.2012.01.030>.
162. Wang J, Shanmugam A, Markand S, Zorrilla E, Ganapathy V, Smith SB. Sigma 1 receptor regulates the oxidative stress response in primary retinal Müller glial cells via NRF2 signaling and system xc⁻, the Na⁺-independent glutamate–cystine exchanger. *Free Radic Biol Med*. 2015;86:25–36. <https://doi.org/10.1016/j.freeradbiomed.2015.04.009>.
163. Prasanth MI, Tencomnao T, Brimson JM. The Sigma-1 receptors role in neuroprotection: Comment on Nrf2 as a therapeutic target in ischemic stroke. *Expert Opin Ther Targets*. 2021;25(7):613–4. <https://doi.org/10.1080/14728222.2021.1948016>.
164. Brimson JM, Prasanth MI, Isidoro C, Sukprasansap M, Tencomnao T. *Cleistocalyx nervosum* var. paniala seed extracts exhibit sigma-1 antagonist sensitive neuroprotective effects in PC12 cells and protect *C. elegans* from stress via the SKN-1/NRF-2 pathway. *Nutr Healthy Aging*. 2021;6(2):131–46. <https://doi.org/10.3233/NHA-200108>.
165. Brimson JM, Prasanth MI, Malar DS, Verma K, Plaingam W, Tencomnao T. Bacopa monnieri protects neuronal cell line and *Caenorhabditis elegans* models of Alzheimer's disease through sigma-1 receptor antagonist sensitive and antioxidant pathways. *Nutr Healthy Aging*. 2022;7(3):173–96. <https://doi.org/10.3233/NHA-220161>.
166. Betzer C, Movius AJ, Shi M, Gai W-P, Zhang J, Jensen PH. Identification of synaptosomal proteins binding to monomeric and oligomeric α -synuclein. *PLoS ONE*. 2015;10(2):e0116473. <https://doi.org/10.1371/journal.pone.0116473>.
167. Schneider JL, Cuervo AM. Autophagy and human disease: emerging themes. *Curr Opin Genet Dev*. 2014;26:16–23. <https://doi.org/10.1016/j.gde.2014.04.003>.
168. Vogiatzi T, Xilouri M, Vekrellis K, Stefanis L. Wild type α -synuclein is degraded by chaperone-mediated autophagy and macroautophagy in neuronal cells. *J Biol Chem*. 2008;283(35):23542–56. <https://doi.org/10.1074/jbc.M801992200>.
169. Galvagnion C. The role of lipids interacting with α -synuclein in the pathogenesis of Parkinson's disease. *J Parkinsons Dis*. 2017;7(3):433–50. <https://doi.org/10.3233/JPD-171103>.
170. Galvagnion C, Brown JW, Ouberai MM, Flagmeier P, Vendruscolo M, Buell AK, et al. Chemical properties of lipids strongly affect the kinetics of the membrane-induced aggregation of α -synuclein. *Proc Natl Acad Sci USA*. 2016;113(26):7065–70. <https://doi.org/10.1073/pnas.1601899113>.
171. Plotegher N, Berti G, Ferrari E, Tessari I, Zanetti M, Lunelli L, et al. DOPAL derived alpha-synuclein oligomers impair synaptic vesicles physiological function. *Sci Rep*. 2017;7:40699. <https://doi.org/10.1038/srep40699>.
172. Limegrover CS, Yurko R, Izzo NJ, LaBarbera KM, Rehak C, Look G, et al. Sigma-2 receptor antagonists rescue neuronal dysfunction induced by Parkinson's patient brain-derived α -synuclein. *J Neurosci Res*. 2021;99(4):1161–76. <https://doi.org/10.1002/jnr.24782>.
173. Gold WA, Krishnaraj R, Ellaway C, Christodoulou J. Rett syndrome: a genetic update and clinical review focusing on comorbidities. *ACS Chem Neurosci*. 2018;9(2):167–76. <https://doi.org/10.1021/acscchemneuro.7b00346>.
174. Vidal S, Xioli C, Pascual-Alonso A, O'Callaghan M, Pineda M, Armstrong J. Genetic landscape of Rett syndrome spectrum: improvements and challenges. *Int J Mol Sci*. 2019;20(16):3925. <https://doi.org/10.3390/ijms20163925>.
175. Chahrouh M, Jung SY, Shaw C, Zhou X, Wong ST, Qin J, et al. MeCP2, a key contributor to neurological disease, activates and represses transcription. *Science*. 2008;320(5880):1224–9. <https://doi.org/10.1126/science.1153252>.
176. Lyst MJ, Bird A. Rett syndrome: a complex disorder with simple roots. *Nat Rev Genet*. 2015;16(5):261–75. <https://doi.org/10.1038/nrg3897>.
177. Kriacionis S, Paterson A, Curtis J, Guy J, Macleod N, Bird A. Gene expression analysis exposes mitochondrial abnormalities in a mouse model of Rett syndrome. *Mol Cell Biol*. 2006;26(13):5033–42. <https://doi.org/10.1128/mcb.01665-05>.
178. Ricciardi S, Boggio EM, Grosso S, Lonetti G, Forlani G, Stefanelli G, et al. Reduced AKT/mTOR signaling and protein synthesis dysregulation in a Rett syndrome animal model. *Hum Mol Genet*. 2011;20(6):1182–96. <https://doi.org/10.1093/hmg/ddq563>.
179. Palmer A, Qayumi J, Ronnett G. MeCP2 mutation causes distinguishable phases of acute and chronic defects in synaptogenesis and maintenance, respectively. *Mol Cell Neurosci*. 2008;37(4):794–807. <https://doi.org/10.1016/j.mcn.2008.01.005>.
180. Fonzo M, Sirico F, Corrado B. Evidence-based physical therapy for individuals with Rett syndrome: a systematic review. *Brain Sci*. 2020;10(7):410. <https://doi.org/10.3390/brainsci10070410>.
181. Ehinger Y, Matagne V, Villard L, Roux JC. Rett syndrome from bench to bedside: recent advances. *F1000Res*. 2018;7:398. <https://doi.org/10.12688/f1000research.14056.1>.
182. Chahil G, Bollu PC. Rett Syndrome. In: Aboubakr S, Abu-Ghosh A, Acharya A, Adibi Sedeh P, Aebi T, Aeddula N, Et al. Editors. StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing. 2023. <https://www.ncbi.nlm.nih.gov/books/NBK482252/>.
183. Ohno K, Saito Y, Ueda R, Togawa M, Ohmae T, Matsuda E, et al. Effect of serotonin 1A agonists and selective serotonin reuptake inhibitors on behavioral and nighttime respiratory symptoms in Rett syndrome. *Pediatr Neurol*. 2016;60:54–9.e1. <https://doi.org/10.1016/j.pediatrneurol.2016.03.016>.
184. Neul JL, Lane JB, Lee H-S, Geerts S, Barrish JO, Annesse F, et al. Developmental delay in Rett syndrome: data from the natural history study. *J Neurodev Disord*. 2014;6(1):20. <https://doi.org/10.1186/1866-1955-6-20>.

185. Luikenhuis S, Giacometti E, Beard CF, Jaenisch R. Expression of MeCP2 in postmitotic neurons rescues Rett syndrome in mice. *Proc Natl Acad Sci USA*. 2004;101(16):6033–8. <https://doi.org/10.1073/pnas.0401626101>.
186. Cohen DR, Matarazzo V, Palmer AM, Tu Y, Jeon O-H, Pevsner J, et al. Expression of MeCP2 in olfactory receptor neurons is developmentally regulated and occurs before synaptogenesis. *Mol Cell Neurosci*. 2003;22(4):417–29. [https://doi.org/10.1016/S1044-7431\(03\)00026-5](https://doi.org/10.1016/S1044-7431(03)00026-5).
187. Chen WG, Chang Q, Lin Y, Meissner A, West AE, Griffith EC, et al. Derepression of BDNF transcription involves calcium-dependent phosphorylation of MeCP2. *Science*. 2003;302(5646):885–9. <https://doi.org/10.1126/science.1086446>.
188. Kikuchi-Utsumi K, Nakaki T. Chronic treatment with a selective ligand for the sigma-1 receptor chaperone, SA4503, up-regulates BDNF protein levels in the rat hippocampus. *Neurosci Lett*. 2008;440(1):19–22. <https://doi.org/10.1016/j.neulet.2008.05.055>.
189. Mysona BA, Zhao J, Smith S, Bollinger KE. Relationship between Sigma-1 receptor and BDNF in the visual system. *Exp Eye Res*. 2018;167:25–30. <https://doi.org/10.1016/j.exer.2017.10.012>.
190. Hashimoto K. Sigma-1 receptor chaperone and brain-derived neurotrophic factor: emerging links between cardiovascular disease and depression. *Prog Neurobiol*. 2013;100:15–29. <https://doi.org/10.1016/j.pneurobio.2012.09.001>.
191. Nakano M, Osada K, Misonoo A, Fujiwara K, Takahashi M, Ogawa Y, et al. Fluvoxamine and sigma-1 receptor agonists dehydroepiandrosterone (DHEA)-sulfate induces the Ser473-phosphorylation of Akt-1 in PC12 cells. *Life Sci*. 2010;86(9–10):309–14. <https://doi.org/10.1016/j.lfs.2009.11.017>.
192. Christ MG, Clement AM, Behl C. The sigma-1 receptor at the crossroad of proteostasis, neurodegeneration, and autophagy. *Trends Neurosci*. 2020;43(2):79–81. <https://doi.org/10.1016/j.tins.2019.12.002>.
193. Urano F. Wolfram syndrome: diagnosis, management, and treatment. *Curr Diab Rep*. 2016;16(1):6. <https://doi.org/10.1007/s11892-015-0702-6>.
194. Abreu D, Urano F. Current landscape of treatments for Wolfram syndrome. *Trends Pharmacol Sci*. 2019;40(10):711–4. <https://doi.org/10.1016/j.tips.2019.07.011>.
195. Delprat B, Maurice T, Delettre C. Wolfram syndrome: MAMs' connection? *Cell Death Dis*. 2018;9(3):364. <https://doi.org/10.1038/s41419-018-0406-3>.
196. Mori T, Hayashi T, Hayashi E, Su TP. Sigma-1 receptor chaperone at the ER-mitochondrion interface mediates the mitochondrion-ER-nucleus signaling for cellular survival. *PLoS ONE*. 2013;8(10):e76941. <https://doi.org/10.1371/journal.pone.0076941>.
197. Crouzier L, Danese A, Yasui Y, Richard EM, Liévens J-C, Patergnani S, et al. Activation of the sigma-1 receptor chaperone alleviates symptoms of Wolfram syndrome in preclinical models. *Sci Transl Med*. 2022;14(631):eabh3763. <https://doi.org/10.1126/scitranslmed.abh3763>.
198. Kornhuber J, Bormann J, Hübers M, Rusche K, Riederer P. Effects of the 1-amino-adamantanes at the MK-801-binding site of the NMDA-receptor-gated ion channel: a human postmortem brain study. *Eur J Pharmacol*. 1991;206(4):297–300. [https://doi.org/10.1016/0922-4106\(91\)90113-v](https://doi.org/10.1016/0922-4106(91)90113-v).
199. Blanpied TA, Clarke RJ, Johnson JW. Amantadine inhibits NMDA receptors by accelerating channel closure during channel block. *J Neurosci*. 2005;25(13):3312–22. <https://doi.org/10.1523/jneurosci.4262-04.2005>.
200. Peeters M, Romieu P, Maurice T, Su TP, Maloteaux JM, Hermans E. Involvement of the sigma1 receptor in the modulation of dopaminergic transmission by amantadine. *Eur J Neurosci*. 2004;19(8):2212–20. <https://doi.org/10.1111/j.0953-816X.2004.03297.x>.
201. Abbas H. Expression of sigma receptors in human cancer cell lines and effects of novel sigma-2 ligands on their proliferation [Doctoral Thesis]. In: The University of Wolverhampton open repository. 2018. <http://hdl.handle.net/2436/621768>. Accessed 1 Dec 2022.
202. Hubsher G, Haider M, Okun MS. Amantadine: the journey from fighting flu to treating Parkinson disease. *Neurology*. 2012;78(14):1096–9. <https://doi.org/10.1212/WNL.0b013e31824e8f0d>.
203. Chang C, Ramphul K. Amantadine. In: Aboubakr S, Abu-Ghosh A, Acharya A, Adibi Sedeh P, Aeby T, Aeddula N, et al. Editors. StatPearls. Treasure Island (FL): StatPearls Publishing. 2023. <https://www.ncbi.nlm.nih.gov/books/NBK499953/>.
204. Schwab RS, England AC, Poskanzer DC, Young RR. Amantadine in the treatment of Parkinson's disease. *JAMA*. 1969;208(7):1168–70.
205. Rascol O, Fabbri M, Poewe W. Amantadine in the treatment of Parkinson's disease and other movement disorders. *Lancet Neurol*. 2021;20(12):1048–56. [https://doi.org/10.1016/S1474-4422\(21\)00249-0](https://doi.org/10.1016/S1474-4422(21)00249-0).
206. Kornhuber J, Schoppmeyer K, Riederer P. Affinity of 1-amino-adamantanes for the σ binding site in post-mortem human frontal cortex. *Neurosci Lett*. 1993;163(2):129–31. [https://doi.org/10.1016/0304-3940\(93\)90362-O](https://doi.org/10.1016/0304-3940(93)90362-O).
207. Crosby NJ, Deane KH, Clarke CE. Amantadine for dyskinesia in Parkinson's disease. *Cochrane Database Syst Rev*. 2003;2003(2):CD003467. <https://doi.org/10.1002/14651858.Cd003467>.
208. Aoki FY, Sitar DS. Clinical pharmacokinetics of amantadine hydrochloride. *Clin Pharmacokinet*. 1988;14(1):35–51. <https://doi.org/10.2165/00003088-198814010-00003>.
209. Al-Salama ZT. Amantadine extended release capsules (GOCOVRI) in Parkinson's disease: a profile of its use in the USA. *Drugs Ther Perspect*. 2022;38(5):203–14. <https://doi.org/10.1007/s40267-022-00912-y>.
210. Hauser RA, Mehta SH, Kremens D, et al. Effects of Gocovri (amantadine) extended-release capsules on motor aspects of experiences of daily living in people with Parkinson's disease and dyskinesia. *Neurol Ther*. 2021;10(2):739–51. <https://doi.org/10.1007/s40120-021-00256-1>.
211. Elmer LW, Juncos JL, Singer C, et al. Pooled analyses of phase III studies of ADS-5102 (amantadine) extended-release capsules for dyskinesia in Parkinson's disease. *CNS Drugs*. 2018;32:387–98. <https://doi.org/10.1007/s40263-018-0498-4>.
212. Isaacson SH, Fahn S, Pahwa R, et al. Parkinson's patients with dyskinesia switched from immediate release amantadine to open-label ADS-5102. *Mov Disord Clin Pract*. 2018;5:183–90. <https://doi.org/10.1002/mdc3.12595>.
213. Oertel W, Eggert K, Pahwa R, Tanner CM, Hauser RA, Trenkwalder C, et al. Randomized, placebo-controlled trial of ADS-5102 (amantadine) extended-release capsules for levodopa-induced dyskinesia in Parkinson's disease (EASE LID 3). *Mov Disord*. 2017;32(12):1701–9. <https://doi.org/10.1002/mds.27131>.
214. Pahwa R, Tanner CM, Hauser RA, Isaacson SH, Nausieda PA, Truong DD, et al. ADS-5102 (amantadine) extended-release capsules for levodopa-induced dyskinesia in Parkinson disease (EASE LID study): a randomized clinical trial. *JAMA Neurol*. 2017;74(8):941–9. <https://doi.org/10.1001/jamaneurol.2017.0943>.
215. Tanner CM, Pahwa R, Hauser RA, Oertel WH, Isaacson SH, Jankovic J, et al. EASE LID 2: a 2-year open-label trial of gocovri (amantadine) extended release for dyskinesia in

- Parkinson's disease. *J Parkinsons Dis.* 2020;10(2):543–58. <https://doi.org/10.3233/JPD-191841>.
216. Rosenberg GA, Appenzeller O. Amantadine, fatigue, and multiple sclerosis. *Arch Neurol.* 1988;45(10):1104–6. <https://doi.org/10.1001/archneur.1988.00520340058012>.
 217. Shaygannejad V, Janghorbani M, Ashtari F, Zakeri H. Comparison of the effect of aspirin and amantadine for the treatment of fatigue in multiple sclerosis: a randomized, blinded, crossover study. *Neurol Res.* 2012;34(9):854–8. <https://doi.org/10.1179/1743132812Y.0000000081>.
 218. Ledinek AH, Sajko MC, Rot U. Evaluating the effects of amantadin, modafinil and acetyl-L-carnitine on fatigue in multiple sclerosis—result of a pilot randomized, blind study. *Clin Neurol Neurosurg.* 2013;115:S86–9. <https://doi.org/10.1016/j.clineuro.2013.09.029>.
 219. Khazaei M, Karevan A, Taheri M, Ghafouri-Fard S. Comparison of the effects of amantadine and ondansetron in treatment of fatigue in patients with multiple sclerosis. *Clin Transl Med.* 2019;8(1):20. <https://doi.org/10.1186/s40169-019-0239-4>.
 220. Puca F, Bricolo A, Turella G. Effect of L-dopa or amantadine therapy on sleep spindles in Parkinsonism. *Electroencephalogr Clin Neurophysiol.* 1973;35(3):327–30. [https://doi.org/10.1016/0013-4694\(73\)90245-9](https://doi.org/10.1016/0013-4694(73)90245-9).
 221. Nourbakhsh B, Revirajan N, Morris B, Cordano C, Creasman J, Manguinao M, et al. Safety and efficacy of amantadine, modafinil, and methylphenidate for fatigue in multiple sclerosis: a randomised, placebo-controlled, crossover, double-blind trial. *Lancet Neurol.* 2021;20(1):38–48. [https://doi.org/10.1016/s1474-4422\(20\)30354-9](https://doi.org/10.1016/s1474-4422(20)30354-9).
 222. Villard V, Espallergues J, Keller E, Vamvakides A, Maurice T. Anti-amnesic and neuroprotective potentials of the mixed muscarinic receptor/sigma1 (σ 1) ligand ANAVEX2-73, a novel aminotetrahydrofuran derivative. *J Psychopharmacol.* 2011;25(8):1101–17. <https://doi.org/10.1177/0269881110379286>.
 223. Lahmy V, Meunier J, Malmström S, Naert G, Givalois L, Kim SH, et al. Blockade of Tau hyperphosphorylation and A β 1–42 generation by the aminotetrahydrofuran derivative ANAVEX2-73, a mixed muscarinic and σ 1 receptor agonist, in a nontransgenic mouse model of Alzheimer's disease. *Neuropsychopharmacology.* 2013;38(9):1706. <https://doi.org/10.1038/npp.2013.70>.
 224. Lahmy V, Long R, Morin D, Villard V, Maurice T. Mitochondrial protection by the mixed muscarinic/ σ 1 ligand ANAVEX2-73, a tetrahydrofuran derivative, in A β 25–35 peptide-injected mice, a nontransgenic Alzheimer's disease model. *Front Cell Neurosci.* 2015;8:463. <https://doi.org/10.3389/fncel.2014.00463>.
 225. Voges O, Weigmann I, Bitterlich N, Missling C, Schindler C. A Phase I dose escalation study to investigate safety, tolerability, and pharmacokinetics of ANAVEX 2-73 in Healthy Male Subjects. *Innov Clin Neurosci.* 2014;12(Suppl B), 1-20; CNS Summit 2014, Boca Raton, Florida.
 226. Hampel H, Williams C, Etcheto A, Goodsaid F, Parmentier F, Sallantin J, et al. A precision medicine framework using artificial intelligence for the identification and confirmation of genomic biomarkers of response to an Alzheimer's disease therapy: Analysis of the blarcamesine (ANAVEX2-73) Phase 2a clinical study. *Alzheimers Dement.* 2020;6(1):e12013. <https://doi.org/10.1002/trc2.12013>.
 227. Lisak RP, Nedelkoska L, Benjamins JA. Sigma-1 receptor agonists as potential protective therapies in multiple sclerosis. *J Neuroimmunol.* 2020;342:577188. <https://doi.org/10.1016/j.jneuroim.2020.577188>.
 228. Kaufmann WE, Sprouse J, Rebowe N, Hanania T, Klamer D, Missling CU. ANAVEX 2–73 (blarcamesine), a Sigma-1 receptor agonist, ameliorates neurologic impairments in a mouse model of Rett syndrome. *Pharmacol Biochem Behav.* 2019;187:172796. <https://doi.org/10.1016/j.pbb.2019.172796>.
 229. Reyes ST, Deacon RM, Guo SG, Altamiras FJ, Castillo JB, van der Wildt B, et al. Effects of the sigma-1 receptor agonist blarcamesine in a murine model of fragile X syndrome: neurobehavioral phenotypes and receptor occupancy. *Sci Rep.* 2021;11(1):17150. <https://doi.org/10.1038/s41598-021-94079-7>.
 230. Anavex-Life-Sciences-Corp. ANAVEX2-73 (Blarcamesine) AVATAR phase 3 trial met primary and secondary efficacy endpoints. In: Anavex press release. 2022. <https://www.anavex.com/post/anavex-2-73-blarcamesine-avator-phase-3-trial-met-primary-and-secondary-efficacy-endpoints>. Accessed 1 Nov 2022.
 231. Hall H, Iulita MF, Gubert P, Flores Aguilar L, Ducatenzeiler A, Fisher A, et al. AF710B, an M1/sigma-1 receptor agonist with long-lasting disease-modifying properties in a transgenic rat model of Alzheimer's disease. *Alzheimers Dement.* 2018;14(6):811–23. <https://doi.org/10.1016/j.jalz.2017.11.009>.
 232. Anavex-life-Sciences-Corp. Anavex Life Sciences reports positive results from phase 1 clinical trial of ANAVEX3-71. In: Anavex press release. 2022. <https://www.anavex.com/post/anavex-life-sciences-reports-positive-results-from-phase-1-clinical-trial-of-anavex-3-71>. Accessed 1 Nov 2022.
 233. Rishton GM, Look GC, Ni ZJ, Zhang J, Wang Y, Huang Y, et al. Discovery of investigational drug CT1812, an antagonist of the Sigma-2 receptor complex for Alzheimer's disease. *ACS Med Chem Lett.* 2021;12(9):1389–95. <https://doi.org/10.1021/acsmcchemlett.1c00048>.
 234. Izzo NJ, Xu J, Zeng C, Kirk MJ, Mozzoni K, Silky C, et al. Alzheimer's therapeutics targeting amyloid beta 1–42 oligomers II: sigma-2/PGRMC1 receptors mediate A β 24 oligomer binding and synaptotoxicity. *PLoS ONE.* 2014;9(11):e111899. <https://doi.org/10.1371/journal.pone.0111899>.
 235. Izzo NJ, Yuede CM, LaBarbera KM, Limegrover CS, Rehak C, Yurko R, et al. Preclinical and clinical biomarker studies of CT1812: a novel approach to Alzheimer's disease modification. *Alzheimers Dement.* 2021;17(8):1365–82. <https://doi.org/10.1371/journal.pone.0111899>.
 236. Harper AR, Nayee S, Topol EJ. Protective alleles and modifier variants in human health and disease. *Nat Rev Genet.* 2015;16(12):689–701. <https://doi.org/10.1038/nrg4017>.
 237. Grundman M, Morgan R, Lickliter JD, Schneider LS, DeKosky S, Izzo NJ, et al. A phase 1 clinical trial of the sigma-2 receptor complex allosteric antagonist CT1812, a novel therapeutic candidate for Alzheimer's disease. *Alzheimers Dement.* 2019;5:20–6. <https://doi.org/10.1016/j.trci.2018.11.001>.
 238. Catalano S, Grundman M, Schneider LS, DeKosky S, Morgan R, Guttendorf R, et al. [P4–567]: a phase 1 safety trial of the A β oligomer receptor antagonist CT1812. *Alzheimers Dement.* 2017;13(7S):P1523–80. <https://doi.org/10.1016/j.jalz.2017.07.730>.
 239. Schwartz AR, Pizon AF, Brooks DE. Dextromethorphan-induced serotonin syndrome. *Clin Toxicol.* 2008;46(8):771–3. <https://doi.org/10.1080/15563650701668625>.
 240. Shin EJ, Nah SY, Chae JS, Bing G, Shin SW, Yen TP, et al. Dextromethorphan attenuates trimethyltin-induced neurotoxicity via sigma1 receptor activation in rats. *Neurochem Int.* 2007;50(6):791–9. <https://doi.org/10.1016/j.neuint.2007.01.008>.
 241. Nguyen L, Thomas KL, Lucke-Wold BP, Cavendish JZ, Crowe MS, Matsumoto RR. Dextromethorphan: an update on its utility for neurological and neuropsychiatric disorders. *Pharmacol Ther.* 2016;159:1–22. <https://doi.org/10.1016/j.pharmthera.2016.01.016>.
 242. Sauvé WM. Recognizing and treating pseudobulbar affect. *CNS Spectr.* 2016;21(S1):34–44. <https://doi.org/10.1017/S1092852916000791>.

243. Garnock-Jones KP. Dextromethorphan/quinidine. *CNS Drugs*. 2011;25(5):435–45. <https://doi.org/10.2165/11207260-0000000000000000>.
244. Cummings JL, Lyketsos CG, Peskind ER, Porsteinsson AP, Mintzer JE, Scharre DW, et al. Effect of dextromethorphan-quinidine on agitation in patients with Alzheimer disease dementia: a randomized clinical trial. *JAMA*. 2015;314(12):1242–54. <https://doi.org/10.1001/jama.2015.10214>.
245. Senanarong V, Cummings J, Fairbanks L, Mega M, Masterman D, O'connor S, et al. Agitation in Alzheimer's disease is a manifestation of frontal lobe dysfunction. *Dement Geriatr Cogn Disord*. 2004;17(1–2):14–20. <https://doi.org/10.1159/000074080>.
246. O'Gorman C, Jones A, Cummings JL, et al. Efficacy and safety of AXS-05, a novel, oral, NMDA-receptor antagonist with multimodal activity, in agitation associated with Alzheimer's disease: results from ADVANCE-1, a phase 2/3, double-blind, active and placebo-controlled trial: developments in clinical trials and cognitive assessment. *Alzheimer Dement*. 2020;16:e047684. <https://doi.org/10.1002/alz.047684>.
247. Chou Y-C, Liao J-F, Chang W-Y, Lin M-F, Chen C-F. Binding of dimemorfan to sigma-1 receptor and its anticonvulsant and locomotor effects in mice, compared with dextromethorphan and dextrorphan. *Brain Res*. 1999;821(2):516–9. [https://doi.org/10.1016/s0006-8993\(99\)01125-7](https://doi.org/10.1016/s0006-8993(99)01125-7).
248. Verhagen Metman L, Del Dotto P, Natté R, van den Munckhof P, Chase TN. Dextromethorphan improves levodopa-induced dyskinesias in Parkinson's disease. *Neurology*. 1998;51(1):203–6. <https://doi.org/10.1212/wnl.51.1.203>.
249. Starr MS, Starr BS, Kaur S. Stimulation of basal and L-DOPA-induced motor activity by glutamate antagonists in animal models of Parkinson's disease. *Neurosci Biobehav Rev*. 1997;21(4):437–46. [https://doi.org/10.1016/s0149-7634\(96\)00039-5](https://doi.org/10.1016/s0149-7634(96)00039-5).
250. Paquette MA, Brudney EG, Putterman DB, Meshul CK, Johnson SW, Berger SP. Sigma ligands, but not N-methyl-D-aspartate antagonists, reduce levodopa-induced dyskinesias. *NeuroReport*. 2008;19(1):111–5. <https://doi.org/10.1097/WNR.0b013e3282f3b0d1>.
251. Liu C-T, Kao L-T, Shih J-H, Chien W-C, Chiu C-H, Ma K-H, et al. The effect of dextromethorphan use in Parkinson's disease: a 6-hydroxydopamine rat model and population-based study. *Eur J Pharmacol*. 2019;862:172639. <https://doi.org/10.1016/j.ejphar.2019.172639>.
252. Fox SH, Metman LV, Nutt JG, Brodsky M, Factor SA, Lang AE, et al. Trial of dextromethorphan/quinidine to treat levodopa-induced dyskinesia in Parkinson's disease. *Mov Disord*. 2017;32(6):893–903. <https://doi.org/10.1002/mds.26976>.
253. Verhagen Metman L, Blanchet PJ, van den Munckhof P, Del Dotto P, Natté R, Chase TN. A trial of dextromethorphan in parkinsonian patients with motor response complications. *Mov Disord*. 1998;13(3):414–7. <https://doi.org/10.1002/mds.870130307>.
254. Wilhelm R, Ahl B, Anghelescu I-G. Dextromethorphan-paroxetine, but not dronabinol, effective for treatment-resistant aggression and agitation in an elderly patient with Lewy body dementia. *J Clin Psychopharmacol*. 2017;37(6):745–7. <https://doi.org/10.1097/jcp.0000000000000805>.
255. Logan BK, Yeakel JK, Goldfogel G, Frost MP, Sandstrom G, Wickham DJ. Dextromethorphan abuse leading to assault, suicide, or homicide. *J Forensic Sci*. 2012;57(5):1388–94. <https://doi.org/10.1111/j.1556-4029.2012.02133.x>.
256. Smith R, Piore E, Myers K, Sirdofsky M, Goslin K, Meekins G, et al. Enhanced bulbar function in amyotrophic lateral sclerosis: the Nuedexta treatment trial. *Neurotherapeutics*. 2017;14(3):762–72. <https://doi.org/10.1007/s13311-016-0508-5>.
257. Green JR, Allison KM, Cordella C, Richburg BD, Pattee GL, Berry JD, et al. Additional evidence for a therapeutic effect of dextromethorphan/quinidine on bulbar motor function in patients with amyotrophic lateral sclerosis: a quantitative speech analysis. *Br J Clin Pharmacol*. 2018;84(12):2849–56. <https://doi.org/10.1111/bcp.13745>.
258. Sancho J, Ferrer S, Burés E, Díaz JL, Torrecilla T, Signes-Costa J, et al. Effect of one-year dextromethorphan/quinidine treatment on management of respiratory impairment in amyotrophic lateral sclerosis. *Respir Med*. 2021;186:106536. <https://doi.org/10.1016/j.rmed.2021.106536>.
259. Chechneva OV, Mayrhofer F, Daugherty DJ, Pleasure DE, Hong J-S, Deng W. Low dose dextromethorphan attenuates moderate experimental autoimmune encephalomyelitis by inhibiting NOX2 and reducing peripheral immune cells infiltration in the spinal cord. *Neurobiol Dis*. 2011;44(1):63–72. <https://doi.org/10.1016/j.nbd.2011.06.004>.
260. Lisak RP, Nedelkoska L, Benjamins JA. Effects of dextromethorphan on glial cell function: proliferation, maturation, and protection from cytotoxic molecules. *Glia*. 2014;62(5):751–62. <https://doi.org/10.1002/glia.22639>.
261. Piore EP, Brooks BR, Cummings J, Schiffer R, Thisted RA, Wynn D, et al. Dextromethorphan plus ultra low-dose quinidine reduces pseudobulbar affect. *Ann Neurol*. 2010;68(5):693–702. <https://doi.org/10.1002/ana.22093>.
262. McGrane I, VandenBerg A, Munjal R. Treatment of pseudobulbar affect with fluoxetine and dextromethorphan in a woman with multiple sclerosis. *Ann Pharmacother*. 2017;51(11):1035–6. <https://doi.org/10.1177/1060028017720746>.
263. Ehde DM, Osborne TL, Jensen MP. Chronic pain in persons with multiple sclerosis. *Phys Med Rehabil Clin N Am*. 2005;16(2):503–12. <https://doi.org/10.1016/j.pmr.2005.01.001>.
264. Merlos M, Romero L, Zamanillo D, Plata-Salamán C, Vela JM. Sigma-1 receptor and pain. *Handb Exp Pharmacol*. 2017;244:131–61. https://doi.org/10.1007/164_2017_9.
265. Panitch HS, Thisted RA, Smith RA, Wynn DR, Wymer JP, Achiron A, et al. Randomized, controlled trial of dextromethorphan/quinidine for pseudobulbar affect in multiple sclerosis. *Ann Neurol*. 2006;59(5):780–7. <https://doi.org/10.1002/ana.20828>.
266. Nelson KA, Park KM, Robinovitz E, Tsigos C, Max MB. High-dose oral dextromethorphan versus placebo in painful diabetic neuropathy and postherpetic neuralgia. *Neurology*. 1997;48(5):1212–8. <https://doi.org/10.1212/wnl.48.5.1212>.
267. Abraham RB, Marouani N, Kollender Y, Meller I, Weinbroum AA. Dextromethorphan for phantom pain attenuation in cancer amputees: a double-blind crossover trial involving three patients. *Clin J Pain*. 2002;18(5):282–5. <https://doi.org/10.1097/00002508-200209000-00002>.
268. Carlsson K, Hoem N, Moberg E, Mathisen L. Analgesic effect of dextromethorphan in neuropathic pain. *Acta Anaesthesiol Scand*. 2004;48(3):328–36. <https://doi.org/10.1111/j.0001-5172.2004.0325.x>.
269. Gilron I, Booher S, Rowan J, Smoller B, Max M. A randomized, controlled trial of high-dose dextromethorphan in facial neuralgias. *Neurology*. 2000;55(7):964–71. <https://doi.org/10.1212/wnl.55.7.964>.
270. McQuay HJ, Carroll D, Jadad AR, Glynn CJ, Jack T, Moore RA, et al. Dextromethorphan for the treatment of neuropathic pain: a double-blind randomised controlled crossover trial with integral n-of-1 design. *Pain*. 1994;59(1):127–33. [https://doi.org/10.1016/0304-3959\(94\)90056-6](https://doi.org/10.1016/0304-3959(94)90056-6).
271. Smith-Hicks CL, Gupta S, Ewen JB, Hong M, Kratz L, Kelley R, Tierney E, Vaurio R, Bibat G, Sanyal A, Yenokyan G, Brereton N, Johnston MV, Naidu S. Randomized open-label trial of dextromethorphan in Rett syndrome. *Neurology*. 2017;89(16):1684–90. <https://doi.org/10.1212/WNL.0000000000004515>.

272. Kumagai K, Shono K, Nakayama H, Ohno Y, Sajai I. (2R-trans)-2-butyl-5-heptylpyrrolidine as a potent sigma receptor ligand produced by *Streptomyces longispororuber*. *Journal Antibiot*. 2000;53(5):467–73. <https://doi.org/10.7164/antibiotics.53.467>.
273. Moebius FF, Reiter RJ, Hanner M, Glossmann H. High affinity of sigma1-binding sites for sterol isomerization inhibitors: evidence for a pharmacological relationship with the yeast sterol C8–C7 isomerase. *Br J Pharmacol*. 1997;121(1):1–6. <https://doi.org/10.1038/sj.bjp.0701079>.
274. Ramachandran S, Chu UB, Mavlyutov TA, Pal A, Pyne S, Ruoho AE. The sigma1 receptor interacts with N-alkyl amines and endogenous sphingolipids. *Eur J Pharmacol*. 2009;609(1–3):19–26. <https://doi.org/10.1016/j.ejphar.2009.03.003>.
275. Li J, Satyshur K, Guo LW, Ruoho AE. Sphingoid bases regulate the Sigma-1 receptor—sphingosine and N,N'-dimethylsphingosine are endogenous agonists. *Int J Mol Sci*. 24(4):3103; [Doi: https://doi.org/10.3390/ijms24043103](https://doi.org/10.3390/ijms24043103)
276. National-Institute-for-Health-and-Care-Excellence. Donepezil, galantamine, rivastigmine and memantine for the treatment of Alzheimer's disease. In: National Institute for Health Care Excellence. 2011;TA217:1-77; <https://www.nice.org.uk/guidance/ta217>. Accessed 1 Nov 2022.
277. Kato K, Hayako H, Ishihara Y, Marui S, Iwane M, Miyamoto M. TAK-147, an acetylcholinesterase inhibitor, increases choline acetyltransferase activity in cultured rat septal cholinergic neurons. *Neurosci Lett*. 1999;260(1):5–8. [https://doi.org/10.1016/S0304-3940\(98\)00943-4](https://doi.org/10.1016/S0304-3940(98)00943-4).
278. Ramakrishnan NK, Visser AK, Schepers M, Luurtsema G, Nyakas CJ, Elsinga PH, et al. Dose-dependent sigma-1 receptor occupancy by donepezil in rat brain can be assessed with 11 C-SA4503 and microPET. *Psychopharmacology*. 2014;231(20):3997–4006. <https://doi.org/10.1007/s00213-014-3533-2>.
279. Sato K, Urbano R, Yu C, Yamasaki F, Sato T, Jordan J, et al. The effect of donepezil treatment on cardiovascular mortality. *Clin Pharmacol Ther*. 2010;88(3):335–8. <https://doi.org/10.1038/clpt.2010.98>.
280. Nordström P, Religa D, Wimo A, Winblad B, Eriksson M. The use of cholinesterase inhibitors and the risk of myocardial infarction and death: a nationwide cohort study in subjects with Alzheimer's disease. *Eur Heart J*. 2013;34(33):2585–91. <https://doi.org/10.1093/eurheartj/ehb182>.
281. Bhuiyan MS, Fukunaga K. Targeting sigma-1 receptor signaling by endogenous ligands for cardioprotection. *Expert Opin Ther Targets*. 2011;15(2):145–55. <https://doi.org/10.1517/14728222.2011.546350>.
282. Ishima T, Nishimura T, Iyo M, Hashimoto K. Potentiation of nerve growth factor-induced neurite outgrowth in PC12 cells by donepezil: role of sigma-1 receptors and IP3 receptors. *Prog Neuropsychopharmacol Biol Psychiatry*. 2008;32(7):1656–9. <https://doi.org/10.1016/j.pnpbp.2008.06.011>.
283. Zhang X, Lian S, Zhang Y, Zhao Q. Efficacy and safety of donepezil for mild cognitive impairment: a systematic review and meta-analysis. *Clin Neurol Neurosurg*. 2022;213:107134. <https://doi.org/10.1016/j.clineuro.2022.107134>.
284. Birks JS, Harvey RJ. Donepezil for dementia due to Alzheimer's disease. *Cochrane Database Syst Rev*. 2018;6(6):CD001190. <https://doi.org/10.1002/14651858.CD001190.pub3>.
285. Wang H, Zong Y, Han Y, Zhao J, Liu H, Liu Y. Compared of efficacy and safety of high-dose donepezil vs standard-dose donepezil among elderly patients with Alzheimer's disease: a systematic review and meta-analysis. *Expert Opin Drug Saf*. 2022;21(3):407–15. <https://doi.org/10.1080/14740338.2022.2027905>.
286. Howard R, McShane R, Lindesay J, Ritchie C, Baldwin A, Barber R, et al. Donepezil and memantine for moderate-to-severe Alzheimer's disease. *N Engl J Med*. 2012;366(10):893–903. <https://doi.org/10.1056/NEJMoa1106668>.
287. Tariot PN, Cummings JL, Katz IR, Mintzer J, Perdomo CA, Schwam EM, et al. A randomized, double-blind, placebo-controlled study of the efficacy and safety of donepezil in patients with Alzheimer's disease in the nursing home setting. *J Am Geriatr Soc*. 2001;49(12):1590–9.
288. Ye CY, Lei Y, Tang XC, Zhang HY. Donepezil attenuates A β -associated mitochondrial dysfunction and reduces mitochondrial A β accumulation in vivo and in vitro. *Neuropharmacology*. 2015;95:29–36. <https://doi.org/10.1016/j.neuropharm.2015.02.020>.
289. Dong H, Yuede CM, Coughlan CA, Murphy KM, Csernansky JG. Effects of donepezil on amyloid- β and synapse density in the Tg2576 mouse model of Alzheimer's disease. *Brain Res*. 2009;1303:169–78. <https://doi.org/10.1016/j.brainres.2009.09.097>.
290. Ishibashi K, Miura Y, Wagatsuma K, Ishiwata K, Ishii K. Changes in brain amyloid- β accumulation after donepezil administration. *J Clin Neurosci*. 2017;45:328–9. <https://doi.org/10.1016/j.jocn.2017.08.025>.
291. Howard RJ, Juszcak E, Ballard CG, Bentham P, Brown RG, Bullock R, et al. Donepezil for the treatment of agitation in Alzheimer's disease. *N Engl J Med*. 2007;357(14):1382–92. <https://doi.org/10.1056/NEJMoa066583>.
292. De Keyser J, Sulter G, Luiten PG. Clinical trials with neuroprotective drugs in acute ischaemic stroke: are we doing the right thing? *Trends Neurosci*. 1999;22(12):535–40. [https://doi.org/10.1016/s0166-2236\(99\)01463-0](https://doi.org/10.1016/s0166-2236(99)01463-0).
293. Modell S, Naber D, Holzbach R. Efficacy and safety of an opiate sigma-receptor antagonist (SL 82.0715) in schizophrenic patients with negative symptoms: an open dose-range study. *Pharmacopsychiatry*. 1996;29(02):63–6. <https://doi.org/10.1055/s-2007-979546>.
294. Hashimoto K, London ED. Interactions of erythro-ifenprodil, threo-ifenprodil, erythro-iodoifenprodil, and eliprodil with subtypes of σ receptors. *Eur J Pharmacol*. 1995;273(3):307–10. [https://doi.org/10.1016/0014-2999\(94\)00763-w](https://doi.org/10.1016/0014-2999(94)00763-w).
295. Brooks S, Kaur S, Starr BS, Starr MS. Motor actions of eliprodil in the normal and monoamine-depleted mouse: a role in the treatment of Parkinson's disease? *J Neural Transm (Vienna)*. 1996;103(6):737–48. <https://doi.org/10.1007/bf01271233>.
296. Demerens C, Stankoff B, Zalc B, Lubetzki C. Eliprodil stimulates CNS myelination: new prospects for multiple sclerosis? *Neurology*. 1999;52(2):346. <https://doi.org/10.1212/wnl.52.2.346>.
297. Silkina I, Gan'shina T, Seredin S, Mirzoiyan R. Gabaergic mechanism of cerebrovascular and neuroprotective effects of afobazole and picamilon. *E Eksp Klin Farmakol*. 2005;68(1):20–4.
298. Antipova T, Sapozhnikova D, Liu B, Seredenin S. Selective anxiolytic afobazole increases the content of BDNF and NGF in cultured hippocampal HT-22 line neurons. *Eksp Klin Farmakol*. 2009;72(1):12–4.
299. Cuevas J, Rodriguez A, Behensky A, Katnik C. Afobazole modulates microglial function via activation of both σ -1 and σ -2 receptors. *J Pharmacol Exp Ther*. 2011;339(1):161–72. <https://doi.org/10.1124/jpet.111.182816>.
300. Behensky AA, Yasny IE, Shuster AM, Seredenin SB, Petrov AV, Cuevas J. Stimulation of sigma receptors with afobazole blocks activation of microglia and reduces toxicity caused by amyloid- β 25–35. *J Pharmacol Exp Ther*. 2013;347(2):458–67. <https://doi.org/10.1124/jpet.113.208348>.
301. Behensky AA, Yasny IE, Shuster AM, Seredenin SB, Petrov AV, Cuevas J. Afobazole activation of σ -1 receptors modulates neuronal responses to amyloid- β 25–35. *J Pharmacol Exp Ther*. 2013;347(2):468–77. <https://doi.org/10.1124/jpet.113.208330>.

302. Martin P, de-Witte PAM, Maurice T, Gammaitoni A, Farfel G, Galer B. Fenfluramine acts as a positive modulator of sigma-1 receptors. *Epilepsy Behav.* 2020;105:106989. <https://doi.org/10.1016/j.yebeh.2020.106989>.
303. Maurice T. Bi-phasic dose response in the preclinical and clinical developments of sigma-1 receptor ligands for the treatment of neurodegenerative disorders. *Expert Opin Drug Discov.* 2021;16(4):373–89. <https://doi.org/10.1080/17460441.2021.1838483>.
304. Knupp KG, Scheffer IE, Ceulemans B, Sullivan JE, Nickels KC, Lagae L, et al. Efficacy and safety of fenfluramine for the treatment of seizures associated with Lennox-Gastaut syndrome: a randomized clinical trial. *JAMA Neurol.* 2022;79(6):554–64. <https://doi.org/10.1001/jamaneurol.2022.0829>.
305. Lagae L, Sullivan J, Knupp K, Laux L, Polster T, Nikanorova M, et al. Fenfluramine hydrochloride for the treatment of seizures in Dravet syndrome: a randomised, double-blind, placebo-controlled trial. *Lancet.* 2019;394(10216):2243–54. [https://doi.org/10.1016/S0140-6736\(19\)32500-0](https://doi.org/10.1016/S0140-6736(19)32500-0).
306. Specchio N, Pietrafusa N, Doccini V, Trivisano M, Darra F, Ragona F, et al. Efficacy and safety of Fenfluramine hydrochloride for the treatment of seizures in Dravet syndrome: a real-world study. *Epilepsia.* 2020;61(11):2405–14. <https://doi.org/10.1111/epi.16690>.
307. Chomchoei C, Brimson J, Brimson S. Repurposing fluoxetine to treat lymphocytic leukaemia: apoptosis induction, sigma-1 receptor upregulation, inhibition of IL-2 cytokine production, and autophagy induction. *Expert Opin Ther Targets.* 2023. <https://doi.org/10.1080/14728222.2022.2166829>.
308. Németh ZK, Szűcs A, Vitrai J, Juhász D, Németh JP, Holló A. Fluoxetine use is associated with improved survival of patients with COVID-19 pneumonia: a retrospective case-control study. *Ideggyogy Sz.* 2021;74(11–12):389–96. <https://doi.org/10.18071/isz.74.0389>.
309. Emilia M, Rosso D, Laura M, María A. Immunomodulatory effects of fluoxetine: a new potential pharmacological action for a classic antidepressant drug? *Pharmacol Res.* 2016;109:101–7. <https://doi.org/10.1016/j.phrs.2015.11.021>.
310. Safrany ST, Brimson JM. Are fluoxetine's effects due to sigma-1 receptor agonism? *Pharmacol Res.* 2016;113:707–8. <https://doi.org/10.1016/j.phrs.2016.05.031>.
311. Mostert JP, Admiraal-Behloul F, Hoogduin JM, Luyendijk J, Heersema DJ, Van Buchem MA, et al. Effects of fluoxetine on disease activity in relapsing multiple sclerosis: a double-blind, placebo-controlled, exploratory study. *J Neurol Neurosurg Psychiatry.* 2008;79(9):1027–31. <https://doi.org/10.1136/jnnp.2007.139345>.
312. Lee JY, Kang SR, Yune TY. Fluoxetine prevents oligodendrocyte cell death by inhibiting microglia activation after spinal cord injury. *J Neurotrauma.* 2015;32(9):633–44. <https://doi.org/10.1089/neu.2014.3527>.
313. Sijens PE, Mostert JP, Irwan R, Potze JH, Oudkerk M, De Keyser J. Impact of fluoxetine on the human brain in multiple sclerosis as quantified by proton magnetic resonance spectroscopy and diffusion tensor imaging. *Psychiatry Res.* 2008;164(3):274–82. <https://doi.org/10.1016/j.psychres.2007.12.014>.
314. Cambon M, Mostert J, D'Hooghe M, Nagels G, Willekens B, Debruyne J, et al. Fluoxetine in progressive multiple sclerosis: the FLUOX-PMS trial. *Mult Scler J.* 2019;25(13):1728–35. <https://doi.org/10.1177/1352458519843051>.
315. Connick P, De Angelis F, Parker RA, Plantone D, Doshi A, John N, et al. Multiple Sclerosis-Secondary Progressive Multi-Arm Randomisation Trial (MS-SMART): a multiarm phase IIb randomised, double-blind, placebo-controlled clinical trial comparing the efficacy of three neuroprotective drugs in secondary progressive multiple sclerosis. *BMJ Open.* 2018;8(8):e021944. <https://doi.org/10.1136/bmjopen-2018-021944>.
316. De Angelis F, Connick P, Parker RA, Plantone D, Doshi A, John N, et al. Amiloride, fluoxetine or riluzole to reduce brain volume loss in secondary progressive multiple sclerosis: the MS-SMART four-arm RCT. *Efficacy Mech Eval.* 2020;7(3):1–72. <https://doi.org/10.3310/eme07030>.
317. Chataway J, De Angelis F, Connick P, Parker RA, Plantone D, Doshi A, et al. Efficacy of three neuroprotective drugs in secondary progressive multiple sclerosis (MS-SMART): a phase 2b, multiarm, double-blind, randomised placebo-controlled trial. *Lancet Neurol.* 2020;19(3):214–25. [https://doi.org/10.1016/S1474-4422\(19\)30485-5](https://doi.org/10.1016/S1474-4422(19)30485-5).
318. Brimson JM, Prasanth MI, Malar DS, Brimson S, Thitilertdech P, Tencomnao T. Drugs that offer the potential to reduce hospitalization and mortality from SARS-CoV-2 infection: the possible role of the Sigma-1 receptor and autophagy. *Expert Opin Ther Targets.* 2021;25(6):35–449. <https://doi.org/10.1080/14728222.2021.1952987>.
319. Nishimura T, Ishima T, Iyo M, Hashimoto K. Potentiation of nerve growth factor-induced neurite outgrowth by fluvoxamine: role of sigma-1 receptors, IP3 receptors and cellular signaling pathways. *PLoS ONE.* 2008;3(7):e2558. <https://doi.org/10.1371/journal.pone.0002558>.
320. Benedetti F, Campori E, Colombo C, Smeraldi E. Fluvoxamine treatment of major depression associated with multiple sclerosis. *J Neuropsychiatry Clin Neurosci.* 2004;16(3):364–6. <https://doi.org/10.1176/jnp.16.3.364>.
321. Ghareghani M, Zibara K, Sadeghi H, Dokoohaki S, Sadeghi H, Aryanpour R, et al. Fluvoxamine stimulates oligodendrogenesis of cultured neural stem cells and attenuates inflammation and demyelination in an animal model of multiple sclerosis. *Sci Rep.* 2017;7(1):4923. <https://doi.org/10.1038/s41598-017-04968-z>.
322. Robson MJ, Elliott M, Seminerio MJ, Matsumoto RR. Evaluation of sigma (σ) receptors in the antidepressant-like effects of ketamine in vitro and in vivo. *Eur Neuropsychopharmacol.* 2012;22(4):308–17. <https://doi.org/10.1016/j.euroneuro.2011.08.002>.
323. Nikiforuk A, Popik P. Ketamine prevents stress-induced cognitive inflexibility in rats. *Psychoneuroendocrinology.* 2014;40:119–22. <https://doi.org/10.1016/j.psyneuen.2013.11.009>.
324. Shibakawa YS, Sasaki Y, Goshima Y, Echigo N, Kamiya Y, Kurahashi K, et al. Effects of ketamine and propofol on inflammatory responses of primary glial cell cultures stimulated with lipopolysaccharide. *Br J Anaesth.* 2005;95(6):803–10. <https://doi.org/10.1093/bja/aei256>.
325. Hollinger A, Rüst CA, Riegger H, Gysi B, Tran F, Brügger J, et al. Ketamine vs haloperidol for prevention of cognitive dysfunction and postoperative delirium: a phase IV multicentre randomised placebo-controlled double-blind clinical trial. *J Clin Anesth.* 2021;68:110099. <https://doi.org/10.1016/j.jclinane.2020.110099>.
326. Lara DR, Bisol LW, Munari LR. Antidepressant, mood stabilizing and procognitive effects of very low dose sublingual ketamine in refractory unipolar and bipolar depression. *Int J Neuropsychopharmacol.* 2013;16(9):2111–7. <https://doi.org/10.1017/S1461145713000485>.
327. Zhang MW, Ho RC. Controversies of the effect of ketamine on cognition. *Front Psychiatry.* 2016;7:47. <https://doi.org/10.3389/fpsy.2016.00047>.
328. Ownby RL, Crocco E, Acevedo A, John V, Loewenstein D. Depression and risk for Alzheimer disease: systematic review, meta-analysis, and metaregression analysis. *Arch Gen Psychiatry.* 2006;63(5):530–8. <https://doi.org/10.1001/archpsyc.63.5.530>.

329. Aan Het Rot M, Zarate CA, Charney DS, Mathew SJ. Ketamine for depression: where do we go from here? *Biol Psychiatry*. 2012;72(7):537–47. <https://doi.org/10.1016/j.biopsych.2012.05.003>.
330. Rocha FL, de Vasconcelos Cunha UG, Paschoalin RC, Hara C, Thomaz DP. Use of subcutaneous ketamine to rapidly improve severe treatment-resistant depression in a patient with Alzheimer's disease. *Int Clin Psychopharmacol*. 2021;36(2):104–5. <https://doi.org/10.1097/YIC.0000000000000334>.
331. He Y, Li H, Huang J, Huang S, Bai Y, Li Y, et al. Efficacy of antidepressant drugs in the treatment of depression in Alzheimer disease patients: a systematic review and network meta-analysis. *J Psychopharmacol*. 2021;35(8):901–9. <https://doi.org/10.1177/02698811211030181>.
332. Yeung L, Wai MS, Fan M, Mak Y, Lam W, Li Z, et al. Hyperphosphorylated tau in the brains of mice and monkeys with long-term administration of ketamine. *Toxicol Lett*. 2010;193(2):189–93. <https://doi.org/10.1016/j.toxlet.2010.01.008>.
333. Ferro M, Angelucci M, Anselmo-Franci J, Canteras N, Da Cunha C. Neuroprotective effect of ketamine/xylazine on two rat models of Parkinson's disease. *Braz J Med Biol Res*. 2007;40:89–96. <https://doi.org/10.1590/s0100-879x2007000100012>.
334. Vecchia DD, Kanazawa LKS, Wendler E, Hocayen PdAS, Vital MABF, Takahashi RN, et al. Ketamine reversed short-term memory impairment and depressive-like behavior in animal model of Parkinson's disease. *Brain Res Bull*. 2021;168:63–73. <https://doi.org/10.1016/j.brainresbull.2020.12.011>.
335. Fan J-C, Song J-J, Wang Y, Chen Y, Hong D-X. Neuron-protective effect of subanesthetic-dosage ketamine on mice of Parkinson's disease. *Asian Pac J Trop Med*. 2017;10(10):1007–10. <https://doi.org/10.1016/j.apjtm.2017.09.014>.
336. Wright JJ, Goodnight PD, McEvoy MD. The utility of ketamine for the preoperative management of a patient with Parkinson's disease. *Anesth Analg*. 2009;108(3):980–2. <https://doi.org/10.1213/ane.0b013e3181924025>.
337. Salazar G, Motamed C. A remifentanyl/ketamine sedation in surgical cancer patients having severe Parkinson's disease: two case reports. *J Opioid Manag*. 2012;8(2):133–4. <https://doi.org/10.5055/jom.2012.0106>.
338. Barenboim I, Lafer B. Maintenance use of ketamine for treatment-resistant depression: an open-label pilot study. *Braz J Psychiatry*. 2018;40(1):110. <https://doi.org/10.1590/1516-4446-2017-2380>.
339. Mathew SJ, Shah A, Lapidus K, Clark C, Jarun N, Ostermeyer B, et al. Ketamine for treatment-resistant unipolar depression. *CNS Drugs*. 2012;26(3):189–204. <https://doi.org/10.2165/11599770-000000000-00000>.
340. Feifel D. Breaking sad: unleashing the breakthrough potential of ketamine's rapid antidepressant effects. *Drug Dev Res*. 2016;77(8):489–94. <https://doi.org/10.1002/ddr.21347>.
341. Shiroma PR, Johns B, Kuskowski M, Wels J, Thuras P, Albott CS, et al. Augmentation of response and remission to serial intravenous subanesthetic ketamine in treatment resistant depression. *J Affect Disord*. 2014;155:123–9. <https://doi.org/10.1016/j.jad.2013.10.036>.
342. Wang X, Chang L, Tan Y, Qu Y, Shan J, Hashimoto K. (R)-ketamine ameliorates the progression of experimental autoimmune encephalomyelitis in mice. *Brain Res Bull*. 2021;177:316–23. <https://doi.org/10.1016/j.brainresbull.2021.10.013>.
343. Wang X, Chang L, Wan X, Tan Y, Qu Y, Shan J, et al. (R)-ketamine ameliorates demyelination and facilitates remyelination in cuprizone-treated mice: a role of gut–microbiota–brain axis. *Neurobiol Dis*. 2022;165:105635. <https://doi.org/10.1016/j.nbd.2022.105635>.
344. Sakai T, Tomiyasu S, Ono T, Yamada H, Sumikawa K. Multiple sclerosis with severe pain and allodynia alleviated by oral ketamine. *Clin J Pain*. 2004;20(5):375–6. <https://doi.org/10.1097/00002508-200409000-00016>.
345. Messer MM, Haller IV. Ketamine therapy for treatment-resistant depression in a patient with multiple sclerosis: a case report. *Innov Clin Neurosci*. 2017;14(1–2):56–9.
346. Patrizi A, Picard N, Simon AJ, Gunner G, Centofante E, Andrews NA, et al. Chronic administration of the N-methyl-D-aspartate receptor antagonist ketamine improves Rett syndrome phenotype. *Biol Psychiatry*. 2016;79(9):755–64. <https://doi.org/10.1016/j.biopsych.2015.08.018>.
347. Kron M, Howell CJ, Adams IT, Ransbottom M, Christian D, Ogier M, et al. Brain activity mapping in Mecp2 mutant mice reveals functional deficits in forebrain circuits, including key nodes in the default mode network, that are reversed with ketamine treatment. *J Neurosci*. 2012;32(40):13860–72. <https://doi.org/10.1523/jneurosci.2159-12.2012>.
348. Jolly T, McLean HS. Use of ketamine during procedural sedation: indications, controversies, and side effects. *J Infus Nurs*. 2012;35(6):377–82. <https://doi.org/10.1097/NAN.0b013e31827068c1>.
349. Mount C, Downton C. Alzheimer disease: progress or profit? *Nat Med*. 2006;12(7):780–4. <https://doi.org/10.1038/nm0706-780>.
350. Guo J, Wang Z, Liu R, Huang Y, Zhang N, Zhang R. Memantine, donepezil, or combination therapy—what is the best therapy for Alzheimer's disease? A network meta-analysis. *Brain Behav*. 2020;10(11):e01831. <https://doi.org/10.1002/brb3.1831>.
351. Schmitt HP. On the paradox of ion channel blockade and its benefits in the treatment of Alzheimer disease. *Med Hypotheses*. 2005;65(2):259–65. <https://doi.org/10.1016/j.mehy.2005.03.011>.
352. Parsons C, Danysz W, Quack G. Memantine is a clinically well tolerated N-methyl-D-aspartate (NMDA) receptor antagonist—a review of preclinical data. *Neuropharmacol*. 1999;38(6):735–67. [https://doi.org/10.1016/s0028-3908\(99\)00019-2](https://doi.org/10.1016/s0028-3908(99)00019-2).
353. Rogawski MA, Wenk GL. The neuropharmacological basis for the use of memantine in the treatment of Alzheimer's disease. *CNS Drug Rev*. 2003;9(3):275–308. <https://doi.org/10.1111/j.1527-3458.2003.tb00254.x>.
354. Martina M, Turcotte MEB, Halman S, Bergeron R. The sigma-1 receptor modulates NMDA receptor synaptic transmission and plasticity via SK channels in rat hippocampus. *J Physiol*. 2007;578(1):143–57. <https://doi.org/10.1113/jphysiol.2006.116178>.
355. Chaki S, Okuyama S, Ogawa S-i, Tomisawa K. Regulation of NMDA-induced [3H] dopamine release from rat hippocampal slices through sigma-1 binding sites. *Neurochem Int*. 1998;33(1):29–34. <https://doi.org/10.1113/jphysiol.2006.116178>.
356. Sha S, Qu WJ, Li L, Lu ZH, Chen L, Yu WF, et al. Sigma-1 receptor knockout impairs neurogenesis in dentate gyrus of adult hippocampus via down-regulation of NMDA receptors. *CNS Neurosci Ther*. 2013;19(9):705–13. <https://doi.org/10.1111/cns.12129>.
357. Pabba M, Wong AY, Ahlskog N, Hristova E, Biscaro D, Nassrallah W, et al. NMDA receptors are upregulated and trafficked to the plasma membrane after sigma-1 receptor activation in the rat hippocampus. *J Neurosci*. 2014;34(34):11325–38. <https://doi.org/10.1523/JNEUROSCI.0458-14.2014>.
358. Stahl SM. Mechanism of action of ketamine. *CNS Spectr*. 2013;18(4):171–4. <https://doi.org/10.1017/S109285291300045X>.
359. Winblad B, Jones RW, Wirth Y, Stöfler A, Möbius HJ. Memantine in moderate to severe Alzheimer's disease: a meta-analysis of randomised clinical trials. *Dement Geriatr Cogn Disord*. 2007;24(1):20–7. <https://doi.org/10.1159/000102568>.
360. Atri A, Molinuevo JL, Lemming O, Wirth Y, Pulte I, Wilkinson D. Memantine in patients with Alzheimer's disease receiving donepezil: new analyses of efficacy and safety for combination

- therapy. *Alzheimers Res Ther.* 2013;5(1):6. <https://doi.org/10.1186/alzrt160>.
361. Schmitt FA, van Dyck CH, Wichems CH, Olin JT, Group MM-M-S. Cognitive response to memantine in moderate to severe Alzheimer disease patients already receiving donepezil: an exploratory reanalysis. *Alzheimer Dis Assoc Disord.* 2006;20(4):255–62. <https://doi.org/10.1097/01.wad.0000213860.35355.d4>.
 362. Lopez OL, Becker JT, Wahed AS, Saxton J, Sweet RA, Wolk DA, et al. Long-term effects of the concomitant use of memantine with cholinesterase inhibition in Alzheimer disease. *J Neurol Neurosurg Psychiatry.* 2009;80(6):600–7. <https://doi.org/10.1136/jnnp.2008.158964>.
 363. Tariot PN, Farlow MR, Grossberg GT, Graham SM, McDonald S, Gergel I, et al. Memantine treatment in patients with moderate to severe Alzheimer disease already receiving donepezil: a randomized controlled trial. *JAMA.* 2004;291(3):317–24. <https://doi.org/10.1001/jama.291.3.317>.
 364. Hendrix S, Ellison N, Stanworth S, Otcheretko V, Tariot PN. Post Hoc Evidence for an additive effect of memantine and donepezil: consistent findings from DOMINO-AD Study and Memantine Clinical Trial Program. *J Prev Alzheimers Dis.* 2015;2(3):165–71. <https://doi.org/10.14283/jpad.2015.66>.
 365. Porsteinsson AP, Grossberg GT, Mintzer J, Olin JT. Memantine treatment in patients with mild to moderate Alzheimer's disease already receiving a cholinesterase inhibitor: a randomized, double-blind, placebo-controlled trial. *Curr Alzheimer Res.* 2008;5(1):83–9. <https://doi.org/10.2174/156720508783884576>.
 366. Leroi I, Overshott R, Byrne EJ, Daniel E, Burns A. Randomized controlled trial of memantine in dementia associated with Parkinson's disease. *Mov Disord.* 2009;24(8):1217–21. <https://doi.org/10.1002/mds.22495>.
 367. Emre M, Tsolaki M, Bonuccelli U, Destée A, Tolosa E, Kutzelnigg A, et al. Memantine for patients with Parkinson's disease dementia or dementia with Lewy bodies: a randomised, double-blind, placebo-controlled trial. *Lancet Neurol.* 2010;9(10):969–77. [https://doi.org/10.1016/S1474-4422\(10\)70194-0](https://doi.org/10.1016/S1474-4422(10)70194-0).
 368. Ondo WG, Shinawi L, Davidson A, et al. Memantine for non-motor features of Parkinson's disease: a double-blind placebo controlled exploratory pilot trial. *Parkinsonism Relat Disord.* 2011;17(3):156–9. <https://doi.org/10.1016/j.parkreldis.2010.12.003>.
 369. Moreau C, Delval A, Tiffreau V, et al. Memantine for axial signs in Parkinson's disease: a randomised, double-blind, placebo-controlled pilot study. *J Neurol Neurosurg Psychiatry.* 2013;84(5):552–5. <https://doi.org/10.1136/jnnp-2012-303182>.
 370. Wang HF, Yu JT, Tang SW, Jiang T, Tan CC, Meng XF, et al. Efficacy and safety of cholinesterase inhibitors and memantine in cognitive impairment in Parkinson's disease, Parkinson's disease dementia, and dementia with Lewy bodies: systematic review with meta-analysis and trial sequential analysis. *J Neurol Neurosurg Psychiatry.* 2015;86(2):135–43. <https://doi.org/10.1136/jnnp-2014-307659>.
 371. Sanjay M, Amir G, John R, Scott O, Stephen D, et al. Short-term tolerability of a nonazapirone selective serotonin 1A agonist in adults with generalized anxiety disorder: a 28-day, open-label study. *Clin Ther.* 2008;30:9. <https://doi.org/10.1016/j.clinthera.2008.09.006>.
 372. Rickels K, Mathew S, Banov MD, Zimbroff DL, Oshana S, Parsons EC Jr, et al. Effects of PRX-00023, a novel, selective serotonin 1A receptor agonist on measures of anxiety and depression in generalized anxiety disorder: results of a double-blind, placebo-controlled trial. *J Clin Psychopharmacol.* 2008;28(2):235–9. <https://doi.org/10.1097/JCP.0b013e31816774de>.
 373. PRX-00023 Therapy in localization-related epilepsy. In: NIH U.S. National Library of Medicine clinical trials web site; 2011. <https://clinicaltrials.gov/ct2/show/NCT01281956>. Accessed 1 Dec 2022.
 374. Merritt HH, Putnam TJ. Sodium diphenyl hydantoinate in the treatment of convulsive disorders. *J Am Med Assoc.* 1938;111(12):1068–73. <https://doi.org/10.1001/jama.1938.02790380010004>.
 375. Rogawski MA, Löscher W. The neurobiology of antiepileptic drugs. *Nat Rev Neurosci.* 2004;5(7):553–64. <https://doi.org/10.1038/nrn1430>.
 376. Tchedre KT, Huang RQ, Dibas A, Krishnamoorthy RR, Dillon GH, Yorio T. Sigma-1 receptor regulation of voltage-gated calcium channels involves a direct interaction. *Invest Ophthalmol Vis Sci.* 2008;49(11):4993–5002. <https://doi.org/10.1167/iov.08-1867>.
 377. Mueller BH II, Park Y, Daudt DR III, Ma H-Y, Akopova I, Stankowska DL, et al. Sigma-1 receptor stimulation attenuates calcium influx through activated L-type voltage gated calcium channels in purified retinal ganglion cells. *Exp Eye Res.* 2013;107:21–31. <https://doi.org/10.1016/j.exer.2012.11.002>.
 378. Zhang K, Zhao Z, Lan L, Wei X, Wang L, Liu X, et al. Sigma-1 receptor plays a negative modulation on N-type calcium channel. *Front Pharmacol.* 2017;8:302. <https://doi.org/10.3389/fphar.2017.00302>.
 379. Zhang H, Cuevas J. Sigma receptors inhibit high-voltage-activated calcium channels in rat sympathetic and parasympathetic neurons. *J Neurophysiol.* 2002;87(6):2867–79. <https://doi.org/10.1152/jn.2002.87.6.2867>.
 380. DeHaven-Hudkins D, Ford-Rice F, Allen J, Hudkins R. Allosteric modulation of ligand binding to [3H](+) pentazocine-defined σ recognition sites by phenytoin. *Life Sci.* 1993;53(1):41–8. [https://doi.org/10.1016/0024-3205\(93\)90609-7](https://doi.org/10.1016/0024-3205(93)90609-7).
 381. Braakman HM, Verhoeven JS, Erasmus CE, Haaxma CA, Willemssen MH, Schelhaas HJ. Phenytoin as a last-resort treatment in SCN 8A encephalopathy. *Epilepsia Open.* 2017;2(3):343–4. <https://doi.org/10.1002/epi4.12059>.
 382. Dilena R, Striano P, Gennaro E, Bassi L, Olivotto S, Tadini L, et al. Efficacy of sodium channel blockers in SCN2A early infantile epileptic encephalopathy. *Brain Dev.* 2017;39(4):345–8. <https://doi.org/10.1016/j.braindev.2016.10.015>.
 383. Su TP, Wu XZ, Cone EJ, Shukla K, Gund TM, Dodge AL, et al. Sigma compounds derived from phencyclidine: identification of PRE-084, a new, selective sigma ligand. *J Pharmacol Exp Ther.* 1991;259(2):543–50.
 384. Motawe ZY, Abdelmaboud SS, Cuevas J, Breslin JW. PRE-084 as a tool to uncover potential therapeutic applications for selective sigma-1 receptor activation. *Int J Biochem Cell Biol.* 2020;126:105803; [Doi: https://doi.org/10.1016/j.biocel.2020.105803](https://doi.org/10.1016/j.biocel.2020.105803)
 385. Motawe ZY, Farsaei F, Abdelmaboud SS, Cuevas J, Breslin JW. Sigma-1 receptor activation-induced glycolytic ATP production and endothelial barrier enhancement. *Microcirculation.* 2020;27(6):e12620. <https://doi.org/10.1111/micc.12620>.
 386. An Y, Qi Y, Li Y, Li Z, Yang C, Jia D. Activation of the sigma-1 receptor attenuates blood-brain barrier disruption by inhibiting amyloid deposition in Alzheimer's disease mice. *Neurosci Lett.* 2022;774:136528. <https://doi.org/10.1016/j.neulet.2022.136528>.
 387. Penas C, Pascual-Font A, Mancuso R, Forés J, Casas C, Navarro X. Sigma receptor agonist 2-(4-morpholinethyl) 1 phenylcyclohexanecarboxylate (Pre084) increases GDNF and BiP expression and promotes neuroprotection after root avulsion injury. *J Neurotrauma.* 2011;28(5):831–40. <https://doi.org/10.1089/neu.2010.1674>.

388. Francardo V, Bez F, Wieloch T, Nissbrandt H, Ruscher K, Cenci MA. Pharmacological stimulation of sigma-1 receptors has neurorestorative effects in experimental parkinsonism. *Brain*. 2014;137(7):1998–2014. <https://doi.org/10.1093/brain/awu107>.
389. Rao YL, Ganaraja B, Murlimanju BV, Joy T, Krishnamurthy A, Agrawal A. Hippocampus and its involvement in Alzheimer's disease: a review. *3 Biotech*. 2022;12(2):55; Doi: <https://doi.org/10.1007/s13205-022-03123-4>
390. Eriksson PS, Perfilieva E, Björk-Eriksson T, Alborn A-M, Nordborg C, Peterson DA, et al. Neurogenesis in the adult human hippocampus. *Nat Med*. 1998;4(11):1313–7. <https://doi.org/10.1038/3305>.
391. Borbély E, Varga V, Szögi T, Schuster I, Bozsó Z, Penke B, et al. Impact of two neuronal Sigma-1 receptor modulators, PRE084 and DMT, on neurogenesis and neuroinflammation in an Aβ1–42-injected, wild-type mouse model of AD. *Int J Mol Sci*. 2022;23(5):2514. <https://doi.org/10.3390/ijms23052514>.
392. Li L, Xu B, Zhu Y, Chen L, Sokabe M, Chen L. DHEA prevents Aβ25–35-impaired survival of newborn neurons in the dentate gyrus through a modulation of PI3K-Akt-mTOR signaling. *Neuropharmacol*. 2010;59(4–5):323–33. <https://doi.org/10.1016/j.neuropharm.2010.02.009>.
393. Estévez-Silva HM, Cuesto G, Romero N, Brito-Armas JM, Acevedo-Aroza A, Acebes Á, et al. Pridopidine promotes synaptogenesis and reduces spatial memory deficits in the Alzheimer's disease APP/PS1 mouse model. *Neurotherapeutics*. 2022;19(5):1566–87. <https://doi.org/10.1007/s13311-022-01280-1>.
394. Guzmán-Lenis MS, Navarro X, Casas C. Selective sigma receptor agonist 2-(4-morpholinethyl) 1-phenylcyclohexanecarboxylate (PRE084) promotes neuroprotection and neurite elongation through protein kinase C (PKC) signaling on motoneurons. *Neuroscience*. 2009;162(1):31–8. <https://doi.org/10.1016/j.neuroscience.2009.03.067>.
395. Mancuso R, Oliván S, Rando A, Casas C, Osta R, Navarro X. Sigma-1R agonist improves motor function and motoneuron survival in ALS mice. *Neurotherapeutics*. 2012;9(4):814–26. <https://doi.org/10.1007/s13311-012-0140-y>.
396. Mancuso R, Del Valle J, Morell M, Pallás M, Osta R, Navarro X. Lack of synergistic effect of resveratrol and sigma-1 receptor agonist (PRE-084) in SOD1G93A ALS mice: overlapping effects or limited therapeutic opportunity? *Orphanet J Rare Dis*. 2014;9(1):1–11. <https://doi.org/10.1186/1750-1172-9-78>.
397. Gaja-Capdevila N, Hernández N, Navarro X, Herrando-Grabulosa M. Sigma-1 receptor is a pharmacological target to promote neuroprotection in the SOD1G93A ALS mice. *Front Pharmacol*. 2021;12:780588. <https://doi.org/10.3389/fphar.2021.780588>.
398. Peviani M, Salvaneschi E, Bontempi L, Petese A, Manzo A, Rossi D, et al. Neuroprotective effects of the Sigma-1 receptor (S1R) agonist PRE-084, in a mouse model of motor neuron disease not linked to SOD1 mutation. *Neurobiol Dis*. 2014;62:218–32. <https://doi.org/10.1016/j.nbd.2013.10.010>.
399. Barwick SR, Siddiq MS, Wang J, Xiao H, Marshall B, Perry E, et al. Sigma 1 receptor co-localizes with NRF2 in retinal photoreceptor cells. *Antioxidants*. 2021;10(6):981. <https://doi.org/10.3390/antiox10060981>.
400. Weng TY, Hung DT, Su TP, Tsai SYA. Loss of Sigma-1 receptor chaperone promotes astrocytosis and enhances the Nrf2 antioxidant defense. *Oxid Med Cell Longev*. 2017;2017:4582135. <https://doi.org/10.1155/2017/4582135>.
401. Lasbleiz C, Peyrel A, Tarot P, Sarniguet J, Cruzier L, Cubedo N, et al. Sigma-1 receptor agonist PRE-084 confers protection against TAR DNA-binding protein-43 toxicity through NRF2 signalling. *Redox Biol*. 2022;58:102542. <https://doi.org/10.1016/j.redox.2022.102542>.
402. Hyrskyluoto A, Pulli I, Törnqvist K, Ho H, Korhonen L, Lindholm D. Sigma-1 receptor agonist PRE084 is protective against mutant huntingtin-induced cell degeneration: involvement of calpastatin and the NF-κB pathway. *Cell Death Dis*. 2013;4(5):e646. <https://doi.org/10.1038/cddis.2013.170>.
403. Bol'Shakova A, Kraskovskaya N, Gainullina A, Kukanova E, Vlasova O, Bezprozvanny I. Neuroprotective effect of σ1-receptors on the cell model of Huntington's disease. *Bull Exp Biol Med*. 2017;164:252–8. <https://doi.org/10.1007/s10517-017-3968-7>.
404. Johnston TH, Geva M, Steiner L, Orbach A, Papapetropoulos S, Savola JM, et al. Pridopidine, a clinic-ready compound, reduces 3, 4-dihydroxyphenylalanine-induced dyskinesia in Parkinsonian macaques. *Mov Disord*. 2019;34(5):708–16. <https://doi.org/10.1002/mds.27565>.
405. Grachev ID, Meyer PM, Becker GA, Bronzel M, Marsteller D, Pastino G, et al. Sigma-1 and dopamine D2/D3 receptor occupancy of pridopidine in healthy volunteers and patients with Huntington disease: a [(18)F] fluspidine and [(18)F] fallypride PET study. *Eur J Nucl Med Mol Imaging*. 2021;48(4):1103–15. <https://doi.org/10.1007/s00259-020-05030-3>.
406. Sahlholm K, Sijbesma JWA, Maas B, Kwizera C, Marcellino D, Ramakrishnan NK, et al. Pridopidine selectively occupies sigma-1 rather than dopamine D2 receptors at behaviorally active doses. *Psychopharmacology*. 2015;232(18):3443–53. <https://doi.org/10.1007/s00213-015-3997-8>.
407. Bourne J, Harris KM. Do thin spines learn to be mushroom spines that remember? *Curr Opin Neurobiol*. 2007;17(3):381–6. <https://doi.org/10.1016/j.conb.2007.04.009>.
408. Bourne JN, Harris KM. Balancing structure and function at hippocampal dendritic spines. *Annu Rev Neurosci*. 2008;31:47–67. <https://doi.org/10.1146/annurev.neuro.31.060407.125646>.
409. Hayashi-Takagi A, Yagishita S, Nakamura M, Shirai F, Wu YI, Loshbaugh AL, et al. Labelling and optical erasure of synaptic memory traces in the motor cortex. *Nature*. 2015;525(7569):333–8. <https://doi.org/10.1038/nature15257>.
410. Ionescu A, Gradus T, Altman T, Maimon R, Avraham NS, Geva M, et al. Targeting the sigma-1 receptor via pridopidine ameliorates central features of ALS pathology in a SOD1G93A model. *Cell Death Dis*. 2019;10(3):210. <https://doi.org/10.1038/s41419-019-1451-2>.
411. Estévez-Silva HM, Mediavilla T, Giacobbo BL, Liu X, Sultan FR, Marcellino DJ. Pridopidine modifies disease phenotype in a SOD1 mouse model of amyotrophic lateral sclerosis. *Eur J Neurosci*. 2022;55(5):1356–72. <https://doi.org/10.1111/ejn.15608>.
412. Wang S-M, Wu H-E, Yasui Y, Geva M, Hayden M, Maurice T, et al. Nucleoporin POM121 signals TFEB-mediated autophagy via activation of SIGMAR1/sigma-1 receptor chaperone by pridopidine. *Autophagy*. 2023;19(1):126–51. <https://doi.org/10.1080/15548627.2022.2063003>.
413. Garcia-Miralles M, Geva M, Tan JY, Yusof NABM, Cha Y, Kusko R, et al. Early pridopidine treatment improves behavioral and transcriptional deficits in YAC128 Huntington disease mice. *JCI insight*. 2017;2:23. <https://doi.org/10.1172/jci.insight.95665>.
414. Nguyen KQ, Rymar VV, Sadikot AF. Impaired TrkB signaling underlies reduced BDNF-mediated trophic support of striatal neurons in the R6/2 mouse model of Huntington's disease. *Front Cell Neurosci*. 2016;10:37. <https://doi.org/10.3389/fncel.2016.00037>.
415. Kusko R, Dreyman J, Ross J, Cha Y, Escalante-Chong R, Garcia-Miralles M, et al. Large-scale transcriptomic analysis reveals that pridopidine reverses aberrant gene expression and activates neuroprotective pathways in the YAC128 HD mouse. *Mol Neurodegener*. 2018;13(1):25. <https://doi.org/10.1186/s13024-018-0259-3>.

416. Xie Y, Hayden MR, Xu B. BDNF overexpression in the fore-brain rescues Huntington's disease phenotypes in YAC128 mice. *J Neurosci*. 2010;30(44):14708–18. <https://doi.org/10.1523/JNEUROSCI.1637-10.2010>.
417. Geva M, Kusko R, Soares H, Fowler KD, Birnberg T, Barash S, et al. Pridopidine activates neuroprotective pathways impaired in Huntington disease. *Hum Mol Genet*. 2016;25(18):3975–87. <https://doi.org/10.1093/hmg/ddw238>.
418. Yagasaki Y, Numakawa T, Kumamaru E, Hayashi T, Su TP, Kunugi H. Chronic antidepressants potentiate via sigma-1 receptors the brain-derived neurotrophic factor-induced signaling for glutamate release. *J Biol Chem*. 2006;281(18):12941–9. <https://doi.org/10.1074/jbc.M508157200>.
419. Huntington Study Group HART Investigators. A randomized, double-blind, placebo-controlled trial of pridopidine in Huntington's disease. *Mov Disord*. 2013;28(10):1407–15. <https://doi.org/10.1002/mds.25362>.
420. De Yebenes JG, Landwehrmeyer B, Squitieri F, Reilmann R, Rosser A, Barker RA, et al. Pridopidine for the treatment of motor function in patients with Huntington's disease (MermaiHD): a phase 3, randomised, double-blind, placebo-controlled trial. *Lancet Neurol*. 2011;10(12):1049–57. [https://doi.org/10.1016/S1474-4422\(11\)70233-2](https://doi.org/10.1016/S1474-4422(11)70233-2).
421. Reilmann R, McGarry A, Grachev ID, Savola J-M, Borowsky B, Eyal E, et al. Safety and efficacy of pridopidine in patients with Huntington's disease (PRIDE-HD): a phase 2, randomised, placebo-controlled, multicentre, dose-ranging study. *Lancet Neurol*. 2019;18(2):165–76. [https://doi.org/10.1016/S1474-4422\(18\)30391-0](https://doi.org/10.1016/S1474-4422(18)30391-0).
422. McGarry A, Auinger P, Kieburz K, Geva M, Mehra M, Ablner V, et al. Additional safety and exploratory efficacy data at 48 and 60 months from Open-HART, an open-label extension study of pridopidine in Huntington disease. *J Huntingtons Dis*. 2020;9(2):173–84. <https://doi.org/10.3233/JHD-190393>.
423. Ponten H, Kullingsjö J, Sonesson C, Waters S, Waters N, Tedroff J. The dopaminergic stabilizer pridopidine decreases expression of l-DOPA-induced locomotor sensitisation in the rat unilateral 6-OHDA model. *Eur J Pharmacol*. 2013;698(1–3):278–85. <https://doi.org/10.1016/j.ejphar.2012.10.039>.
424. McFarthing K, Prakash N, Simuni T. Clinical trial highlights – dyskinesia. *J Parkinsons Dis*. 2019;9:449–65. <https://doi.org/10.3233/JPD-199002>.
425. Nadjar A, Gerfen CR, Bezdard E. Priming for l-dopa-induced dyskinesia in Parkinson's disease: a feature inherent to the treatment or the disease? *Prog Neurobiol*. 2009;87(1):1–9. <https://doi.org/10.1016/j.pneurobio.2008.09.013>.
426. Khoury R, Rajamanickam J, Grossberg GT. An update on the safety of current therapies for Alzheimer's disease: focus on rivastigmine. *Ther Adv Drug Saf*. 2018;9(3):171–8. <https://doi.org/10.1177/2042098617750555>.
427. Terada K, Migita K, Matsushima Y, Sugimoto Y, Kamei C, Matsumoto T, et al. Cholinesterase inhibitor rivastigmine enhances nerve growth factor-induced neurite outgrowth in PC12 cells via sigma-1 and sigma-2 receptors. *PLoS ONE*. 2018;13(12):e0209250. <https://doi.org/10.1371/journal.pone.0209250>.
428. Corey-Bloom J. A randomized trial evaluating the efficacy and safety of ENA 713 (rivastigmine tartrate), a new acetylcholinesterase inhibitor, in patients with mild to moderately severe Alzheimer's disease. *Int J Geriatr Psychiatry*. 1998;13:55–65.
429. Rösler M, Anand R, Cicin-Sain A, Gauthier S, Agid Y, Dal-Bianco P, et al. Efficacy and safety of rivastigmine in patients with Alzheimer's disease: international randomised controlled trial. *BMJ*. 1999;318(7184):633–8. <https://doi.org/10.1136/bmj.318.7184.633>.
430. Finkel SI. Effects of rivastigmine on behavioral and psychological symptoms of dementia in Alzheimer's disease. *Clin Ther*. 2004;26(7):980–90. [https://doi.org/10.1016/s0149-2918\(04\)90172-5](https://doi.org/10.1016/s0149-2918(04)90172-5).
431. Yi B, Sahn JJ, Ardestani PM, Evans AK, Scott LL, Chan JZ, et al. Small molecule modulator of sigma 2 receptor is neuroprotective and reduces cognitive deficits and neuroinflammation in experimental models of Alzheimer's disease. *J Neurochem*. 2017;140(4):561–75. <https://doi.org/10.1111/jnc.13917>.
432. Yano T, Tanabe H, Kobayashi K, Kobayashi H, Nabetani A, Sakai Y, et al. Sigma-1 receptor is a molecular target for novel neuroprotectant T-817MA. *Alzheimer Dement*. 2015;11(7):P861. <https://doi.org/10.1016/j.jalz.2015.08.038>.
433. Nguyen PTH, Kimura T, Ho SA, Tran AH, Ono T, Nishijo H. Ameliorative effects of a neuroprotective agent, T-817MA, on place learning deficits induced by continuous infusion of amyloid- β peptide (1–40) in rats. *Hippocampus*. 2007;17(6):443–55. <https://doi.org/10.1002/hipo.20281>.
434. Kimura T, Hong Nguyen PT, Ho SA, Tran AH, Ono T, Nishijo H. T-817MA, a neurotrophic agent, ameliorates the deficits in adult neurogenesis and spatial memory in rats infused icv with amyloid- β peptide. *Br J Pharmacol*. 2009;157(3):451–63. <https://doi.org/10.1111/j.1476-5381.2009.00141.x>.
435. Schneider LS, Thomas RG, Hendrix S, Rissman RA, Brewer JB, Salmon DP, et al. Safety and efficacy of edonergic maleate for patients with mild to moderate Alzheimer disease: a phase 2 randomized clinical trial. *JAMA Neurol*. 2019;76(11):1330–9. <https://doi.org/10.1001/jamaneurol.2019.1868>.
436. Schneider L, Porsteinsson A, Farlow M, Shimakura A, Nakagawa M, Iwakami N. The neuroprotective and neurotrophic agent T-817MA for Alzheimer's disease: randomized, double-blind, placebo-controlled proof-of-concept trial outcomes. *Alzheimers Dement*. 2013;9(4):P530–1. <https://doi.org/10.1016/j.jalz.2013.04.272>.
437. Audronytė E, Kaubrys G. Odor identification and discrimination as markers of early Alzheimer's disease. *Alzheimers Dement*. 2022;18:e063101. <https://doi.org/10.1002/alz.063101>.
438. Hudd F, Shiel A, Harris M, Bowdler P, Wood B, Tsivos D, et al. novel blood biomarkers that correlate with cognitive performance and hippocampal volumetry: potential for early diagnosis of Alzheimer's disease. *J Alzheimers Dis*. 2019;67(3):931–47. <https://doi.org/10.3233/JAD-180879>.
439. Pinto JM, Wroblewski KE, Kern DW, Schumm LP, McClintock MK. Olfactory dysfunction predicts 5-year mortality in older adults. *PLoS ONE*. 2014;9(10):e107541. <https://doi.org/10.1371/journal.pone.0107541>.
440. Adams DR, Kern DW, Wroblewski KE, McClintock MK, Dale W, Pinto JM. Olfactory dysfunction predicts subsequent dementia in older US adults. *J Am Geriatr Soc*. 2018;66(1):140–4. <https://doi.org/10.1111/jgs.15048>.
441. Wang W, Lee J, Harrou F, Sun Y. Early detection of Parkinson's disease using deep learning and machine learning. *IEEE Access*. 2020;8:147635–46. <https://doi.org/10.1109/ACCESS.2020.3016062>.
442. Lin C-H, Wang F-C, Kuo T-Y, Huang P-W, Chen S-F, Fu L-C. Early detection of Parkinson's disease by neural network models. *IEEE Access*. 2022;10:19033–44. <https://doi.org/10.1109/ACCESS.2022.3150774>.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.