



Correction to: Comparing Acute Kidney Injury Reports Among Antibiotics: A Pharmacovigilance Study of the FDA Adverse Event Reporting System (FAERS)

Taylor M. Patek^{1,2} · Chengwen Teng^{1,2} · Kaitlin E. Kennedy^{1,2} · Carlos A. Alvarez³ · Christopher R. Frei^{1,2,4,5}

Published online: 26 June 2020
© Springer Nature Switzerland AG 2020

Correction to: Drug Safety (2020) 43:17–22
<https://doi.org/10.1007/s40264-019-00873-8>

In the original publication of the article, the second sentence in the figure 2 caption should read as below:

A total of 2,042,801 reports (including 20,138 AKI reports) were included in the study. This is an example demonstrating how to calculate RORs: AKI RORs for vancomycin plus cefepime = $[22/(113 - 22)/((20,138 - 22)/(2,042,801 - 20,138 - (113 - 22)))] = 24.31$.

The original article can be found online at <https://doi.org/10.1007/s40264-019-00873-8>.

✉ Chengwen Teng
tengc3@livemail.uthscsa.edu

- ¹ Pharmacotherapy Division, College of Pharmacy, The University of Texas at Austin, San Antonio, TX, USA
- ² Pharmacotherapy Education and Research Center, Long School of Medicine, The University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Dr., MSC-6220, San Antonio, TX 78229, USA
- ³ Division of Clinical/Translational Research, Hodge School of Pharmacy, Texas Tech University Health Sciences Center, Dallas, TX, USA
- ⁴ South Texas Veterans Health Care System, San Antonio, TX, USA
- ⁵ University Health System, San Antonio, TX, USA