Sourcing and Using Appropriate Health State Utility Values in Economic Models in Health Care

Roberta Ara¹ · Tessa Peasgood¹ · Clara Mukuria¹ · Helene Chevrou-Severac² · Donna Rowen¹ · Ismail Azzabi-Zouraq² · Suzy Paisley¹ · Tracey Young¹ · Ben van Hout¹ · John Brazier¹

Published online: 19 October 2017
© Springer International Publishing AG 2017

1 Background

Decision analytic models (DAMs) in health care are generally used to assess different interventions to determine which provides the best value for money [1]. These models explore both the costs and benefits accrued by patients receiving the interventions, and results are presented in terms of the incremental cost per benefit of the intervention under evaluation [2]. While not all policy decision makers and reimbursement authorities conform to the same methodology [3], there has been a substantial growth in the use of incremental cost per quality-adjusted life-year (QALY) over the last two decades [4]. As the incremental QALY is the denominator in the outputs of the model, results can be sensitive to both the health state utility values (HSUVs) used and the methods and techniques used to deploy these values within the model [5].

The different components or stages involved in sourcing appropriate HSUVs (Fig. 1) are rarely independent and the process is generally iterative and often challenging [6]. In addition to identifying an appropriate preference-based measure for the particular condition (Fig. 1), and satisfying any associated reimbursement agency requirements, factors such as the advantages (or disadvantages) of collecting utility evidence in randomised clinical trials and other sources [7], the potential need to use a mapping function to predict the required HSUVs [8], and satisfying the exact

and often evolving health states used within the DAM all require consideration.

Once the most appropriate HSUVs available have been identified, these may not match the requirements of the DAM exactly and analysts may then need to make a series of methodological decisions or assumptions related to the practicalities of using this evidence within the DAM. These may include adjustments to account for age, sex or adverse events, evidence for the baseline or counterfactual (i.e. the trajectory of HSUVs for people who do not have a particular health condition or clinical event), estimate HSUVs for comorbidities (when a second health condition is present concurrently with the primary health condition), and characterising uncertainty appropriately (when using summary statistics sourced from different studies) (Fig. 1).

2 Existing Recommendations on Best Practice

2.1 Recommendations for Decision Analytic Models

Guidance on appropriate DAM approaches are available in the literature [1, 2, 9–11] and cover the choice of model structure [12]; measuring and estimating costs [13, 14]; identifying, reviewing and synthesising evidence on health-related quality of life [6, 15]; and methods for characterising uncertainty [16]. In addition, a number of checklists have been published over the years to encourage good practice in this area [1, 2, 9, 17, 18].

It is worth noting that there are issues where best practice can be defined and issues for which countries may have legitimately different preferences (such as the use of public or patient preferences). Both the US Panel on Cost-Effectiveness in Health and Medicine, and the UK National Institute for Health and Care Excellence (NICE) have

✉ Roberta Ara
r.m.ara@sheffield.ac.uk

¹ School of Health and Related Research, University of Sheffield, Regent Court, 30 Regent Street, Sheffield, UK

² Takeda Pharmaceuticals International AG, Thurgauerstrasse 130, 8152 Glattpark-Opfikon (Zurich), Switzerland

Fig. 1 The iterative process involved in identifying, analysing and using health state utility values used in decision analytic models. *HRQoL* health-related quality of life, *CSPBM* condition-specific

preference-based measure, *PBM* preference-based measure, *RCT* randomised controlled trial

developed guidelines with a view to standardising the methodological approaches in health technology appraisals to facilitate comparability of submissions [19, 20]; however, these and similar guidelines in other settings do not provide detailed guidance on good practice for HSUVs other than stating a preference for a particular preference-based measure (e.g. EQ-5D), source of evidence (e.g. patient), methodology (e.g. time trade-off) and source of preference weights (e.g. general population or patient), or specifying a specific output from the models (e.g. cost per QALY).

2.2 Recommendations for Health State Utility Values

A series of technical support documents commissioned via the NICE's Decision Support Unit in 2010 provide some recommendations of good practice when using HSUVs [6, 21–23], and a recent textbook provides an up-to-date review of preference-based measures of health [24]. More recently, their use in DAMs has formed the focus of two International Society for Pharmacoeconomics and Outcomes Research (ISPOR) Task Force reports [7, 8]. The first describes a framework for collecting HSUVs within

clinical studies and covers a broad spectrum from early planning of requirements, inclusion in clinical trial protocols, the advantages and design of alternative sources such as prospective and cross-sectional observational studies, statistical analyses and reporting standards [7]. The second provides recommendations relating to conducting high-quality mapping studies involving HSUVs and includes good practices for the selection of datasets used to obtain the mapping function, model selection and performance assessment, the use of results (including characterising variability and uncertainty appropriately), and reporting standards [8].

3 Standardising Good Practices

Good practice recommendations and checklists have evolved over time to address the developments in methodological research and the needs associated with the increasing complexity of DAMs. Consequently, there is no common source that covers the state-of-the-art recommendations across the full spectrum of considerations associated with the use of HSUVs informing DAMs, and there are still areas where recommendations for good

practice are absent. The result of this is there is no standardisation across current practice and policy decisions informed by the results from these models may be suboptimal.

The articles in this special issue provide a comprehensive compilation of the most robust recommendations in this area and are intended as a reference source for state-of-the-art evidence on good practice across the full spectrum of needs associated with HSUVs that inform DAMs.

Acknowledgements The authors would like to thank Prof. Jon Karnon, Ph.D., University of Adelaide, and Dr. Andrew Lloyd, Ph.D., Bladen Associates Ltd, for their editorial review.

Compliance with Ethical Standards

Disclosures This article is published in a special edition journal supplement wholly funded by Takeda Pharmaceutical International AG, Zurich, Switzerland.

References

- Gold MR. Panel on cost-effectiveness in health and medicine. *Med Care*. 1996;34(12 Suppl.):DS197–9.
- Drummond MF, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. *Methods for the economic evaluation of health care programmes*. Oxford: Oxford University Press; 2015.
- Barbieri M, Drummond M, Rutten F, Cook J, Glick HA, Lis J, et al. ISPOR Good Research Practices Economic Data Transferability Task Force. What do international pharmacoeconomic guidelines say about economic data transferability? *Value Health*. 2010;13(8):1028–37.
- Teja T, Michael C. Skills of the trade: the tufts cost-effectiveness analysis registry. *J Benefit Cost Anal*. 2012;3(1):1–9.
- Ara R, Brazier JE. Populating an economic model with health state utility values: moving toward better practice. *Value Health*. 2010;13(5):509–18.
- Papaioannou D, Brazier J, Paisley S. NICE DSU Technical Support Document 9: the identification, review and synthesis of health state utility values from the literature. University of Sheffield. 2011. <http://www.nicedsu.org.uk>. Accessed 7 Aug 2016.
- Wolowacz SE, Briggs A, Belozeroff V, Clarke P, Doward L, Goeree R, et al. Estimating health-state utility for economic models in clinical studies: an ISPOR Good Research Practices Task Force Report. *Value Health*. 2016;19(6):704–19.
- Wailoo AJ, Hernandez-Alava M, Manca A, Mejia A, Ray J, Crawford B, et al. Mapping to estimate health-state utility from non-preference-based outcome measures: an ISPOR Good Practices for Outcomes Research Task Force Report. *Value Health*. 2017;20(1):18–27.
- Eddy DM. Technology assessment: the role of mathematical modeling. In: *Assessing Medical Technologies*. Washington DC: National Academies Press; 1985. pp. 144–75.
- Weinstein MC, O'Brien B, Hornberger J, Jackson J, Johannesson M, McCabe C, et al. Principles of good practice for decision analytic modeling in health-care evaluation: report of the ISPOR Task Force on Good Research Practices—modeling studies. *Value Health*. 2003;6(1):9–17.
- Caro JJ, Briggs AH, Siebert U, Kuntz KM. Modeling good research practices—overview a report of the ISPOR-SMDM modeling good research practices task force–1. *Med Decis Mak*. 2012;32(5):667–77.
- Brennan A, Chick SE, Davies R. A taxonomy of model structures for economic evaluation of health technologies. *Health Econ*. 2006;15(12):1295–310.
- Young TA. Estimating mean total costs in the presence of censoring. *Pharmacoeconomics*. 2005;23(12):1229–42.
- Lee CY, Wand MP. Variational methods for fitting complex Bayesian mixed effects models to health data. *Stat Med*. 2016;35(2):165–88.
- Kaltenthaler E, Tappenden P, Paisley S. Reviewing the evidence to inform the population of cost-effectiveness models within health technology assessments. *Value Health*. 2013;16(5):830–6.
- Briggs AH. Handling uncertainty in cost-effectiveness models. *Pharmacoeconomics*. 2000;17(5):479–500.
- Husereau D, Drummond M, Petrou S, Carswell C, Moher D, Greenberg D, et al. Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement. *Cost Eff Resour Alloc*. 2013;11(1):6.
- Drummond MF, Jefferson TO. Guidelines for authors and peer reviewers of economic submissions to the BMJ. The BMJ Economic Evaluation Working Party. *BMJ*. 1996;313(7052):275.
- Sanders GD, Neumann PJ, Basu A, Brock DW, Feeny D, Krahn M, et al. Recommendations for conduct, methodological practices, and reporting of cost-effectiveness analyses: second panel on cost-effectiveness in health and medicine. *JAMA*. 2016;316(10):1093–103.
- Guide to the methods of technology appraisal 2013. <http://nice.org.uk>. Accessed 30 Nov 2016.
- Longworth L, Rowen D. NICE DSU technical support document 10: the use of mapping methods to estimate health state utility values. Sheffield: Decision Support Unit, ScHARR, University of Sheffield; 2011.
- Brazier J, Longworth L. NICE DSU Technical Support Document 8: an introduction to the measurement and valuation of health for NICE submissions. NICE Decision Support Unit, London. 2011. <http://www.nicedsu.org.uk>. Accessed 7 Aug 2016.
- Ara R, Wailoo AJ. NICE DSU Technical Support Document 12: the use of health state utility values in decision models. 2011. <http://www.nicedsu.org.uk>. Accessed 7 Aug 2016.
- Brazier J, Ratcliffe J, Salomon J, Tsuchiya A. *Measuring and valuing health benefits for economic evaluation*. Oxford: Oxford University Press; 2016.