



Correction to: PAIT-Survey Follow-Up: Changes in Albuminuria in Hypertensive Diabetic Patients with Mild-Moderate Chronic Kidney Disease

Francesco Fici¹ · Elif Ari Bakir² · Elif Ilkay Yüce³ · Serdal Kanuncu⁴ · Wim Makel⁵ · Bahar Arican Tarim⁶ · Nicolás Roberto Robles^{1,7,8} 

Published online: 12 January 2021
© Italian Society of Hypertension 2021

Correction to:
High Blood Press Cardiovasc Prev (2020) 27(1):43–49
<https://doi.org/10.1007/s40292-020-00358-1>

Table 3 should appear as follows:

* $p < 0.001$ vs baseline, [McNemar test]

** $p < 0.05$ vs baseline, [McNemar test]

[§] $p < 0.05$ vs other groups, [Chi-square test]

Table 3 Changes in prevalence of patients with albuminuria with different treatment strategies

Treatment	Amlodipine–valsartan	Amlodipine–perindopril	Lercanidipine–enalapril	Verapamil–trandolapril
Microalbuminuria				
Before (n)	77	68	92	26
After (n)	65	60	54	21
Δ (n)	– 12	– 8	– 38	– 5
%	– 15.6	– 11.8	– 41.3	– 19.2**
Macroalbuminuria				
Before	18	10	16	3
After	12	10	12	3
Δ (n)	– 6	0	– 4	0
%	– 27.8	–	– 25.0	–
Normoalbuminuria				
Before	122	75	133	28
After	140	83	175	33
Δ (n)	+ 18	+ 8	+ 42	+ 5
%	+ 14.8	+ 10.7	+ 28.6 [§]	+ 17.8**

The original article can be found online at <https://doi.org/10.1007/s40292-020-00358-1>.

✉ Nicolás Roberto Robles
nrrobles@yahoo.es

¹ Catedra de Riesgo Cardiovascular, Universidad de Salamanca, Salamanca, Spain

² Department of Nephrology, Bahcesehir University Hospital, Istanbul, Turkey

³ Kelkit State Hospital, Gumushane, Turkey

⁴ Family Health Center, Defne-Hatay, Turkey

⁵ Clinical Research Facilities International B.V, Schaijk, The Netherlands

⁶ Kartal Research and Education Hospital, Istanbul, Turkey

⁷ Hospital Universitario de Badajoz, Badajoz, Spain

⁸ Unidad de Hipertensión Arterial, Hospital Infanta Cristina, Carretera de Portugal s/n, 06080 Badajoz, Spain