The Logarithmic Sobolev Inequality Along the Ricci Flow: The Case $\lambda_0(g_0) = 0$

Rugang Ye

Received: 26 August 2014 / Accepted: 6 February 2015 / Published online: 19 March 2015
© School of Mathematical Sciences, University of Science and Technology of China and Springer-Verlag Berlin Heidelberg 2015

Abstract A uniform logarithmic Sobolev inequality, a uniform Sobolev inequality and a uniform κ -noncollapsing estimate along the Ricci flow are established in the situation that a certain smallest eigenvalue for the initial metric is zero.

Keywords Uniform · Logarithmic Sobolev inequality · Sobolev inequality · Ricci flow · Eigenvalue

Mathematics Subject Classification 53C44 · 35K55

1 Introduction

In [5] and [6], logarithmic Sobolev inequalities along the Ricci flow in all dimensions $n \geq 2$ were obtained using Perelman's entropy monotonicity, which lead to Sobolev inequalities and κ -noncollapsing estimates. In particular, a uniform logarithmic Sobolev inequality, a uniform Sobolev inequality and a uniform κ -noncollapsing estimate were obtained without any restriction on time, provided that the smallest eigenvalue $\lambda_0(g_0)$ of the operator $-\Delta + \frac{R}{4}$ for the initial metric is positive. In this paper, we extend these uniform results to the case $\lambda_0(g_0) = 0$.

Consider a compact manifold M of dimension $n \geq 2$. Let $g = g(t)$ be a smooth solution of the Ricci flow

$$\frac{\partial g}{\partial t} = -2\text{Ric} \quad (1.1)$$

R. Ye (✉)
Department of Mathematics, University of California, Santa Barbara, USA
e-mail: yer@math.ucsb.edu

on $M \times [0, T)$ for some (finite or infinite) $T > 0$ with a given initial metric $g(0) = g_0$.

Theorem 1.1 *Assume that $\lambda_0(g_0) = 0$. For each $t \in [0, T)$ and each $\sigma > 0$ there holds*

$$\int_M u^2 \ln u^2 d\text{vol} \leq \sigma \int_M \left(|\nabla u|^2 + \frac{R}{4} u^2 \right) d\text{vol} - \frac{n}{2} \ln \sigma + C \quad (1.2)$$

for all $u \in W^{1,2}(M)$ with $\int_M u^2 d\text{vol} = 1$, where C depends only on (M, g_0) .

We would like to emphasize the key feature that no constant in this inequality depends on T . The same is true in the subsequent theorems. Note that as in [5], a log gradient version of the logarithmic Sobolev inequality follows as a consequence of (1.2). We omit its statement. Our next result provides a dependence of the above C on g_0 in terms of rudimentary geometric data. To simplify the statements, we assume that $|Rm| \leq 1$ for g_0 , which can always be achieved by a rescaling. Then we can also assume $T \geq 2\alpha(n)$ for a positive constant $\alpha(n)$ depending only on n such that $|Rm| \leq 2$ on $[0, \alpha(n)]$. (Namely the maximal possible T such that the solution $g = g(t)$ can be extended to a smooth solution of the Ricci flow on $[0, T)$ has this property.)

Theorem 1.2 *There are for each $v_0 > 0$, each $D_0 > 0$, each $\epsilon > 0$ and each integer $l \geq 3$ a positive number $C = C(v_0, D_0, \epsilon, l, n)$ with the following properties. Assume $\lambda_0(g_0) = 0$, $\text{vol}_{g_0}(M) \geq v_0$, $\text{diam}_{g_0} \leq D_0$, and the normalization conditions $|Rm|_{g_0} \leq 1$ and $T \geq 2\alpha(n)$. $\text{vol}_{g_0}(M) \geq v_0$. Then one of the following two cases must occur:*

- 1) $g(\alpha(n))$ lies in the ϵ -neighborhood of a Ricci flat metric on M in the C^l norm,
- 2) the logarithmic Sobolev inequality (1.2) holds true for each $t \in [0, T)$, each $\sigma > 0$ and all $u \in W^{1,2}(M)$ with $\int_M u^2 d\text{vol} = 1$, where $C = C(v_0, D_0, \epsilon, l, n)$.

It turns out that we have a better result in dimension $n = 3$. The same holds true in dimension $n = 2$. But Theorem 1.5 and Theorem 3.7 in [6] provide a stronger result in this dimension.

Theorem 1.3 *Assume that $n = 3$. There is for each $v_0 > 0$ and each $D_0 > 0$ a positive number $C = C(v_0, D_0)$ with the following properties. Assume $\lambda_0(g_0) = 0$, $\text{vol}_{g_0}(M) \geq v_0$, $\text{diam}_{g_0} \leq D_0$, and the normalization conditions $|Rm|_{g_0} \leq 1$ and $T \geq 2\alpha(n)$. Then the logarithmic Sobolev inequality (1.2) holds true for each $t \in [0, T)$, each $\sigma > 0$ and all $u \in W^{1,2}(M)$ with $\int_M u^2 d\text{vol} = 1$, where $C = C(v_0, D_0)$.*

As in [5], Theorems 1.1–1.3 lead to Sobolev inequalities along the Ricci flow, which in turn lead to κ -noncollapsing estimates. We consider only the case $n \geq 3$ although the methods also work for $n = 2$, because the case $n = 2$ is covered by the results in [6].

Theorem 1.4 Assume that $n \geq 3$ and $\lambda_0(g_0) = 0$. Then there holds for each $t \in [0, T)$

$$\left(\int_M |u|^{\frac{2n}{n-2}} d\text{vol} \right)^{\frac{n-2}{n}} \leq A \int_M \left(|\nabla u|^2 + \frac{R}{4} u^2 \right) d\text{vol} + B \int_M u^2 d\text{vol} \quad (1.3)$$

for all $u \in W^{1,2}(M)$, where A and B depend only on (M, g_0) . We have $A = A(v_0, D_0)$ and $B = B(v_0, D_0)$ for given $v_0 > 0$ and $D_0 > 0$, if $n = 3$ and g_0 satisfies the conditions in Theorem 1.3. We also have $A = A(v_0, D_0, n)$ and $B = B(v_0, D_0, n)$ for given $v_0 > 0$ and $D_0 > 0$, provided that g_0 satisfies the conditions in Theorem 1.2 and $g(\alpha(n))$ does not lie in the ϵ -neighborhood of any Ricci flat metric on M in the C^3 norm.

Theorem 1.5 Assume that $n = 3$ and $\lambda_0(g_0) = 0$. Let $L > 0$ and $t \in [0, T)$. Consider the Riemannian manifold (M, g) with $g = g(t)$. Assume $R \leq \frac{1}{r^2}$ on a geodesic ball $B(x, r)$ with $0 < r \leq L$. Then there holds

$$\text{vol}(B(x, r)) \geq \left(\frac{1}{2^{n+3}A + 2BL^2} \right)^{\frac{n}{2}} r^n, \quad (1.4)$$

where A and B are from Theorem 1.4.

As in [5], the above results extend to various versions of the modified Ricci flows. Moreover, the κ -noncollapsing estimates ensure that we can obtain smooth blow-up limits at the time infinity under the assumption that $\lambda_0(g_0) = 0$. As in [5], the above results also extend to the Ricci flow with surgeries. We omit the statements of those results because they are completely analogous to the corresponding ones in [5].

2 The Proofs

Proof of Theorem 1.1 Consider a fixed $t_1 \in (0, T)$. Let u_1 be a positive eigenfunction for the eigenvalue $\lambda_0(g(t_1))$ associated with the metric $g(t_1)$, such that $\int_M u_1^2 d\text{vol} = 1$ with respect to $g(t_1)$. Let $f = f(t)$ be the smooth solution of the equation

$$\frac{\partial f}{\partial t} = -\Delta f + |\nabla f|^2 - R \quad (2.1)$$

on $[0, t_1]$ with $f(t_1) = -2 \ln u_1$. Note that (2.1) is equivalent to

$$\frac{\partial v}{\partial t} = -\Delta v + Rv, \quad (2.2)$$

where $v = e^{-f}$. So the solution $f(t)$ exists. We also infer $\frac{d}{dt} \int_M v d\text{vol} = 0$, and hence $\int_M v d\text{vol} = 1$ for all $t \in [0, t_1]$.

We set $u = e^{-\frac{f}{2}}$. By [4, (1.4)] we then have

$$\begin{aligned} \frac{d}{dt} \int_M \left(|\nabla u|^2 + \frac{R}{4} u^2 \right) d\text{vol} &= \frac{1}{4} \frac{d}{dt} \int_M \left(|\nabla f|^2 + R \right) e^{-f} d\text{vol} \\ &\geq \frac{1}{2} \int_M |Ric + \nabla^2 f|^2 e^{-f} d\text{vol}. \end{aligned} \quad (2.3)$$

It follows that

$$\lambda_0(g(t_1)) \geq \lambda_0(g_0) + \frac{1}{2} \int_0^{t_1} \int_M |Ric + \nabla^2 f|^2 e^{-f} d\text{vol} dt. \quad (2.4)$$

We choose $t_1 = \min\{\frac{T}{2}, 1\}$. If $\lambda_0(g(t_1)) > 0$, we first apply Theorem 1.1 in [5] or Theorem 1.1 in [6] to obtain (1.2) for $g(t)$ on $[0, t_1]$. Then we apply Theorem 4.2 in [5] to obtain (1.2) on $[t_1, T)$ with a larger C . If $\lambda_0(g(t_1)) = 0$, we deduce from (2.4)

$$Ric + \nabla^2 f = 0 \quad (2.5)$$

on $[0, t_1]$. It follows that $g = g(t)$ is a steady Ricci soliton for all t . Hence the logarithmic Sobolev inequality for g_0 provided by Theorem 3.3 in [5] holds true for all $g(t)$. Actually, Proposition 5.20 in [1] implies that g_0 is Ricci flat, hence $g(t) = g_0$ for all t . $\square$

Lemma 2.1 *For given $v_0 > 0$, $D_0 > 0$, $\epsilon > 0$ and $l \geq 3$ there is a positive constant $\mu_0 = \mu_0(v_0, D_0, \epsilon, l, n)$ with the following properties. Let $g = g(t)$ be a smooth solution of the Ricci flow on $M \times [0, T)$ which satisfies the normalization conditions in Theorem 1.2 and the conditions $\text{vol}_{g(0)} \geq v_0$ and $\text{diam}_{g(0)} \leq D_0$. Assume $\lambda_0(g_0) = 0$. If $\lambda(g(\alpha(n))) < \mu_0$, then $g(\alpha(n))$ lies in the ϵ -neighborhood of a Ricci flat metric with respect to the C^l -norm.*

Proof Assume that μ_0 does not exist. Then we can find a sequence of manifolds M_k of a fixed dimension n and a sequence of smooth solutions $g_k = g_k(t)$ on $M_k \times [0, T_k)$ satisfying the normalization conditions and the conditions $\text{vol}_{g_k(0)} \geq v_0$ and $\text{diam}_{g_k(0)} \leq D_0$, such that $\lambda_0(g_k(0)) = 0$, $\lambda_0(g_k(\alpha(n))) \rightarrow 0$, and $g_k(\alpha(n))$ does not lie in the ϵ -neighborhood of any Ricci flat metric with respect to the C^l -norm. By Gromov–Cheeger–Hamilton compactness theorem [3], we can find a subsequence of (M_k, g_k) , which we still denote by (M_k, g_k) , such that $(M_k, g_k, [\frac{\alpha(n)}{2}, \alpha(n)])$ converge smoothly to a limit Ricci flow $(M, g, [\frac{\alpha(n)}{2}, \alpha(n)])$. There holds $\lambda_0(g(\alpha(n))) = 0$. By the monotonicity of λ_0 along the Ricci flow (see [4] or the above proof of Theorem 1.1), we have $\lambda_0(g_k(t)) \geq 0$ for all $t \in [0, T_k)$. Hence $\lambda_0(g(t)) \geq 0$ for all $t \in [\frac{\alpha(n)}{2}, \alpha(n)]$. Now the argument in the proof of Theorem 1.1 implies that $g(\alpha(n))$ is Ricci flat. But $(M_k, g_k(\alpha(n)))$ converge smoothly to $(M, g(\alpha(n)))$, so $g_k(\alpha(n))$ lies in the ϵ -neighborhood of a Ricci flat metric with respect to the C^l -norm whenever k is large enough. This is a contradiction. $\square$

Proof of Theorem 1.2 Assume that $g(\alpha(n))$ does not lie in the ϵ -neighborhood of any Ricci flat metric with respect to the C^l -norm. Then $\lambda_0(g(\alpha(n))) \geq \mu_0$ by

Lemma 2.1. Now we obtain a desired logarithmic Sobolev inequality for $g(t)$ on $[0, \alpha(n)]$ by Theorem 1.1 in [5]. Alternatively, we can apply the arguments in [7] for controlling the evolution of the Sobolev constant to bound the Sobolev constant for $g(t)$ on $[0, \alpha(n)]$, and then apply Theorem 3.3 in [5] to infer the desired logarithmic Sobolev inequality. Next we apply the bound for the Sobolev constant at $t = \alpha(n)$, the bound $\lambda_0 \geq \mu_0$ at $t = \alpha(n)$ and the arguments in the proof of Theorem 3.5 in [5] to deduce a logarithmic Sobolev inequality of the kind [5, (3.11)] at $t = \alpha(n)$. Then we apply Theorem 1.2 in [5] on $[\alpha(n), T)$ and combine it with Theorem 1.1 in [5]. Then we arrive at the desired logarithmic Sobolev inequality on $[\alpha(n), T)$. $\square$

Proof of Theorem 1.3 By [2], there is for each given flat metric g an ϵ -neighborhood of g with respect to the C^6 -norm, such that the Ricci flow starting at any metric in the neighborhood converges smoothly to a Ricci flat metric at a fixed exponential rate as $t \rightarrow \infty$. Moreover, the limit Ricci flat metric lies in the 2ϵ -neighborhood of g with respect to the C^6 -norm. We call such a neighborhood a Ricci contraction ϵ -neighborhood.

Now for given $v_0 > 0$, $D_0 > 0$ and $K_0 > 0$ the moduli space $\mathcal{M}^0(v_0, D_0, K_0)$ of flat metrics on M with $\text{vol} \geq v_0$, $\text{diam} \leq D_0$ and $|Rm| \leq K_0$ is C^∞ compact modulo diffeomorphisms by Gromov–Cheeger compactness theorem and the Einstein equation. So there is a uniform $\epsilon = \epsilon(v_0, D_0, K_0)$ such that each $g \in \mathcal{M}^0(v_0, D_0, K_0)$ has a Ricci contraction ϵ -neighborhood. Moreover, there is a uniform upper bound $C(v_0, D_0, K_0)$ for the Sobolev constant for $g \in \mathcal{M}^0(v_0, D_0, K_0)$.

Now consider for given $v_0 > 0$ and $D_0 > 0$ a smooth solution of the Ricci flow $g = g(t)$ satisfying the normalization conditions and the conditions $\text{vol}_{g_0}(M) \geq v_0$ and $\text{diam}_{g_0}(M) \leq D_0$. Let $\epsilon > 0$ and $l = 6$, where ϵ is to be determined. Assume that $g(\alpha(n))$ lies in the ϵ -neighborhood of a Ricci flat metric $\bar{g}$ with respect to the C^6 norm. Since $n = 3$, $\bar{g}$ is flat. There is a positive number $\epsilon_0 = \epsilon_0(v_0, D_0)$ such that $\text{vol}_{\bar{g}}(M) \geq \frac{1}{2}v_0$, $\text{diam}_{\bar{g}}(M) \leq 2D_0$ and $|Rm|_{\bar{g}} \leq 3$, i.e., $\bar{g} \in \mathcal{M}^0(\frac{1}{2}v_0, 2D_0, 3)$, whenever $\epsilon \leq \epsilon_0$. Now we choose $\epsilon = \min\{\epsilon_0(v_0, D_0), \epsilon(\frac{1}{2}v_0, 2D_0, 4)\}$. Then the maximally extended $g(t)$ converges smoothly to a flat metric g^* at exponential rate where the rate depends only on v_0 and D_0 . We can choose ϵ_0 sufficiently small such that $g^* \in \mathcal{M}^0(\frac{1}{3}v_0, 3D_0, 4)$. We can also make the C^6 norm of $g(t) - g^*$ sufficiently small for all $t \geq \alpha(n)$ such that the Sobolev constant of $g(t)$ is bounded above by $2C(\frac{1}{3}v_0, 3D_0, 4)$ for all $t \geq \alpha(n)$. A desired logarithmic Sobolev inequality then follows for $t \in [\alpha(n), T)$.

A desired logarithmic Sobolev inequality for $g(t)$ on $[0, \alpha(n)]$ follows from Theorem 1.1 in [5]. It also follows from the arguments for controlling the evolution of the Sobolev constant in [7]. $\square$

References

1. Chow, B., Knopf, D.: The Ricci flow: An introduction. Mathematical surveys and monographs. Am. Math. Soc. **110** (2004)
2. Guenther, C., Isenberg, J., Knopf, D.: Stability of the Ricci flow at Ricci flat metrics. Commun. Anal. Geom. **10**, 741–777 (2002)
3. Hamilton, R.S.: A compactness property for solutions of the Ricci flow. Am. J. Math. **117**, 545–572 (1995)

4. Perelman, G.: The entropy formula for the Ricci flow and its geometric applications. November (2002), <http://arxiv.org/abs/math.DG/0211159v1>
5. Ye, R.: The logarithmic Sobolev inequality along the Ricci flow. [arXiv:0707.2424](https://arxiv.org/abs/0707.2424)
6. Ye, R.: The logarithmic Sobolev inequality along the Ricci flow in dimension 2. July (2007) [arXiv:0708.2003](https://arxiv.org/abs/0708.2003)
7. Ye, R.: Ricci flow, Einstein metrics and space forms. *Trans. Am. Math. Soc.* **338**, 871–895 (1993)