



On Bloch's "Principle of Topological Continuity"

Walter Bergweiler¹ · Alexandre Eremenko²

Received: 26 September 2023 / Revised: 4 January 2024 / Accepted: 5 January 2024
© The Author(s) 2024

Abstract

We discuss to what extent certain results about totally ramified values of entire and meromorphic functions remain valid if one relaxes the hypothesis that some value is totally ramified by assuming only that all islands over some Jordan domain are multiple. In particular, we prove a result suggested by Bloch which says that an entire function of order less than 1 has a simple island over at least one of two given Jordan domains with disjoint closures.

Keywords Meromorphic function · Entire function · Ramified value · Island · Ahlfors theory · Covering surface · Quasiconformal deformation · Line complex

Mathematics Subject Classification 30D30 · 30D35 · 30D45

1 Introduction

About one hundred years ago, André Bloch [14] wrote a paper consisting mainly of heuristic speculations based on two philosophical principles. In this paper, he anticipated several important results of 20th century geometric function theory—and even today reading this paper can still be rewarding.

Dedicated to the memory of Larry Zalcman.

Communicated by Mark Agranovsky.

✉ Walter Bergweiler
bergweiler@math.uni-kiel.de

Alexandre Eremenko
eremenko@math.purdue.edu

¹ Mathematisches Seminar, Christian-Albrechts-Universität zu Kiel, Ludwig-Meyn-Str. 4, 24098 Kiel, Germany

² Department of Mathematics, Purdue University, West Lafayette, IN 47907, USA

The first of these principles Bloch phrased as “Nihil est in infinito quod non prius fuerit in finito”. This principle has often been interpreted as meaning that if all entire functions with a certain property are constant, then the family of functions which are holomorphic in some domain and have this property is normal. This is an important guideline in the theory of normal families; see [9, 35, 36] for a discussion. We will call it *Bloch’s normal family principle* in the sequel. Another interpretation of “Nihil est in infinito quod non prius fuerit in finito” given by Bloch is that features of transcendental entire and meromorphic functions are in some form already present in polynomials and rational functions.

The second principle that Bloch discusses he calls the *principle of topological continuity*. It says that certain true statements remain true if one modifies the data from a metric point of view, but not from the topological point of view.

To illustrate this principle, he quotes the following theorem.

Theorem A *Let D be a domain and let $a_1, a_2, a_3 \in \mathbb{C}$ be distinct. Let \mathcal{F} be the family of all functions holomorphic in D which do not have a simple a_j -point, for all $j \in \{1, 2, 3\}$. Then \mathcal{F} is normal.*

He then argues [14, p. 87] that the principle of topological continuity should give the following result.

Theorem A’ *Let D be a domain and let D_1, D_2 and D_3 be three disks in \mathbb{C} which have pairwise disjoint closures. Let \mathcal{F} be the family of all functions holomorphic in D such that there does not exist a domain U with $\bar{U} \subset D$ such that U is mapped univalently onto one of the disks D_j . Then \mathcal{F} is normal.*

Theorem A was known when Bloch wrote this. Theorem A’ was not known then. It was first proved by Ahlfors six years later, see [1–4], his definitive account being [5]. Theorem A’, and its generalization Theorem D’ below, are among the principal results of his theory of covering surfaces which earned him one of the two first Fields medals in 1936.

Ahlfors did not use Theorem A in his proof of Theorem A’, nor do the more recent proofs in [17, 18, 30]. Ahlfors does, however, give an interesting discussion of Bloch’s principle of topological continuity in [3, pp. 202–203]. A deduction of Theorem A’ from Theorem A—and thus in some sense a confirmation of Bloch’s principle of topological continuity—was given in [8]. This deduction was based on a rescaling principle of Zalcman [35].

Zalcman’s lemma has become a major tool in the theory of normal families by giving a rigorous formulation of *Bloch’s normal family principle*. For example, it shows that Theorems A and A’ can be deduced from the following corresponding results about entire functions.

Theorem B *Let f be entire and let $a_1, a_2, a_3 \in \mathbb{C}$ be distinct. Suppose that all a_j -points are multiple, for all $j \in \{1, 2, 3\}$. Then f is constant.*

Theorem B’ *Let f be entire and let D_1, D_2 and D_3 be three disks in \mathbb{C} which have pairwise disjoint closures. Suppose that there does not exist $j \in \{1, 2, 3\}$ and a bounded domain U in \mathbb{C} which is mapped univalently onto D_j . Then f is constant.*

As a second example for his principle of topological continuity Bloch considers the following result.

Theorem C *Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be a non-constant entire function of order less than 1 and let $a_1, a_2 \in \mathbb{C}$ be distinct. Then there exists $j \in \{1, 2\}$ such that f has a simple a_j -point.*

This theorem can be proved using Nevanlinna theory, and Bloch was aware of this proof [12, 13]. We will sketch a proof after Theorem D' below.

Bloch argues [14, p. 88] that Theorem C together with his principle of topological continuity should yield the following result.

Theorem C' *Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be a non-constant entire function of order less than 1 and let D_1 and D_2 be two disks in \mathbb{C} with disjoint closures. Then there exists $j \in \{1, 2\}$ and a domain U in \mathbb{C} which is mapped univalently onto D_j .*

So far as we know, there is no proof of Theorem C' in the literature, neither by Bloch nor by someone else. We will give a proof of (a generalization of) Theorem C' in Sect. 3, using some ideas of Goldberg and Tairova [19].

Theorems B and B' are actually special cases of more general results. The generalization of Theorem B (which Bloch also knew) is the following result due to Nevanlinna [25, Chapitre IV, no. 51].

Theorem D *Let $f: \mathbb{C} \rightarrow \bar{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ be meromorphic and $q \in \mathbb{N}$. Let $a_1, \dots, a_q \in \mathbb{C}$ be distinct and $m_1, \dots, m_q \in \mathbb{N} \cup \{\infty\}$. If, for all $j \in \{1, \dots, q\}$, all a_j -points have multiplicity at least m_j , then*

$$\sum_{j=1}^q \left(1 - \frac{1}{m_j}\right) \leq 2 \tag{1.1}$$

or f is constant.

Here $m_j = \infty$ means that f does not take the value a_j at all, and we put $1/m_j = 1/\infty = 0$ in this case.

To state the corresponding generalization of Theorem B' we introduce some terminology. Let $f: \mathbb{C} \rightarrow \bar{\mathbb{C}}$ be meromorphic and let D be a Jordan domain in $\bar{\mathbb{C}}$. A connected component U of $f^{-1}(D)$ is called an *island* of f over D if it is bounded and simply-connected. Then $f: U \rightarrow D$ is a proper mapping. The degree of this proper mapping is called the *multiplicity* of the island U . An island of multiplicity 1 is called a *simple island*.

Our terminology differs slightly from that in the classical literature, as we require that islands are simply-connected. Of course, this does not make a difference for islands of entire functions over domains in \mathbb{C} , or for simple islands.

The following generalization of Theorem B' is due to Ahlfors, who called it "Scheibensatz" [5, p. 190].

Theorem D' *Let $f: \mathbb{C} \rightarrow \bar{\mathbb{C}}$ be meromorphic and $q \in \mathbb{N}$. Let D_1, \dots, D_q be Jordan domains in $\bar{\mathbb{C}}$ with pairwise disjoint closures and let $m_1, \dots, m_q \in \mathbb{N} \cup \{\infty\}$. If, for*

all $j \in \{1, \dots, q\}$, all islands over D_j have multiplicity at least m_j , then (1.1) holds or f is constant.

As mentioned, Ahlfors did not require that islands are simply-connected, but his theory yields the above result also with this additional condition; see [28, Chapter X, §2, Theorem 29] or [31, Theorem VI.5 or VI.13]. Theorems B and B' are obtained from Theorems D and D' by choosing $q = 4, m_1 = m_2 = m_3 = 2, m_4 = \infty$ and $a_4 = \infty$.

There are also normal family analogues of Theorems D and D' according to Bloch's normal family principle. These can be obtained from Theorems D and D' via Zalcman's lemma.

Suppose now that we have equality in (1.1); that is,

$$\sum_{j=1}^q \left(1 - \frac{1}{m_j}\right) = 2. \tag{1.2}$$

Then, apart from permutation of the m_j , we have one of the following six cases:

- (i) $q = 2, (m_1, m_2) = (\infty, \infty)$.
- (ii) $q = 3, (m_1, m_2, m_3) = (2, 2, \infty)$.
- (iii) $q = 4, (m_1, m_2, m_3, m_4) = (2, 2, 2, 2)$.
- (iv) $q = 3, (m_1, m_2, m_3) = (2, 3, 6)$.
- (v) $q = 3, (m_1, m_2, m_3) = (2, 4, 4)$.
- (vi) $q = 3, (m_1, m_2, m_3) = (3, 3, 3)$.

Selberg [27, Satz II] determined all transcendental meromorphic functions f of finite order such that f has only finitely many a_j -points of multiplicity less than m_j , with the m_j chosen such that (1.2) holds. He used this to determine the possible orders of these functions [27, Satz IV–VII]. Let $\rho(f)$ denote the order of a meromorphic function f ; see [20, Chapter 2, Section 1] for the definition of the order and other concepts from the theory of meromorphic functions used in the sequel.

We briefly summarize Selberg's reasoning and results. First, in case (i) we assume that $(a_1, a_2) = (0, \infty)$ and find that the function f has the form $f = Qe^P$ with a rational function Q and a polynomial P . Thus $\rho(f) \in \mathbb{N}$ in this case.

In case (ii) we assume that $a_1 = 1, a_2 = -1$ and $a_3 = \infty$. Then

$$R(z) = \frac{f'(z)^2}{f(z)^2 - 1} \tag{1.3}$$

has poles only at the simple ± 1 -points of f . Thus R has only finitely many poles. Since f has finite order, the lemma on the logarithmic derivative [20, Chapter 3, Section 1] yields that Nevanlinna's proximity functions $m(r, R)$ satisfies $m(r, R) = O(\log r)$. We conclude that R is a rational function. Hence f has the form

$$f(z) = \cosh\left(\int \sqrt{R(z)} dz\right). \tag{1.4}$$

This implies that $\rho(f)$ is an integer multiple of $1/2$. (Equation (1.4) and its implication for $\rho(f)$ were actually already obtained by Valiron [32, p. 77], before Selberg's work and in fact before Nevanlinna developed his theory.)

If all ± 1 -points are multiple, then R has no poles and is thus a polynomial. In this case we find that $\rho(f) \geq 1$. Note that this proves Theorem C.

In cases (iii)–(vi), assume that all a_j are in \mathbb{C} and that M is the least common multiple of the m_j . We then find that

$$R(z) = \frac{f'(z)^M}{\prod_{j=1}^q (f(z) - a_j)^{(m_j-1)M/m_j}}$$

is a rational function and f has the form

$$f(z) = E \left(\int \sqrt[M]{R(z)} dz \right).$$

where E is an elliptic function.

We only state the conclusions about the order of f that Selberg drew from this.

Theorem E *Let $f: \mathbb{C} \rightarrow \overline{\mathbb{C}}$ be a transcendental meromorphic function of finite order, $q \in \mathbb{N}$, $a_1, \dots, a_q \in \overline{\mathbb{C}}$ distinct and $m_1, \dots, m_q \in \mathbb{N} \cup \{\infty\}$. Suppose that (1.2) holds so that we have one of the cases (i)–(vi) listed above.*

If, for all $j \in \{1, \dots, q\}$, all but finitely many a_j -points of f have multiplicity at least m_j , then $\rho(f) \in \mathbb{N}$ in case (i), $2\rho(f) \in \mathbb{N}$ in case (ii), $\rho(f) \in \mathbb{N}_0$ in case (iii), $3\rho(f) \in \mathbb{N}_0$ in case (iv), $2\rho(f) \in \mathbb{N}_0$ in case (v) and $3\rho(f)/2 \in \mathbb{N}_0$ in case (vi).

If all (and not only all but finitely many) a_j -points have multiplicity at least m_j , then in addition $\rho(f) \geq 1$ in cases (i) and (ii) while $\rho(f) \geq 2$ in cases (iii)–(vi).

The question that motivated this paper is whether there exists a Theorem E' which corresponds to Theorem E in the same way that Theorems A', B', C' and D' correspond to Theorems A, B, C and D. In other words, we ask to what extent the conclusion of Theorem E remains valid if instead of the hypotheses of Theorem D we assume the hypotheses of Theorem D'. Note that Theorem C', which was envisaged by Bloch using his principle of topological continuity, says that this holds in case (ii) if f is entire. In contrast, we will see that the corresponding result does not hold in cases (i) and (iii).

The following result corresponds to case (i) of Theorem D.

Theorem 1.1 *Let f be a transcendental entire function and let D be a Jordan domain in \mathbb{C} . If f has only finitely many islands over D , then $\rho(f) \geq 1/2$.*

Conversely, for every $\rho \in [1/2, \infty)$ and every Jordan domain D there exists an entire function f of order ρ such that f has no island over D .

Our next result corresponds to case (ii) of Theorem D. It has Theorem C' as a corollary. Here and in the following we will denote the multiplicity of an island U over some Jordan domain by $\mu(U)$.

Theorem 1.2 *Let f be a transcendental entire function of finite order and let D_1 and D_2 be Jordan domains in \mathbb{C} with disjoint closures. Suppose that f has only finitely many simple islands over D_1 and D_2 . Let N be the number (counting multiplicity) of critical points c of f such that $f(c) \notin D_1 \cup D_2$. Put*

$$p = 2N + 2 + \sum_U (\mu(U) - 2), \tag{1.5}$$

where the sum is taken over all islands U over D_1 or D_2 . Then $1 \leq p < \infty$ and there exists $c > 0$ such that

$$\log M(r, f) \sim c r^{p/2} \tag{1.6}$$

as $r \rightarrow \infty$. In particular, $\rho(f) = p/2$.

Remark In the proof of Theorem 1.2 the hypothesis that f is entire and of finite order will be used only to conclude that f^{-1} has finitely many transcendental singularities. So actually we prove a more general theorem:

Let U be a simply-connected domain in \mathbb{C} and let $f : U \rightarrow \mathbb{C}$ be holomorphic. Suppose that f^{-1} has only finitely many transcendental singularities over $\overline{\mathbb{C}}$ and that f has only finitely many simple islands over two Jordan domains in \mathbb{C} with disjoint closures. Then $U = \mathbb{C}$, and f is either a polynomial or an entire function satisfying (1.6).

Under the hypothesis that there exist $a_1, a_2 \in \mathbb{C}$ such that f has only finitely many simple a_j -points, the conclusions of Theorem 1.2 and the above remark were obtained by Goldberg and Tairova [19]. Their proof was based on topological arguments and it probably can be extended to a proof of Theorem 1.2.

Note that we have $\mu(U) \geq 1$ and hence $\mu(U) - 2 \geq -1$ for all islands U , with $\mu(U) - 2 = -1$ only for the at most finitely many simple islands. Thus $p < \infty$ implies in particular that $N < \infty$ and that $\mu(U) > 2$ for at most finitely many islands U .

Note also that if all islands over D_1 and D_2 are multiple, then $\mu(U) \geq 2$ for all islands U and thus $p \geq 2$. Thus Theorem C' follows from Theorem 1.2.

Theorems 1.1 and 1.2 concern entire functions. The analogous results do not hold if instead of entire functions we consider meromorphic functions which have no island over a domain containing ∞ .

Theorem 1.3 *Let D_1 and D_2 be Jordan domains in $\overline{\mathbb{C}}$ with disjoint closures. Then, given $\rho \in [0, \infty)$, there exists a meromorphic function f of order ρ which has no island over D_1 and D_2 .*

Theorem 1.4 *Let D be a Jordan domain in $\overline{\mathbb{C}}$ and let $a_1, a_2 \in \overline{\mathbb{C}} \setminus D$ be distinct. Then, given $\rho \in [0, \infty)$, there exists a meromorphic function f of order ρ which has no island over D and for which all a_1 -points and a_2 -points are multiple.*

The functions f constructed in the proofs of Theorems 1.3 and 1.4 have the property that

$$T(r, f) \geq c (\log r)^2 \tag{1.7}$$

for some $c > 0$ and all large r .

Question 1.1 Do we have (1.7) for every meromorphic function f satisfying the conditions of Theorems 1.3 or 1.4?

Remark After submission of this paper, Jim Langley communicated to us the following argument showing that the answer to Question 1.1 is positive in the case of Theorem 1.3. Thus suppose that D_1 and D_2 are as in Theorem 1.3 and that f is a meromorphic function satisfying $\liminf_{r \rightarrow \infty} T(r, f)/(\log r)^2 = 0$. It is shown in [23, Proof of Theorem 6] that then there exists $w \in \overline{\mathbb{C}}$ and a sequence (t_n) tending to ∞ such that $f(z)$ tends to w uniformly on the circles $\{z: |z| = t_n\}$ as $n \rightarrow \infty$, with respect to the spherical metric. Let $j \in \{1, 2\}$ such that $w \notin D_j$. This implies that for large n the disk $\{z: |z| < t_n\}$ contains a component of $f^{-1}(D_j)$. This component need not be simply-connected. If it is multiply-connected, then any bounded complementary component contains a preimage of $f^{-1}(D_k)$ where $k \in \{1, 2\}, k \neq j$. We thus obtain a "nested" chain of alternating preimages of D_1 and D_2 . Such a chain must be finite, and its terminal member is simply-connected.

The following result says that in case (iii) there is no analogue of Theorem E if instead of multiple a_j -points one considers multiple islands over certain Jordan domains. In fact, it suffices to replace a_j -points by multiple islands over some Jordan domain for one j .

Theorem 1.5 Let $a_1, a_2, a_3 \in \overline{\mathbb{C}}$ be distinct and $\rho \in (0, \infty)$. Then there exists a meromorphic function f of order ρ and a Jordan domain D whose closure is contained in $\overline{\mathbb{C}} \setminus \{a_1, a_2, a_3\}$ such that all a_j -points of f are multiple for $j \in \{1, 2, 3\}$ and such that f has no simple island over D .

Question 1.2 Does the conclusion of Theorem 1.5 also hold for $\rho = 0$?

Theorem 1.5 deals with case (iii). We do not know whether there are analogous results for the cases (iv)–(vi).

Question 1.3 In cases (iv)–(vi), is there an analogue of Theorem E if instead of multiple a_j -points we consider multiple islands over Jordan domains D_j with disjoint closures?

The answer may depend on whether we replace multiple a_j -points by multiple islands over the D_j for all $j \in \{1, \dots, q\}$ or only for some j .

There exist functions of order 0 for which all but finitely many islands over D_j have multiplicity m_j . In fact, this is the case already in the situation of Theorem E. Given a_j and m_j as there there exists a function of order 0 for which all but finitely many a_j -points have multiplicity at least m_j . Such a function f was considered already by Teichmüller [29, p. 734]. It satisfies

$$T(r, f) \sim c (\log r)^2$$

as $r \rightarrow \infty$, for some $c > 0$. This leads to the following question analogous to Question 1.1.

Question 1.4 Let $f: \mathbb{C} \rightarrow \overline{\mathbb{C}}$ be meromorphic and $q \in \mathbb{N}$. Let D_1, \dots, D_q be Jordan domains in $\overline{\mathbb{C}}$ with pairwise disjoint closures and let $m_1, \dots, m_q \in \mathbb{N} \cup \{\infty\}$ satisfy (1.2). Suppose that, for all $j \in \{1, \dots, q\}$, all but finitely many islands over D_j have multiplicity at least m_j . Does there exist $c > 0$ such that (1.7) holds for all large r ?

2 Results Used in the Proofs

We begin with a classical result of Wiman [20, Chapter 5, Theorem 1.3].

Lemma 2.1 *Let f be a non-constant entire function. Suppose that there exists $C > 0$ such that $\min_{|z|=r} |f(z)| < C$ for all $r > 0$. Then $\rho(f) \geq 1/2$.*

The next result [20, Chapter 5, Theorem 1.2] is a version of the Denjoy–Carleman–Ahlfors theorem.

Lemma 2.2 *Let f be an entire function of finite order and let $C > 0$. Then $\{z \in \mathbb{C}: |f(z)| > C\}$ has at most $\max\{1, 2\rho(f)\}$ connected components.*

The following result [20, Chapter 5, Theorem 1.4] is a consequence of Lemma 2.2 and a result of Lindelöf. It is also called Denjoy–Carleman–Ahlfors theorem.

Lemma 2.3 *Let f be a non-constant entire function of finite order. Then f has at most $2\rho(f)$ finite asymptotic values.*

We will use some results about quasiconformal and quasiregular mappings. We refer to [24] for the definition and basic properties, noting that quasiregular mappings are called *quasiconformal functions* there. Let D be a domain and $f: D \rightarrow \overline{\mathbb{C}}$ be quasiregular. We use the notation

$$\mu_f(z) = \frac{f_{\bar{z}}(z)}{f_z(z)}, \quad K_f(z) = \frac{1 + |\mu_f(z)|}{1 - |\mu_f(z)|} \quad \text{and} \quad K(f) = \operatorname{ess\,sup}_{z \in D} |K_f(z)|.$$

The Hölder continuity of quasiconformal mappings [24, § II.4.2] yields the following result.

Lemma 2.4 *Let $\phi: \mathbb{C} \rightarrow \mathbb{C}$ be quasiconformal. Then $|\phi(z)| = O(|z|^{K(\phi)})$ as $|z| \rightarrow \infty$.*

A basic result [24, § V.1] in the theory of quasiconformal mappings says that there exist quasiconformal mappings with prescribed dilatation. One consequence of this is the following result.

Lemma 2.5 *Let $f: \mathbb{C} \rightarrow \overline{\mathbb{C}}$ be quasiregular. Then there exists a quasiconformal mapping $\phi: \mathbb{C} \rightarrow \mathbb{C}$ such that $f \circ \phi$ is meromorphic.*

The next result is known as the Teichmüller–Wittich–Belinskii theorem [24, § V.6].

Lemma 2.6 *Let $\phi: \mathbb{C} \rightarrow \mathbb{C}$ be quasiconformal. Suppose that*

$$\int_{\{z \in \mathbb{C}: |z| > 1\}} \frac{K_\phi(z) - 1}{x^2 + y^2} dx dy < \infty. \tag{2.1}$$

Then there exists $c \in \mathbb{C} \setminus \{0\}$ such that

$$\phi(z) \sim cz \text{ as } z \rightarrow \infty. \tag{2.2}$$

Remark The condition (2.1) is satisfied in particular if the set A of all z satisfying $|z| > 1$ where ϕ is not conformal satisfies

$$\int_A \frac{dx dy}{|z|^2} < \infty. \tag{2.3}$$

We denote the open and closed disk of radius r around a point $a \in \mathbb{C}$ by $D(a, r)$ and $\overline{D}(a, r)$. We also put $\mathbb{D} := D(0, 1)$.

Beurling and Ahlfors [11] characterized the homeomorphisms of $\partial\mathbb{D}$ which admit a quasiconformal extension to \mathbb{D} ; see [24, § II.7.1]. Such homeomorphisms are called quasisymmetric.

We will use the following sufficient condition for quasisymmetry. It is surely known, but we did not find a reference.

Lemma 2.7 *Let $h: \partial\mathbb{D} \rightarrow \partial\mathbb{D}$ be an orientation-preserving homeomorphism. Suppose that there exists a finite subset A of $\partial\mathbb{D}$ such that h is continuously differentiable with non-zero derivative in $\partial\mathbb{D} \setminus A$. Suppose also that for all $a \in A$ there exists $\gamma_a > 0$ such that*

$$\frac{|h(ae^{it}) - h(a)|}{t^{\gamma_a}}$$

has one-sided, non-zero limits as $t \rightarrow 0^\pm$. Then h is quasisymmetric.

Sketch of proof The restriction of h to a closed arc which contains no point of A is clearly quasisymmetric. To see that h is quasisymmetric on an arc which has a point of A as one of its endpoints it is convenient to consider quasisymmetric mappings on \mathbb{R} rather than $\partial\mathbb{D}$. Here this claim follows since $t \mapsto t^\gamma$ is quasisymmetric on $[0, 1]$ and since the composition of quasisymmetric mappings is again quasisymmetric [21, Theorem 9]. Finally, quasisymmetry in the union of these intervals (or arcs) follows from [21, Theorem 3]. □

Let $B: \mathbb{D} \rightarrow \mathbb{D}$ be a Blaschke product of degree $d \geq 2$ fixing 0. Thus B has the form

$$B(z) = e^{i\alpha} z \prod_{k=1}^{d-1} \frac{z - a_k}{1 - \overline{a_k}z}$$

with $\alpha \in \mathbb{R}$ and $a_1, \dots, a_{d-1} \in \mathbb{D}$.

Suppose that $r \in (0, 1)$ is such that all zeros of B are in $D(0, r)$. By a result of Walsh [33, Theorem 1], all critical points of B in \mathbb{D} are also contained in $D(0, r)$. Since $B(\overline{D}(0, r)) \subset D(0, r)$ by Schwarz's lemma we see that $B^{-1}(D(0, r)) \setminus \overline{D}(0, r)$ is a ring domain.

Branner and Fagella [16, p. 163] showed that there exists a quasiregular mapping $A: \mathbb{D} \rightarrow \mathbb{D}$ such that $A(z) = B(z)$ for $z \in \mathbb{D} \setminus B^{-1}(D(0, r))$ while $A(z) = z^d$ for $z \in \overline{D}(0, r)$. Moreover, $A(z) \neq 0$ for $z \in \mathbb{D} \setminus \{0\}$.

They state this only for the case $d = 2$, but their proof extends to the general case. Note that in order to prove this result one has to define the mapping in the ring domain $B^{-1}(D(0, r)) \setminus \overline{D}(0, r)$. The construction of this mapping is done in [16, Exercise 2.3.3] for arbitrary degree.

Branner and Fagella also note that we may choose A to depend continuously on B . This implies that the dilatation $K(A)$ of A also depends continuously on B . Restricting to a compact set of Blaschke products B we find that the dilatation of the corresponding quasiregular maps A is uniformly bounded.

We summarize the above discussion in the following result.

Lemma 2.8 *Let $0 < r_1 < r_2 < 1$ and let $B: \mathbb{D} \rightarrow \mathbb{D}$ be a Blaschke product of degree $d \geq 2$. Suppose that $B(0) = 0$ and that the zeros of B are contained in $D(0, r_1)$. Then there exists a quasiregular mapping $A: \mathbb{D} \rightarrow \mathbb{D}$ and a neighborhood W of $\partial\mathbb{D}$ such that $A(z) = z^d$ for $z \in D(0, r_2)$, $A(z) = B(z)$ for $z \in W \cap \mathbb{D}$ and $A(z) \neq 0$ for $z \in \mathbb{D} \setminus \{0\}$.*

Moreover, there exists a constant C depending only on r_1, r_2 and d such that A may be chosen to satisfy $K(A) \leq C$.

We will also need the following result.

Lemma 2.9 *Let U be a simply-connected, unbounded domain in \mathbb{C} which is bounded by piecewise analytic curves. Suppose that each disk $D(0, t)$ intersects only finitely many of these boundary curves.*

Let $f: \overline{U} \rightarrow \mathbb{C}$ be a bounded, continuous function which is holomorphic in U . Suppose that there exist $r, R > 0$ such that $|f(z)| > r$ for all $z \in U$ while $|f(z)| = r$ for all $z \in \partial U$ satisfying $|z| > R$. Then there exists a curve γ tending to ∞ in U and $a \in \partial D(0, r)$ such that $f(z) \rightarrow a$ as $z \rightarrow \infty$ on γ .

Proof Let $\phi: \mathbb{D} \rightarrow U$ be a conformal mapping. The hypotheses imply that the boundary of U in $\overline{\mathbb{C}}$ is locally connected. Thus ϕ has a continuous extension $\phi: \overline{\mathbb{D}} \rightarrow \overline{U} \cup \{\infty\}$.

Put $E = \phi^{-1}(\infty)$. Then E is a compact subset of $\partial\mathbb{D}$. By a result of Beurling [26, Theorem 9.19], E has logarithmic capacity zero. Put $g = f \circ \phi$. There exists an open arc A containing E such that $|g(z)| = r$ for $z \in A \setminus E$. Noting that $|g(z)| > r$ and thus $g(z) \neq 0$ for $z \in \mathbb{D}$, we deduce from the Schwarz reflection principle that g can be extended to a function holomorphic in $\mathbb{D} \cup (A \setminus E) \cup (\mathbb{C} \setminus \overline{\mathbb{D}})$.

Since E has logarithmic capacity zero, it also has analytic capacity zero; see, e.g., [34, Proposition 3.5]. Since g is bounded this yields that g has a holomorphic extension to $\mathbb{D} \cup A \cup (\mathbb{C} \setminus \overline{\mathbb{D}})$; see [34, Appendix II].

Next we note that U is unbounded and that ∞ is accessible in U . Thus $E \neq \emptyset$. Taking a point $\xi \in E$ we have $\phi(t\xi) \rightarrow \infty$ as $t \rightarrow 1$. The conclusion follows for the curve $\gamma: [0, 1) \rightarrow \mathbb{C}$, $\gamma(t) = \phi(t\xi)$, and $a = g(\xi)$. \square

We will also use the following result about the growth of composite meromorphic functions; see [10, Satz 2.3 and Satz 5.7] and [7, Corollary 4].

Lemma 2.10 *Let f be a meromorphic function and g be an entire function. Then*

$$\rho(g) \liminf_{r \rightarrow \infty} \frac{\log T(r, f)}{\log \log r} \leq \rho(f \circ g) \leq \rho(g) \limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log \log r}. \tag{2.4}$$

In [10, Satz 5.7] the left inequality in (2.4) is proved only under the additional hypothesis that

$$\limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log \log r} < \infty.$$

This additional hypothesis is removed in [7, Corollary 4]. It will be satisfied, however, in our applications. In fact, we will consider only the case where $(\log T(r, f))/\log \log r$ tends to a finite limit. In this case we deduce from (2.4) that

$$\rho(f \circ g) = \rho(g) \lim_{r \rightarrow \infty} \frac{\log T(r, f)}{\log \log r}. \tag{2.5}$$

3 Proofs of Theorems 1.1–1.5

Proof of Theorem 1.1 Let f be a transcendental entire function which has only finitely many islands over some Jordan domain D . Then there exists a connected component U of $f^{-1}(D)$ which is unbounded. This implies that the minimum modulus $\min_{|z|=r} |f(z)|$ is bounded. Wiman's theorem (Lemma 2.1) now yields that $\rho(f) \geq 1/2$.

For the converse result, let D be a Jordan domain in \mathbb{C} . Without loss of generality we may assume that $0 \in D$. Thus there exists $\varepsilon > 0$ such that $D(0, \varepsilon) \subset D$. For $\rho \in [1/2, 1)$ we consider the function g defined by

$$g(z) = \prod_{n=1}^{\infty} \left(1 - \frac{z}{n^{1/\rho}}\right)$$

Then g is a function of order ρ which has only positive zeros and satisfies $g(x) \rightarrow 0$ as $x \rightarrow \infty$. For $\rho \in (1/2, 1)$ the last statement follows from, e.g., [15, Theorem 4.1.8] while for $\rho = 1/2$ it follows from the explicit representation

$$g(z) = \frac{\sin(\pi\sqrt{z})}{\pi\sqrt{z}}.$$




Fig. 1 The graphs Γ (right) and Γ^* (left)

We conclude that for sufficiently small $\delta > 0$ the function $f := \delta g$ has no island over $D(0, \varepsilon)$ and hence no island over D .

To obtain functions of order $\rho \in [1, \infty)$ we write $\rho = p\rho_0$ with $p \in \mathbb{N}$ and $\rho_0 \in [1/2, 1)$. Choosing g of order ρ_0 and $\delta > 0$ as above we find that the function f given by $f(z) = \delta g(z^p)$ has order ρ and that f has no island over D . \square

Proof of Theorem 1.2 Let $r > 0$ be such that the closures of D_1 and D_2 and all finite asymptotic values are contained in $D(0, r)$. Note that by the Denjoy–Carleman–Ahlfors theorem (Lemma 2.3) there are only finitely many asymptotic values. We also assume that $\partial D(0, r)$ contains no critical value. This can be achieved since the set of critical values is countable.

We consider the following graph Γ on the sphere $\overline{\mathbb{C}}$. It has two vertices, which we denote by \times and \circ and which lie on $\partial D(0, r)$, and three edges, two of which are arcs on the circle $\partial D(0, r)$ connecting \times and \circ , while the third edge is a crosscut of $D(0, r)$ connecting \times and \circ which separates $\overline{D_1}$ and $\overline{D_2}$ and which contains no critical or asymptotic value. The components of the complement of the set of vertices and edges are called faces. We then have three faces. The face $\overline{\mathbb{C}} \setminus \overline{D(0, r)}$ will be denoted by F_∞ and, for $j \in \{1, 2\}$, the face containing $\overline{D_j}$ will be denoted by F_j .

We consider the preimage $\Gamma^* = f^{-1}(\Gamma)$ of Γ . It yields a partition of the plane into faces, edges and vertices. It is similar to a *line complex*; see [20, Chapter 7, Section 4]. One difference is that a line complex is always connected, while Γ^* need not be connected. We will see, however, that Γ^* is connected if r is chosen sufficiently large.

For our purposes only the topology of Γ^* is relevant. Thus we do not distinguish between the preimage Γ^* and its image under a homeomorphism of the plane. In figures like Fig. 1 we usually draw only a homeomorphic image of $\Gamma^* = f^{-1}(\Gamma)$, not the true preimage. In such figures we use the labels ①, ② and \otimes for the faces F_1, F_2 and F_∞ as well as for their preimages in Γ^* . The same remark applies to the vertices \times and \circ .

Figure 1 shows Γ and Γ^* for a function having only islands of multiplicity 2 over D_1 and D_2 , and no critical or finite asymptotic values outside D_1 and D_2 . An example is given by the sine or cosine function if $-1 \in D_1$ and $1 \in D_2$.

Fig. 2 Example of a graph Γ^*




Figure 2 shows Γ^* for a function having one simple island and one island of multiplicity 4 over D_1 , one island of multiplicity 3 over D_2 , while all other islands over D_1 and D_2 have multiplicity 2.


Clearly, Γ^* is a bipartite, properly embedded graph. (Here “properly embedded” means that it not only lies in the plane, but also that its vertices and edges do not accumulate to a point in the plane.) We list some properties of this graph.

- (a) Each vertex has degree 3 and lies on the boundaries of three faces with labels ①, ② and ∞ .
- (b) There are only finitely many digons labeled ① or ②. (A digon is a face with only two boundary vertices and two boundary edges.)
- (c) Each face labeled ∞ is unbounded and there are only finitely many such faces.
- (d) Each face labeled ① or ② is bounded.
- (e) If r is large enough, then Γ^* is connected.

Property (a) is obvious from the definition of Γ^* . To prove (b) we note that for a digon V labeled ① with $j \in \{1, 2\}$ the mapping $f: V \rightarrow F_j$ is bijective and hence V contains a simple island over D_j . Thus (b) follows from the hypothesis that there are only finitely many simple islands over D_1 and D_2 .

To prove (c) we note that in a face labeled ∞ the function f is unbounded. Thus such a face is unbounded and by the Denjoy–Carleman–Ahlfors theorem (Lemma 2.2) there are only finitely many such faces.

Fig. 3 The graph Δ corresponding to the graph Γ^* in Fig. 2



To prove (d), let V be an unbounded face labeled $\textcircled{1}$ with $j \in \{1, 2\}$. Then ∂V contains an infinite chain $\cdots - o - \times - o - \times - \cdots$. Since every vertex lies on the boundary of a face labeled $\textcircled{\otimes}$, and since there are only finitely many faces labeled $\textcircled{\otimes}$, there exist a face V' labeled $\textcircled{\otimes}$ such that that this chain contains infinitely many vertices which lie on both ∂V and $\partial V'$. Let v_1 and v_2 be two such vertices and let v_0 be a vertex between them. Then there exists a face V'' labeled $\textcircled{\otimes}$ such that $v_0 \in \partial V''$. Connecting v_1 and v_2 by a crosscut in V' we see that V'' must intersect this crosscut. Thus $V'' = V'$. We conclude that there are infinitely many triplets of adjacent vertices which are on the boundary of both V and V' . As the middle vertex of such a triplet has degree 3, it must be connected to one of the other two vertices of the triplet by a double edge. In other words, such a triplet leads to a digon. Since there are only finitely many digons by (b), this is a contradiction, completing the proof of (d).

As a preparation for the proof of (e), we note that there is a one-to-one correspondence between the components of Γ^* and the components of $f^{-1}(\overline{D}(0, r))$. In fact, given a component C of Γ^* , the corresponding component of $f^{-1}(\overline{D}(0, r))$ is obtained by “filling” those faces labeled $\textcircled{1}$ or $\textcircled{2}$ whose boundaries are contained in C . Note that these faces are all bounded by (d). Reversing this process, one obtains a component of Γ^* from a component of $f^{-1}(\overline{D}(0, r))$. We conclude from this that the number of components of Γ^* is a non-increasing function of r .

For the proof of (e), as well as some subsequent arguments, it will be convenient to consider a graph Δ which in some sense is dual to Γ^* : To each bounded face V we associate a point $v \in V$. (Recall that by (c) and (d) the bounded faces are those labeled $\textcircled{1}$ or $\textcircled{2}$.) These points v are the vertices of Δ . Two vertices are connected by an edge if the two faces of Γ^* that contain these vertices share a common edge in Γ^* . We take this edge in Δ to be in the union of the closures of the two faces in Γ^* , crossing the edge in Γ^* which separates these faces once. So f is bounded on the set of edges of Δ . Figure 3 shows the graph Δ corresponding to the graph Γ^* in Fig. 2. (Again we only consider a homeomorphic image.)

Similarly as above we see that there is a one-to-one correspondence between the components of Δ and the components of $f^{-1}(\overline{D}(0, r))$, and hence to the components of Γ^* . To pass from a component of Δ to a component of $f^{-1}(\overline{D}(0, r))$ consider for a vertex v the closure \overline{V} of the face V that contains v . Taking the union of the closures \overline{V} over all v in a component of Δ yields a component of $f^{-1}(\overline{D}(0, r))$. This process can be reversed, so indeed there is the one-to-one correspondence mentioned. Bounded components of Γ^* correspond to bounded components of Δ and unbounded components of Γ^* correspond to unbounded components of Δ .

Next we show that every component of Δ is a tree. Otherwise there exists a closed curve and in fact a Jordan curve in Δ . We thus have a Jordan curve γ contained in the union of the faces labeled ① or ② which crosses each edge of Γ^* at most once. The interior of γ contains some vertex, and this vertex is on the boundary of a face V labeled \otimes . Since V is unbounded, it must intersect γ , which is a contradiction. Thus every component of Δ is a tree.

For a face V labeled ① or ② the degree of the vertex v of Δ such that $v \in V$ is given by $\mu(V)$. We will also write $\mu(v)$ instead of $\mu(V)$. By hypothesis, there are only finitely many vertices of degree 1. Since each bounded tree contains a vertex of degree 1, we conclude that Δ has only finitely many bounded components. This implies that Δ also has an unbounded component.

We connect the finitely many bounded components of Δ by some paths to an unbounded component. Then there exists $C > 0$ such that $|f(z)| \leq C$ for z on one of these paths. This implies that $r < C$ and hence shows that if r is chosen large enough at the beginning, then all components of Δ are unbounded. It follows that all components of Γ^* are unbounded.

To complete the proof of (e), suppose that Γ^* is disconnected. Then there exists a face V whose boundary contains two (unbounded) components of Γ^* . We connect these two components by a crosscut γ in V . This crosscut separates V into two domains V_1 and V_2 .

Since Γ^* has no bounded components, V and hence V_1 and V_2 are simply-connected. By (c), the face V is labeled \otimes and hence f is unbounded there. We claim that f is unbounded in each of the domains V_1 and V_2 . In fact, suppose that f is bounded in V_j where $j \in \{1, 2\}$. Lemma 2.9 yields that f has an asymptotic value of modulus r with asymptotic path contained in V_j . This contradicts our assumption made at the beginning that all asymptotic values are contained in $D(0, r)$. Hence f is unbounded in both V_1 and V_2 .

Again there exists $C > 0$ such that $|f(z)| \leq C$ for z on the crosscut γ . Increasing r to some value greater than C thus increases the number of components of $\{z \in \mathbb{C} : |f(z)| > r\}$. Since, by Lemma 2.2, the number of these components is at most $\max\{1, 2\rho(f)\}$ we conclude that if r is large enough, then Γ^* has only one component; that is, Γ^* is connected. This yields (e) and completes the proof of statements (a)–(e).

As mentioned, a line complex is always connected, while Γ^* and Δ need not be. But (e) says that this holds for large r .

Let p be the number of unbounded faces of Γ^* . Thus p is the number of components of $\{z \in \mathbb{C} : |f(z)| > r\}$. An argument similar to the one above about the one-to-one

correspondence between the components of Γ^* and the components of Δ yields that p equals the number of complementary components of Δ .

Next we show that

$$p = 2 + \sum_v (\mu(v) - 2), \tag{3.1}$$

with the sum taken over all vertices v of Δ . The proof of (3.1) will only use that Δ is an infinite properly embedded tree and that Δ has only finitely many vertices of degree 1. To prove (3.1) suppose first that the number of vertices v for which $\mu(v) \neq 2$ is finite. If there are no such vertices, then Δ is an infinite chain $\cdots - \circ - \times - \circ - \times - \cdots$. In this case $\sum_v (\mu(v) - 2) = 0$ and thus (3.1) holds.

Let now $n \in \mathbb{N}$ and suppose that (3.1) has been proved if the number of vertices v with $\mu(v) \neq 2$ is less than n . Let Δ be a graph for which there are n such vertices. If Δ has a vertex of degree 1, we replace this vertex by an infinite half-chain $\circ - \times - \circ - \cdots$ (or $\times - \circ - \times - \cdots$). The new graph Δ' obtained has only $n - 1$ vertices v with $\mu(v) \neq 2$ and both left and right side of (3.1) differ for Δ and Δ' by 1. Thus (3.1) holds for Δ since it holds for Δ' by induction hypothesis.

If Δ does not have a vertex of degree 1, then Δ has a vertex of degree at least 3, and there exists such a vertex v_0 with $\mu(v_0) \geq 3$ such that v_0 bounds $\mu(v_0) - 1$ half-chains $\circ - \times - \circ - \cdots$ (or $\times - \circ - \times - \cdots$), while all other vertices of Δ are on the remaining part of Δ . Replacing the $\mu(v_0) - 1$ half-chains by only one such half-chain yields a graph Δ' which has only $n - 1$ vertices v with $\mu(v) \neq 2$. The left and right side of (3.1) differ for Δ and Δ' by $\mu(v_0) - 2$. Again (3.1) holds for Δ since it holds for Δ' by induction hypothesis.

This proves (3.1) if the number of vertices v for which $\mu(v) \neq 2$ is finite. However, minor modifications of the argument show that if the number of vertices v for which $\mu(v) \neq 2$ is infinite, then the number of complementary components of Δ is infinite. Thus we also see that if p is finite, then the number of vertices v with $\mu(v) \neq 2$ is finite. This means that the sum in (3.1) is finite.

To pass from (3.1) to (1.5), let v be a vertex of Δ and let V be the face of Γ^* containing v . Then V is labeled \textcircled{j} with $j \in \{1, 2\}$ and it is an island over F_j . Its boundary contains $\mu(V)$ vertices \times and $\mu(V)$ vertices \circ . So it is an n -gon with $n = 2\mu(V)$.

Also, V contains at least one island over D_j . Let U_1, \dots, U_m be the islands over D_j that are contained in V . Then

$$\mu(V) = \sum_{k=1}^m \mu(U_k). \tag{3.2}$$

Let $N(V)$ be the number of critical points of f contained in V that are not mapped to D_j . Since V contains $\mu(V) - 1$ critical points and each U_k contains $\mu(U_k) - 1$ critical points, we deduce from (3.2) that

$$N(V) = \mu(V) - 1 - \sum_{k=1}^m (\mu(U_k) - 1) = m - 1.$$

This allows to rewrite (3.2) in the form

$$\mu(V) - 2 = 2m - 2 + \sum_{k=1}^m (\mu(U_k) - 2) = 2N(V) + \sum_{k=1}^m (\mu(U_k) - 2). \tag{3.3}$$

To obtain (1.5) we want to sum this over all V . Note that there may be (finitely many) vertices v in Δ of degree 2 for which the corresponding face V contains one critical point and two simple islands over D_j . For such a face V both the left and right hand side of (3.3) are equal to 0. Whether we include such a face or not will not affect the sum. Similarly, the sum in (1.5) is unchanged if we remove this critical point and the two simple islands.

Summing (3.3) over all bounded faces V and combining this with (3.1) thus yields (1.5) with $\sum_V N(V)$ instead of N , where the sum is taken over all bounded faces V of Γ^* .

To complete the proof of (1.5) we thus need to show that

$$N = \sum_V N(V). \tag{3.4}$$

In other words, we have to show that the unbounded faces of Γ^* contain no critical points of f . In order to do so, let V be an unbounded face of Γ^* . Since V is simply-connected and bounded by a single curve, unbounded in both directions, there exists a conformal mapping ϕ from \mathbb{D} to V such that ϕ has a continuous extension to $\overline{\mathbb{D}} \setminus \{1\}$, mapping $\partial\mathbb{D} \setminus \{1\}$ to ∂V and satisfying $\phi(z) \rightarrow \infty$ as $z \rightarrow 1$. Then

$$u(z) = \log \frac{|f(\phi(z))|}{r}$$

defines a positive harmonic function in \mathbb{D} which extends continuously to $\overline{\mathbb{D}} \setminus \{1\}$, with $u(z) = 0$ for $z \in \partial\mathbb{D} \setminus \{1\}$. This yields that u is a constant multiple of the Poisson kernel; see, e.g., [6, Theorem 6.19]. But this implies that $\log f \circ \phi$ has no critical points in \mathbb{D} . Hence f has no critical point in V . This completes the proof of (3.4) and hence of (1.5).

For $j \in \{1, 2\}$ we choose $a_j \in D_j$ and a conformal mapping $\psi_j: F_j \rightarrow \mathbb{D}$ satisfying $\psi_j(a_j) = 0$. For a face V labeled \textcircled{j} we choose a conformal mapping $\tau_V: \mathbb{D} \rightarrow V$ with $\tau_V(0) \in f^{-1}(a_j)$. Then $B := \psi_j \circ f \circ \tau_V$ is a Blaschke product and $B(0) = 0$. The zeros of B are contained in $\tau_V^{-1}(f^{-1}(D_j))$. We will show that there exists $r_1 \in (0, 1)$, depending only on the choice of the F_j but not on V , such $\tau_V^{-1}(f^{-1}(D_j))$ and hence the zeros of B are contained in $D(0, r_1)$.

In order to do so we note that since $p < \infty$, for all but finitely many faces V labeled \textcircled{j} there is exactly one island U of multiplicity 2 over D_j contained in V , but

there are no further islands over D_j contained in V , and there are no critical points in $V \setminus U$. For such a face V and island U we then have $f^{-1}(D_j) \cap V = U$ and the mapping $f: V \setminus \overline{U} \rightarrow F_j \setminus \overline{D_j}$ is a covering map of degree 2. Denoting by $\text{mod}(\Omega)$ the modulus of a ring domain Ω we find that

$$\begin{aligned} \text{mod}\left(\mathbb{D} \setminus \overline{\tau_V^{-1}(f^{-1}(D_j))}\right) &= \text{mod}\left(V \setminus \overline{f^{-1}(D_j)}\right) \\ &= \text{mod}(V \setminus \overline{U}) = \frac{1}{2} \text{mod}(F_j \setminus \overline{D_j}). \end{aligned}$$

Thus the modulus of $\mathbb{D} \setminus \overline{\tau_V^{-1}(f^{-1}(D_j))}$ is bounded below. This implies that there exists $r_1 \in (0, 1)$ such that $\tau_V^{-1}(f^{-1}(D_j)) \subset D(0, r_1)$ for all such faces V . Increasing r_1 if necessary we may assume that this inclusion also holds for the finitely many faces V where $f: V \setminus \overline{f^{-1}(D_j)} \rightarrow F_j \setminus \overline{D_j}$ is not a covering of degree 2.

We now choose $r_2 \in (r_1, 1)$ and apply Lemma 2.8 to B . With the function A obtained from this lemma we define $f_V: V \rightarrow F_j$ by $f_V = \psi_j^{-1} \circ A \circ \tau_V^{-1}$. Thus f_V is a quasiregular mapping having one a_j -point of multiplicity $\mu(V)$ and no other a_j -point in V , and there exists a neighborhood W of ∂V with $f_V(z) = f(z)$ for $z \in V \cap W$. Since there are only finitely many faces V labeled \textcircled{j} with $j \in \{1, 2\}$ for which the degree of the mapping $f_V: V \rightarrow F_j$ is greater than 2, Lemma 2.8 also yields that we may choose the mappings f_V with uniformly bounded dilatation.

We now define a mapping $g: \mathbb{C} \rightarrow \mathbb{C}$ by putting $g(z) = f_V(z)$ if $z \in V$ for such a face V , and $g(z) = f(z)$ otherwise. Then g is quasiregular. By Lemma 2.5 there exists a quasiconformal homeomorphism $\phi: \mathbb{C} \rightarrow \mathbb{C}$ such that the mapping $h := g \circ \phi$ is entire. It follows that all except possibly finitely many a_j -points are multiple, for $j \in \{1, 2\}$. Moreover, Lemma 2.4 yields that $\rho(h) \leq K(\phi)\rho(f) < \infty$.

As in (1.3) we now consider

$$R(z) = \frac{h'(z)^2}{(h(z) - a_1)(h(z) - a_2)}$$

and deduce from the lemma on the logarithmic derivative that R is a rational function. Assuming without loss of generality that $a_{1,2} = \pm 1$ we find as in (1.4) that h has the form

$$h(z) = \cosh\left(\int \sqrt{R(z)} dz\right). \tag{3.5}$$

We saw above that $\{z \in \mathbb{C}: |f(z)| > r\}$ has p components if r is sufficiently large. This implies that $\{z \in \mathbb{C}: |h(z)| > r\}$ has p components for large r . Together with (3.5) this yields that

$$\int \sqrt{R(z)} dz \sim \alpha z^{p/2} \quad \text{and hence} \quad R(z) \sim \beta z^{p-2}$$

as $z \rightarrow \infty$, for certain $\alpha, \beta \in \mathbb{C} \setminus \{0\}$. It follows that

$$\log M(r, h) \sim \gamma r^{\rho/2} \tag{3.6}$$

for some $\gamma > 0$.

The function g agrees with f in $\mathbb{C} \setminus f^{-1}(F_1 \cup F_2)$. Thus g is holomorphic in a neighborhood of any point $z \in \mathbb{C}$ for which $|g(z)| > r$. Hence for large r the function ϕ is conformal in a neighborhood of z if $|h(z)| = |g(\phi(z))| > r$. Thus the set of points in $\mathbb{C} \setminus \mathbb{D}$ where ϕ is not conformal is contained in

$$A := \{z \in \mathbb{C} : |z| > 1 \text{ and } |h(z)| < r\},$$

provided r is sufficiently large. It can be deduced from (3.5) that A satisfies (2.3). Lemma 2.6 and the remark following it now imply that there exists $a \in \mathbb{C}$ such that $\phi(z) \sim az$ as $z \rightarrow \infty$. Now (1.6) follows from (3.6). \square

Proof of Theorem 1.3 Without loss of generality we may assume that $0 \in D_1$ and $\infty \in D_2$. Then there exists $\varepsilon > 0$ such that $D(0, \varepsilon) \subset D_1$ and $\{z \in \mathbb{C} : |z| > 1/\varepsilon\} \cup \{\infty\} \subset D_2$. We put $a = 1 + \varepsilon$ and

$$f(z) = \prod_{k=1}^{\infty} \frac{1 - \frac{z}{a^k}}{1 + \frac{z}{a^k}}. \tag{3.7}$$

It is easy to see that the infinite product converges and thus defines a function f meromorphic in \mathbb{C} .

For $x \geq a$ there exists $n \in \mathbb{N}$ and $\eta \in [0, 1)$ such that $x = a^{n+\eta}$. Hence

$$|f(x)| \leq \left| \frac{1 - \frac{x}{a^n}}{1 + \frac{x}{a^n}} \right| = \frac{a^\eta - 1}{a^\eta + 1} \leq \frac{a - 1}{a + 1} < \varepsilon.$$

This implies that f has no island over $D(0, \varepsilon)$ and hence no island over D_1 . An analogous argument shows that f has no island over $\{z \in \mathbb{C} : |z| > 1/\varepsilon\} \cup \{\infty\}$ and hence no island over D_2 .

Standard arguments show that the function f defined by (3.7) has order 0 and in fact that

$$T(r, f) \sim c (\log r)^2 \tag{3.8}$$

for some $c > 0$ as $r \rightarrow \infty$.

This completes the proof for the case that $\rho = 0$. To deal with the general case we note that if g is any entire function, then $f \circ g$ has no island over D_1 and D_2 . Since

$$\lim_{r \rightarrow \infty} \frac{\log T(r, f)}{\log \log r} = 2 \tag{3.9}$$

by (3.8) we deduce from (2.5) that $\rho(f \circ g) = 2\rho(g)$. Thus we can achieve that $f \circ g$ has the preassigned order ρ by choosing g with $\rho(g) = \rho/2$. \square

Proof of Theorem 1.4 Without loss of generality we may assume that $a_{1,2} = \pm 2i$ and that $\infty \in D$. Let $\delta \in (0, 1)$ and $\beta \in (0, 1/2)$ and let U be the domain which contains the imaginary axis and which is bounded by the curve

$$\gamma : [0, \infty) \rightarrow \mathbb{C}, \quad \gamma(t) = \delta + e^{i\beta\pi}t,$$

and the curves $-\gamma$, $\bar{\gamma}$ and $-\bar{\gamma}$ obtained from γ by reflections. The idea is to glue the restriction (of a modification) of $2 \sinh(\arcsin z)$ to the domain U and the restriction of the function given by (3.7) to a half-plane.

In order to do so we note that $\arcsin \gamma$ is an injective curve in the first quadrant which connects $\arcsin \delta$ with ∞ . A computation shows that

$$\arcsin \gamma(t) = \left(\frac{1}{2} + \beta\right)\pi - i(\log t + \log 2) + O\left(\frac{1}{t}\right) \tag{3.10}$$

as $t \rightarrow \infty$.

Next we consider $G(z) = f(-z)$, with the function f from (3.7); that is,

$$G(z) = \prod_{k=1}^{\infty} \frac{1 + \frac{z}{a^k}}{1 - \frac{z}{a^k}}, \tag{3.11}$$

with some $a > 1$ to be determined later. Denoting by $\log G$ the branch of the logarithm with $\log G(0) = 1$ we find that

$$\log G(it) = \sum_{k=1}^{\infty} \log \frac{1 + \frac{it}{a^k}}{1 - \frac{it}{a^k}} = 2i \sum_{k=1}^{\infty} \arg\left(1 + \frac{it}{a^k}\right) = 2ih(t) \tag{3.12}$$

with

$$h(t) := \sum_{k=1}^{\infty} \arctan\left(\frac{t}{a^k}\right). \tag{3.13}$$

We have

$$\int_1^{\infty} \arctan\left(\frac{t}{a^s}\right) ds \leq h(t) \leq \int_0^{\infty} \arctan\left(\frac{t}{a^s}\right) ds. \tag{3.14}$$

Now

$$\int_0^{\infty} \arctan\left(\frac{t}{a^s}\right) ds = \frac{1}{\log a} \int_0^t \frac{\arctan u}{u} du \sim \frac{\pi}{2 \log a} \log t$$

as $t \rightarrow \infty$. Together with (3.14) this yields that

$$h(t) = \frac{\pi}{2 \log a} \log t + O(1) \tag{3.15}$$

as $t \rightarrow \infty$. It follows easily from (3.13) that h is increasing and concave. This implies that (3.15) can be improved to

$$h(t) = \frac{\pi}{2 \log a} \log t + \eta + o(1)$$

for some $\eta \in \mathbb{R}$ as $t \rightarrow \infty$. Hence

$$\log G(it) = i \frac{\pi}{\log a} \log t + 2i\eta + o(1) \tag{3.16}$$

as $t \rightarrow \infty$ by (3.12).

Let now V be the domain bounded by the curves γ and $\bar{\gamma}$ which contains the interval (δ, ∞) . Thus

$$V = \{z \in \mathbb{C} : |\arg(z - \delta)| < \beta\pi\}.$$

Let $b > 0$ and define

$$H: V \rightarrow \mathbb{C}, \quad H(z) = e^b G((z - \delta)^{1/(2\beta)}).$$

It follows from (3.16) that

$$\log H(\gamma(t)) = b + \log G(it^{1/(2\beta)}) = b + i \left(\frac{\pi}{2\beta \log a} \log t + 2\eta \right) + o(1) \tag{3.17}$$

as $t \rightarrow \infty$. This holds for any choice of a, b and β . We choose $\beta = \pi/(2 \log a)$ and $b = (1/2 + \beta)\pi$. Note that this still leaves the possibility to choose a later. Then the right hand sides of (3.10) and (3.17) have the same asymptotics as $t \rightarrow \infty$, apart from an additive constant.

Using interpolation it can now be shown that there exists an odd quasiconformal mapping $\psi: U \rightarrow \mathbb{C}$, symmetric with respect to \mathbb{R} , which agrees with the arcsine in a neighborhood of the imaginary axis and which satisfies

$$\psi(\gamma(t)) = \log H(\gamma(t)) \quad \text{for all } t \geq 0. \tag{3.18}$$

Note that $\log H$ and hence ψ map the curve γ , and hence the boundary of V , to the line $\{z \in \mathbb{C} : \operatorname{Re} z = b\}$.

Next we consider the function $S: \{z \in \mathbb{C} : |\operatorname{Re} z| \leq b\} \rightarrow \mathbb{C}$,

$$S(z) = \begin{cases} \frac{2}{b}(\operatorname{Re} z + b)e^z - e^{-z} & \text{if } -b \leq \operatorname{Re} z \leq -\frac{b}{2}, \\ 2 \sinh z & \text{if } |\operatorname{Re} z| < \frac{b}{2}, \\ e^z + \frac{2}{b}(\operatorname{Re} z - b)e^{-z} & \text{if } \frac{b}{2} \leq \operatorname{Re} z \leq b. \end{cases} \tag{3.19}$$

Thus $S(z) = \pm e^{\pm z}$ if $\operatorname{Re} z = \pm b$. It is easy to see that S is quasiregular. Finally we define $F: \mathbb{C} \rightarrow \overline{\mathbb{C}}$ by

$$F(z) = \begin{cases} S(\psi(z)) & \text{if } z \in \overline{U}, \\ H(z) & \text{if } z \in V, \\ -H(-z) & \text{if } z \in -V. \end{cases}$$

Note that by (3.18) and (3.19) we $S(\psi(z)) = \exp \psi(z) = H(z)$ for $z \in \partial V$. Thus F defines a quasiregular mapping. Lemma 2.5 yields that there exists a quasiconformal homeomorphism $\phi: \mathbb{C} \rightarrow \mathbb{C}$ such that $f := F \circ \phi$ is meromorphic.

Noting that the hyperbolic sine has the totally ramified values $\pm i$ and recalling that we have assumed that $a_{1,2} = \pm 2i$ we see that all a_j -points are multiple for $j \in \{1, 2\}$. A similar argument as in the proof of Theorem 1.3 shows that given $\varepsilon > 0$, we can choose a in (3.11) such that f has no island over $\{z \in \mathbb{C}: |z| > 1/\varepsilon\} \cup \{\infty\}$. Choosing ε sufficiently small we conclude that f has no island over D .

Finally, we have

$$n(r, F) \sim \frac{1}{\beta \log a} \log r = \frac{2}{\pi} \log r \quad \text{and} \quad n\left(r, \frac{1}{F - a_j}\right) \sim \frac{2}{\pi} \log r$$

as $r \rightarrow \infty$, for $j \in \{1, 2\}$. Lemma 2.4 together with standard arguments now shows that f has order 0 and in fact that (3.9) holds.

This proves the theorem for $\rho = 0$. As at the end of the proof of Theorem 1.3 we can use this to obtain the result for any preassigned order $\rho \in (0, \infty)$ by considering $f \circ g$ instead of f for an entire function g satisfying $\rho(g) = \rho/2$. □

Proof of Theorem 1.5 The idea behind the construction is due to Künzi [22]. The details are somewhat different though.

An outline of the construction is as follows. We consider two elliptic functions g_1 and g_2 , both having periods 2 and $2i\tau$, where $\tau > 0$. We restrict g_1 and g_2 to the sectors

$$S_1 := \{z \in \mathbb{C}: |\operatorname{Im} z| \leq \tau \operatorname{Re} z\} \quad \text{and} \quad S_2 := \{z \in \mathbb{C}: |\operatorname{Im} z| \leq -\tau \operatorname{Re} z\}.$$

We will modify the g_j near ∂S_j to obtain quasiregular mappings $f_j: S_j \rightarrow \mathbb{C}$ satisfying

$$f_1(t(1 \pm i\tau)) = f_2(t(-1 \pm i\tau)). \tag{3.20}$$

With $\alpha := \arctan \tau$ this yields that

$$f_0(z) = \begin{cases} f_1(z^{2\alpha/\pi}) & \text{if } \operatorname{Re} z \geq 0, \\ f_2(-(-z)^{2\alpha/\pi}) & \text{if } \operatorname{Re} z \leq 0, \end{cases} \tag{3.21}$$

defines a quasiregular mapping $f_0: \mathbb{C} \rightarrow \overline{\mathbb{C}}$. By Lemma 2.5 there exists a quasiconformal mapping $\phi: \mathbb{C} \rightarrow \mathbb{C}$ such that $F := f_0 \circ \phi$ is meromorphic. The mapping f we

want to construct is then given by $f(z) = F(z^p)$ for some $p \geq 2$. Using Lemma 2.6 we will see that $\rho(f) = 4p\alpha/\pi$ so that we can achieve any preassigned positive order for f .

Moreover, both f_1 and f_2 will have the critical values a_1, a_2 and a_3 . The fourth critical value of f_1 will be different from that of f_2 . The domain D will be such that it contains these fourth critical values.

We now come to the details of the construction. Without loss of generality we may assume that $\{a_1, a_2, a_3\} = \{0, 1, \infty\}$. Let

$$R = \{x + iy : 0 \leq x \leq 1, 0 \leq y \leq \tau\}$$

and let g_1 be a conformal mapping from the interior of R onto the upper half-plane. The mapping g_1 extends continuously to the boundary of R and we may normalize it to satisfy $g_1(0) = 1, g_1(1) = \infty$ and $g_1(1 + i\tau) = -1$. Then $a := g_1(i\tau) \in (-1, 1)$. The mapping g_1 can be extended by reflections to an elliptic function with periods 2 and $2i\tau$.

The mapping g_1 can be expressed in terms of the Weierstrass \wp -function with these periods. In fact, if L is the fractional linear transformation satisfying $L(\infty) = 1, L(e_1) = \infty$ and $L(e_3) = -1$, then $g_1 = L \circ \wp$.

To define the quasiregular mapping $f_1 : S_1 \rightarrow \overline{\mathbb{C}}$ we put, for $m, n \in \mathbb{Z}$,

$$R_{m,n} = m + in\tau + R = \{x + iy : m \leq x \leq m + 1, n\tau \leq y \leq (n + 1)\tau\}.$$

If $R_{m,n} \subset S_1$, we put $f_1(z) = g_1(z)$ for $z \in R_{m,n}$. We also put $f_1(x) = g_1(x)$ for $0 \leq x < 1$.

It remains to define f_1 in $\Delta_{m,n} := S_1 \cap R_{m,n}$ for those $m, n \in \mathbb{Z}$ for which $R_{m,n} \not\subset S_1$, but the interior of $R_{m,n}$ intersects S_1 . (This is the case if $m \geq 0$ and $n = m$ or $n = -m - 1$.) Note that $\Delta_{m,n}$ is a triangle for such m and n . We begin by defining f_1 on $\Delta := \Delta_{0,0}$. In fact, we will first define f_1 on $\partial\Delta$. This will be done in such a way that it can be extended quasiconformally to the interior of Δ using Lemma 2.7.

Given that f_1 is defined already on $\partial\Delta \cap S_1$, it remains to define f_1 on $\partial\Delta \cap \partial S_1 = \{t(1 + i\tau) : 0 \leq t \leq 1\}$. To motivate the definition we note that g_1 maps $\partial\Delta \cap \partial S_1$ to a curve in the upper half-plane which connects 1 to -1 . We want to define f_1 such that it maps $\partial\Delta \cap \partial S_1$ to the semicircle $\{e^{it} : 0 \leq t \leq \pi\}$ which also connects 1 and -1 . The quasiconformal extension of f_1 will then map Δ to the domain $\Omega_1 := \{z \in \mathbb{C} : \text{Im } z > 0, |z| > 1\}$. Thus we make the ansatz

$$f_1(t(1 + i\tau)) = \exp(i\pi H(t)) \tag{3.22}$$

with a homeomorphism $H : [0, 1] \rightarrow [0, 1]$ satisfying $H(0) = 0$ and $H(1) = 1$.

We want to choose H such that the resulting mapping $f_1 : \partial\Delta \rightarrow \partial\Omega_1$ can be extended quasiconformally to Δ . Let $\sigma : \mathbb{D} \rightarrow \Delta$ and $\tau : \Omega_1 \rightarrow \mathbb{D}$ be conformal mappings. These mappings have continuous extensions to the boundaries so that we have a mapping $h := \tau \circ f_1 \circ \sigma : \partial\mathbb{D} \rightarrow \partial\mathbb{D}$. We thus want to choose H such that h satisfies the hypotheses of Lemma 2.7.

Since g_1 has (simple) critical points at 0 and $1 + i\tau$ and $f_1(z) = g_1(z)$ for $z \in \partial\Delta \cap S_1$, we find that this is the case if $H \in C^2[0, 1]$ with $H'(0) = H'(1) = 0$, $H''(0) \neq 0$, $H''(1) \neq 0$, and $H'(x) > 0$ for $0 < x < 1$. So we fix any such mapping H . We extend the mapping h to a quasiconformal self-mapping of \mathbb{D} . The corresponding extension of the mapping $f_1 : \partial\Delta \rightarrow \partial\Omega_1$ is then given by $f_1 := \tau^{-1} \circ h \circ \sigma^{-1} : \Delta \rightarrow \Omega_1$.

Next we define f_1 on $\Delta_{1,1}$. Again we define it first on $\partial\Delta_{1,1}$. As f_1 is defined on $\partial\Delta_{1,1} \cap S_1$ already, we have to define it only on $\partial\Delta_{1,1} \cap \partial S_1 = \{t(1+i\tau) : 1 \leq t \leq 2\}$. We do so by putting $f_1(t(1+i\tau)) = f_1((2-t)(1+i\tau))$ for $1 \leq t \leq 2$. As before the mapping $f_1 : \partial\Delta_{1,1} \rightarrow \overline{\mathbb{C}}$ can be extended quasiconformally to the interior of $\Delta_{1,1}$.

We have thus defined f_1 on $\Delta_{0,0} \cup \Delta_{1,1}$. We extend the definition to $\Delta_{m,m}$ with $m \geq 2$ by periodicity; that is, we put $f_1(z) = f_1(z - 2\lfloor m/2 \rfloor(1+i\tau))$ for $z \in \Delta_{m,m}$.

It remains to define f_1 in the still missing triangles in the lower half-plane, which are of the form $\Delta_{m,-m-1}$ with $m \in \mathbb{N}_0$. This we do by reflection in the real axis; that is, we put $f_1(z) = \overline{f_1(\bar{z})}$ for z in such a triangle.

We have thus defined the quasiregular mapping $f_1 : S_1 \rightarrow \overline{\mathbb{C}}$. To define the quasiregular mapping $f_2 : S_2 \rightarrow \overline{\mathbb{C}}$, we put

$$g_2(z) = -\overline{g_1(\bar{z} + 1 + i\tau)}.$$

Note that g_2 maps the rectangle $R_{-1,0}$ onto the upper half-plane and satisfies $g_2(0) = 1$, $g_2(i\tau) = \infty$, $g_2(-1+i\tau) = -1$ and $g_2(-1) = -g_1(i\tau) = -a$.

For $m, n \in \mathbb{Z}$ such that $R_{m,n} \subset S_2$ and $z \in \mathbb{R}_{m,n}$ we put $f_2(z) = g_2(z)$. We also put $f_2(x) = g_2(x)$ for $-1 < x \leq 0$ and define f_2 on ∂S_2 by (3.20). This defines f_2 on the sector S_2 except for the interior of the triangles $R_{-m,m-1} \cap S_2$ and $R_{-m,-m} \cap S_2$ with $m \in \mathbb{N}$. As before we can extend f_2 quasiconformally to these triangles. Here the triangle $R_{-m,m-1}$ is mapped onto the half-disk $\Omega_2 := \{z \in \mathbb{C} : \text{Im } z > 0, |z| < 1\}$ while $R_{-m,-m}$ is mapped onto $\overline{\Omega_2}$.

Thus for $j \in \{1, 2\}$ we have defined a quasiregular mapping $f_j : S_j \rightarrow \overline{\mathbb{C}}$ such that (3.20) holds. This implies that the mapping f_0 defined by (3.21) is quasiregular.

By construction, all (-1) -points and all poles of f_0 are multiple, and except for the origin all 1-points of f_0 are also multiple. Moreover, all a -points in the right half-plane are multiple and all $(-a)$ -points in the left half-plane are multiple. It follows from (3.20) and (3.22) that f_0 maps the imaginary axis to $\partial\mathbb{D}$.

Let D be a Jordan domain which contains a and $-a$ and whose closure is contained in \mathbb{D} . Then an island over D cannot intersect the imaginary axis. Thus all islands over D are contained in the right or left half-plane. Those contained in the right half-plane contain a multiple a -point while those contained in the left half-plane contain a multiple $(-a)$ -point. We deduce that there are no simple islands over D .

As mentioned above, Lemma 2.5 yields that there exists a quasiconformal mapping $\phi : \mathbb{C} \rightarrow \mathbb{C}$ such that $F := f_0 \circ \phi$ is meromorphic. It is easy to see that the set A of all z satisfying $|z| > 1$ where f_1 and f_2 are not meromorphic satisfies (2.3). Lemma 2.6 now yields that ϕ satisfies (2.2). Since an elliptic function has order 2 this implies that $\rho(F) = 4\alpha/\pi$.

As it is the case for f_0 , the function F has no simple island over D and all poles and all (± 1) -points of F are multiple, except for the simple 1-point at the origin. We finally put $f(z) = F(z^p)$ for some $p \in \mathbb{N}$ with $p \geq 2$. Then the origin is a multiple

1-point of f and we conclude that f has no simple island over D and that all poles and all (± 1) -points of f are multiple. Moreover, $\rho(f) = 4\rho\alpha/\pi$.

Since $\alpha = \arctan \tau$ we can achieve $\rho(f) = \rho$ for any given $\rho \in (0, \infty)$ by a suitable choice of τ and p . \square

Acknowledgements We thank Jim Langley and the referee for valuable comments and suggestions.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Ahlfors, L.V.: Sur une généralisation du théorème de Picard. C. R. Acad. Sci. Paris **194**, 245–247 (1932); Collected Papers, Birkhäuser, Boston, Basel, Stuttgart, 1982, Vol. I, pp. 145–147
2. Ahlfors, L.V.: Sur les fonctions inverses des fonctions méromorphes. C. R. Acad. Sci. Paris **194**, 1145–1147 (1932); Collected Papers, Vol. I, pp. 149–151
3. Ahlfors, L.V.: Quelques propriétés des surfaces de Riemann correspondant aux fonctions méromorphes. Bull. Soc. Math. Fr. **60**, 197–207 (1932); Collected Papers, Vol. I, pp. 152–162
4. Ahlfors, L.V.: Über die Kreise die von einer Riemannschen Fläche schlicht überdeckt werden. Comment. Math. Helv. **5**, 28–38 (1933); Collected Papers, Vol. I, pp. 163–173
5. Ahlfors, L.V.: Zur Theorie der Überlagerungsflächen. Acta Math. **65**, 157–194 (1935); Collected Papers, Vol. I, pp. 214–251
6. Axler, S., Bourdon, P., Ramey, W.: Harmonic function theory, 2nd edn. Graduate Texts in Mathematics, vol. 137. Springer, New York (2001)
7. Bergweiler, W.: On the growth rate of composite meromorphic functions. Complex Var. Theory Appl. **14**(1–4), 187–196 (1990)
8. Bergweiler, W.: A new proof of the Ahlfors five islands theorem. J. Anal. Math. **76**, 337–347 (1998)
9. Bergweiler, W.: Bloch's principle. Comput. Methods Funct. Theory **6**, 77–108 (2006)
10. Bergweiler, W., Jank, G., Volkmann, L.: Wachstumsverhalten zusammengesetzter Funktionen. Results Math. **7**(1), 35–53 (1984)
11. Beurling, A., Ahlfors, L.: The boundary correspondence under quasiconformal mappings. Acta Math. **96**, 125–142 (1956); The collected works of Arne Beurling, Birkhäuser Boston, Inc., Boston, MA, 1989, Vol. I, pp. 231–248; Collected Papers (of Ahlfors), Vol. II, pp. 104–121
12. Bloch, A.: Quelques théorèmes sur les fonctions entières et méromorphes d'une variable. C. R. Acad. Sci. Paris **181**, 1123–1125 (1925)
13. Bloch, A.: Quelques théorèmes sur les fonctions entières et méromorphes d'une variable. C. R. Acad. Sci. Paris **182**, 367–369 (1926)
14. Bloch, A.: La conception actuelle de la théorie des fonctions entières et méromorphes. Enseign. Math. **25**, 83–103 (1926)
15. Boas, R.P., Jr.: Entire functions. Academic Press, Inc., New York (1954)
16. Branner, B., Fagella, N.: Quasiconformal surgery in holomorphic dynamics. Cambridge Studies in Advanced Mathematics, vol. 141. Cambridge University Press, Cambridge (2014)
17. de Thélin, H.: Une démonstration du théorème de recouvrement de surfaces d'Ahlfors. Enseign. Math. (2) **51**(3–4), 203–209 (2005)
18. Duval, J.: Sur la théorie d'Ahlfors des surfaces. Enseign. Math. (2) **60**(3–4), 417–420 (2014)

19. Goldberg, A.A., Tairova, V.G.: On entire functions with two finite completely ramified values (in Russian). *Zapiski Meh-mat. fakulteta Kharkovskogo gos. universiteta i Harkovskogo mat. obshchestva*, 29(Ser. 4), 67–78 (1963)
20. Goldberg, A.A., Ostrovskii, I.V.: Value distribution of meromorphic functions. *Translations of Mathematical Monographs*, vol. 236. American Mathematical Society, Providence (2008)
21. Kelingos, J.A.: Boundary correspondence under quasiconformal mappings. *Mich. Math. J.* **13**, 235–249 (1966)
22. Künzi, H.: Zur Theorie der Viertelsenden Riemannscher Flächen. *Comment. Math. Helv.* **30**, 107–115 (1956)
23. Langley, J.K.: On differential polynomials, fixpoints and critical values of meromorphic functions. *Results Math.* **35**(3–4), 284–309 (1999)
24. Lehto, O., Virtanen, K.I.: Quasiconformal mappings in the plane, 2nd edn. *Die Grundlehren der mathematischen Wissenschaften*, vol. 126. Springer, New York (1973)
25. Nevanlinna, R.: *Le théorème de Picard-Borel et la théorie des fonctions méromorphes*. Gauthiers-Villars, Paris (1929)
26. Pommerenke, Ch.: Boundary behaviour of conformal maps. *Die Grundlehren der Mathematischen Wissenschaften*, vol. 299. Springer, Berlin (1992)
27. Selberg, H.L.: Über einige Eigenschaften bei der Werteverteilung der meromorphen Funktionen endlicher Ordnung. *Avh. Det Norske Videnskaps-Akademi i Oslo I. Matem.-Naturvid. Kl.* **7**, 17pp (1928)
28. Stoilow, S. (in collaboration with C. A. Cazacu): *Teoria funcțiilor de o variabilă complexă. Vol. 2. Funcții armonice. Suprafețe Riemanniene*. Editura Academiei Republicii Populare Romîne, Bucharest (1958)
29. Teichmüller, O.: Einfache Beispiele zur Wertverteilungslehre. *Deutsche Math.* **7**, 360–368 (1944); *Gesammelte Abhandlungen—Collected Papers*. Springer, Berlin, 1982, pp. 728–736
30. Tôki, Y.: Proof of Ahlfors principal covering theorem. *Rev. Math. Pures Appl.* **2**, 277–280 (1957)
31. Tsuji, M.: *Potential theory in modern function theory*. Maruzen Co., Ltd., Tokyo (1959)
32. Valiron, G.: *Lectures on the general theory of integral functions*. Édouard Privat, Toulouse 1923; reprint: Chelsea, New York (1949)
33. Walsh, J.L.: Note on the location of zeros of the derivative of a rational function whose zeros and poles are symmetric in a circle. *Bull. Am. Math. Soc.* **45**, 462–470 (1939)
34. Zalcman, L.: Analytic capacity and rational approximation. *Lecture Notes in Mathematics*, vol. 50. Springer, Berlin (1968)
35. Zalcman, L.: A heuristic principle in complex function theory. *Am. Math. Mon.* **82**, 813–817 (1975)
36. Zalcman, L.: Normal families: new perspectives. *Bull. Am. Math. Soc. (N. S.)* **35**, 215–230 (1998)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.