Physical activity, Sedentary Behaviour and Their Correlates in Adults with Autism Spectrum Disorder: a Systematic Review

C. Thompson^{1,2} · M. Brook¹ · S. Hick¹ · C. Miotti¹ · R. Toong¹ · J.A. McVeigh^{1,2,3}

Received: 27 May 2021 / Accepted: 5 February 2022
© The Author(s) 2022

Abstract

The aim of this systematic review was to investigate what is currently known about autistic adults' participation in physical activity and sedentary behaviour and explore any associated correlates of these behaviours. The databases EMBASE, MEDLINE, PsycINFO, PubMed, and SPORTDiscus were searched for relevant literature. The review was registered with PROSPERO (No. 189187). The search identified 2834 papers, with 12 meeting the criteria for inclusion. Correlates of physical activity and/or sedentary behaviour were reported in eight included studies. The limited evidence identified suggest that autistic adults are unlikely to meet Australia's physical activity guidelines. Future research should use device-derived measurement to more accurately quantify activity levels within the autistic adult population.

Keywords Autism spectrum disorder · Physical activity · Sedentary behaviour · Adults

Introduction

Physical activity is defined as bodily movements produced by skeletal muscles requiring energy expenditure (World Health Organization, 2017). An individual's daily physical activity pattern consists of various activities along a continuum of intensities, from sedentary behaviour (e.g. sitting, lying) to moderate and vigorous physical activity (MVPA) (e.g. running, swimming) (Department of Health, 2019). Australia's physical activity and sedentary behaviour guidelines recommend that adults perform sufficient physical activity (i.e. participate in > 150 min MVPA/week) and minimise and break up their sedentary behaviour (Department of Health, 2019). Health outcomes associated with sufficient physical activity participation include decreased mortality, reduced risk of developing chronic conditions (e.g. cardiovascular disease) (Lear et al., 2017), maintaining healthy blood pressure and cholesterol levels, preventing

obesity and promoting physical and mental health (Kokinos, 2012; World Health Organization, 2017). Despite the benefits of sufficient participation in physical activity, approximately half of Australian adults were considered insufficiently active based on self-reports in 2017–2018 (Australian Bureau of Statistics, 2019). The patterns and volumes of physical activity and sedentary behaviour are influenced by a wide variety of factors or correlates, including a diagnosis of autism spectrum disorder (ASD).

Autism spectrum disorder is a lifelong, neurodevelopmental condition characterised by deficits in social communication and restricted, repetitive behaviours and interests (American Psychiatric Association, 2013). It is essential to continue to develop knowledge of ASD across the lifespan due to the condition's lifelong nature, and the unique challenges adulthood presents for autistic individuals (Howlin et al., 2015). Autistic adults are reported to have lower participation in domains of life, including education, employment (Knüppel et al., 2019), social relationships (Tobin et al., 2014), leisure and physical activity compared to neurotypical peers (García-Villamisar & Dattilo, 2010; Hillier et al., 2020). Autistic adults may also experience motor and social skill deficits, difficulties with sensory processing, poor sleep and restrictive behaviour patterns (Nichols et al., 2018). These characteristics may act as unique correlates limiting autistic adults' involvement in a range of occupational roles and activities,

✉ C. Thompson
craig.thompson@curtin.edu.au

¹ Curtin School of Allied Health, Curtin University, Perth, Australia

² enAble Institute, Curtin University, Perth, Australia

³ Movement Physiology Laboratory, School of Physiology, University of Witwatersrand, Johannesburg, South Africa

including participating in recommended physical activity levels (Menear & Neumeier, 2015).

The research examining correlates of time spent in physical activity and sedentary behaviour among autistic adults is scant. However, several studies have reported on time spent in physical activity and sedentary behaviour (and the correlates thereof) in autistic children. Autistic children spend more time in sedentary behaviour than neurotypical children (Jones et al., 2017; Must et al., 2014) and have lower levels of physical activity (Jones et al., 2017). Access to an outdoor environment, social interactions and being of male gender demonstrated to positively influence physical activity participation in autistic children (Healy & Garcia, 2019; Pan et al., 2011). A larger number of close friends and lower parental rating of prosocial behaviour negatively impact sedentary behaviour participation in this population (Healy & Garcia, 2019). Moreover, there are consistent findings that suggest a decline in physical activity participation and lesser adherence to Australia's physical activity and sedentary behaviour guidelines as autistic children age (Healy et al., 2019; Jones et al., 2017; MacDonald et al., 2011; Simpson et al., 2019). It is important to understand the factors impacting upon and health implications of physical activity and sedentary behaviour participation in order to support autistic adults to achieve positive health outcomes (Simpson et al., 2019; Tanguay & Lohr, 2015).

The usefulness of the evidence for physical activity levels is influenced by the measurement method. Self-report and device-derived measurement are two common methods used to measure physical activity and sedentary behaviour (Yano et al., 2019). Self-report measures of physical activity and sedentary behaviour vary according to a range of factors, including the variables being quantified (e.g. frequency and/or duration of participation), unit of measurement (e.g. length of time, categorical ranking) and method of application (e.g. self-completed questionnaire, interview). The use of measurement devices (i.e. fitness trackers and accelerometers) to quantify physical activity and sedentary behaviour is growing in popularity (Yano et al., 2019). Device-derived measurement of physical activity and sedentary behaviour enables the identification of, and ability to discriminate between characteristics of activity patterns including frequency, duration, type and intensity (Strath et al., 2012).

The current evidence base in autistic children has increasingly used device derived measures of physical activity, as demonstrated by a recent systematic review where 31 of the 35 studies used device derived methods. In the autistic adult population, most studies have used questionnaires, such as the paper by Engelhardt et al. (2017). The limited use of device derived measurement has also meant that studies in the autistic adult population have mainly focused on either physical activity, such as Hamm and Yun (2019), or

sedentary behaviours, such as Finke et al. (2018), resulting in a limited understanding of the continuum of activity.

Currently, there is limited evidence examining time spent in physical activity and sedentary behaviour among the autistic adult population, and the evidence regarding correlates of these behaviours is scarce. Therefore, the aim of this review was to systematically investigate what is known about the time that autistic adults spend in physical activity and sedentary behaviour and to explore any associated correlates of these behaviours.

Methods

Design

A systematic review of the literature regarding the physical activity and sedentary behaviour prevalence of autistic adults and associated correlates was conducted. The guidelines outlined in the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statement were followed (Moher et al., 2009). Prior to commencing this systematic review, the proposed methodology was registered with PROSPERO (No. 189187) (National Institute for Health Research, 2020).

Search Strategy

A literature search was conducted using five selected online databases: EMBASE, MEDLINE, PsycINFO, PubMed, and SPORTDiscus. The search strategy included keywords within four categories: (1) population, (2) physical activity, (3) sedentary behaviour and (4) Autism spectrum disorder. Table 1 (insert Table 1 after this paragraph) provides the search terms used for all databases. In order to broaden the scope of the review and ensure all papers with potential for inclusion were captured, no limitations (e.g. year of publication) were applied to initial searches of each respective database. Searches were designed to capture papers published from the establishment of the databases to April 2020.

Papers were included if they were (1) peer-reviewed, (2) written in English, (3) accessible by researchers in full-text and (4) reported a measure of individuals' physical activity and/or sedentary behaviour. Academic theses, commentaries, secondary reviews and papers omitting baseline or habitual physical activity and/or sedentary behaviour data were excluded. In order to further broaden the scope for the review, reference lists of short-listed papers were screened manually to identify any other potentially relevant papers. Following initial database searches and manual review of references, the title and reference of short-listed papers were entered into a shared Microsoft ExcelTM, 2020 spreadsheet for review by all researchers.

Table 1 Search terms used in all databases

Population		Outcome	
Age	Diagnosis	Physical activity	Sedentary behaviour
Adult*	Autis*	"Physical* activ*"	"Sedentary behavio*"
Old* adult*	"Autism spectrum disorder"	Exercis*	Sedentar*
Elder*	ASD	"Motor activit*"	"Physical inactiv*"
	Asperger*	"Locomotor activit*"	Inactiv*
	"Autistic disorder"	Sport*	Gaming
	"Pervasive developmental disorder"	Acceleromet*	"Television view*"
	PDDNOS	Recreation*	"TV"
	PDD		Television
	"Pervasive developmental disorder not otherwise specified"		"Video game*"
			"Screen time"
			"Electronic game*"
			Computer*
			E-game*
			Video*
			"Screen based media"
			"Electronic media"

A systematic screening process (Figure 1) was conducted to ensure selective inclusion of relevant papers: (1) All four authors screened paper titles and discussed their potential suitability for inclusion; (2) two authors (MB, RT) screened remaining papers' abstracts, with third author (CM) being consulted if a unanimous decision regarding inclusion was not achieved; (3) two authors (MB, RT) screened remaining papers' full text, with a third author (CM) being consulted if a unanimous decision regarding inclusion was not achieved; and (4) all four authors reviewed and agreed upon final papers deemed to have met the inclusion criteria. Authors' individual decisions for each phase of data screening were recorded on a shared Microsoft ExcelTM, 2020 spreadsheet, with individual columns to input 'N' to represent 'No', 'Y' to represent 'Yes' and '?' to represent 'Maybe' next to each respective study.

Quality Assessment

The Standard Quality Assessment Criteria was used to critically appraise each of the final 12 studies to be included in the review to determine the quality of the studies (Kmet et al., 2004). Four authors completed the quantitative or qualitative checklists for each of the 12 included studies dependent on respective study design (Kmet et al., 2004). To determine the individual study quality, criteria assessed included, but was not exclusive to: study design (e.g. appropriateness), sample selection, size and description (e.g. adequate and representative), random allocation and blinding (e.g. appropriately used), outcome measures, analysis methods (e.g. described and justified), confounding variables (e.g. if they were controlled for) and results (e.g. supporting hypothesis). Each study was awarded a score of '2' for 'Yes', '1' for 'Partial' or '0' for 'No' across each criterion

on the checklist according to the assessment guidelines (Kmet et al., 2004). For a summary of categorical ratings for each included study, see Appendix A. From the summed categorical ratings, percentage scores were calculated for the quality of each study. All four authors discussed individual scoring and negotiated to full agreement. Studies were then given a categorical rating of quality according to the following protocol: > 80% = strong, 70–80% = good, 50–70% = adequate, < 50% = limited (Lee et al., 2008). Study quality across the 12 studies was categorically ranked 'good' overall, with an average score of 74% in accordance with the Standard Quality Assessment Criteria (Table 2) (insert Table 2 at the beginning of page 8) (Kmet et al., 2004).

Data Extraction

Data regarding participant sample, study design, ASD diagnosis, outcome of interest, instrument employed to collect data, results and correlates were extracted from each of the 12 included studies. The studies were screened by three authors (CM, MB and RT) to complete data extraction independently. To increase accuracy, each study was examined by two of the three screening authors. All data were manually extracted and entered into a shared Microsoft ExcelTM, 2020 spreadsheet. Following initial data extraction, information recorded for all studies was manually reviewed and re-assessed by a fourth author (SH). Following the final review, all authors negotiated the 5% discrepancy to reach full agreement.

Data Standardisation

In the 12 included studies, physical activity and sedentary behaviour data were reported on in various formats

Fig. 1 PRISMA flow diagram of search results

dependent on the study design and outcomes of interest. Prevalence of physical activity was reported as MVPA minutes per day, number of steps per day, MET minutes per

week and Godin-Shephard Leisure-Time Physical Activity Questionnaire (GSLTPAQ) scores or self-reported activity level (e.g. number of participations per week). For sedentary

Table 2 Characteristics of included studies ($N = 12$)

No	Study	Sample	Study design	Diagnosis verification	Target outcome (PA/SB)	Measurement tool(s)	Measure(s)	Results	Correlates	Quality assessment score
1	Benson et al	$N = 32$ (23 M; 71.9%), aged 18–35 yrs ASD: $n = 15$ (11 M; 73.3%), mean age 22.8 ± 4.5 yrs, presence of co-occurring ID not reported NT: $n = 17$ (12 M; 70.6%), mean age 23.7 ± 4.3 yrs	Cross-sectional	Psychiatric assessment based on the ADOS ($n = 13$) Current ASD diagnoses by medical doctor obtained from parents ($n = 2$)	PA	GSLTPAQ Motionlogger™ actigraph (trix-ial piezoelectric accelerometer): non-dominant wrist 24 h/day for 14 consecutive days; cut off point of $> 22,000$ cpm for MVPA (specific PA threshold has not been validated for the Motionlogger™ actigraph)	GSLTPAQ score MVPA	Self-reported: ASD MVPA: 28 min/day NT MVPA: 40 min/day Objective: ASD MVPA: 16 min/day NT MVPA: 37 min/day	Sleep, day of the week (weekday vs weekend)	Good 17/22 (77%)
2	Bishop-Fitzpatrick et al. (2017)	$N = 60$ (46 M; 76.7%), aged 24–55 yrs ASD: $N = 60$ (46 M; 76.7%), mean age 31.48 ± 6.75 yrs, co-occurring ID $n = 16$ (26.7%). Sample also included mothers of participants No control group	Correlation	Administration of the ADI-R	PA and SB	Separate in-person interviews with ASD adults and their mothers	Self-reported participation in recreational activities (participations/week; engagements/year)	Self-Reported: MVPA: 15 min/day SB: 72 min/day Mother-reported: MVPA: 8 min/day SB: 37 min/day	Perceived stress	Strong 17/20 (85%)

Table 2 (continued)

No	Study	Sample	Study design	Diagnosis verification	Target outcome (PA/SB)	Measurement tool(s)	Measure(s)	Results	Correlates	Quality assessment score
3	Engelhardt, Mazurek, & Hilgard (2017)	N = 119 (103 M; 87%, mean age 20.48 yrs ASD: <i>n</i> = 59 (51 M; 86%), mean age 20.42 yrs, IQ scores $\geq$ to 85. ASD sample recruited from participants concurrently engaged in a larger primary study NT: <i>n</i> = 60 (52 M; 87%), mean age 20.54 yrs	Cross-sectional	ADOS and/or ADI-R	SB	Self-report questionnaire	Time spent playing video games (hours/day)	ASD SB: 168 min/day NT SB: 63 min/day	Day of the week, pathological game use symptoms	Strong 16/20 (80%)
4	Finke, Hickerson, & Kremkow (2018)	N = 10 (9 M; 90%), aged 18–24 yrs ASD: <i>N</i> = 10 (9 M; 90%); mean age 21 yrs, presence of co-occurring ID not reported No control group; purposive sampling method used	Case-series	No verification of ASD diagnoses completed; participants self-reported ASD diagnosis, health professional who made the initial diagnosis and the age of diagnosis	SB	Semi-structured interview	Time spent playing video games (hours/week; days/week)	SB: 266 min/day	Gender	Strong 16/20 (80%)

Table 2 (continued)

No	Study	Sample	Study design	Diagnosis verification	Target outcome (PA/SB)	Measurement tool(s)	Measure(s)	Results	Correlates	Quality assessment score
5	Garcia-Pastor, Salinero, & Ruiz-Vicente (2019)	N=78 (56 M; 71.8%), aged 7–48yrs ASD Adults (> 19 yrs); n=34 (23 M; 67.6%), mean age 31.09 ± 6.85 yrs, presence of co-occurring ID not reported No NT control group	Cross-sectional	ADOS-2 and/or ADI-R	PA and SB	GTIM accelerometer (Actigraph): ≥ 10 h/day for a minimum of 4 consecutive days including 1 weekend day; counts recorded over periods of 60 s; applied cut-off points for adults MVPA obtained from GTIM accelerometer calibration studies Count ranges for activity intensities in adults: - 0–99: sedentary - 100–1951: light - 1952–5724: moderate - 5725–9498: vigorous - 9599 and above: very vigorous Periods of 20 min or greater with intensity counts as 0 were considered non-wearing time and excluded from analysis	Sedentary time, MVPA, steps	ASD adults MVPA: 40 min/day SB: 1100 min/day	Age, day of the week (weekend), vs obesity	Strong 18/22 (82%)

Table 2 (continued)

No	Study	Sample	Study design	Diagnosis verification	Target outcome (PA/SB)	Measurement tool(s)	Measure(s)	Results	Correlates	Quality assessment score
6	Hamm and Yun (2019)	N=320 (138 M; 43.1%, aged 18–35 yrs ASD: $n = 143$; mean age 25.39 ± 4.5 yrs, presence of co-occurring ID not reported NT: $n = 177$; mean age 24.22 ± 5.1 yrs)	Cross-sectional	No verification of ASD diagnoses completed; participants self-reported ASD diagnosis	PA	GSLTPAQ	GSLTPAQ score	ASD MVPA: 39 min/day NT MVPA: 55 min/day	Health-related quality of life	Strong 16/20 (80%)
7	Jozkowski and Cermak (2020)	N=36 (19 M; 52.8%), aged 18–25 yrs ASD: $n = 18$ (16 M; 88.9%), mean age 21.35 yrs, presence of co-occurring ID not reported NT: $n = 18$ (3 M; 16.7%), mean age 19.25 yrs Convenience sampling method used resulted in non-homogenous sample groups	Pre-post	Researcher review of documentation including ICD-9 diagnostic code, or written diagnosis of 'autism', 'autistic disorder', or 'ASD' on a physician signed form or verified individualised education plan	PA and SB	Demographic and Physical Activity Questionnaire	Self-described activity level (frequency/week), time spent playing videogames (hours/day)	ASD MVPA: 18 min/day SB: 150 min/day NT MVPA: 33 min/day SB: 45 min/day	Fitness level, social interaction	Good 19/24 (79%)

Table 2 (continued)

No	Study	Sample	Study design	Diagnosis verification	Target outcome (PA/SB)	Measurement tool(s)	Measure(s)	Results	Correlates	Quality assessment score
8	LaLonde, MacNeill, Wolfe, Eversole, Ragozy, & Poling (2014)	N=5 (4 M; 80%), aged 21–26yrs ASD: N=5 (4 M; 80%), presence of co-occurring ID not reported No NT control group	Pre-post	No verification of ASD diagnoses completed; participants were sampled from an ASD-specific classroom at a post-secondary educational facility that provides transitional services to young adults with disabilities	PA	Fitbit Zip™ device (pedometer that uses a 3-axis accelerometer), 6 h/day, number of days of wear varied depending on participants school attendance (range: 25–42 days)	Steps/day	ASD MVPA: 73 min/day	Nil reported	Adequate 12/20 (60%)
9	Li, Curiel, Ragozy, & Poling (2019)	N=4 (2 M; 50%), aged 21–25 yrs ASD: n=2 (1 M; 50%), mean age 23 yrs, presence of co-occurring ID not reported No NT control group	Pre-post	No verification of ASD diagnoses completed; participants were sampled from an ASD-specific classroom at a post-secondary educational facility that provides transitional services to young adults with disabilities	PA	Fitbit Zip™ device (pedometer that uses a 3-axis accelerometer), 6 h/day, number of days of wear varied depending on participants school attendance (range: 32–64 days)	Steps/day	ASD MVPA: 69 min/day	Nil reported	Adequate 12/20 (60%)
10	Nastasi, Shepard, & Raiff (2020)	N=4 (2 M; 50%), aged 23–55 yrs ASD: n=1 (0 M; 0%), mean age 23 yrs, presence of co-occurring ID not reported No NT control group	Pre-post	No verification of ASD diagnoses completed; diagnoses for the participants were provided by clinicians not involved in this research	PA	Fitbit Flex™ (pedometer that uses a 3-axis accelerometer), wrist-worn 24 h/day (excluding charging approximately 30 min every 2–3 days)	Steps/day	ASD MVPA: 61 min/day	Nil reported	Adequate 10/20 (50%)

Table 2 (continued)

No	Study	Sample	Study design	Diagnosis verification	Target outcome (PA/SB)	Measurement tool(s)	Measure(s)	Results	Correlates	Quality assessment score
11	Takeshi and Chisato (2014)	N = 26 (26 M; 100%), aged 18–20 ASD: n = 7 (7 M; 100%), mean age 18.4 ± 2.1 yrs, presence of co-occurring ID not reported. Sample recruited from ASD-specific sporting organisations NT: n = 9 (9 M; 100%), mean age 18.9 ± 0.4 yrs. More active NT group used for comparison	Correlation	No verification of ASD diagnoses completed; reported diagnosis by physicians (n = 5), no formal diagnosis of ASD (n = 1), unable to confirm formal ASD diagnosis (n = 1)	PA	IPAQ	MET min/week	ASD MVPA: 95 min/day NT MVPA: 95 min/day	Arterial stiffness	Good 17/22 (77%)
12	Stacey et al. (2019)	N = 249 (81 M; 32.5%), aged 25–85 yrs ASD: n = 145 (58 M; 40%), presence of co-occurring ID not reported NT: n = 104 (23 M; 22.1%)	Cross-sectional	No verification of ASD diagnoses completed; all autistic participants reported having received a formal autism diagnosis from a health professional as per the DSM-5 or an equivalent diagnosis from the DSM-III or DSM-IV	PA and SB	Self-report questionnaire	Frequency of participation in leisure activities (participations/year)	ASD MVPA: 13 min/day SB: 155 min/day NT MVPA: 14 min/day SB: 147 min/day	Nil reported	Strong 16/20 (80%)

ADI-R Autism Diagnostic Interview-Revised; *ADOS* Autism Diagnostic Observation Schedule; *ASD* Autism Spectrum Disorder; *DSM* Diagnostic and Statistical Manual of Mental Disorders; *GSL/TPAQ* Godin-Shepherd Leisure-Time Physical Activity Questionnaire; *ICD-9* International Classification of Diseases (Ninth Revision); *ID* Intellectual disability; *IPAQ* International Physical Activity Questionnaire; *MVPA* moderate-vigorous physical activity; *NT* neurotypical; *PA* physical activity; *SB* sedentary behaviour; *Yrs* years

behaviour, the prevalence was reported as sedentary time (e.g. minutes/day) or time spent playing video games (e.g. hours/day, hours per week, days/week).

Prior to data synthesis and comparison, estimates of time spent in MVPA and sedentary behaviour were standardised to minutes per day to produce a comparable data set. The included studies were divided among two authors (SH, RT) to complete standardisation calculations. The data standardisation process required some assumptions be made in order to convert the available reported data into the form of MVPA and sedentary behaviour minutes per day. For a full explanation of the standardisation process and any assumptions made for each data format, see Appendix B. All calculations were initially completed manually on paper with the assistance of scientific calculators. All completed calculations were transcribed onto an electronic document by one author (SH) and cross-checked by two authors (SH, RT). Where discrepancies occurred, resolution was achieved through discussion. For full details of all completed calculations for each included study, see Appendix C. Final results of MVPA and sedentary behaviour minutes per day were rounded to the nearest whole number as per standard mathematical protocol.

Results

Study Characteristics

A total of 12 studies met the inclusion criteria for this review with six studies reporting physical activity (Benson et al., 2019; Hamm & Yun, 2019; LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020; Takeshi & Chisato, 2014), two studies reporting sedentary behaviour (Engelhardt et al., 2017; Finke et al., 2018) and four studies including data on both physical activity and sedentary behaviour (Bishop-Fitzpatrick et al., 2017; Garcia-Pastor et al., 2019; Jozkowski & Cermak, 2020; Stacey et al., 2019) (Figure 1). Table 2 provides a summary of the characteristics of the included studies.

The average age of participants in the included studies was approximately 27 years (*range* = 18–85 years). Of the 12 studies, one study used a sample comprising solely of males (Takeshi & Chisato, 2014), and another included one single female participant (Finke et al., 2018). All remaining studies gathered data from a sample with a higher proportion of male compared to female participants (Garcia-Pastor et al., 2019; LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020; Stacey et al., 2019). Within the included studies, six made use of a neurotypical control group for comparative purposes (Benson et al., 2019; Engelhardt et al., 2017; Hamm & Yun, 2019; Jozkowski & Cermak, 2019; Otsuki & Ohashi, 2014; Stacey et al.,

2019). A range of measurement tools were used across the 12 studies, with seven studies using self-report measures (Benson et al., 2019; Bishop-Fitzpatrick et al., 2017; Engelhardt et al., 2017; Garcia-Pastor et al., 2019; Hamm & Yun, 2019; Jozkowski & Cermak, 2020; LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020; Stacey et al., 2019), four studies using a measurement device (Nastasi et al., 2020; Pinder-Amaker, 2014) and one study using both self-report and measurement device (Benson et al., 2019). All studies included in Table 2 describe their research objective(s) in sufficient detail and are published in peer-reviewed journals. Sample sizes ranged from 4 to 320 participants, with one study obtaining data from one autistic adult participant (Nastasi et al., 2020). Of the 12 studies, four collected data from a sample of 10 participants or less (Finke et al., 2018; LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020). Several sampling methods were used across the studies, with one recruiting autistic participants exclusively from ASD-specific sporting organisations (Takeshi & Chisato, 2014).

Autistic Adults' Physical Activity

Self-Report Findings

Of the 12 studies, six reported data of time spent in physical activity obtained from a form of self-report measure (*mean* = 35 MVPA min/day, *range* = 13–95) (Benson et al., 2019; Bishop-Fitzpatrick et al., 2017; Hamm & Yun, 2019; Jozkowski & Cermak, 2020; Stacey et al., 2019; Takeshi & Chisato, 2014). Estimates of autistic adults' physical activity levels from questionnaires commonly used to measure physical activity in the general population (e.g. International Physical Activity Questionnaire, GSLTPAQ) were higher (*mean* = 54 MVPA min/day, *range* = 28–95) than estimates from questionnaires tailored to meet specific study objectives and for use with autistic adults (*mean* = 16 MVPA min/day, *range* = 13–18) (Jozkowski & Cermak, 2020; Stacey et al., 2019). Estimates obtained from interviews were more similar to the tailored questionnaire estimates (15 MVPA min/day) (Bishop-Fitzpatrick et al., 2017). In one study using both self- and proxy-report, the mother-reported value of daily physical activity was lower than that reported by the autistic adult participants themselves (mother-report *mean* = 8 MVPA min/day vs self-report *mean* = 15 MVPA min/day) (Bishop-Fitzpatrick et al., 2017). Overall, within the six studies measuring physical activity participation using self-report, three of the autistic adult samples would be deemed insufficiently active (i.e. not participate in > 150 MVPA min/week) (Bishop-Fitzpatrick et al., 2017; Jozkowski & Cermak, 2020; Stacey et al., 2019).

Objective Findings

Within the included studies, five reported estimates of time spent in PA from measurement devices (*mean* = 52 MVPA min/day, *range* = 16–73) (Benson et al., 2019; Garcia-Pastor et al., 2019; LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020). The two categories of these measurement devices used within these five studies were wearable activity trackers (i.e. Fitbit, Garmin) and research-grade accelerometers (i.e. Motionlogger and GT1M ActiGraph).

Within this review, three studies used wearable activity trackers to obtain data on autistic adults' PA (*mean* = 68 MVPA min/day, *range* = 61–73) (LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020). All three studies using wearable activity trackers for physical activity measurement scored less than 60% when assessed with the Standard Quality Assessment Criteria and hence were classified to be of 'adequate' quality (Kmet et al., 2004; LaLonde et al., 2014; Lee et al., 2008; Li et al., 2019; Nastasi et al., 2020). All the participants in these three studies would be considered to participate in sufficient physical activity (i.e. participate in > 150 MVPA min/week).

Of the 12 studies, two used accelerometers to estimate autistic adults' time spent in physical activity (*mean* = 28 MVPA min/day, *range* = 16–40) (Benson et al., 2019; Garcia-Pastor et al., 2019). These studies scored 77 (Benson et al., 2019) and 82% (Garcia-Pastor et al., 2019) when assessed with the Standard Quality Assessment Criteria and hence were classified to be of 'good' and 'strong' quality, respectively (Kmet et al., 2004; Lee et al., 2008). Estimates of physical activity obtained from accelerometers were lower (*mean* = 28 MVPA min/day, *range* = 16–40) (Benson et al., 2019; Garcia-Pastor et al., 2019) compared to physical activity estimates from wearable activity trackers (*mean* = 68 MVPA min/day, *range* = 61–73) (LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020).

Autistic Adults' Sedentary Behaviour

Self-Report Findings

Within the included studies, five reported estimates of time spent in sedentary behaviour from a form of self-report measure (*mean* = 162 sedentary behaviour min/day, *range* = 72–266) (Bishop-Fitzpatrick et al., 2017; Engelhardt et al., 2017; Finke et al., 2018; Jozkowski & Cermak, 2020; Stacey et al., 2019). Estimates of sedentary behaviour levels gathered from questionnaires were lower (*mean* = 158 sedentary behaviour min/day, *range* = 150–168) (Engelhardt et al., 2017; Jozkowski & Cermak, 2020; Stacey et al., 2019) than estimates from in-person interviews (*mean* = 169 sedentary behaviour min/day, *range* = 72–266) (Bishop-Fitzpatrick et al., 2017; Finke et al., 2018). In one study,

mother-reported SB estimates were lower than that obtained from interviews with the autistic adult (mother-report *mean* = 37 sedentary behaviour min/day vs self-report *mean* = 72 sedentary behaviour min/day) (Bishop-Fitzpatrick et al., 2017).

Objective Findings

Out of the 12 studies, one reported device-derived estimates of autistic adults' time spent in sedentary behaviour obtained using the GITM accelerometer (1100 sedentary behaviour min/day) (Garcia-Pastor et al., 2019).

Studies Including Neurotypical Adults

A total of six studies included a neurotypical control group alongside autistic adult samples for comparison (Benson et al., 2019; Engelhardt et al., 2017; Hamm & Yun, 2019; Jozkowski & Cermak, 2020; Stacey et al., 2019; Takeshi & Chisato, 2014). Another two studies included participants without ASD. However, due to the diagnosis of other neurological conditions (e.g. cognitive impairments, mental health conditions), they did not fit the criteria for being neurotypical participants (Li et al., 2019; Nastasi et al., 2020). All six studies, including neurotypical adult participants, relied on self-report for measurement of physical activity and/or sedentary behaviour participation (Benson et al., 2019; Engelhardt et al., 2017; Hamm & Yun, 2019; Jozkowski & Cermak, 2020; Stacey et al., 2019; Takeshi & Chisato, 2014). In addition to a self-report questionnaire, one study also obtained estimates regarding physical activity participation of autistic and neurotypical adults via accelerometry (Benson et al., 2019).

Physical Activity Findings

Of the six studies with autistic and neurotypical participants, five reported estimates of physical activity participation (ASD *mean* = 35 MVPA min/day, *range* = 13–95; NT *mean* = 46 MVPA min/day, *range* = 14–95) (Benson et al., 2019; Hamm & Yun, 2019; Jozkowski & Cermak, 2020; Stacey et al., 2019; Takeshi & Chisato, 2014). Of these studies, autistic adults' level of physical activity participation was lower than the neurotypical adult sample in four of the studies (Benson et al., 2019; Hamm & Yun, 2019; Jozkowski & Cermak, 2020; Stacey et al., 2019). The difference in autistic adults' MVPA minutes per day ranged from 1 to 21 min less than that of the neurotypical adults (Benson et al., 2019; Stacey et al., 2019). The greatest difference in daily physical activity participation between groups was in the one study that used accelerometry for physical activity measurement (Benson et al., 2019). In one study, MVPA minutes per day

were the same for autistic and neurotypical adults (Takeshi & Chisato, 2014).

Sedentary Behaviours Findings

Of the six included studies with both autistic and neurotypical participants, three studies reported data about time spent in sedentary behaviour (ASD *mean* = 158 sedentary behaviour min/day, *range* = 150–168; neurotypical *mean* = 85 sedentary behaviour min/day, *range* = 45–147) (Engelhardt et al., 2017; Jozkowski & Cermak, 2020; Stacey et al., 2019). Within all three of these studies, autistic adults' level of sedentary behaviour participation was reported to be higher than the neurotypical adults (Engelhardt et al., 2017; Jozkowski & Cermak, 2020; Stacey et al., 2019). The difference in autistic adults' sedentary behaviour minutes per day ranged from 8 to 105 min greater than the neurotypical adults (Engelhardt et al., 2017; Jozkowski & Cermak, 2020; Stacey et al., 2019).

Correlates

Within this review, eight studies reported correlates of physical activity, sedentary behaviour or both among autistic adults (Table 2) (Benson et al., 2019; Bishop-Fitzpatrick et al., 2017; Engelhardt et al., 2017; Finke et al., 2018; Garcia-Pastor et al., 2019; Hamm & Yun, 2019; Jozkowski & Cermak, 2020; Takeshi & Chisato, 2014). A total of ten potential correlates were identified within the studies. The most commonly described correlate for time spent in physical activity and sedentary behaviour was day of the week (i.e. weekday vs weekend) (Benson et al., 2019; Engelhardt et al., 2017; Garcia-Pastor et al., 2019). In two studies, the day of the week was suggested to have no influence on autistic adults' physical activity and sedentary behaviour levels (Benson et al., 2019; Finke et al., 2018). Alternatively, the remaining study reported a larger number of hours per day spent in sedentary behaviour, specifically videogames, on weekends compared to weekdays for both autistic and neurotypical adults (Engelhardt et al., 2017).

Participant's fitness level was positively associated with levels of physical activity (Jozkowski & Cermak, 2020). Sleep, health-related quality of life and social interaction were all also shown to be positively correlated with physical activity levels (Benson et al., 2019; Hamm & Yun, 2019; Jozkowski & Cermak, 2020). Conversely, obesity was negatively associated with autistic adults' level of physical activity, and older participants were observed to have lower levels of physical activity declined (Garcia-Pastor et al., 2019). In males, arterial stiffness was negatively associated with physical activity level (Takeshi & Chisato, 2014). Additionally, levels of perceived stress were reported to be negatively

associated with participants' physical activity and sedentary behaviour levels (Bishop-Fitzpatrick et al., 2017).

Higher levels of sedentary behaviour participation were reported to be positively associated with being of male gender compared to female (Finke et al., 2018). As participants aged, their sedentary behaviour participation was also found to increase (Garcia-Pastor et al., 2019). Furthermore, pathological game use symptoms were reported to have positive associations with time spent in sedentary behaviour (Engelhardt et al., 2017; Garcia-Pastor et al., 2019).

Discussion

This systematic review investigated what is currently known about autistic adults' physical activity and sedentary behaviour levels, along with associated correlates to these behaviours. Of the 2834 studies identified through database searching, only 12 met the inclusion criteria for analysis. Findings of this review highlight the scarcity of available evidence on the physical activity and sedentary behaviour of autistic adults. Additionally, various data collection methods were used within the primary studies, some of which may not be best suited for use within the autistic adult population. Of the six studies using self-report measures of physical activity, three of the autistic adult samples would be deemed to be insufficiently active (i.e. not participate in > 150 MVPA min/week) (Bishop-Fitzpatrick et al., 2017; Jozkowski & Cermak, 2020; Stacey et al., 2019). The results of one of the two studies using accelerometry identified that their sample as not meeting the physical activity guidelines (i.e. not participate in > 150 MVPA min/week) (Benson et al., 2019). These differing data collection methods also yielded inconsistent estimates of autistic adults' physical activity and sedentary behaviour participation. Therefore, estimates obtained from this review are not likely to be a true representation of autistic adults' physical activity and sedentary behaviour participation. This signals the need for additional studies using more accurate and reliable methods to quantify these behaviours.

The primary data collection method used by the studies included in this review was self-report. Self-report may be prone to inaccuracies regarding the frequency, duration and intensity of activities retrospectively recalled (Dyrstad et al., 2014). This can result in an overestimation of activity levels when compared to device-derived measures (Benson et al., 2019). Self-reporting accuracy limitations may be amplified among autistic adults due to their difficulties with social communication and interactions and/or potential co-occurring intellectual disability (American Psychiatric Association, 2013; Benson et al., 2019). In addition, most questionnaires only allow for reporting of time spent in sedentary leisure activities (e.g. videogaming, watching

television). This does not take into consideration time spent sedentary during productive activities (e.g. office work, university lectures), which are likely to be a large proportion of autistic adults' daily activity. So evidenced by the wide range of estimates obtained, relying solely on self-report measures to estimate autistic adults' physical activity and sedentary behaviour may not be a reliable data collection method.

Measurement devices, such as accelerometers or wearable activity trackers, are not reliant on participants' ability to understand questions and/or respectively recall activity. Therefore, measurement devices are more likely to capture accurate estimates of physical activity and sedentary behaviour throughout the day. Accelerometry has been validated across multiple populations and is currently viewed as the 'gold standard' measurement of activity levels (Sylvia et al., 2014; Verceles & Hager, 2015). In the current review, accelerometers were only used in two studies (Benson et al., 2019; Garcia-Pastor et al., 2019), and one of these did not separate sleep time from daytime sedentary behaviour in their results (Benson et al., 2019). This makes the interpretation of sedentary behaviour estimates difficult and may have skewed results. Wearable activity trackers may be useful in studies wanting to objectively measure time spent in autistic adults' physical activity and sedentary behaviour, due to their lower cost, accessibility, and low burden of wear (Semanik et al., 2020). Wearable activity trackers were used in three studies. However, their small sample sizes and low ecological validity should also be considered in the interpretation of results (LaLonde et al., 2014; Li et al., 2019; Nastasi et al., 2020). Due to the limited number of high-quality studies using measurement devices, the proportion of the autistic adult population meeting the recommended levels of physical activity is unknown.

The quantification of physical activity and sedentary behaviour has been inconsistent between neurotypical adults and autistic adults. The ability to accurately quantify physical activity and sedentary behaviour is important due to their association with chronic health conditions and mental well-being (Kokkinos, 2012; World Health Organization, 2017). The secondary sequelae of insufficient physical activity may be exaggerated for autistic adults due to a greater incidence of poor health outcomes, such as lower life expectancy and higher rates of mental health conditions compared to neurotypical adults (Hirvikoski et al., 2016; van Heijst & Geurts, 2015). Current research has also shown that physical activity participation can provide additional benefits specific to the autistic adult population, such as reduced stereotypic behaviours, anxiety, and stress levels, as well as providing valuable opportunities for social interaction (Lang et al., 2010). Factors reported to be associated with autistic adults' physical activity and sedentary behaviour participation were limited within the included studies yet provide initial insights for future studies to consider.

Correlates to autistic adults' physical activity and sedentary behaviour were reported in eight studies, with 10 potential correlates defined. Day of the week was the only correlate to be reported across multiple studies, yet findings were inconsistent (Benson et al., 2019; Engelhardt et al., 2017; Garcia-Pastor et al., 2019). In one study, autistic adults were demonstrated to spend greater time in sedentary behaviour on weekends compared to weekdays (Engelhardt et al., 2017), which is consistent with findings with prior research on neurotypical adults (Marshall et al., 2015). Additionally, it can be presumed that individuals would also participate in lower levels of physical activity on weekends compared to weekdays (Chaput et al., 2014). This is similar to what has been shown in autistic adolescents, whereby time spent in physical activity was significantly lower on weekends compared to weekdays (Stanish et al., 2017). These differences may be attributed to having more free time available on weekends than weekdays and providing a basis for research to increase weekend physical activity participation in the autistic population. However, due to scarce and inconsistent reporting of the impact that the day of the week has on autistic adults' activity levels, it is unclear how this can be targeted by interventions until clearer measurements of these behaviours on days of the week are obtained. Furthermore, studies on autistic children and neurotypical adults report levels of physical activity participation to be negatively associated with increasing age and poorer health, and positively associated with social interaction and being of male gender (Bennie et al., 2016; Healy & Garcia, 2019; Jones et al., 2017; Pan et al., 2011). These correlates show similarities to those reported across the studies of this review and may also apply to autistic adults.

The interpretations of this review should be considered in the context of the limitations present within the primary studies. Such limitations included the poor quality of reported data and questionable reliability and accuracy of data collection methods. Due to the differing nature of the primary studies, a series of assumptions had to be made to allow for standardisation (Appendix B) and, thus, limit the accuracy of numerical values derived. Furthermore, the potential occurrence of common methodological flaws, such as inaccuracies in assessing the quality of included studies, cannot be ruled out (Garg et al., 2008). Biases may have also occurred as there is inevitably subjectivity in decision-making to determine study eligibility. Finally, the authors manually extracted data from the studies and did not contact primary researchers for verification of accuracy or to attain additional research information (e.g. missing/unreported data). This may have impacted the reliability of reported results and subsequent interpretations, further limiting extrapolation of findings.

Future research examining autistic adults' physical activity and sedentary behaviour participation should use more

accurate and reliable methods to capture the information. In particular, measurement devices are recommended over self-report measures for this population as they eliminate potential inaccuracies due to the possibility of autistic adults having difficulty interpreting questions. Studies including neurotypical control groups and large samples representative of the autistic adult population are also warranted. Future studies should also examine a range of potential correlates to allow for the directionality and relative strength of associations with physical activity and sedentary behaviour to be examined. This may assist in the development of tailored interventions to promote physical activity in the autistic adult population.

Currently, it is not possible to accurately estimate autistic adults' physical activity and sedentary behaviour levels due to the poor quality of the current literature and the limited number of studies. However, based on the available evidence, it appears that autistic adults may be less physically active and more sedentary than neurotypical adults and that correlates that influence these behaviours may exist. Hence, this population may be at greater risk of numerous physical and mental health consequences. Accurate estimates of autistic adults' physical activity and sedentary behaviour are urgently needed in order to determine the true extent of inactivity and the proportion of the autistic adult population participating in sufficient physical activity.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s40489-022-00305-x>.

Acknowledgements This work would be impossible without the contributions of the autistic individuals across the world. We are grateful to the Curtin School of Allied Health for the support of the research.

Funding Open Access funding enabled and organized by CAUL and its Member Institutions.

Declarations

Conflict of Interest All authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Australian Bureau of Statistics. (2019). *National health survey: First results, 2017–18* (No. 4364.055.001). <https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/4364.0.55.001~2017-18~Main%20Features~Physical%20activity~115>
- Department of Health. (2019). Australia's physical activity and sedentary behaviour guidelines and the Australian 24-hour movement guidelines. <https://www1.health.gov.au/internet/main/publishing.nsf/Content/health-pubhlth-strateg-phys-act-guidelines>
- Bennie, J. A., Pedisic, Z., van Uffelen, J. G. Z., Gale, J., Banting, L. K., Vergeer, I., Stamatakis, E., Bauman, A. E., & Biddle, S. J. H. (2016). The descriptive epidemiology of total physical activity, muscle-strengthening exercises and sedentary behaviour among Australian adults – results from the National Nutrition and Physical Activity Survey. *BMC Public Health*, 16(1), 73. <https://doi.org/10.1186/s12889-016-2736-3>
- Benson, S., Bender, A. M., Wickenheiser, H., Naylor, A., Clarke, M., Samuels, C. H., & Werthner, P. (2019). Differences in sleep patterns, sleepiness, and physical activity levels between young adults with autism spectrum disorder and typically developing controls. *Developmental Neurorehabilitation*, 22(3), 164–173 (2019). <https://doi.org/10.1080/17518423.2018.1501777>
- Bishop-Fitzpatrick, L., Smith DaWalt, L., Greenberg, J. S., & Mailick, M. R. (2017). Participation in recreational activities buffers the impact of perceived stress on quality of life in adults with autism spectrum disorder. *Autism Research*, 10(5), 973–982. <https://doi.org/10.1002/aur.1753>
- Chaput, J.-P., Carson, V., Gray, C. E., & Tremblay, M. S. (2014). Importance of all movement behaviors in a 24 hour period for overall health. *International Journal of Environmental Research and Public Health*, 11(12), 12575–12581. <https://www.mdpi.com/1660-4601/11/12/12575>
- Dyrstad, S. M., Hansen, B. H., Holme, I. M., & Anderssen, S. A. (2014). Comparison of self-reported versus accelerometer-measured physical activity. *Medicine & Science in Sports & Exercise*, 46(1), 99–106. <https://doi.org/10.1249/MSS.0b013e3182a0595f>
- Engelhardt, C. R., Mazurek, M. O., & Hilgard, J. (2017). Pathological game use in adults with and without autism spectrum disorder. *PeerJ*, 5, e3393. <https://doi.org/10.7717/peerj.3393>
- Finke, E. H., Hickerson, B. D., & Kremkow, J. M. D. (2018). To be quite honest, if it wasn't for videogames i wouldn't have a social life at All”: Motivations of young adults with autism spectrum disorder for playing videogames as leisure. *American Journal of Speech-Language Pathology*, 27(2), 672–689 (2018). doi:https://doi.org/10.1044/2017_AJSLP-17-0073
- Garcia-Pastor, T., Salinero, J. J., Theirs, C. I., & Ruiz-Vicente, D. (2019). Obesity status and physical activity level in children and adults with autism spectrum disorders: A pilot study. *Journal of Autism Developmental Disorders*, 49(1), 165–172. <https://doi.org/10.1007/s10803-018-3692-9>
- García-Villamizar, D. A., & Dattilo, J. (2010). Effects of a leisure programme on quality of life and stress of individuals with ASD. *Journal of Intellectual Disability Research*, 54(7), 611–619. <https://doi.org/10.1111/j.1365-2788.2010.01289.x>
- Garg, A. X., Hackam, D., & Tonelli, M. (2008). Systematic review and meta-analysis: When one study is just not enough. *Clinical Journal of the American Society of Nephrology*, 3(1), 253–260. <https://doi.org/10.2215/cjn.01430307>
- Hamm, J., & Yun, J. (2019). Influence of physical activity on the health-related quality of life of young adults with and without autism spectrum disorder. *Disability and Rehabilitation*, 41(7), 763–769. <https://doi.org/10.1080/09638288.2017.1408708>

- Healy, S., & Garcia, J. M. (2019). Psychosocial correlates of physical activity participation and screen-time in typically developing children and children on the autism spectrum. *Journal of Developmental and Physical Disabilities*, 31(3), 313–328. <https://doi.org/10.1007/s10882-018-9642-9>
- Healy, S., Aigner, C. J., Haegele, J. A., & Patterson, F. (2019). Meeting the 24-hr movement guidelines: An update on US youth with autism spectrum disorder from the 2016 National Survey of Children's Health. *Autism Research*, 12(6), 941–951. <https://doi.org/10.1002/aur.2095>
- Hillier, A., Buckingham, A., & Schena, D. (2020). Physical activity among adults with autism: Participation, attitudes, and barriers. *Perceptual and Motor Skills*, 127(5), 874–890. <https://doi.org/10.1177/0031512520927560>
- Hirvikoski, T., Mittendorf-Rutz, E., Boman, M., Larsson, H., Lichtenstein, P., & Bölte, S. (2016). Premature mortality in autism spectrum disorder. *British Journal of Psychiatry*, 208(3), 232–238. <https://doi.org/10.1192/bjp.bp.114.160192>
- Howlin, P., Moss, P., Savage, S., Bolton, P., & Rutter, M. (2015). Outcomes in adult life among siblings of individuals with autism. *Journal of Autism and Developmental Disorders*, 45(3), 707–718. <https://doi.org/10.1007/s10803-014-2224-5>
- Jones, R. A., Downing, K., Rinehart, N. J., Barnett, L. M., May, T., McGillivray, J. A., Papadopoulos, N. V., Skouteris, H., Timperio, A., & Hinkley, T. (2017). Physical activity, sedentary behavior and their correlates in children with autism spectrum disorder: A systematic review. *PLoS ONE*, 12(2), e0172482. <https://doi.org/10.1371/journal.pone.0172482>
- Jozkowski, A. C., & Cermak, S. A. (c). (2019). Moderating effect of social interaction on enjoyment and perception of physical activity in young adults with autism spectrum disorders. *International Journal of Developmental Disabilities*, 66(3), 222–234. <https://doi.org/10.1080/20473869.2019.1567091>
- Kmet, L., Lee, R., & Cook, L. (2004). Standard quality assessment criteria for evaluating primary research papers from a variety of fields. <https://www.ihe.ca/advanced-search/standard-quality-assessment-criteria-for-evaluating-primary-research-papers-from-a-variety-of-fields>
- Knüppel, A., Telléus, G. K., Jakobsen, H., & Lauritsen, M. B. (2019). Characteristics of young adults with autism spectrum disorder performing different daytime activities. *Journal of Autism and Developmental Disorders*, 49(2), 542–555. <https://doi.org/10.1007/s10803-018-3730-7>
- Kokkinos, P. (2012). Physical activity, health benefits, and mortality risk. *ISRN Cardiology*, 2012, 718789. <https://doi.org/10.5402/2012/718789>
- LaLonde, K. B., MacNeill, B. R., Eversole, L. W., Ragotzy, S. P., & Poling, A. (2014). Increasing physical activity in young adults with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 8(12), 1679–1684. <https://doi.org/10.1016/j.rasd.2014.09.001>
- Lang, R., Koegel, L. K., Ashbaugh, K., Regester, A., Ence, W., & Smith, W. (2010). Physical exercise and individuals with autism spectrum disorders: A systematic review. *Research in Autism Spectrum Disorders*, 4(4), 565–576 (2010). doi:<https://doi.org/10.1016/j.rasd.2010.01.006>
- Lear, S. A., Hu, W., Rangarajan, S., Gasevic, D., Leong, D., Iqbal, R., Casanova, A., Swaminathan, S., Anjana, R. M., Kumar, R., Rosengren, A., Wei, L., Yang, W., Chuangshi, W., Huaxing, L., Nair, S., Diaz, R., Swidon, H., Gupta, R., Mohammadifard, N., Lopez-Jaramillo, P., Oguz, A., Zatonska, K., Seron, P., Avezum, A., Poirier, P., Teo, K., & Yusuf, S. (2017). The effect of physical activity on mortality and cardiovascular disease in 130 000 people from 17 high-income, middle-income, and low-income countries: the PURE study. *The Lancet*, 390(10113), 2643–2654 (2017). doi:[https://doi.org/10.1016/S0140-6736\(17\)31634-3](https://doi.org/10.1016/S0140-6736(17)31634-3)
- Lee, L., Packer, T. L., Tang, S. H., & Girdler, S.. Self-management education programs for age-related macular degeneration: A systematic review. *Australasian Journal on Ageing*, 27(4), 170–176. <https://doi.org/10.1111/j.1741-6612.2008.00298.x>
- Li, A., Curiel, H., Ragotzy, S. P., & Poling, A. (2019). Using a lottery to promote physical activity by young adults with developmental disabilities. *Behavior Analysis in Practice*, 12(3), 612–616. <https://doi.org/10.1007/s40617-018-00292-8>
- MacDonald, M., Esposito, P., & Ulrich, D. (2011). The physical activity patterns of children with autism. *BMC Research Notes*, 4(1), 422 <https://doi.org/10.1186/1756-0500-4-422>
- Marshall, S., Kerr, J., Carlson, J., Cadmus-Bertram, L., Patterson, R., Wasilenko, K., Crist, K., Rosenberg, D., & Natarajan, L. (2015). Patterns of weekday and weekend sedentary behavior among older adults. *Journal of Aging and Physical Activity*, 23(4), 534. <https://doi.org/10.1123/japa.2013-0208>
- Menear, K. S., & Neumeier, W. H. (2015). Promoting physical activity for students with autism spectrum disorder: Barriers, benefits, and strategies for success. *Journal of Physical Education, Recreation & Dance*, 86(3), 43–48. <https://doi.org/10.1080/07303084.2014.998395>
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *BMJ*, 339, b2535. <https://doi.org/10.1136/bmj.b2535>
- Must, A., Phillips, S. M., Curtin, C., Anderson, S. E., Maslin, M., Lividini, K., & Bandini, L. G. (2014). Comparison of sedentary behaviors between children with autism spectrum disorders and typically developing children. *Autism*, 18(4), 376–384. <https://doi.org/10.1177/1362361313479039>
- Nastasi, J. A., Sheppard, R. D., & Raiff, B. R. (2020). Token-economy-based contingency management increases daily steps in adults with developmental disabilities. *Behavioral Interventions*, 35(2), 315–324. <https://doi.org/10.1002/bin.1711>
- National Institute for Health Research. (2020). Welcome to prospero – International prospective register of systematic reviews. <https://www.crd.york.ac.uk/prospero>
- Nichols, C., Block, M. E., Bishop, J. C., & McIntire, B. (2018). Physical activity in young adults with autism spectrum disorder: Parental perceptions of barriers and facilitators. *Autism*, 23(6), 1398–1407. <https://doi.org/10.1177/1362361318810221>
- Pan, C.-Y., Tsai, C.-L., & Hsieh, K.-W. (2011). Physical activity correlates for children with autism spectrum disorders in middle school physical education. *Research Quarterly for Exercise and Sport*, 82(3), 491–498. <https://doi.org/10.1080/02701367.2011.10599782>
- Pinder-Amaker, S. (2014). Identifying the unmet needs of college students on the autism spectrum. *Harvard Review of Psychiatry*, 22(2), 125–137. <https://doi.org/10.1097/hrp.0000000000000032>
- American Psychiatric Association. (2013). The diagnostic and statistical manual of mental disorders: DSM 5. *American Psychiatric Publishing*.
- Semanik, P., Lee, J., Pellegrini, C. A., Song, J., Dunlop, D. D., & Chang, R. W. (2020). Comparison of physical activity measures derived from the fitbit flex and the ActiGraph GT3X+ in an employee population with chronic knee symptoms. *ACR Open Rheumatology*, 2(1), 48–52. <https://doi.org/10.1002/acr2.11099>
- Simpson, K., Adams, D., Bruck, S., & Keen, D. (2019). Investigating the participation of children on the autism spectrum across home, school, and community: A longitudinal study. *Child: Care, Health and Development*, 45(5), 681–687. <https://doi.org/10.1111/cch.12679>
- Stacey, T.-L., Froude, E. H., Trollor, J., & Foley, K.-R. (2019). Leisure participation and satisfaction in autistic adults and neurotypical adults. *Autism*, 23(4), 993–1004. <https://doi.org/10.1177/1362361318791275>

- Stanish, H. I., Curtin, C., Must, A., Phillips, S., Maslin, M., & Bandini, L. G.. Physical activity levels, frequency, and type among adolescents with and without autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 47(3), 785–794. <https://doi.org/10.1007/s10803-016-3001-4>
- Strath, S. J., Pfeiffer, K. A., & Whitt-Glover, M. C. (2012). Accelerometer use with children, older adults, and adults with functional limitations. *Medicine & Science in Sports & Exercise*, 44(1S), S77–S85. <https://doi.org/10.1249/MSS.0b013e3182399eb1>
- Sylvia, L. G., Bernstein, E. E., Hubbard, J. L., Keating, L., & Anderson, E. J. (2014). Practical guide to measuring physical activity. *Journal of the Academy of Nutrition and Dietetics*, 114(2), 199–208. <https://doi.org/10.1016/j.jand.2013.09.018>
- Takeshi, O., & Chisato, O. (2014). Participation in physical activity and arterial stiffness in males with autism spectrum disorder. *Artery Research*, 8(3), 110–114. <https://doi.org/10.1016/j.artres.2014.04.001>
- Tanguay, P. E., & Lohr, D. W. (2015). Autism spectrum disorders. In M. K. Duncan (Ed.), *Dulcan's Textbook of Child and Adolescent Psychiatry* (2nd ed., pp. 83–94). <https://doi.org/10.1176/appi.books.9781615370306.md08>
- Tobin, M. C., Drager, K. D. R., & Richardson, L. F. (2014). A systematic review of social participation for adults with autism spectrum disorders: Support, social functioning, and quality of life. *Research in Autism Spectrum Disorders*, 8(3), 214–229. <https://doi.org/10.1016/j.rasd.2013.12.002>
- van Heijst, B. F., & Geurts, H. M. (2015). Quality of life in autism across the lifespan: A meta-analysis. *Autism*, 19(2), 158–167. <https://doi.org/10.1177/1362361313517053>
- Verceles, A. C., & Hager, E. R. (2015). Use of accelerometry to monitor physical activity in critically ill subjects: A systematic review. *Respiratory Care*, 60(9), 1330–1336. <https://doi.org/10.4187/respcare.03677>
- World Health Organization. (2017). *Physical activity*. <https://www.who.int/dietphysicalactivity/pa/en/>
- Yano, S., Koohsari, M. J., Shibata, A., Ishii, K., Mavoa, S., & Oka, K. (2019). Assessing physical activity and sedentary behavior under free-living conditions: Comparison of active style Pro HJA-350IT and ActiGraphTM GT3X+. *International journal of environmental research and public health*, 16(17), 3065. <https://www.mdpi.com/1660-4601/16/17/3065>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.