Correction to: Measuring physical performance in highly active older adults: associations with age and gender?

Gro Gujord Tangen^{1,2,3} · Hilde Stendal Robinson¹

Published online: 17 August 2019
© Springer Nature Switzerland AG 2019

Correction to: Aging Clinical and Experimental Research
<https://doi.org/10.1007/s40520-019-01190-x>

In the published article, the Table 2 was published incorrectly. The correct Table 2 is given below.

The original article has been updated.

The original article can be found online at <https://doi.org/10.1007/s40520-019-01190-x>.

✉ Gro Gujord Tangen
g.g.tangen@medisin.uio.no
Hilde Stendal Robinson
h.s.robinson@medisin.uio.no

¹ Department of Interdisciplinary Health Sciences, Institute of Health and Society, University of Oslo, P.O. Box 1089, Blindern, 0317 Oslo, Norway

² Norwegian National Advisory Unit on Ageing and Health, Vestfold Hospital Trust, Tonsberg, Norway

³ Department of Geriatric Medicine, Oslo University Hospital, Oslo, Norway

Table 2 Results on performance-based tests of physical function

Test	All (<i>n</i> = 105)	Men (<i>n</i> = 57)	Women (<i>n</i> = 48)	<i>B</i> ^a / <i>OR</i> ^b (95% CI)	<i>p</i>
General mobility					
SPPB (0–12)					
Median (IQR)	12 (2)	12 (1)	12 (2)		
Min–max	4–12	6–12	4–12		
> 10 points, <i>n</i> (%)	79 (75)	44 (77)	35 (73)	1.8 (0.7, 4.9) ^a	0.227 ^c
Gait speed, m/s, mean (SD)	1.14 (0.2)	1.11 (0.2)	1.18 (0.2)	− 0.4 (− 1.1, 0.04) ^b	0.292 ^d
TUG, s, mean (SD)	8.2 (1.5)	8.5 (1.5)	7.8 (1.5)	0.5 (− 0.1, 1.0) ^b	0.075 ^d
Balance					
OLS-EO, s median (IQR)					
Median (IQR)	24.7 (20.2)	24.7 (20.6)	26.5 (19.8)		
Min–max	2.2–30.0	3.0–30.0	2.2–30.0		
30 s, <i>n</i> (%)	49 (46.7)	25 (43.9)	24 (50)	1.1 (0.5, 2.6) ^a	0.839 ^c
OLS-EC, median (IQR)					
Median (IQR)	3.0 (3.4)	3.0 (3.6)	3.0 (3.3)		
Min–max	0.7–30.0	0.7–30.0	1.1–14.5		
< 2 s, <i>n</i> (%)	78 (74.3)	43 (75.4)	35 (72.9)	1.6 (0.6, 4.1) ^a	0.346 ^c
Mini-BESTest (0–28)					
Median (IQR)	24.0 (4.0)	25.0 (4)	23.6 (3.2)	1.4 (0.4, 2.3) ^b	0.009 ^d
Min–max	14–28	15–28	14–28		
Muscle strength					
30sSTS, mean (SD)					
Mean (SD)	16.9 (5.1)	17.6 (5.1)	16.2 (5.0)	2.1 (0.3, 4.0) ^b	0.026 ^d
Grip strength, kg, mean (SD)					
Dominant hand	32.7 (9.5)	39.1 (7.6)	25.1 (4.6)	15.1 (12.8, 17.3) ^b	< 0.001 ^d
Non-dominant hand	32.0 (9.7)	38.9 (6.9)	23.7 (4.8)	16.0 (13.7, 18.2) ^b	< 0.001 ^d
Endurance					
Stair test, s, median (IQR)					
Median (IQR)	51.3 (16.1)	50.8 (14.7)	51.5 (20.6)	− 9.3 (1.7, 2.9) ^b	0.002 ^d
Min–max	33.9–125.9	33.9–103.1	37.9–125.9		

OR odds ratio, *SPPB* short physical performance battery, *OLS-EO* one leg standing-eyes open, *OLS-EC* one leg standing-eyes closed, *30sSTS* 30-seconds sit-to-stand test, *Mini-BESTest* mini balance evaluations systems test, *TUG* timed up and go

^a*OR*

^bUnstandardized coefficient

^cLogistic regression, using dichotomized variable as dependent variable, with age as covariate

^dLinear regression analysis with age as covariate

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.