



Biological Frailty Index in centenarians

Beatrice Arosio¹ · Annalisa Geraci² · Evelyn Ferri² · Daniela Mari³ · Matteo Cesari^{1,4}

Received: 22 July 2021 / Accepted: 28 September 2021 / Published online: 16 October 2021
© The Author(s) 2021

Abstract

This study measured the subclinical frailty of centenarians by looking at the accumulation of their biological abnormalities. For this aim, a biological Frailty Index (FI) was computed in centenarians living in Northern Italy. The median value of the biological FI was 0.33 (interquartile range, IQR 0.28–0.41). The biological FI did not significantly differ between women (0.34, IQR 0.31–0.39) and men (0.32, IQR 0.26–0.43). The biological FI seems to have a narrower distribution compared to clinical FI we previously computed in the same cohort. In conclusion, our study suggests that centenarians benefit from exceptional biological reserves that might be underestimated by clinical appearances.

Keywords Centenarians · Biological frailty index · Biological reserves · Longevity

Introduction

The amount of people reaching old age has been growing exponentially in the last decades, and centenarians represent the fastest-growing group (World Population Prospects 2019: Highlights; <https://population.un.org/wpp/>). Centenarians are persons with an extraordinary adaptive capacity, probably thanks to unusual functional reserves. Centenarians may live with debilitating disease, but still present an advantage in terms of incident disability and death [1, 2].

They constitute a very heterogeneous population as result of lifestyle habits, environmental factors, and histories that have differently affected their biological and clinical profile over the life course [3, 4]. Thus, centenarians may be subjects with not only good but also very poor health status, as

demonstrated by the different degrees of frailty we previously reported [5].

Aging occurs at molecular and cellular levels [6]. Interestingly, centenarians seem to express molecular signatures suggestive of a slower process compared to other persons [7, 8].

Recently, it has been explained that Frailty Index (FI) exclusively based on biological parameters may define the biological age of the individual, potentially capturing variations in the health status before the manifestation of clinical deficits [6, 9].

The aim of this study was to measure the subclinical frailty of centenarians by looking at the accumulation of their biological abnormalities. Since available measures of biological age are not optimized to disentangle the heterogeneity that characterizes centenarians [10], in this study, we have computed a biological FI by the means of blood tests in a cohort of well-characterized centenarians living in Northern Italy.

Study design

The participants belonged to a large cohort enrolled during a study conducted between 2007 and 2014 and funded by the Italian Ministry of University and Scientific Research. The cohort was composed by 125 centenarians. Forty-six registry offices in Northern Italy were contacted to collect dates of birth of living people close to 100 years at the

✉ Beatrice Arosio
beatrice.arosio@unimi.it

¹ Laboratorio di Geriatria, Department of Clinical Sciences and Community Health, University of Milan, Via Pace 9, 20122 Milan, Italy

² Geriatric Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Via Pace 9, 20122 Milan, Italy

³ Laboratorio Sperimentale di Ricerche di Neuroendocrinologia Geriatrica ed Oncologica, Istituto Auxologico Italiano, IRCCS, Via Zucchi 18, 20095 Cusano Milanino, Italy

⁴ Geriatric Unit, IRCCS Istituti Clinici Scientifici Maugeri, Via Camaldoli 64, 20138 Milan, Italy

enrolment. Sixty-five out of 125 centenarians with all available variables needed for the computation of the biological FI were included. All these persons had a clinical FI already described [5].

Briefly, a trained multidisciplinary team went to each centenarian's house or nursing home to administer a standard structured questionnaire and collect blood samples [11].

The biological FI was computed considering a total of 42 variables including routine blood tests [6], telomere length [12] and Apolipoprotein E genotype [9]. The 20th and 80th percentiles of each variable were considered as cut-points. The values under the 20th percentile and over the 80th percentile were considered abnormal. These biomarkers and their cut-points are presented in Table 1.

Each biomarker was categorized to assume the value of 0 if its value fell within the range of normality or 1 if abnormal. The biological FI was then calculated as the ratio between the number of biomarkers presenting abnormal values and the number of considered biomarkers ($n = 42$).

Results

Overall, a total of 65 centenarians (46 women and 19 men) were included in this study. The mean age of the sample was 101.3 (standard deviation, SD 2.0) years. The age was similar between women and men (101.2, SD 2.1 and 101.6, SD 2.0, respectively). As expected, the prevalence of women was higher than men (71% and 29%, respectively).

The median value of the biological FI was 0.33 (interquartile range, IQR 0.28–0.41). The biological FI did not significantly differ between women (0.34, IQR 0.31–0.39) and men (0.32, IQR 0.26–0.43). Figure 1 shows the distribution of the biological FI, which ranged between 0.11 and 0.69. Age was weakly correlated with the biological FI (Spearman's $r = 0.26$, $p = 0.04$).

Discussion

To our knowledge, this is the first study measuring a biological FI in a cohort of well-characterized centenarians. Interestingly, it seems to have a more narrow distribution compared to the clinical FI we previously computed [5].


In fact, in the same cohort, we reported a higher clinical FI (median 0.50, IQR 0.40–0.58), and a wider spectrum of values (ranging between 0.13 and 0.73) [5].

In a cohort of persons aged 80 years and older, it has been reported that the clinically fittest persons (FI values between 0 and 0.02) had a mean biological FI of 0.33, indicating that this latter is able to detect the subclinical accumulation of deficits and anticipate the clinical phenotype [9]. Similarly, community-dwelling men aged 40–79 showed a higher

Table 1 Biomarkers and cut-points of the biological FI

Biomarkers	20th–80th percentile	
	Men	Women
Glycemia (mg/dl)	80–111	78–97
Insulin (μ U/ml)	2.8–11.8	3.6–12.0
Albumin (g/dl)	3.4–4.2	3.2–4.1
Urea (mg/dl)	39.8–74.6	37.2–71.0
Creatinine (mg/dl)	0.9–1.3	0.6–1.3
Uric Acid (mg/dl)	4.7–7.4	4.2–6.3
Cholesterol (mg/dl)	152–220	152–221
HDL (mg/dl)	41.0–62.0	39.8–59.0
Triglycerides (mg/dl)	78–154	73.4–149.6
Direct Bilirubin (mg/dl)	0.06–0.18	0.05–0.14
Total Bilirubin (mg/dl)	0.3–0.7	0.2–0.6
AST (U/L)	11–19	14–20
ALT (U/L)	5–10	5–13
GGT (U/L)	11.2–29.4	11.0–41.4
ALP (U/L)	65.0–133.4	58.4–135.6
Calcium (mg/dl)	9.5–10.2	9.4–10.3
Iron (μ g/dl)	42.6–98.0	47.2–101.8
Phosphorus (mg/dL)	2.7–3.4	3.0–4.0
hs-CRP (mg/dl)	1.0–12.7	0.9–11.9
Lymphocytes ($\times 10^3/\mu$ l)	1.14–1.78	1.06–1.86
Leukocytes ($\times 10^3/\mu$ l)	5.8–7.8	5.2–7.4
Monocytes ($\times 10^3/\mu$ l)	0.3–0.5	0.2–0.4
Haemoglobin (g/dl)	11.9–13.9	10.9–13.1
MCV (fl)	83–93	81–91
MCH (pg)	27.4–31.4	28.1–31.0
MCHC (g/dl)	32.6–35.6	33.1–36.1
Platelets ($\times 10^3/\mu$ l)	175.0–254.0	162.4–296.2
CMV	Negativity	
PAI-1 Act (ng/ml)	1.0–5.1	1.0–3.9
Fibrinogen Antigen (mg/ml)	2.8–5.6	3.1–6.3
VWF Antigen (%)	165.6–294.4	186.0–318.4
Adams-13 Antigen (%)	30.1–52.0	37.5–49.6
IGF-1 (ng/ml)	41.3–108.4	42.9–103.5
FT3 (pg/ml)	2.0–2.9	2.2–2.8
FT4 (ng/ml)	9.8–14.3	9.9–14.0
TSH (μ U/ml)	1.4–6.7	1.1–3.0
PTH (ng/l)	40.7–130.4	53.6–220.8
SHBG (nmol/l)	65.0–108.8	70.6–137.2
Testosterone (nmol/l)	4.7–15.4	0.2–0.9
25-OH Vitamin D (μ g/l)	3.0–9.5	3.0–8.6
Telomere Length	> 0.76	> 0.87
Apolipoprotein E e4	Negativity	

HDL High Density Lipoprotein, *AST* Aspartate Transaminase, *ALT* Alanine Transferase, *GGT* γ -Glutamyl Transpeptidase, *ALP* Alkaline Phosphatase, *hs-CRP* High Sensitivity C-reactive Protein, *MCV* Mean Corpuscular Volume, *MCH* Mean Corpuscular Hemoglobin, *MCHC* Mean Corpuscular Hemoglobin Concentration, *CMV* Cytomegalovirus, *PAI-1 Act* Plasminogen Activator Inhibitor-1 Activity, *VWF* Von Willebrand Factor, *IGF-1* Insulin-like Growth Factor-1, *FT3* Free Triiodothyronine, *FT4* Free Thyroxine, *TSH* Thyroid-Stimulating Hormone, *PTH* Parathyroid Hormone, *SHBG* Sex Hormone Binding Globulin

Fig. 1 Distribution of the biological FI in centenarians

biological FI (based on routine blood tests) compared to the clinical one and a significant association with mortality and adverse health outcomes [6].

Nevertheless, our findings suggest that centenarians benefit from exceptional biological reserves that might be underestimated by clinical appearances. Indeed, in our cohort of centenarians, we got the counterintuitive finding of a biological FI lower than the clinical FI we previously reported.

This result may suggest that, at very advanced age, the biology of the system might be “better” than what clinically manifested. The hypothesis might be explained by the lower relevance that clinical constructs (e.g., definition of the diagnoses) may have with increasing age, especially if compared to the biological substratum feeding them [13]. After all, it is possible that several clinical deficits could be overestimated in centenarians. For example, some tools (e.g., Mini-Mental State Examination) are not validated for extremely old persons [14] and do not often consider peculiar characteristics (e.g., fatigue) potentially affecting their results.

We found a weak association between age and biological FI in centenarians probably because of the narrow range of chronological age and the similar biological FI observed in men and women. This last result is apparently in contrast with the so-called “sex-frailty paradox”, describing women as frailer than men but, at the same time, presenting longer life expectancy [15].

It is possible that, at an extremely advanced age (as in centenarians), the paradox may lose value because of the

ceiling effect determined by the exceptional age and the favourable biology that allows it.

The main limitation of our study resides in the relatively low number of participants, which might have affected the statistical power of our analyses. We cannot also exclude that our sample does not represent the population of centenarians, and that third factors not considered in the study may differently explain our findings. For all these reasons, this study has to be considered an exploratory analysis that needs to be confirmed in a larger population.

In conclusion, our study suggests that centenarians benefit from exceptional biological reserves that might be underestimated by clinical appearances. Further studies are needed to disentangle the relationship between chronological age, biological age, and clinical complexity in older persons, especially at a very advanced age.

Funding This work was supported by the Italian Ministry of University and Research (PRIN 2006 and PRIN 2009 to DM).

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval The protocol received approval from the Ethical Committee of the Fondazione IRCCS Ca’ Granda Ospedale Maggiore Poli-

clinico, Milan (Prot. n. 2035, amendment 30/11/2011). An informed consent was obtained from all participants.

Human and animal participants All procedures involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Informed consent The informed consent was obtained for all individual participants included in the study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Andersen SL (2020) Centenarians as Models of Resistance and Resilience to Alzheimer's Disease and Related Dementias. *Adv Geriatr Med Res* 2:e200018. <https://doi.org/10.20900/agmr20200018>
- Andersen-Ranberg K, Schroll M, Jeune B (2001) Healthy centenarians do not exist, but autonomous centenarians do: a population-based study of morbidity among Danish centenarians. *J Am Geriatr Soc* 49:900–908. <https://doi.org/10.1046/j.1532-5415.2001.49180.x>
- Ostan R, Monti D, Mari D, Arosio B, Gentilini D, Ferri E et al (2019) Heterogeneity of Thyroid Function and Impact of Peripheral Thyroxine Deiodination in Centenarians and Semi-Supercentenarians: Association With Functional Status and Mortality. *J Gerontol A Biol Sci Med Sci* 74:802–810. <https://doi.org/10.1093/gerona/gly194>
- Salvioli S, Capri M, Bucci L, Lanni C, Racchi M, Uberti D et al (2009) Why do centenarians escape or postpone cancer? The role of IGF-1, inflammation and p53. *Cancer Immunol Immunother* 58:1909–1917. <https://doi.org/10.1007/s00262-008-0639-6>
- Arosio B, Ferri E, Casati M, Mari D, Vitale G, Cesari M (2019) The Frailty Index in centenarians and their offspring. *Aging Clin Exp Res* 31:1685–1688. <https://doi.org/10.1007/s40520-019-01283-7>
- Blodgett JM, Theou O, Howlett SE, Rockwood K (2017) A frailty index from common clinical and laboratory tests predicts increased risk of death across the life course. *Geroscience* 39:447–455. <https://doi.org/10.1007/s11357-017-9993-7>
- Tedone E, Arosio B, Gussago C, Casati M, Ferri E, Ogliaeri G et al (2014) Leukocyte telomere length and prevalence of age-related diseases in semisupercentenarians, centenarians and centenarians' offspring. *Exp Gerontol* 58:90–95. <https://doi.org/10.1016/j.exger.2014.06.018>
- Sebastiani P, Federico A, Morris M, Gurinovich A, Tanaka T, Chandler KB et al (2021) Protein signatures of centenarians and their offspring suggest centenarians age slower than other humans. *Aging Cell* 20:e13290. <https://doi.org/10.1111/ace1.13290>
- Mitnitski A, Collerton J, Martin-Ruiz C, Jagger C, von Zglinicki T, Rockwood K et al (2015) Age-related frailty and its association with biological markers of ageing. *BMC Med* 13:161. <https://doi.org/10.1186/s12916-015-0400-x>
- Bacalini MG, Gentilini D, Monti D et al (2021) No association between frailty index and epigenetic clocks in Italian semi-supercentenarians. *Mech Ageing Dev* 197:111514. <https://doi.org/10.1016/j.mad.2021.111514>
- Bucci L, Ostan R, Giampieri E, Cevenini E, Pini E, Scurti M et al (2014) Immune parameters identify Italian centenarians with a longer five-year survival independent of their health and functional status. *Exp Gerontol* 54:14–20. <https://doi.org/10.1016/j.exger.2014.01.023>
- Tedone E, Arosio B, Gussago C et al (2014) Leukocyte telomere length and prevalence of age-related diseases in semisupercentenarians, centenarians and centenarians' offspring. *Exp Gerontol* 58:90–95. <https://doi.org/10.1016/j.exger.2014.06.018>
- Cesari M, Perez-Zepeda MU, Marzetti E (2017) Frailty and Multimorbidity: Different Ways of Thinking About Geriatrics. *J Am Med Dir Assoc* 18:361–364. <https://doi.org/10.1016/j.jamda.2016.12.086>
- Arosio B, Ostan R, Mari D et al (2017) Cognitive status in the oldest old and centenarians: a condition crucial for quality of life methodologically difficult to assess. *Mech Ageing Dev* 165(Pt B):185–194. <https://doi.org/10.1016/j.mad.2017.02.010>
- Gordon EH, Hubbard RE (2018) The Pathophysiology of Frailty: Why Sex Is So Important. *J Am Med Dir Assoc* 19:4–5. <https://doi.org/10.1016/j.jamda.2017.10.009>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.