Racial/Ethnic Disparities in HPV Vaccine Uptake Among a Sample of College Women

Chukwuemeka Okafor · Xingdi Hu · Robert L Cook

Received: 9 August 2014 / Revised: 20 October 2014 / Accepted: 3 November 2014 / Published online: 4 December 2014
© W. Montague Cobb-NMA Health Institute 2014

Abstract

Objective The aim of this study is to determine the association between racial/ethnic status and uptake and completion of the HPV vaccine series in college women.

Methods Participants were recruited from a large university in North Central Florida. Young women between 18 and 26 years of age who were currently enrolled in a college course comprised the study sample. Participants completed an anonymous online survey that assessed sociodemographic characteristics, sexual behaviors, gynecological healthcare utilization, and perception of risk to HPV-associated diseases. Multivariable analysis was conducted to determine the relationship between racial/ethnic status and HPV vaccination status. **Results** Of the 835 with complete data (51.0 % white, 16.5 % black, 13.8 % Hispanic, 8.3 % Asian, and 9.9 % other), 53 % had initiated (receipt of at least one dose) the three-dose HPV vaccine series. Of those who initiated, 70 % indicated that they had completed all three doses. In adjusted analysis, blacks were significantly less likely to report initiation [adjusted prevalence ratio (aPR) = 0.78; 95 % confidence interval (CI), 0.63, 0.97] and completion (aPR=0.64; 95 % CI: 0.48, 0.84) of the three dose HPV vaccine as compared to whites. Although completion rates were lower in all other racial/ethnic groups as compared to whites, these rates did not reach statistical significance.

Conclusions These findings are consistent with research from other types of settings and demonstrate lower initiation and completion rates of HPV vaccine among black women attending college as compared to their white counterparts. Additional research is needed to understand why black college women have low initiation and completion rates.

Keywords Human papillomavirus · HPV vaccine · Vaccination · College women · Racial/ethnic disparities

Introduction

Human papillomavirus (HPV) is the most prevalent sexually transmitted infection in the USA [1]. Persistent infection with high-risk HPV sub-types 16 and 18 is associated with 99.7 % of all cervical cancers [2]. Two HPV vaccines, the quadravalent HPV-4 (Gardasil, Merck, Whitehouse Station, NJ) [3] and the bivalent HPV-2 (Cervarix, GlaxoSmithKline, Philadelphia, PA) vaccine [4], are currently recommended by the Advisory Committee on Immunization Practices (ACIP) [5]. Current guidelines by the ACIP recommend routine vaccination of females aged 11 to 12 years and catch-up vaccination for females aged 13–26 years old [5]. Both HPV vaccines administered as a three-dose series have demonstrated between 70 and 95 % efficacy in reducing the rate of HPV-associated diseases [3, 4]. Despite the vaccine's benefits, rates of vaccine uptake among eligible women in the USA have been low to moderate. Rates of receipt of ≥ 1 , ≥ 2 , and ≥ 3 doses of the vaccine among adolescent girls aged 13 to 17 years were approximately 54, 43, and 33 %, respectively, in 2012 [6]. Whereas, among females aged 18–26 years, initiation rates (receipt of ≥ 1 dose) range from 43 to 45 % [8–10], while completion rates (receipt of three doses) among those who start the series range from 13 to 58 % [11–13].

Racial/ethnic minorities are disproportionately affected by the burden of HPV-associated cervical cancer incidence and mortality [14], yet racial/ethnic disparities in uptake of the HPV vaccine continue to be reported. In studies of non-college women, some studies show black and Hispanic adolescent women have similar or even higher rates of initiating HPV vaccination than their white counterparts; however, nearly all studies show that black women are less likely to

C. Okafor (✉) · X. Hu · R. L. Cook
Department of Epidemiology, College of Public Health & Health Professions, College of Medicine, University of Florida, 2004 Mowry Road, P.O Box 100231, Gainesville, FL 32610, USA
e-mail: emekaokafor@phhp.ufl.edu

complete the series as compared to white women [12, 15–19]. Several factors have been associated with HPV vaccine uptake and could help explain any racial/ethnic disparities. These include factors such as sociodemographic characteristics, cost or lack of health insurance, provider recommendation, negative attitudes toward vaccination (e.g., concerns about side effects), lack of knowledge about HPV infection, and a woman's perceived risk for HPV infection (e.g., a family or friends experience with a HPV-associated disease) [7, 12, 20–24]. Also, women who seek sexual health services such as Pap tests or prescription contraception (e.g., hormonal contraception) may be more proactive to their sexual health, engage in regular contact with healthcare services and therefore, more likely to be made aware of and receive HPV vaccination. Additionally, parental concerns that HPV vaccination may give young women a false sense of security and license for increased sexual activity may also influence HPV vaccination rates [7].

Female college students have demonstrated increased knowledge of HPV infection and vaccination as compared to a community sample of similar age [25] along with access to a college's health care center that provides HPV vaccination. Whether racial/ethnic disparity in HPV vaccination persists in a sample of female college students has not been thoroughly described. As college women are exposed to known risk factors for HPV infection such as early initiation of sexual activity and multiple sexual partners [20, 26], it is therefore important to continuously monitor the progress of HPV vaccine coverage and to delineate factors influencing vaccination. Therefore, the aims of this analysis were to (1) determine the rates of initiation and completion of the three-dose HPV vaccine series among a sample of college women, and (2) to determine the association of racial/ethnic status with vaccine initiation and completion.

Methods

Participant Recruitment

As part of a larger study related to oral HPV infection, participants for this study include a convenience sample of young women recruited from the campus of the University of Florida, a large US southeastern university. From April 2011 to September 2011, women were recruited within and around the university Student Health Care Center (SHCC). The sample included women seeking care at the center, as well as those just passing by, referred by friends, or responding to brochures or online advertisements about the study. Women were eligible to participate if they were at least 18 years of age and currently enrolled in a college course. Among women who inquired about the study and declined to participate, the primary reason for non-participation was lack of time. The study

was approved by the institutional review boards at the University of Florida and the Moffitt Cancer Research Center. All participants provided written informed consent and received a \$20 incentive.

Data Collection Procedure

Participants completed an anonymous 10–15 min web-based survey in a private room. Survey responses were entered directly by participants using a netbook computer.

Measures

The primary outcome measure “HPV vaccination status” was assessed with the following question: “How many HPV vaccine shots have you received in your lifetime (Cervarix or Gardasil)? (0, 1, 2, or 3). In this analysis, we defined HPV vaccine initiation as receipt of any ($n \geq 1$) HPV vaccine dose and vaccine completion as receipt of all three doses. Participants who reported “0” number of shots in the vaccine question were defined as unvaccinated. Participants with missing data or who answered “don't know” to the HPV vaccination question ($n=68$) were excluded from the analysis yielding a final study sample of 835 college women aged 18–26 years. There were no significant differences in the distribution of the sociodemographic variables, and all other covariates between those included in the final analytic sample and those that were excluded (data not shown).

Participant's sociodemographic characteristics, sexual behavior, gynecological health care utilization, and perceptions of risk for HPV-associated diseases were assessed for their association with HPV vaccination and for possible inclusion in multivariable analysis. A single question was used to ask participants to describe their racial/ethnic status. We categorized racial/ethnic status as white, black, Hispanic, Asian, and other. Age was dichotomized to 18–20 versus 21–26 years. Participants' relationship status was categorized as single, in a relationship, and other.

Sexual behavior was measured as number of vaginal sex partners in the past year, which was an ordinal variable categorized as none, 1–2, or 3+ partners. We included variables to assess gynecological health care utilization. Participants were asked whether they had ever received a Pap test, dichotomized as yes versus no, and to indicate through a check list which contraceptive methods they had used in the past year. Hormonal contraception which must be prescribed by a healthcare provider was dichotomized as yes versus no. The survey included items related to perceptions of risk of HPV-associated diseases in which participants were asked whether they ever had a physician or health care provider diagnosis of an abnormal Pap smear/colposcopy/biopsy, genital warts, or genital HPV, and whether they had a family history of head and/or neck cancer.

Data Analysis

We compared sociodemographic characteristics, sexual behaviors, gynecological health care utilization, and perceived risk of HPV infection by racial/ethnic status using chi-square or Fishers exact test. Similar methods were used to determine bivariate associations of factors associated with HPV vaccination. Covariates with *p* values at or below 0.2 were included in a multivariable analysis. We used log-binomial regression models to calculate crude and adjusted prevalence ratios with 95 % confidence intervals [27]. All data analyses were conducted using SAS version 9.3 (Cary, NC).

Results

Sample Characteristics

The final study sample included 835 college women aged 18–26 years. Approximately, half of the female college students in this sample self-identified as white (51 %), followed by black (16.5 %), Hispanic (13.8 %), Asian (8.3 %), and other (9.9 %). Table 1

displays the sociodemographic, sexual behavior, gynecological health care utilization, and perceptions of risk of HPV-associated disease by racial/ethnic status. The results show that there were several significant differences across racial/ethnic groups for several variables related to sociodemographic characteristics, sexual behavior, and gynecological health care utilization (Table 1).

HPV Vaccine Coverage

Table 2 displays the HPV vaccination status of the study sample by racial/ethnic groups. Overall, 53 % of the study participants had initiated the vaccine (receipt of at least one dose); of these, 70 % had received all three doses. The rates of initiating the HPV vaccine were similar between whites and Hispanics (57 and 57 %, respectively) and higher than all other racial groups (blacks: 42 %, Asians: 51 %, other: 47 %). Whites had the highest completion rate, whereas blacks had the lowest (whites: 78 %, blacks: 45 %, Hispanics: 66 %, Asians: 65 %, other: 69 %).

Table 1 Sociodemographic characteristics, sexual behaviors, sexual health care utilization, and perceptions of risk, by racial/ethnic status

Variable	White (<i>n</i> =425) (%)	Black (<i>n</i> =138) (%)	Hispanic (<i>n</i> =119) (%)	Asian (<i>n</i> =69) (%)	Other ^a (<i>n</i> =83) (%)	<i>p</i> Value
Total	51.0	16.5	13.8	8.3	9.9	
Age						
18–20	47.5	61.6	52.1	56.5	59.0	.032
21–26	52.5	38.4	47.9	43.5	41.0	
Relationship status						
Single	43.3	72.5	47.9	60.9	43.4	<.001
In a relationship	10.1	24.6	39.5	37.7	54.2	
Other ^b	46.6	2.9	12.6	1.5	2.4	
Vaginal sex partners—past year						
None in the past year	23.9	45.6	19.5	45.5	27.2	<.001
1–2	59.8	40.4	61.0	43.9	61.7	
3+	16.3	14.0	19.5	10.6	11.1	
Ever had a Pap test						
Yes	62.4	38.4	49.6	28.4	53.0	<.001
No	37.7	61.6	50.4	71.6	47.0	
Hormonal contraception—past year						
Yes	64.5	21.0	57.1	30.4	56.6	<.001
No	35.5	79.0	42.9	69.6	43.4	
F/H of head/neck cancer ^c						
Yes	6.4	3.3	3.9	13.5	4.5	.089
No	93.6	96.7	96.1	86.5	95.5	
History of HPV-associated disease ^d						
Yes	6.4	6.7	5.9	4.6	3.7	.873
No	93.6	93.3	94.1	95.4	96.3	

^a Includes American Indian/Alaskan Native, and Hawaiian/Pacific Islander

^b Includes cohabiting or living together and married

^c Family history of head and neck cancer

^d Includes abnormal Pap test, colposcopy, or genital warts

HPV Vaccine Initiation

Table 2 shows results from the unadjusted and adjusted prevalence ratios of HPV vaccine initiation. After adjusting for age and hormonal contraception use in the past year, the prevalence of HPV vaccine initiation was significantly lower among blacks (aPR=0.80, 95 % CI=0.65, 0.99; Table 2) than whites. Asides from Hispanics, who had similar prevalence rates as compared to whites (aPR=1.00, 95 % CI=0.85, 1.18), the prevalence for vaccine initiation among all other racial/ethnic groups were lower than their white counterparts, although these associations were not statistically significant.

HPV Vaccine Completion

In unadjusted analysis, blacks had significantly lower rates of HPV vaccine completion (aPR=0.58, 95 % CI=0.43, 0.77) than whites. All other racial/ethnic groups also had lower rates of completion than white women but not statistically significant. In adjusted analyses that controlled for hormonal contraception use in the past year, the prevalence of HPV vaccine completion remained significantly lower for blacks (aPR=0.64, 95 % CI=0.48, 0.84) than whites. All other racial/ethnic groups had lower but non-significant HPV vaccine completion rates as compared to white women.

Discussion

The findings from this analysis indicate that female college students in this sample had modest rates of HPV vaccine initiation (53 %), whereas, among those who initiated, completion rate was relatively high (70 %). The rates reported in this study are slightly higher than those reported in previous studies of women aged 18–26 years old which have ranged from 25 to 50 % for initiation [8, 22, 28–31] and 2 to 53 % for completion [8, 15, 25, 31, 32], with one study reporting a completion rate of 70 % [25].

In our multivariable models, we found younger-aged college women to report significantly higher prevalence of initiating HPV vaccination. This likely reflects increased healthcare provider recommendation for HPV vaccine as well as the eligibility of this age group for the vaccine for children program (VFC), which provides free vaccines for children and adolescents through 18 years of age who are uninsured. Our finding shows that past year hormonal contraception use is predictive of HPV vaccine initiation. This finding is not surprising, as women who seek sexual health promotion services may be more concerned about their sexual health and more likely to seek other sexual health preventive services including HPV vaccination. In this analysis, black women had significantly lower prevalence rates of initiating and completing the HPV vaccine, while no other racial/ethnic group had initiation or completion rates that were significantly different from whites. We are aware of only one study that has investigated racial/ethnic differences in HPV vaccine uptake in

Table 2 Unadjusted and adjusted prevalence ratios of HPV vaccine initiation and completion

Variables	Initiated (1 to 3 doses) ^a			Completed (3 doses) ^b		
	Vaccine initiation (%)	PR (95 % CI)	aPR (95 % CI)	Vaccine completion (%)	PR (95 % CI)	aPR (95 % CI)
Total	53.3			70.1		
Age						
18–20	59.8	1.30 (1.14, 1.48)***	1.34 (1.18, 1.52)***	68.7	0.95 (0.84, 1.08)	–
21–26	46.1	Ref	Ref	72.1	Ref	
Racial/ethnic						
White	57.7	Ref	Ref	77.8	Ref	Ref
Black	42.0	0.73 (0.59, 0.90)**	0.78 (0.63, 0.97)*	44.8	0.58 (0.43, 0.77)***	0.64 (0.48, 0.84)***
Hispanic	57.1	0.99 (0.83, 1.18)	1.00 (0.85, 1.18)	66.2	0.85 (0.71, 1.02)	0.86 (0.71, 1.03)
Asian	50.7	0.88 (0.69, 1.13)	0.96 (0.76, 1.20)	65.7	0.84 (0.66, 1.08)	0.87 (0.67, 1.13)
Other	47.0	0.82 (0.64, 1.04)	0.79 (0.63, 1.00)	69.2	0.88 (0.71, 1.11)	0.93 (0.75, 1.15)
Hormonal contraception—past year						
No	44.6	Ref	Ref	63.6	Ref	Ref
Yes	61.1	1.37 (1.20, 1.57)***	1.33 (1.14, 1.54)***	74.4	1.17 (1.02, 1.33)*	1.24 (0.91, 1.70)

* $p < .05$; ** $p < .01$; *** $p < .001$

^a Initiation rate defined as the percentage of college women with ≥ 1 dose of the HPV vaccine, among the study sample

^b Completion rate defined as the percentage of college women with three dose of the HPV vaccine, among those who had ≥ 1 HPV vaccine dose

college women. Bednarczyk et al. [33] found that black women were 33 % less likely to have initiated the HPV vaccination as compared to white women; however, data for completion rate among racial/ethnic group was not reported. Of note in the present analysis is the relatively higher number of single relationship status and lower number of past year vaginal sex partners among black women than other racial/ethnic groups. Therefore, it is possible that black women in this study may have perceived themselves to be at lower risk for an HPV infection and not likely to benefit from receiving the HPV vaccine. Additionally, black women in this analysis reported lower rates of past year hormonal contraception use and Pap tests as compared to most racial/ethnic groups. It is also possible that the black women in this sample may have been less proactive to sexual health, less inclined to seek sexual health preventive services, and thus have missed opportunities to be made aware of and receive HPV vaccine.

The findings from this analysis should be interpreted cautiously. Firstly, this was a cross-sectional analysis, and therefore we are unable to make any inferences about causality. Secondly, for our primary outcome measure, we relied on self-reported HPV vaccination status, and there is a potential for some reporting bias in our estimates of HPV vaccination uptake. In one study, adolescent girls aged 14–17 years were more likely to underestimate than to overestimate their HPV vaccine uptake [34], and therefore, there is a possibility that the actual HPV vaccination rate may be higher. Also, we did not account for other important factors including socioeconomic status and cultural health beliefs about HPV vaccination which could have influenced the propensity to get vaccinated. Thirdly, since our study was from a single university, our results may not be generalizable to all college students.

Despite these limitations, our analyses provide a 5-year update on rates of initiation and completion of the HPV vaccine in a sample of college women. Our findings suggest that racial/ethnic disparity in HPV vaccination persists even among female college students, specifically in black women. Although the completion rates are encouraging, our analysis suggest that more targeted intervention for HPV vaccination may be needed among minority women.

Acknowledgments This research was supported in part by a research grant from the Investigator-Initiated Studies Program of Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc. Funding was provided to the University of Florida with no specific restrictions. The opinions expressed are those of the authors and do not necessarily represent those of Merck Sharp & Dohme Corp. We appreciate the help and support from the staff at the University of Florida Student Health Care Center.

Conflict of Interest Chukwuemeka Okafor, Xingdi Hu, and Robert L. Cook declare that they have no conflict of interest.

Informed Consent All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of

1975, as revised in 2000 (5). Informed consent was obtained from all participants for being included in the study.

Animal Studies No animal studies were carried out by the authors for this article.

References

- Weinstock H, Berman S, Cates Jr W. Sexually transmitted diseases among American youth: incidence and prevalence estimates, 2000. *Perspect Sex Reprod Health*. 2004;36(1):6–10.
- Hariri S, Unger ER, Sternberg M, Dunne EF, Swan D, Patel S, et al. Prevalence of genital human papillomavirus among females in the United States, the National Health And Nutrition Examination Survey, 2003–2006. *J Infect Dis*. 2011;204(4):566–73.
- Garland SM, Hernandez-Avila M, Wheeler CM, Perez G, Harper DM, Leodolter S, et al. Quadrivalent vaccine against human papillomavirus to prevent anogenital diseases. *N Engl J Med*. 2007;356(19):1928–43.
- Paavonen J, Jenkins D, Bosch FX, Naud P, Salmerón J, Wheeler CM, et al. Efficacy of a prophylactic adjuvanted bivalent L1 virus-like-particle vaccine against infection with human papillomavirus types 16 and 18 in young women: an interim analysis of a phase III double-blind, randomised controlled trial. *Lancet*. 2007;369(9580):2161–70.
- Markowitz LE, Dunne EF, Saraiya M, Lawson HW, Chesson H, Unger ER. Quadrivalent human papillomavirus vaccine: recommendations of the advisory committee on immunization practices (ACIP). *MMWR Recomm Rep Morb Mortal Wkly Rep Recomm Rep Cent Dis Control*. 2007;56(RR-2):1–24.
- Human Papillomavirus Vaccination Coverage Among Adolescent Girls, 2007–2012, and Postlicensure Vaccine Safety Monitoring, 2006–2013 — United States [Internet]. [cited 2014 May 25]. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6229a4.htm#tab1>
- Jeudin P, Liveright E, del Carmen MG, Perkins RB. Race, ethnicity, and income factors impacting human papillomavirus vaccination rates. *Clin Ther*. 2014;36(1):24–37.
- Licht AS, Murphy JM, Hyland AJ, Fix BV, Hawk LW, Mahoney MC. Is use of the human papillomavirus vaccine among female college students related to human papillomavirus knowledge and risk perception? *Sex Transm Infect*. 2010;86(1):74–8.
- Lindley LL, Elkind JS, Landi SN, Brandt HM. Receipt of the human papillomavirus vaccine among female college students in the United States, 2009. *J Am Coll Health*. 2013;61(1):18–27.
- Bernat DH, Gerend MA, Chevallier K, Zimmerman MA, Bauermeister JA. Characteristics associated with initiation of the human papillomavirus vaccine among a national sample of male and female young adults. *J Adolesc Health*. 2013;53(5):630–6.
- Williams WW, Lu P-J, Saraiya M, Yankey D, Dorell C, Rodriguez JL, et al. Factors associated with human papillomavirus vaccination among young adult women in the United States. *Vaccine*. 2013;31(28):2937–46.
- Chou B, Krill LS, Horton BB, Barat CE, Trimble CL. Disparities in human papillomavirus vaccine completion among vaccine initiators. *Obstet Gynecol*. 2011;118(1):14–20.
- Tan W, Viera AJ, Rowe-West B, Grimshaw A, Quinn B, Walter EB. The HPV vaccine: are dosing recommendations being followed? *Vaccine*. 2011;29(14):2548–54.
- Howlader N, Noone A, Krapcho N. SEER Cancer Statistics Review, 1975–2008. Bethesda: National Cancer Institute; 2011.
- Cook RL, Zhang J, Mullins J, Kauf T, Brumback B, Steingraber H, et al. Factors associated with initiation and completion of human

- papillomavirus vaccine series among young women enrolled in Medicaid. *J Adolesc Health*. 2010;47(6):596–9.
16. Niccolai LM, Mehta NR, Hadler JL. Racial/ethnic and poverty disparities in human papillomavirus vaccination completion. *Am J Prev Med*. 2011;41(4):428–33.
 17. Dempsey A, Cohn L, Dalton V, Ruffin M. Worsening disparities in HPV vaccine utilization among 19–26 year old women. *Vaccine*. 2011;29(3):528–34.
 18. Gold R, Naleway AL, Jenkins LL, Riedlinger KK, Kurosky SK, Nystrom RJ, et al. Completion and timing of the three-dose human papillomavirus vaccine series among adolescents attending school-based health centers in Oregon. *Prev Med*. 2011;52(6):456–8.
 19. Dorell CG, Yankey D, Santibanez TA, Markowitz LE. Human papillomavirus vaccination series initiation and completion, 2008–2009. *Pediatrics*. 2011;128(5):830–9.
 20. Ratanasiripong NT. A review of human papillomavirus (HPV) infection and HPV vaccine-related attitudes and sexual behaviors among college-aged women in the United States. *J Am Coll Health J ACH*. 2012;60(6):461–70.
 21. Holman DM, Benard V, Roland KB, Watson M, Liddon N, Stokley S. Barriers to human papillomavirus vaccination among us adolescents: a systematic review of the literature. *JAMA Pediatr*. 2014;168(1):76–82.
 22. Scarinci IC, Garcés-Palacio IC, Partridge EE. An examination of acceptability of HPV vaccination among African American women and Latina immigrants. *J Womens Health* 2002. 2007;16(8):1224–33.
 23. Lau M, Lin H, Flores G. Factors associated with human papillomavirus vaccine-series initiation and healthcare provider recommendation in US adolescent females: 2007 National Survey of Children's Health. *Vaccine*. 2012;30(20):3112–8.
 24. Ylitalo KR, Lee H, Mehta NK. Health care provider recommendation, human papillomavirus vaccination, and race/ethnicity in the US National Immunization Survey. *Am J Public Health*. 2013;103(1):164–9.
 25. Wolwa M, Blavo C, Shah R, Fleisher JM, Espinal T. Cervical Cancer Knowledge and Prevention Among College Women. *J Community Health*. 2013 May 29.
 26. Rysavy MB, Kresowik JDK, Liu D, Mains L, Lessard M, Ryan GL. Human papillomavirus vaccination and sexual behavior in young women. *J Pediatr Adolesc Gynecol*. 2014;27(2):67–71.
 27. Spiegelman D, Hertzmark E. Easy SAS calculations for risk or prevalence ratios and differences. *Am J Epidemiol*. 2005;162(3):199–200.
 28. Marchand E, Glenn BA, Bastani R. HPV vaccination and sexual behavior in a community college sample. *J Community Health*. 2013;38(6):1010–4.
 29. Crosby R, Schoenberg N, Hopenhayn C, Moore G, Melhan W. Correlates of intent to be vaccinated against human papillomavirus: an exploratory study of college-aged women. *Sex Health*. 2007;4(1):71–3.
 30. Chao C, Velicer C, Slezak JM, Jacobsen SJ. Correlates for human papillomavirus vaccination of adolescent girls and young women in a managed care organization. *Am J Epidemiol*. 2010;171(3):357–67.
 31. Marchand E, Glenn BA, Bastani R. Low HPV vaccine coverage among female community college students. *J Community Health*. 2012;37(6):1136–44.
 32. Wetherill RR, Neal DJ, Fromme K. Parents, peers, and sexual values influence sexual behavior during the transition to college. *Arch Sex Behav*. 2010;39(3):682–94.
 33. Bednarczyk RA, Birkhead GS, Morse DL, Doleys H, McNutt L-A. Human papillomavirus vaccine uptake and barriers: association with perceived risk, actual risk and race/ethnicity among female students at a New York State university, 2010. *Vaccine*. 2011;29(17):3138–43.
 34. Stupiansky NW, Zimet GD, Cummings T, Fortenberry JD, Shew M. Accuracy of self-reported human papillomavirus vaccine receipt among adolescent girls and their mothers. *J Adolesc Health Off Publ Soc Adolesc Med*. 2012;50(1):103–5.