Correction to: Assessment and Treatment of Challenging Behavior Maintained by a Nonvocal Mands Function

Sara Jeglum^{1,2} • Jonathan D. Schmidt^{1,2} • Morgan Hallgren¹ • Jennifer Vetter^{1,2} • Amanda Goetzel¹

Published online: 17 November 2020

© Association for Behavior Analysis International 2020

Correction to: Behavior Analysis in Practice

<https://doi.org/10.1007/s40617-020-00486-z>

Figure 2 did not have a data path of open circles depicted. Below is the corrected Fig. 2, with a more accurate figure caption.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s40617-020-00486-z>

✉ Jonathan D. Schmidt
schmidtj@kennedykrieger.org

¹ Department of Behavioral Psychology, Kennedy Krieger Institute,
707 N. Broadway, Baltimore, MD 21205, USA

² Johns Hopkins University School of Medicine, Baltimore, MD, USA

Emesis, Head-Banging, Aggression, Disruption, and Biting Others

Fig. 2 Treatment evaluation. In the top panel, rates of challenging behavior are depicted by closed circles during treatment and by open squares during baseline. The bottom panel depicts rates of mands during baseline (open squares), during Mike's Way (open circles) in

treatment, and during Caregiver's Way (closed circles) in treatment. During the first treatment phase and the beginning of the second treatment phase, Mike's Way comprised the full 10-min session