



# A New Extension of the Topp–Leone-Family of Models with Applications to Real Data

Mustapha Muhammad<sup>1</sup> · Lixia Liu<sup>2</sup> · Badamasi Abba<sup>3,4</sup> ·  
Iisyaku Muhammad<sup>5</sup> · Mouna Bouchane<sup>6</sup> · Hexin Zhang<sup>7</sup> · Sani Musa<sup>8</sup>

Received: 13 February 2022 / Revised: 28 July 2022 / Accepted: 26 September 2022 /  
Published online: 18 October 2022

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

## Abstract

In this article, we proposed a new extension of the Topp–Leone family of distributions. Some important properties of the model are developed, such as quantile function, stochastic ordering, model series representation, moments, stress–strength reliability parameter, Renyi entropy, order statistics, and moment of residual life. A particular member called new extended Topp–Leone exponential (NETLE) is discussed. Maximum likelihood estimation (MLE), least-square estimation (LSE), and percentile estimation (PE) are used for the model parameter estimation. Simulation studies were conducted using NETLE to assess the MLE, LSE, and PE performance by examining their bias and mean square error (MSE), and the result was satisfactory. Finally, the applications of the NETLE to two real data sets are provided to illustrate the importance of the NETLG families in practice; the data sets consist of daily new deaths due to COVID-19 in California and New Jersey, USA. The new model outperformed many other existing Topp–Leone’s and exponential related distributions based on the real data illustrations.

**Keywords** Topp–Leone model · Moments · Renyi entropy · Stress–strength parameter · Maximum likelihood estimation · Least square estimation · Percentile estimation

**Mathematics Subject Classification** 62E05 · 62F10 · 62F12

---

✉ Lixia Liu  
liulixia@hebtu.edu.cn

Extended author information available on the last page of the article

## 1 Introduction

In life science and engineering studies such as medicine, biomedical sciences, biostatistics, communication, computer engineering, reliability, survival analysis, and life testing, statistical models play an indispensable role in studying the natural phenomena that occur in such fields of study. Probability models are used to model and characterize natural life phenomena. In probability studies, extending the classical probability models becomes necessary due to the natural events that occur and require higher dimensional data analysis and complex decisions. Whereas the traditional models cannot efficiently model such natural phenomena, especially when the failure rates are non-monotone. Thus, the practitioners proposed many ways of adding new parameter(s) to manipulate the existing traditional model and improve their quality and flexibility to provide higher accuracy so that the exploration of lifetime data can be assessed better. Data exploration in lifetime studies is one of the keys to analyze real-world, enabling us to measure, predict, and explain quantities of interest that arise in the fields of artificial intelligence, big data analysis, data mining, biomedical sciences, business studies, and communication with the aid of several powerful tools in probability and statistical concepts, programming, optimization, algorithms, and computational techniques; for more studies, one can see [1–4].

Topp Leone (TL) model was introduced in [5], the model is defined on finite support  $w \in (0, 1)$  with the cumulative distribution function (cdf),

$$H(w) = w^\alpha(2 - w)^\alpha, \quad \alpha > 0.$$

TL possesses a bathtub failure rate and has only one parameter; it has a closed-form of the cumulative distribution function, unlike beta distribution. This model has received comprehensive study and applications in sciences and social sciences such as biology, economics, ecology, etc. One can see [6] for the moments properties of the TL. [7] provides some stochastic ordering results and reliability studies of the TL. [8] discussed the Bayesian estimation of the TL under trimmed samples. [9] provided the recurrence relations for the moments of order statistics from the TL without any restrictions on the shape parameter and many relations when the shape parameter is an integer. [10] derived some explicit algebraic expressions for the single and product moments of order statistics from TL, and gave an identity about single moments of the order statistics. [11] proposed the use of maximum likelihood and uniformly minimum variance unbiased estimation procedures for the stress–strength reliability parameter from TL for the problem on both complete and left censored data, also considering the interval estimation of the reliability parameter. [12] develops a Bayesian estimation in the context of non-informative priors for the shape parameter of the mixture of TL distributions for a censored data set. Under the assumption of the known scale parameter of the TL, [13] provided Bayes and empirical Bayes estimates of the unknown parameter under non-informative and suitable conjugate priors and under the assumption of squared and linear-exponential error loss functions, also concluded that the proposed estimates are minimax and admissible. [14] constructed the Bayesian point and interval estimation for the shape parameter of the TL under lower  $k$ -record values. [15] discussed the moments of dual generalized order statistics from TL-weighted Weibull and its characterization.

Recently, [16] proposed a new inverted Topp–Leone distribution called the Kies inverted Topp–Leone with its applications to COVID-19 mortality rates in the United Kingdom and Canada. [17] introduced the odd Weibull inverted Topp–Leone distribution with its application to COVID-19 mortality rates in the United Kingdom and Canada. [18] introduced the type I half-logistic Nadarajah-Haghighi distribution and illustrated its good performance in fitting the COVID-19 mortality rates data from California, USA. [19] proposed exponentiated Gumbel–Weibull-Logistic and provided its application to Nigeria’s COVID-19 infections data. [20] estimated the daily recovery cases from COVID-19 in Egypt using power-odd generalized exponential Lomax distribution. [21] provided some statistical inferences based on the exponentiated exponential model that aided in assessing covid-19 cases in Kerala. [22] compared the performance of some lifetime models, including the Weibull model, using COVID-19 data from Pakistan. Among others.

TL has tractable close form properties, but it is not flexible enough to cover a wide range of practical applications, especially in lifetime data analysis. But, the demand for flexible lifetime models in practice is increasing due to the inability of the classical ones and rapid progress in applied statistical studies, biomedical sciences, engineering, computer sciences, reliability, econometrics, etc. Thus, researchers are encouraged to extend classical models to more flexible ones capable of modelling various failure rates that occur in lifetime analysis. TL families of distributions have been extended to a generator of distributions by various authors, and their positive impact has been discussed comprehensively in the literature. For example, [23–25] proposed various Topp–Leone-G (TLG) family, [26] introduced the generalized Topp–Leone-G (GTLG), [27] type II Topp–Leone-G (TIITLG), [28] type II generalized Topp–Leone-G (TIIGTG), [29] power Topp–Leone-G (PwTLG), [30] Type II power Topp–Leone-G (TIPTLG), [31] extended Topp–Leone -G (ETLG), G–fixed–Topp–Leone (GFTL) [32], exponentiated Generalized Topp Leone-G (EGTLG) [33], Marshall–Olkin Topp Leone-G (MOTLG) [34], Topp–Leone odd log-logistic-G (TLOLLG) [35], and Topp–Leone Marshall–Olkin-G (TLMOG) [36], among other.

In probability and statistical studies, the generators of probability models have significantly contributed to the literature regarding distribution theory and led to numerous useful tools in mathematical and statistical theory and practice. Here, we proposed a new extension of the Topp–Leone generator of distributions (NETLG) as an additional tool for statistical studies. This work aims to introduce another flexible generator of distributions from the TL that can accommodate various failure rates and can be used to model different kinds of skewed data. An additional parameter was employed to the usual TLG by applying a function involving exponential and natural logarithm. The additional power parameter allows the tuning of the model functions for better flexibility and provides some new statistical viewpoints for modelling data. Any valid baseline model can be chosen to propose a new flexible member of the NETLG model; the new model can provide more flexible shapes of the density and failure rates in comparison to parent distribution; the new generator’s special models have the capability to provide a better fit than some other existing alternative models. Moreover, different probability models serve different purposes and represent different data generation processes. In addition, we want to investigate some of the important mathematical and statistical properties regarding the new model to present a closed-form and convenient

**Table 1** Some Topp–Leone-G families

Name	CDF
Topp–Leone-G (TL-G) [23]	$H(x) = G^\alpha(x; \xi)(2 - G(x; \xi))^\alpha, \quad \alpha > 0.$
New Topp–Leone-G (NTL-G) [24]	$H(x) = \left(G^\theta(x; \xi) \left(2 - G^\theta(x; \xi)\right)\right)^\alpha, \quad \alpha, \theta > 0.$
Power Topp–Leone-G (PwTL-G) [29]	$H(x) = e^{\alpha\beta\left(1 - \frac{1}{G(x; \xi)}\right)} \left(2 - e^{\beta\left(1 - \frac{1}{G(x; \xi)}\right)}\right)^\alpha, \quad \alpha, \beta > 0.$
Extended Topp–Leone-G (ETL-G) [31]	$H(x) = G^\alpha(x; \xi)(2 - G(x; \xi))^{\alpha\beta}, \quad \alpha > 0, \beta \in \mathbb{R}.$

representation of the model properties with the aid of several mathematical techniques, computational algorithms, and computer packages for numerical assessment. Finally, two applications of the NETLG family to COVID 19 data are used to illustrate its importance in practice.

In Sect. 2, we derived the new model and discussed some important properties. In Sect. 3, maximum likelihood estimation (MLE), least-square estimation (SLE), and percentile estimation (PE) are proposed for the parameter estimation and assessed by simulation studies. Section 4 provides an application of the new model family for illustration. In Sect. 5, the conclusion.

## 2 New Extension of the Topp–Leone-G Models (NETLG) and Its Properties

Let  $G(x; \xi)$  be any valid baseline cumulative distribution function,  $x \in \mathbb{R}$ , and  $\xi$  is a vector of parameters, the cumulative distribution function (cdf) of some Topp–Leone-G families are given in Table 1:

In this work, we proposed the new extended Topp–Leone generator of distributions (NETLG) due to the idea in [37]. The perspective of this model derivation is innovative and will have broader academic value. No doubt, the studies related to this model will provide tools for extending several mathematical results in probability and mathematical statistics. The additional parameter provides more flexibility to the new model than other existing TL families. It is a fact that different distributions serve different purposes and represent different data generation processes; thus, our new model provided a new means of data generating processes and a Monte Carlo simulation process. The cumulative distribution function of the new extended Topp–Leone generator of distributions (NETLG) is given by

$$F(x) = e^{-\alpha(-\log G(x; \xi))^\beta} \left(2 - e^{-(-\log G(x; \xi))^\beta}\right)^\alpha, \quad \alpha, \beta > 0. \quad (1)$$

If  $\beta = 1$  we have the Topp–Leone-G model [23]. The corresponding probability density function (pdf) is given by

$$f(x) = 2\alpha\beta \frac{g(x; \xi)}{G(x; \xi)} (-\log G(x; \xi))^{\beta-1} e^{-\alpha(-\log G(x; \xi))^\beta} \left(1 - e^{-(-\log G(x; \xi))^\beta}\right) \left(2 - e^{-(-\log G(x; \xi))^\beta}\right)^{\alpha-1}, \quad (2)$$

where  $g(x; \xi)$  is the corresponding pdf of the baseline cdf  $G(x; \xi)$ . The survival function and hazard rate function (hrf) of the NETLG for  $\beta, \alpha > 0$  are given by

$$s(x) = 1 - e^{-\alpha(-\log G(x; \xi))^\beta} \left(2 - e^{-(-\log G(x; \xi))^\beta}\right)^\alpha, \quad (3)$$

and

$$h(x) = 2\alpha\beta \frac{g(x; \xi)}{G(x; \xi)} \frac{(-\log G(x; \xi))^{\beta-1} e^{-\alpha(-\log G(x; \xi))^\beta} \left(1 - e^{-(-\log G(x; \xi))^\beta}\right) \left(2 - e^{-(-\log G(x; \xi))^\beta}\right)^{\alpha-1}}{1 - e^{-\alpha(-\log G(x; \xi))^\beta} \left(2 - e^{-(-\log G(x; \xi))^\beta}\right)^\alpha}, \quad (4)$$

respectively.

The asymptotic properties of the NETLG model are discussed below.

**Lemma 1** *Let  $X \sim \text{NETLG}$ ; then we have the following asymptotic.*

$$\begin{aligned} F(x) &\sim 2e^{-\alpha(-\log G(x; \xi))^\beta} \text{ as } G \rightarrow 0, & (5) \\ s(x) &\sim \alpha \left(1 - e^{-(-\log G(x; \xi))^\beta}\right)^2 \sim \alpha (-\log G(x; \xi))^{2\beta} \text{ as } G \rightarrow 1, \\ f(x) &\sim 2\alpha\beta g(x; \xi) (-\log G(x; \xi))^{\beta-1} \left(1 - e^{-(-\log G(x; \xi))^\beta}\right) \text{ as } G \rightarrow 1, \\ h(x) &\sim \frac{2\beta g(x; \xi) (-\log G(x; \xi))^{\beta-1}}{\left(1 - e^{-(-\log G(x; \xi))^\beta}\right)} \text{ as } G \rightarrow 1, \\ f(x) &\sim 4\alpha\beta \frac{g(x; \xi)}{G(x; \xi)} (-\log G(x; \xi))^{\beta-1} e^{-\alpha(-\log G(x; \xi))^\beta} \text{ as } G \rightarrow 0. & (6) \end{aligned}$$

The quantile of distribution has many uses in theoretical studies and statistics applications. Quantile function serves as a tool for parameter estimation and simulation. The quantile function of the NETLG can be derived as

$$Q(u) = G^{-1} \left( e^{-\left[-\log\left(1 - \sqrt{1 - u^{\frac{1}{\alpha}}}\right)\right]^{1/\beta}} \right), \quad 0 \leq u \leq 1. \quad (7)$$

In particular,  $Q(\frac{1}{2})$  is the median. The skewness and kurtosis of the NETLG can be discussed by the Bowley's skewness (B), and Moor's kurtosis (M) defined respectively as

$$B = \frac{Q(\frac{3}{4}) - 2Q(\frac{2}{4}) + Q(\frac{1}{4})}{Q(\frac{3}{4}) - Q(\frac{1}{4})}$$

and

$$M = \frac{Q(\frac{7}{8}) - Q(\frac{5}{8}) + Q(\frac{3}{8}) - Q(\frac{1}{8})}{Q(\frac{6}{8}) - Q(\frac{2}{8})}.$$

## 2.1 Stochastic Ordering

Stochastic ordering is another aspect of probability theory that allows us to discuss some of the relative characters of distributions. There are various means by which we might say that a random variable  $X$  is smaller than a random variable  $Y$ . However, in stochastic ordering, we say that  $X$  is stochastically smaller than  $Y$  denoted as  $X \prec_{st} Y$ . There are many other stochastic orders such as likelihood, hazard, etc., for details one can see [38] among others. Here, we obtained the likelihood ordering result for NETLG under some possible conditions. Let a random variable  $X_1$  with pdf  $f_1(x)$  and  $X_2$  having pdf  $f_2(x)$ , then  $X_1$  is said to be smaller than  $X_2$  in the likelihood ratio order (denoted  $X_1 \prec_{lr} X_2$ ) if  $f_1(x)/f_2(x)$  is decreasing in  $x$ .

**Proposition 1** *Let  $X_1$  be a random variable having  $NETLG_1(\alpha_1, \beta, \xi)$  and random variable  $X_2$  having  $NETLG_2(\alpha_2, \beta, \xi)$ , then,  $X_1 \prec_{lr} X_2$  if  $\alpha_1 \leq \alpha_2$ .*

**Proof** We show that  $f_1(x)/f_2(x)$  is decreasing.

$$\frac{f_1(x; \alpha_1, \beta, \xi)}{f_2(x; \alpha_2, \beta, \xi)} = \frac{\alpha_1}{\alpha_2} e^{-(\alpha_1 - \alpha_2)(-\log G(x; \xi))^\beta} \left(2 - e^{-(\log G(x; \xi))^\beta}\right)^{\alpha_1 - \alpha_2},$$

and

$$\begin{aligned} \frac{\partial}{\partial x} \frac{f_1(x; \alpha_1, \beta, \xi)}{f_2(x; \alpha_2, \beta, \xi)} &= 2 \frac{\alpha_1}{\alpha_2} \beta (\alpha_1 - \alpha_2) \frac{g(x; \xi)}{G(x; \xi)} (-\log G(x; \xi))^{\beta-1} e^{-(\alpha_1 - \alpha_2)(-\log G(x; \xi))^\beta} \\ &\quad \times \left(1 - e^{-(\log G(x; \xi))^\beta}\right) \left(2 - e^{-(\log G(x; \xi))^\beta}\right)^{\alpha_1 - \alpha_2 - 1}, \end{aligned}$$

thus,  $\frac{\partial}{\partial x} \frac{f_1(x; \alpha_1, \beta, \xi)}{f_2(x; \alpha_2, \beta, \xi)} \leq 0$  if  $\alpha_1 \leq \alpha_2$ . □

## 2.2 Series Representation of the pdf

The series representation of the NETLG could make it easier for us to compute some properties of the NETLG. But we need the following expansions. For  $|z| \leq 1$  and  $b > 0$  real and non integer,

$$(1 - z)^{b-1} = \sum_{i=0}^{\infty} \frac{(-1)^i \Gamma(b)}{i! \Gamma(b-i)} z^i. \quad (8)$$

Also, for a power series to the power of  $n \in \mathbb{N}$  we need:

**Lemma 2** ([39]) *For a given power series of the form  $\sum_{k=0}^{\infty} a_k x^k$ , let  $n$  be a positive integer, then,*

$$\left(\sum_{k=0}^{\infty} a_k x^k\right)^n = \sum_{k=0}^{\infty} c_k x^k,$$

where  $c_0 = a_0^n$ ,  $c_m = \frac{1}{m a_0} \sum_{k=1}^m (kn - m + k) a_k c_{m-k}$  for  $m \geq 1$ .

Form the (2); the following expressions can be presented in a series form by using (8) as

$$\begin{aligned} & e^{-\alpha(-\log G(x;\xi))^\beta} \left(1 - e^{-(-\log G(x;\xi))^\beta}\right) \left(2 - e^{-(-\log G(x;\xi))^\beta}\right)^{\alpha-1} \\ &= \sum_{i,k=0}^{\infty} \sum_{j=0}^{i+1} \omega_{i,j,k} (-\log G(x;\xi))^{\beta k}, \end{aligned} \tag{9}$$

where,  $\omega_{i,j,k} = \frac{(-1)^{j+k} \alpha^k (j+1)^k \Gamma(\alpha) \binom{i+1}{j}}{i! k! \Gamma(\alpha-1)}$ , therefore, (2) can be presented as

$$\begin{aligned} f(x) &= \sum_{i,k=0}^{\infty} \sum_{j=0}^{i+1} \omega_{i,j,k}^* \frac{g(x;\xi)}{G(x;\xi)} (-\log G(x;\xi))^{\beta(k+1)-1} \\ &= \sum_{i,k=0}^{\infty} \sum_{j=0}^{i+1} \omega_{i,j,k}^* \tau_k(x; \beta, \xi), \end{aligned} \tag{10}$$

where,  $\omega_{i,j,k}^* = \frac{2\alpha\beta(-1)^{j+k} \alpha^k (j+1)^k \Gamma(\alpha) \binom{i+1}{j}}{i! k! \Gamma(\alpha-1)}$  and  $\tau_k(x; \beta, \xi) = \frac{g(x;\xi)}{G(x;\xi)} (-\log G(x;\xi))^{\beta(k+1)-1}$ . Further, for an integer  $\beta(k+1) - 1$  we can proceed and simplify  $(-\log G(x;\xi))^{\beta(k+1)-1}$  in (10) by some algebra as

$$\begin{aligned} (-\log G(x;\xi))^{\beta(k+1)-1} &= \sum_{l=0}^{\beta(k+1)-1} \binom{\beta(k+1)-1}{l} (-\log G(x;\xi) - 1)^l, \\ &= 1 + \sum_{l=1}^{\beta(k+1)-1} \binom{\beta(k+1)-1}{l} \left(\sum_{n=0}^{\infty} w_n (G(x;\xi) - 1)^n\right)^l, \end{aligned}$$

where  $w_0 = -1$ , and  $w_n = \frac{(-1)^n}{n}$ , therefore, we can apply the lemma 2 in above to get

$$(-\log G(x;\xi))^{\beta(k+1)-1} = 1 + \sum_{l=1}^{\beta(k+1)-1} \binom{\beta(k+1)-1}{l} \sum_{n=0}^{\infty} c_n (G(x;\xi) - 1)^n,$$

where  $c_0 = w_0^n$ , and  $c_m = \frac{1}{mw_0} \sum_{n=1}^m (ln - m + n)w_n c_{m-n}$ , hence,

$$(-\log G(x; \xi))^{\beta(k+1)-1} = 1 + \sum_{l=1}^{\beta(k+1)-1} \sum_{n=0}^{\infty} d_{k,v,n} G^v(x; \xi), \tag{11}$$

where  $d_{k,v,n} = \sum_{v=0}^n \binom{\beta(k+1)-1}{l} \binom{n}{v} c_n (-1)^{n-1}$ , thus, by putting (11) in (10) we have

$$f(x) = \sum_{i,k=0}^{\infty} \sum_{j=0}^{i+1} \omega_{i,j,k}^* \frac{g(x; \xi)}{G(x; \xi)} + \sum_{i,k,n=0}^{\infty} \sum_{l=1}^{\beta(k+1)-1} \sum_{j=0}^{i+1} \omega_{i,j,k}^* d_{k,v,n} g(x; \xi) G^{v-1}(x; \xi), \tag{12}$$

### 2.3 Moments

Let  $X$  follow the NETLG distributions, the  $r^{th}$  moment of  $X$  can be obtained from  $\mu_r = E[X^r] = \int_{-\infty}^{\infty} x^r f(x) dx$ . The moments can be computed directly from the (10) as

$$\mu_r = \sum_{i,k=0}^{\infty} \sum_{j=0}^{i+1} \omega_{i,j,k}^* \int_{-\infty}^{\infty} x^r \tau_k(x; \beta, \xi) dx, \tag{13}$$

where  $\omega_{i,j,k}^*$  and  $\tau_k(x; \beta, \xi)$  are given previously in (10). In other way, we can compute the moment from (12) given by

$$\mu_r = \sum_{i,k=0}^{\infty} \sum_{j=0}^{i+1} \omega_{i,j,k}^* \int_{-\infty}^{\infty} x^r \frac{g(x; \xi)}{G(x; \xi)} dx + \sum_{i,k,n=0}^{\infty} \sum_{l=1}^{\beta(k+1)-1} \sum_{j=0}^{i+1} v^{-1} \omega_{i,j,k}^* d_{k,v,n} E_v[X^r],$$

where  $\omega_{i,j,k}^*$  given in (12),  $d_{k,v,n}$  in (11), and  $E_v[X^r]$  is the  $r^{th}$  moment of the exponentiated baseline distribution  $G^v(x; \xi)$  distribution.

Next, we consider the incomplete moments used in computation the mean deviations, income inequalities, and moments of residual life. The incomplete moment of  $X$  defined as  $I_r(t) = \int_{-\infty}^t x^r f(x) dx$  from (10) can take the following form

$$I_r(t) = \sum_{i,k=0}^{\infty} \sum_{j=0}^{i+1} \omega_{i,j,k}^* \int_{-\infty}^t x^r \tau_k(x; \xi) dx, \tag{14}$$

where  $\tau_k(x; \xi)$  is given in the equation (10). The first incomplete moment  $I_1(t)$ , i.e when  $r = 1$ , is a tool for computing the mean deviations about the mean  $\delta_1(X)$  and the mean deviations about median  $\delta_2(X)$  defined by

$$\delta_1(X) = 2\mu_1 F(\mu_1) - 2I_1(\mu_1) \quad \text{and} \quad \delta_2(X) = \mu_1 - 2I_1(M),$$

where  $\mu_1$  is the first moment of  $X$  i.e,  $\mu_1$ ;  $M = Q(0.5)$  from (7); and  $F(\cdot)$  is the cdf in (1).



The Lorenz and Bonferroni curves are computed using  $I_1(t)$ . For a random variable  $X$  and a given probability  $p$ , the Lorenz and Bonferroni curves are defined by  $L(p) = (\mu_1)^{-1}I_1(q)$  and  $B(q) = (p\mu_1)^{-1}I_1(q)$  respectively, and  $q = Q(p)$  from (7).

Moreover,  $I_1(t)$  can be used to compute the two mean of residual life, i.e. the mean residual life defined by  $M(t) = E(X - t|X > t)$ , or,  $M(t) = \int_0^\infty \frac{s(x+t)}{s(t)} dx$ , where  $s(t)$  is the survival function of  $X$ , and the mean reverse residual life defined by  $m(t) = E(t - X|X \leq t)$ , or  $m(t) = \int_0^t \frac{F(x)}{F(t)} dx$ , where  $F(\cdot)$  is the cdf of  $X$ . These measures are applied in the determination of the distributions of the extreme values.

## 2.4 Stress–Strength Reliability

In practice, a good design of a system is so that the system can resist the assumed stress to be applied. Assume that a component possesses stress  $X$  and is subjected to a strength  $Y$ , then parameter  $R = P(Y < X)$  discusses the system performance. It is called the stress–strength parameter in reliability studies. The system will fail when the applied stress is higher than the assumed system strength. We can find some applications of  $R$  in various fields in [40, chap. 7].  $R$  has been discussed by many researchers in the literature through various perspectives. For example, let  $X$  and  $Y$  be an independent random variables, then: normal distribution (N) [41, 42], Weibull (W) [43–45], exponential (E) [46], generalized exponential (GE) [47], beta-Erlang truncated exponential (BETE) [48], Poisson-odd generalized exponential family (POGE) [49], generalized logistic (GL) [50], Poisson-generalized half logistic (PGHL) [51], generalized exponential Poisson (GEP) [52], Poisson half logistic (PHL) [53], exponentiated sine Weibull (ESW) [54], extended cosine Weibull (ECSW) [55], among others.

Let  $X$  and  $Y$  be independent random variables, having the density  $f_1(x; \alpha_1, \beta, \xi)$  and  $Y$  with cdf  $F_2(y; \alpha_2, \beta, \xi)$  from NETLG, then, the stress–strength reliability parameter is  $R = P(Y < X) = \int_{-\infty}^\infty f_1(x) F_2(x) dx$ . Therefore,

$$R = \int_{-\infty}^\infty 2\alpha_1\beta \frac{g(x; \xi)}{G(x; \xi)} (-\log G(x; \xi))^{\beta-1} e^{-(\alpha_1+\alpha_2)(-\log G(x; \xi))^\beta} \\ \times \left(1 - e^{-(\log G(x; \xi))^\beta}\right) \left(2 - e^{-(\log G(x; \xi))^\beta}\right)^{\alpha_1+\alpha_2-1} dx,$$

hence,

$$R = \frac{\alpha_1}{\alpha_1 + \alpha_2} \int_{-\infty}^\infty f(x; \alpha_1 + \alpha_2, \beta, \xi) dx = \frac{\alpha_1}{\alpha_1 + \alpha_2}.$$

## 2.5 Entropy

Entropy is the degree of disorder or randomness in a system. Here, we computed the Renyi entropy of the NETLG families defined by  $Re(\rho) = (1 - \rho)^{-1} \log \left[ \int_{-\infty}^\infty f^\rho(x) dx \right]$ , where  $\rho > 0$  and  $\rho \neq 1$ . We started by simplify the expression of  $f^\rho(x)$  as:

$$f^\rho(x) = 2^\rho \alpha^\rho \beta^\rho \frac{g^\rho(x; \xi)}{G^\rho(x; \xi)} (-\log G(x; \xi))^{\rho(\beta-1)} e^{-\alpha\rho(-\log G(x; \xi))^\beta} \times \left(1 - e^{-(\log G(x; \xi))^\beta}\right)^\rho \left(2 - e^{-(\log G(x; \xi))^\beta}\right)^{\rho(\alpha-1)}, \tag{15}$$

we can simplify the following expression obtained from the above equation as

$$e^{-\alpha\rho(-\log G(x; \xi))^\beta} \left(1 - e^{-(\log G(x; \xi))^\beta}\right)^\rho \left(2 - e^{-(\log G(x; \xi))^\beta}\right)^{\rho(\alpha-1)} = \sum_{i, j, k=0}^\infty \phi_{i, j, k} (-\log G(x; \xi))^{\beta k}, \tag{16}$$

where,  $\phi_{i, j, k} = \binom{\rho(\alpha-1)}{i} \binom{i+\rho}{j} \frac{(-1)^{j+k} (\alpha\rho+j)^k}{k!}$ , thus, by substituting the above series in (15) we get

$$f^\rho(x) = 2^\rho \alpha^\rho \beta^\rho \sum_{i, j, k=0}^\infty \phi_{i, j, k} \frac{g^\rho(x; \xi)}{G^\rho(x; \xi)} (-\log G(x; \xi))^{\beta(k+\rho)-\rho},$$

hence, the Renyi entropy of the NETLG families can be obtained from

$$Re(\rho) = (1 - \rho)^{-1} \log \left[ 2^\rho \alpha^\rho \beta^\rho \sum_{i, j, k=0}^\infty \phi_{i, j, k} \int_{-\infty}^\infty \frac{g^\rho(x; \xi)}{G^\rho(x; \xi)} (-\log G(x; \xi))^{\beta(k+\rho)-\rho} dx \right].$$

### 2.6 Order Statistics

Let  $X_1, X_2, \dots, X_n, n \geq 1$ , be an ordered sample obtained from NETLG families with cdf  $F(x)$  and pdf  $f(x)$ , then, the density function of the  $i^{th}$ -order statistics is represented by  $f_{i:n}(x)$  and define by

$$f_{i:n}(x) = \frac{f(x)}{B(i, n - i + 1)} \sum_{j=0}^{n-i} \binom{n-i}{j} (-1)^j F^{i+j-1}(x).$$

Simplifying  $F^{i+j-1}(x)$  in above, we have

$$F^{i+j-1}(x) = e^{-\alpha(i+j-1)(-\log G(x; \xi))^\beta} \left(2 - e^{-(\log G(x; \xi))^\beta}\right)^{\alpha(i+j-1)},$$

thus, the density function of the  $i^{th}$ -order statistics be comes the mixture of the NETLG( $\alpha(i + j), \beta, \xi$ ) as

$$f_{i:n}(x) = \frac{1}{(i + j) B(i, n - i + 1)} \sum_{j=0}^{n-i} \binom{n-i}{j} (-1)^j f(x; \alpha(i + j), \beta, \xi). \tag{17}$$

The  $r^{th}$  moments of the  $i^{th}$ –order statistics can be computed from (17) by considering (13) as

$$E[X_{i:n}^r] = \frac{1}{(i+j)B(i, n-i+1)} \sum_{j=0}^{n-i} \binom{n-i}{j} (-1)^j E[X^r]_{f_{[\alpha(i+j), \beta, \xi]}}$$

where  $E[X^r]_{f_{[\alpha(i+j), \beta, \xi]}}$  is the expectation of NETLG families with respect to the density  $f(x; \alpha(i+j), \beta, \xi)$ .

The asymptotic distributions for the extreme order statistics  $X_{1:n}$  and  $X_{n:n}$  from NETLG families can be discussed according to the details in [56, chap. 8], among others.

## 2.7 A Special Member of the NETLG Families

A special member of the NETLG is derived and discussed in this subsection, namely the new extended Topp–Leone exponential (NETLE) distributions.

### 2.7.1 New Extended Topp–Leone Exponential (NETLE)

Let the baseline density in (2) be exponential distribution with parameter  $\lambda > 0$ , having cdf and pdf given by  $G(x; \lambda) = 1 - e^{-\lambda x}$  and  $g(x; \lambda) = \lambda e^{-\lambda x}$  respectively,  $x > 0$ . The density ( $f(x)$ ) and the hazard ( $h(x)$ ) functions of the NETLE are given by

$$f(x) = \frac{2\alpha\beta\lambda e^{-\lambda x}}{(1 - e^{-\lambda x})} (-\log(1 - e^{-\lambda x}))^{\beta-1} e^{-\alpha(-\log(1 - e^{-\lambda x}))^\beta} \left(1 - e^{-(-\log(1 - e^{-x}))^\beta}\right) \times \left(2 - e^{-(-\log(1 - e^{-x}))^\beta}\right)^{\alpha-1}, \quad (18)$$

$$h(x) = \frac{2\alpha\beta\lambda e^{-\lambda x}}{(1 - e^{-\lambda x})} (-\log(1 - e^{-\lambda x}))^{\beta-1} e^{-\alpha(-\log(1 - e^{-\lambda x}))^\beta} \times \frac{\left(1 - e^{-(-\log(1 - e^{-x}))^\beta}\right) \left(2 - e^{-(-\log(1 - e^{-x}))^\beta}\right)^{\alpha-1}}{1 - e^{-(-\log(1 - e^{-\lambda x}))^\beta} \left(2 - e^{-(-\log(1 - e^{-x}))^\beta}\right)^\alpha}.$$

Figures 1 and 2 show the plots of the density and hazard functions of the NETLU for some parameter values.

Let be a random variable  $X$  follow NETLE, then the quantile function of  $X$  is given by

$$Q_{NETLE}(u) = -\lambda^{-1} \log(1 - q(u, \alpha, \beta)), \quad 0 \leq u \leq 1,$$




Fig. 1 Plots of the density and hazard rate function of the NETLE distribution




Fig. 2 Plots of the hazard rate function of the NETLE distribution




Fig. 3 Plots of the B-skewness and M-kurtosis of the NETLE distribution

where  $q(u, \alpha, \beta) = e^{-\left[-\log\left(1-\sqrt{1-u^{\frac{1}{\alpha}}}\right)\right]^{1/\beta}}$ .

Figure 3 illustrated that the B-skewness and M-kurtosis of the NETLE; notice that both the B skewness and M kurtosis are independent of  $\lambda$ . The skewness decreases in  $\alpha$  and unimodal in  $\beta$ , while the kurtosis decreases in  $\alpha$  and increases in  $\beta$ .

**Proposition 2** Let  $X \sim \text{NETLE}$  with pdf in (18), then,

1. For a very large  $t > 0$ , i.e, as  $t \rightarrow \infty$  the asymptotic of the mean residual life is given by

$$M(t) \sim \frac{1}{2\lambda\beta}.$$

2. For a very small  $t > 0$ , i.e, as  $t \rightarrow 0$  the asymptotic of the mean reverse residual life for  $\beta = 1$  is

$$m(t) \sim \frac{t}{\alpha + 1}.$$

**Proof** 1. The asymptotic of the survival function of NETLE can be obtained from (6) as  $s(x) \sim \alpha (-\log(1 - e^{-\lambda x}))^{2\beta} \sim e^{-\lambda x}$  as  $x \rightarrow \infty$ , therefore, as  $t \rightarrow \infty$  we have  $M = \int_0^\infty \frac{s(x+t)}{s(t)} dx \sim \int_0^\infty e^{-2\lambda\beta x} dx = \frac{1}{2\lambda\beta}$ .

2. As  $x \rightarrow 0$  the cdf of NETLE from (5) for  $\beta = 1$  become:  $F(x) \sim 2(1 - e^{-\lambda x})^\alpha \sim 2\lambda^\alpha x^\alpha$ , therefore, as  $t \rightarrow 0$ ,  $m(x) = \int_0^t \frac{F(x)}{F(t)} dx \sim t^{-\alpha} \int_0^t x^\alpha dx = \frac{t}{\alpha+1}$ . □

**Proposition 3** Let  $X_1 \leq X_2 \leq \dots \leq X_n$ , be from NETLE with pdf in (18), let  $B_n = (X_{n:n} - a_n)/b_n$ , then,  $B_n \xrightarrow{d} B$  implies that

$$\lim_{n \rightarrow \infty} P(B_n \leq x) = G(x) = e^{-e^{-x}},$$

for every valid  $x \in \mathfrak{R}$  of  $G(x)$ .  $a_n = F^{-1}(1 - n^{-1})$  and  $b_n = E[X - a_n | X > a_n]$  from theorem 8.3.4 of [56].

**Proof** From theorem 8.3.2 of [56], we used  $\lim_{t \rightarrow \infty} \frac{s(t+x E(X-t|X>t))}{s(t)}$ , based on the theorem 2 number 1.,  $M(t) \sim \frac{1}{2\lambda\beta}$  as  $t \rightarrow \infty$ , thus,  $\lim_{t \rightarrow \infty} \frac{s(t+x E(X-t|X>t))}{s(t)} \sim \lim_{t \rightarrow \infty} \frac{e^{-2\lambda\beta(t + \frac{x}{2\lambda\beta})}}{e^{-\lambda\beta t}} = e^{-x}$ . □

**Proposition 4** Let  $X_1 \leq X_2 \leq \dots \leq X_n$ , be from NETLE with pdf in (18), let  $B_n^* = (X_{1:n} - a_n^*)/b_n^*$ , then, for  $\beta = 1$ ,  $B_n^* \xrightarrow{d} B^*$  is implies that

$$\lim_{n \rightarrow \infty} P(B_n^* \leq x) = G^*(x; \theta) = 1 - e^{-e^x},$$

for every valid  $x \in \mathfrak{R}^+$  of  $G^*(x)$ , thus,  $a_n^* = F^{-1}(\frac{1}{n})$  and  $b_n^* = E(a_n^* - X | X \leq a_n^*)$  from the theorem 8.3.6 of [56].

**Proof** From theorem 8.3.6 of [56] we can consider  $\lim_{t \rightarrow 0} \frac{F(t+x E(t-X|X \leq t))}{F(t)}$ . Based on the Theorem 2 number 2., the asymptotic of  $E(t - X | X \leq t)$  for  $\beta = 1$  as  $\lim_{t \rightarrow 0}$  is  $m(t) \sim \frac{t}{\alpha+1}$ , thus,  $\lim_{t \rightarrow 0} \frac{F(t+x E(t-X|X \leq t))}{F(t)} \sim \lim_{t \rightarrow 0} \frac{2\lambda^\alpha (t + \frac{tx}{\alpha+1})^\alpha}{2\lambda^\alpha t^\alpha} = (1 + \frac{x}{\alpha+1})^\alpha \rightarrow e^x$  as  $\alpha \rightarrow \infty$ . □

### 3 Model Parameter Estimation

The parameters of the NETLG are estimated using the method of maximum likelihood estimation (MLE), least-square estimation (SLE), and percentile estimation (PE). The performance of the techniques is examined by simulation studies using NETLE.

### 3.1 Maximum Likelihood Estimation (MLE)

Let  $X_1, \dots, X_n$  be a random sample of size  $n$  from the NETLG. Let  $\Theta = (\alpha, \beta, \xi)^T$  be a vector of parameters, with the MLEs as  $\hat{\Theta} = (\hat{\alpha}, \hat{\beta}, \hat{\xi})^T$ . The  $\hat{\Theta}$  can be computed by the maximization of the log-likelihood function ( $\ell(\Theta)$ ) in Eq. (19).

$$\begin{aligned} \ell(\theta) &= n \log 2 + n \log \alpha + n \log \beta + \sum_{i=1}^n \log g(x_i; \xi) - \sum_{i=1}^n \log G(x_i; \xi) \\ &+ (\beta - 1) \sum_{i=1}^n \log (-\log G(x_i; \xi)) - \alpha \sum_{i=1}^n (-\log G(x_i; \xi))^\beta \\ &+ \sum_{i=1}^n \log \left(1 - e^{-(\log G(x_i; \xi))^\beta}\right) + (\alpha - 1) \sum_{i=1}^n \log \left(2 - e^{-(\log G(x_i; \xi))^\beta}\right) \end{aligned} \tag{19}$$

In other way, by solving the nonlinear system given below in (20) to (22).

$$\frac{\partial \ell}{\partial \alpha} = \frac{n}{\alpha} - \sum_{i=1}^n (-\log G(x_i; \xi))^\beta + \alpha \sum_{i=1}^n \log \left(2 - e^{-(\log G(x_i; \xi))^\beta}\right) \tag{20}$$

$$\begin{aligned} \frac{\partial \ell}{\partial \beta} &= \frac{n}{\beta} + \sum_{i=1}^n \log (-\log G(x_i; \xi)) - \alpha \sum_{i=1}^n (-\log G(x_i; \xi))^\beta \log (-\log G(x_i; \xi)) \\ &+ \sum_{i=1}^n \frac{e^{-(\log G(x_i; \xi))^\beta} (-\log G(x_i; \xi))^\beta \log (-\log G(x_i; \xi))}{1 - e^{-(\log G(x_i; \xi))^\beta}} \\ &+ (\alpha - 1) \sum_{i=1}^n \frac{e^{-(\log G(x_i; \xi))^\beta} (-\log G(x_i; \xi))^\beta \log (-\log G(x_i; \xi))}{2 - e^{-(\log G(x_i; \xi))^\beta}} \end{aligned} \tag{21}$$

$$\begin{aligned} \frac{\partial \ell}{\partial \xi} &= \sum_{i=1}^n \frac{g^\xi(x_i; \xi)}{g(x_i; \xi)} - \sum_{i=1}^n \frac{G^\xi(x_i; \xi)}{G(x_i; \xi)} - (\beta - 1)\beta \sum_{i=1}^n \frac{(-\log G(x_i; \xi))^{\beta-1} G^\xi(x_i; \xi)}{(-\log G(x_i; \xi))^\beta G(x_i; \xi)} \\ &+ \alpha\beta \sum_{i=1}^n \frac{(-\log G(x_i; \xi))^{\beta-1} G^\xi(x_i; \xi)}{G(x_i; \xi)} \\ &- \beta \sum_{i=1}^n \frac{e^{-(\log G(x_i; \xi))^\beta} (-\log G(x_i; \xi))^{\beta-1} G^\xi(x_i; \xi)}{\left(1 - e^{-(\log G(x_i; \xi))^\beta}\right) G(x_i; \xi)} \\ &- (\alpha - 1)\beta \sum_{i=1}^n \frac{e^{-(\log G(x_i; \xi))^\beta} (-\log G(x_i; \xi))^{\beta-1} G^\xi(x_i; \xi)}{\left(2 - e^{-(\log G(x_i; \xi))^\beta}\right) G(x_i; \xi)}. \end{aligned} \tag{22}$$

Where  $g^\xi(x_i; \xi)$  and  $G^\xi(x_i; \xi)$  are the partial derivative with respect to  $\xi$ . Under the usual condition for the parameters in the interior of the  $(\alpha, \beta, \xi)$  space but not on the boundary, The asymptotic distribution of  $(\hat{\Theta} - \Theta)$  as  $n \rightarrow \infty$  is the multivariate normal distribution with zero means and covariance matrix  $\mathbf{I}^{-1}(\Theta)$ . The asymptotic behavior

is also valid as  $\mathbf{I}(\Theta) = \lim_{n \rightarrow \infty} n^{-1} \mathbf{J}_n(\Theta)$ , where  $\mathbf{J}_n(\Theta)$  is a unit information matrix evaluated at  $\hat{\Theta}$ , and  $J(\Theta) = (\partial^2 \ell(\Theta) / \partial \Theta \partial \Theta^T)$ .

### 3.2 Least Square Method (LSE)

Let  $X_1 < \dots < X_n$  be an ordered random sample of size  $n$  from NETLG families of distributions. The LSEs for the vector of parameters  $\Theta = (\alpha, \beta, \xi)^T$ , i.e.  $\tilde{\Theta} = (\tilde{\alpha}, \tilde{\beta}, \tilde{\xi})^T$  can be obtained by minimizing  $L(\Theta)$  given by,

$$L(\Theta) = \sum_{i=1}^n \left( e^{-\alpha(-\log G(x_i; \xi))^\beta} \left( 2 - e^{-(-\log G(x_i; \xi))^\beta} \right)^\alpha - \frac{i}{n+1} \right)^2,$$

or by the solution of the following nonlinear equations, which can be done numerically using R software, among others.

$$\begin{aligned} \frac{\partial L}{\partial \alpha} &= 2 \sum_{i=1}^n \left( e^{-\alpha(-\log G(x_i; \xi))^\beta} \left( 2 - e^{-(-\log G(x_i; \xi))^\beta} \right)^\alpha - \frac{i}{n+1} \right) \\ &\quad \times F(x_i) \left[ \log \left( 2 - e^{-(-\log G(x_i; \xi))^\beta} \right) - (-\log G(x_i; \xi))^\beta \right] \\ \frac{\partial L}{\partial \beta} &= 2 \sum_{i=1}^n \left( e^{-\alpha(-\log G(x_i; \xi))^\beta} \left( 2 - e^{-(-\log G(x_i; \xi))^\beta} \right)^\alpha - \frac{i}{n+1} \right) \\ &\quad \times F(x_i) (-\log G(x_i; \xi))^\beta \log(-\log G(x_i; \xi)) \left[ \frac{e^{-(-\log G(x_i; \xi))^\beta}}{\left( 2 - e^{-(-\log G(x_i; \xi))^\beta} \right)} - \alpha \right] \\ \frac{\partial L}{\partial \xi} &= 2\alpha\beta \sum_{i=1}^n \left( e^{-\alpha(-\log G(x_i; \xi))^\beta} \left( 2 - e^{-(-\log G(x_i; \xi))^\beta} \right)^\alpha - \frac{i}{n+1} \right) \\ &\quad \times F(x_i) (-\log G(x_i; \xi))^{\beta-1} \frac{G^\xi(x_i; \xi)}{G(x_i; \xi)} \left[ 1 - \frac{e^{-(-\log G(x_i; \xi))^\beta}}{\left( 2 - e^{-(-\log G(x_i; \xi))^\beta} \right)} \right] \end{aligned}$$

### 3.3 Percentile Estimation (PE)

The quantile of the NETLG in (7) can be used for parameter estimation. Let  $X_1, X_2, \dots, X_n$  be an ordered random sample of size  $n$  from NETLG families of distributions, the unknown parameters  $\Theta = (\alpha, \beta, \xi)^T$  can be estimated by equating the sample percentile points to the population percentile points. Let  $u_i$  denotes an estimate of  $F(x_{i:n})$ , then the percentile estimators  $\tilde{\Theta} = (\tilde{\alpha}, \tilde{\beta}, \tilde{\xi})^T$  can be obtained by minimizing (23) or by the solution of the  $\frac{\partial P}{\partial \alpha} = \frac{\partial P}{\partial \beta} = \frac{\partial P}{\partial \xi} = 0$ .




Fig. 4 Plots of the bias and MSE of the simulated data for  $\alpha = 2.5, \beta = 1.5, \lambda = 0.5$

$$P(\Theta) = \sum_{i=1}^n \left( x_{i:n} - G^{-1} \left( e^{-\left[ -\log \left( 1 - \sqrt{1 - u_i^{\frac{1}{\alpha}}} \right) \right]^{1/\beta}} \right) \right)^2. \tag{23}$$

### 3.4 Simulation Studies

Simulation studies are conducted to examine the performances of the different parameter estimation techniques by discussing their biases and mean square error (MSE) of the estimators. A moderate sample size of  $N = 1000$  is generated each of sizes  $n = 30, 60, 90, \dots, 300$ , from the NETLE for some chosen parameters values. The computations were performed using the R3.5.3-software [57]. The resulting simulation studies are given in Fig. 4, 5, 6 and 7. The results from the figures indicated that both the MLE, LSE, and PE performed consistently as expected, and an increase in the sample sizes decreases the MSE, the bias appears negative in some cases. Thus, we can conclude that these three techniques can be enough for the parameter estimation of the NETLE and the other NETLG distributions.

### 4 Real Data Illustration

We illustrated the advantages and flexibility of the NETLG families using NETLE distribution and compared its performance with some other popular models using two real data sets. We estimated the competing models by maximum likelihood and compared fit using the Akaike Information Criterion (AIC), Bayesian Information Criterion (BIC), and Consistent Akaike Information Criterion (CAIC), Kolmogorov Smirnov (KS), Anderson-Darling (AD), and Cramer-von Mises (CvM) measures. The model with the smallest value of these measures fitted the data better. The competing models are the extended Topp–Leone exponential (ExtTLE) [31], extended Erlang-






Fig. 5 Plots of the bias and MSE of the simulated data for  $\alpha = 0.8, \beta = 0.9, \lambda = 0.5$




Fig. 6 Plots of the bias and MSE of the simulated data for  $\alpha = 1.5, \beta = 1.2, \lambda = 1.1$




Fig. 7 Plots of the bias and MSE of the simulated data for  $\alpha = 1.9, \beta = 0.9, \lambda = 1.0$

truncated exponential (ExETE) [58], Poisson Topp–Leone exponential (PTLE) [59], generalized exponential Poisson (GEP) [60], Topp–Leone exponential (TLE) [23], Topp–Leone generalized inverted exponential (TLGIE) [61], generalized exponential (GE) [62], beta exponential (BE) [63], beta Erlang-truncated exponential (BETE) [48], and exponential (E) distribution.


#### 4.1 Data I

The first data consist of the daily new deaths due to COVID-19 in New Jersey, USA, from March 12, 2020 to July 25, 2021, extracted from <https://www.worldometers.info/coronavirus/usa/new-jersey/>. The data: 1, 1, 2, 5, 2, 6, 5, 8, 19, 21, 22, 31, 37, 24, 42, 79, 100, 206, 124, 226, 81, 98, 259, 308, 222, 263, 284, 189, 106, 409, 398, 409, 365, 260, 150, 198, 425, 352, 413, 214, 279, 86, 121, 449, 372, 518, 352, 231, 162, 74, 386, 318, 297, 171, 150, 165, 88, 226, 210, 248, 231, 126, 121, 94, 161, 176, 120, 151, 111, 64, 18, 48, 162, 81, 141, 115, 84, 24, 58, 139, 114, 87, 73, 80, 87, 88, 112, 97, 56, 108, 42, 56, 63, 62, 41, 38, 29, 14, 36, 55, 52, 33, 45, 34, 27, 5, 17, 41, 52, 24, 22, 23, 50, 64, 106, 31, 50, 6, 30, 23, 43, 31, 20, 20, 5, 2, 2, 23, 33, 18, 13, 17, 16, 18, 15, 9, 10, 6, 3, 6, 3, 13, 3, 10, 11, 2, 5, 4, 15, 2, 31, 9, 7, 9, 1, 2, 2, 7, 2, 6, 8, 4, 3, 7, 5, 15, 9, 6, 5, 3, 22, 7, 11, 5, 9, 4, 4, 5, 8, 9, 4, 5, 4, 2, 1, 3, 9, 10, 3, 5, 4, 13, 7, 3, 3, 3, 4, 3, 7, 5, 7, 3, 7, 2, 1, 17, 12, 4, 4, 2, 4, 3, 18, 16, 17, 8, 12, 2, 10, 17, 15, 10, 6, 9, 1, 2, 16, 23, 16, 10, 12, 4, 10, 22, 12, 19, 25, 28, 15, 14, 38, 36, 36, 24, 32, 17, 16, 46, 59, 36, 27, 38, 11, 20, 84, 56, 62, 45, 42, 23, 15, 87, 111, 77, 59, 61, 29, 24, 86, 125, 66, 51, 50, 28, 26, 106, 139, 81, 47, 23, 19, 39, 115, 188, 82, 106, 32, 34, 38, 120, 124, 110, 99, 86, 173, 113, 34, 113, 87, 24, 16, 51, 136, 86, 109, 59, 17, 22, 132, 115, 81, 82, 72, 29, 30, 70, 109, 100, 93, 77, 24, 23, 93, 147, 79, 64, 47, 13, 13, 31, 135, 89, 62, 50, 24, 17, 104, 99, 69, 46, 46, 14, 21, 48, 128, 42, 30, 36, 16, 17, 45, 133, 46, 40, 34, 15, 22, 41, 79, 31, 27, 31, 40, 7, 61, 50, 38, 28, 24, 7, 15, 82, 75, 30, 24, 11, 9, 14, 51, 49, 34, 42, 33, 12, 25, 50, 59, 47, 43, 40, 9, 18, 45, 65, 30, 27, 39, 13, 19, 61, 49, 20, 25, 34, 12, 16, 42, 49, 33, 27, 22, 11, 10, 28, 43, 25, 26, 20, 9, 14, 23, 32, 23, 17, 14, 7, 9, 25, 34, 14, 12, 16, 6, 5, 7, 28, 6, 12, 8, 6, 5, 1, 31, 6, 2, 2, 3, 5, 11, 12, 7, 4, 4, 2, 3, 15, 18, 6, 12, 4, 3, 3, 6, 13, 5, 5, 5, 1, 4, 13, 3, 3, 5, 4, 4, 7, 11, 2, 2, 5, 3, 6, 12, 5, 2, 4, 5, 2, 4, 6, 3, 2, 5, 7, 3, 1, 8, 11, 4, 7, 5, 5, 4, 9, 6, 7, 9.


The results obtained from these measures and the estimators are provided in the Table 2. The results show that NETLE has the smallest value of the AIC, BIC, CAIC, KS, AD and CvM; thus NETLE provides a good representation of the data better than the other competing TL families and extensions exponential. Thus, NETLE can be recommended as a good model for modeling COVID-19 data and other studies in various fields of applied statistics. Figure 8 shows the plots of the (a) histogram with the fitted NETLE, PTLE, and TLE densities and (b) empirical cdf with fitted NETLE, PTLE, and TLE cdfs for the New Jersey data. Figure 9 is quantile-quantile plots of the NETLE, PTLE, and TLE for the New Jersey data set, it can be seen from the quantile-quantile plots that NETLE has more quantiles laying on the straight line.

**Table 2** MLEs, L, AIC, BIC, CAIC, KS, AD, and CvM for the New Jersey data set

Model	$\hat{\alpha}$	$\hat{\beta}$	$\hat{\lambda}$	$\hat{\theta}$	$\hat{a}$	$\hat{b}$	L	AIC	BIC	CAIC	KS	AD	CvM
NETLE	0.0091	3.1599	0.0009	-	-	-	-2446.6	48991.1	4911.8	4899.2	0.0466	0.9117	0.1202
ExTLE	0.7516	0.6289	-	0.0114	-	-	-2482.2	4970.4	4983.1	4970.4	0.0775	5.3609	0.7930
PTLE	0.8942	-	2.9992	-	-	0.0039	-2466.9	4939.8	4952.5	4939.9	0.0534	3.1108	0.4370
TLGIE	36.2636	-	0.0862	0.3732	-	-	-2481.5	4969.1	4981.7	4969.1	0.0825	5.9208	0.9615
TLE	0.6851	-	0.0073	-	-	-	-2491.5	4987.0	4995.5	4987.1	0.0913	6.5934	0.9805
GE	0.6851	-	0.0146	-	-	-	-2491.5	4987.0	4995.5	4987.1	0.0909	6.5934	0.9805
E	0.0190	-	-	-	-	-	-2516.3	5034.6	5038.9	5034.7	0.1612	6.4186	0.9540
BE	-	-	0.0021	-	0.6882	6.3917	-2489.3	4984.6	4997.3	4984.6	0.0856	6.2885	0.9325
BETE	0.0652	-	-	0.0754	0.6868	2.8914	-2489.5	4987.0	5003.9	4987.1	0.0852	6.3208	0.7375
GEP	0.0084	0.9038	2.9405	-	-	-	-2468.9	4943.7	4956.4	4943.8	0.0642	3.3547	0.4759
ExETE	0.6851	0.0183	1.5784	-	-	-	-2491.5	4989.0	5001.7	4989.1	0.0908	6.5929	0.9804



**Fig. 8** Plots of the **a** histogram with the fitted NETLE, PTLE and TLE densities, and **b** empirical cdf with fitted NETLE, PTLE and TLE cdfs for the New Jersey data




**Fig. 9** Plots of the quantile-quantile plots of the NETLE(left), PTLE(middle) and TLE(right) for the New Jersey data set


**4.2 Data II**

The second data is the daily new deaths due to COVID-19 in California, USA, collected from March 12, 2020 to September 30, 2020. The data was extracted from <https://www.worldometers.info/coronavirus/usa/california/>. The data: 1, 1, 1, 5, 4, 3, 4, 1, 10, 6, 11, 14, 17, 12, 25, 12, 14, 35, 30, 24, 41, 44, 28, 33,54, 64, 61, 25, 46, 44, 53, 55, 80, 86, 89, 105, 28, 48, 73, 120, 103, 71, 91, 32, 58, 85, 75, 89, 80, 75, 24, 71, 92, 74, 81, 91, 64, 26, 61, 97, 89, 80, 104, 55, 79, 32, 103, 86, 106, 69, 71, 31, 19, 43, 102, 82, 97, 74, 27, 47, 72, 61, 63, 72, 66, 29, 23, 95, 97, 71, 47, 72, 27, 30, 85, 79, 74, 65, 67, 24, 48, 69, 96, 79, 64, 33, 32, 42, 104, 82, 98, 63, 29, 19, 75, 118, 150, 137, 102, 73, 26, 46, 138, 125, 127, 121, 91, 12, 57, 119, 155, 156, 134, 90, 27, 92, 169, 175, 113, 191, 136, 38, 108, 196, 169, 148, 188, 103, 67, 87, 182, 160, 186, 151, 75, 19, 98, 179, 164, 134, 166, 146, 18, 104, 149, 142, 140, 144, 67, 35, 80, 144, 157, 167, 152, 65, 22, 33, 72, 154, 99, 172, 71, 52, 75, 152, 105, 90, 99, 73, 31, 53, 123, 117, 88, 132, 51, 21, 34, 150, 107.

Table 3 provide the resulting test of the competing models. The NETLE provides a better fit to the data set than the other models because NETLE has the smallest values of all the measures. Figure 10 shows the plots of the (a) histogram with the fitted NETLE, PTLE, and TLE densities and (b) empirical cdf with fitted NETLE, PTLE, and TLE cdfs for the California data set. Figure 11 is quantile-quantile plots of the NETLE, PTLE, and TLE for the California data set. In addition, Fig. 11 displayed that



**Fig. 10** Plots of the **a** histogram with the fitted NETLE, PTLE and TLE densities, and **b** empirical cdf with fitted NETLE, PTLE and TLE cdfs for the California data



**Fig. 11** Plots of the quantile-quantile plots of the NETLE (left), PTLE (middle), and TLE (right) for the California data set

the quantiles of the NETLE are laying on the straight line better than the PTLE and TLE, which indicates the good performance of our new model.

### 5 Conclusion

This paper proposes and studies a new extension of the Topp–Leone family of distributions. The model’s important mathematical and statistical properties are derived such as stochastic ordering, model series representation, moments, stress–strength reliability parameter, Renyi entropy, order statistics, and the moment of residual life. A new member of NETLG called new extended Topp–Leone exponential (NETLE) is derived; we study its skewness and kurtosis, quantile, residual life, reverse residual life, and extreme value distributions. The model parameter estimation is conducted by maximum likelihood estimation (MLE), least-square estimation (LSE), and percentile estimation (PE). The performance of the MLE, LSE, and PE is examined by simulation studies from the NETLE by discussing the estimators’ bias and mean square error (MSE), and the result was very good, as their MSE decreased with the increase in the sample size. In the end, we compare the performance of the NETLE in practice with some other popular models using two sets of daily new deaths due to COVID-19 data from California and New Jersey, USA, and our new model performs better than the other existing models as measured by some model selection criteria and goodness of fit

**Table 3** MLEs, L, AIC, BIC, CAIC, KS, AD, and CvM for the California data set

Model	$\hat{\alpha}$	$\hat{\beta}$	$\hat{\lambda}$	$\hat{\theta}$	$\hat{a}$	$\hat{b}$	L	AIC	BIC	CAIC	KS	AD	CvM
NETLE	6.3169	0.2407	0.0557	–	–	–	–1050.9	2107.4	2117.4	2107.6	0.0651	0.8527	0.1332
ExTLE	1.7785	0.0100	–	0.0177	–	–	–1062.9	2131.8	2141.7	2132.0	0.1221	2.7129	0.4499
PTLE	1.5850	–	–	–	124.60	$2.83 \times 10^{-4}$	–1054.5	2114.9	2124.8	2115.0	0.0845	1.4507	0.2377
TLGIE	158.70	–	0.0250	0.3534	–	–	–1166.8	2339.6	2349.5	2339.7	0.2447	17.2015	2.9973
TLE	1.7750	–	0.0089	–	–	–	–1062.8	2129.6	2136.2	2129.7	0.1222	2.6975	0.4472
GE	1.7750	–	0.0178	–	–	–	–1062.8	2129.6	2136.2	2129.7	0.1219	2.6975	0.4472
E	–	–	0.0127	–	–	–	–1078.0	2158.0	2161.3	2158.8	0.1810	2.4661	0.4078
BE	–	–	0.0028	–	1.7861	7.6492	–1060.9	2127.9	2137.8	2128.0	0.1154	2.4367	0.4032
BETE	0.0348	–	–	0.0883	1.7875	7.3683	–1060.9	2129.9	2143.1	2130.1	0.1153	2.4375	0.4033
GEP	0.0192	2.0100	$2.72 \times 10^{-7}$	–	–	–	–10623.6	2133.3	2143.2	2133.4	0.1171	2.7712	0.4599
ExETE	1.7752	1.2890	0.0139	–	–	–	–1062.8	2131.6	2141.5	2131.8	0.1223	2.6974	0.4472

statistics. For further studies, several special models can be derived and investigated, and different methods of estimation of the models can be considered, such as Bayesian analysis under various prior, regression analysis, survival analysis, the stress–strength reliability estimation under various viewpoints, and other applied studies due to the flexibility of the proposed family. However, the properties and characterizations of some special members of the model may require some complicated mathematical ideas such as special integrals and series representations which may lead to many useful mathematical tools. We hope that the new model will attract wider applications in various fields of studies.

**Acknowledgements** The authors are thankful to the Editorial Board and the reviewers for their valuable comments and suggestions. Conceptualization, M.M, L.L,B.A,I,M; Formal analysis, M.M, L.L,B.A,I,M,M,B,S,M; Funding acquisition, L.L.; Investigation, M.M, L.L,B.A,I,M,M,B,S,M; Methodology, M.M, L.L,B.A,I,M,M,B,S,M; Project administration, M.M, L.L.; Resources, M.M, L.L,B.A,I,M,M,B,S,M; Software, M.M,B.A,I,M,M,B,H.Z,S,M; Supervision, L.L,M,M; Validation, M.M,L.L,B.A,I,M,M,B,H.Z,S,M; Writing—original draft, M.M,S,M; Writing—review & editing, M.M, L.L,B.A,I,M,M,B,H.Z,S,M.

**Funding** Funding is provided by Key Fund of the Department of Education of Hebei Province (Grant No. ZD2018065).

**Data availability** NA.

## Declarations

**Conflict of interest** The authors declare that they have no conflict of interest.

**Ethical statement:** We are hereby declare that this manuscript is the result of our independent creation under the reviewers comments. Except for the quoted contents, this manuscript does not contain any research achievement that have been published or written by other individuals or groups. We are the only authors of this manuscript. The legal responsibility of this statement should be borne by us.

## References

1. Shi Y (2022) *Advances in big data analytics: theory, algorithms and practices*. Springer
2. Olson DL, Shi Y, Shi Y (2007) *Introduction to business data mining*, vol 10. McGraw-Hill, New York
3. Shi Y, Tian Y, Kou G, Peng Y, Li J (2011) *Optimization based data mining: theory and applications*. Springer
4. Tien JM (2017) Internet of things, real-time decision making, and artificial intelligence. *Ann Data Sci* 4(2):149–178
5. Topp CW, Leone FC (1955) A family of j-shaped frequency functions. *J Am Stat Assoc* 50(269):209–219
6. Nadarajah S, Kotz S (2003) Moments of some j-shaped distributions. *J Appl Stat* 30:311–317
7. Ghitany ME, Kotz S, Xie M (2005) On some reliability measures and their stochastic orderings for the Topp–Leone distribution. *J Appl Stat* 32:715–722
8. Sindhu T, Saleem M, Aslam M (2013) Bayesian estimation for Topp–Leone distribution under trimmed samples. *J Basic Appl Sci Res* 3:347–60
9. MirMostafae S (2014) On the moments of order statistics coming from the Topp–Leone distribution. *Stat Prob Lett* 95:85–91
10. Genc AI (2012) Moments of order statistics of Topp–Leone distribution. *Stat Pap* 53:117–131
11. Genc AI (2013) Estimation of  $P(x < y)$  with Topp–Leone distribution. *J Stat Comput Simul* 83:326–339
12. Sindhu TN, Hussain Z, Aslam M (2019) On the bayesian analysis of censored mixture of two Topp–Leone distribution. *Sri Lankan J Appl Stat* 19

13. Bayoud HA (2016) Admissible minimax estimators for the shape parameter of Topp–Leone distribution. *Commun Stat Theor Methods* 45:71–82
14. MirMostafae S, Mahdizadeh M, Aminzadeh M (2016) Bayesian inference for the Topp–Leone distribution based on lower k-record values. *Japan J Ind Appl Math* 33:637–669
15. Singh B, Khan R, and Khan A (2021) Moments of dual generalized order statistics from Topp–Leone weighted weibull distribution and characterization. *Ann Data Sci* 1–20
16. Almetwally EM, Alharbi R, Alnagar D, Hafez EH (2021) A new inverted Topp–Leone distribution: applications to the covid-19 mortality rate in two different countries. *Axioms* 10:25
17. Almetwally EM (2021) The odd Weibull inverse Topp–Leone distribution with applications to covid-19 data. *Ann Data Sci* 9:1–20
18. Shrahili M, Muhammad M, Elbatal I, Muhammad I, Bouchane M, Abba B (2021) Properties and applications of the type I half-logistic Nadarajah–Haghighi distribution. *Austrian J Stat* (Accepted manuscript)
19. Osatohanmwon P, Efe-Eyefia E, Oyegun FO, Osemwenkhai JE, Ogbonmwan SM, Afere BA (2022) The exponentiated Gumbel–Weibull logistic distribution with application to Nigeria’s covid-19 infections data. *Ann Data Sci* 1–35
20. Mohamed SA, Mousa AE, Abo-Hussien M, Ismail M (2022) Estimation of the daily recovery cases in Egypt for covid-19 using power odd generalized exponential lomax distribution. *Ann Data Sci* 9:71–99
21. Pathak A, Kumar M, Singh SK, Singh U (2022) Statistical inferences: Based on exponentiated exponential model to assess novel corona virus (covid-19) Kerala patient data. *Ann Data Sci* 9:101–119
22. Ahsan-ul Haq M, Ahmed M, Zafar J, Ramos PL (2022) Modeling of covid-19 cases in Pakistan using lifetime probability distributions. *Ann Data Sci* 9:141–152
23. Al-Shomrani A, Arif O, Shawky A, Hanif S, Shahbaz MQ (2016) Topp–Leone family of distributions: some properties and application. *Pakistan J Stat Oper Res* 443–451
24. Rezaei S, Sadr BB, Alizadeh M, Nadarajah S (2017) Topp–Leone generated family of distributions: properties and applications. *Commun Stat Theor Methods* 6:2893–2909
25. Sangsanit Y, Bodhisuwan W (2016) The Topp–Leone generator of distributions: properties and inferences. *Songklanakaraj J Sci Technol* 38
26. Mahdavi A (2017) Generalized Topp–Leone family of distributions. *J Biostat Epid* 3:65–75
27. Elgarhy M, Arslan Nasir M, Jamal F, Ozel G (2018) The type II Topp–Leone generated family of distributions: properties and applications. *J Stat Manag Syst* 21:1529–1551
28. Hassan AS, Elgarhy M, Ahmad Z (2019) Type II generalized Topp–Leone family of distributions: properties and applications. *J Data Sci* 17:638–659
29. Bantan RA, Jamal F, Chesneau C, Elgarhy M (2019) A new power Topp–Leone generated family of distributions with applications. *Entropy* 21:1177
30. Bantan RA, Jamal F, Chesneau C, Elgarhy M (2020) Type II power Topp–Leone generated family of distributions with statistical inference and applications. *Symmetry* 12:75
31. Chesneau C, Sharma VK, Bakouch HS (2021) Extended Topp–Leone family of distributions as an alternative to beta and Kumaraswamy type distributions: Application to glycosaminoglycans concentration level in urine. *Int J Biomath* 14(2):2050088
32. Ali Z, Ali A, Ozel G (2020) A modification in generalized classes of distributions: a new Topp–Leone class as an example. *Commun Stat Theor Methods* 1–23
33. Reyad HM, Alizadeh M, Jamal F, Othman S, Hamedani GG (2019) The exponentiated generalized Topp–Leone-g family of distributions: properties and applications. *Pakistan J Stat Oper Res* 1–24
34. Khaleel MA, Oguntunde PE, Al Abbasi JN, Ibrahim NA, AbuJarad MH (2020) The marshallolkin Topp–Leone-g family of distributions: a family for generalizing probability models. *Sci African* 8:00470
35. Brito E, Cordeiro GM, Yousof H, Alizadeh M, Silva G (2017) The Topp–Leone odd log-logistic family of distributions. *J Stat Comput Simul* 87:3040–3058
36. Chipepa F, Oluyede B, Makubate B et al (2020) The Topp–Leone-Marshall–Olkin-g family of distributions with applications. *Int J Stat Prob* 9:15–32
37. Muhammad M, Liu L (2021) A new extension of the beta generator of distributions. *Mathematica Slovaca* (accepted manuscript)
38. Shaked M, Shanthikumar JG (2017) *Stochastic orders*
39. Gradshteyn I, Ryzhik I, Jeffrey A, Zwillinger D (2007) *Table of integrals, series and products*, 7th edn, New York



40. Kotz S, Pensky M (2003) The stress-strength model and its generalizations: theory and applications. World Scientific
41. Guo H, Krishnamoorthy K (2004) New approximate inferential methods for the reliability parameter in a stress-strength model: the normal case. *Commun Stat Theor Methods* 33:1715–1731
42. Barbiero A (2011) Confidence intervals for reliability of stress-strength models in the normal case. *Commun Stat Simul Comput* 40(6):907–925
43. Krishnamoorthy K, Lin Y (2010) Confidence limits for stress-strength reliability involving Weibull models. *J Stat Plan Inf* 140:1754–1764
44. Kundu D, Gupta RD (2006) Estimation of  $P[y < x]$  for Weibull distributions. *IEEE Trans Reliab* 55:270–280
45. Asgharzadeh A, Valiollahi R, Raqab MZ (2011) Stress-strength reliability of Weibull distribution based on progressively censored samples. *SORT-Stat Oper Res Trans* 103–124
46. Baklizi A, El-Masri AEQ (2004) Shrinkage estimation of  $P(x < y)$  in the exponential case with common location parameter. *Metrika* 59(2):163–171
47. Kundu D, Gupta RD (2005) Estimation of  $P[y < x]$  for generalized exponential distribution. *Metrika* 61(3):291–308
48. Shrahili M, Elbatal I, Muhammad I, Muhammad M (2021) Properties and applications of beta erlang-truncated exponential distribution. *J Math Comput Sci* 22(1):16–37
49. Muhammad M (2016) Poisson-odd generalized exponential family of distributions: theory and applications. *Hacet J Math Stat* 47(6):1652–1670
50. Asgharzadeh A, Valiollahi R, Raqab MZ (2013) Estimation of the stress-strength reliability for the generalized logistic distribution. *Stat Methods* 15:73–94
51. Muhammad M, Liu L (2019) A new extension of the generalized half logistic distribution with applications to real data. *Entropy* 21(4):339
52. Nadarajah S, Bagheri S, Alizadeh M, Samani EB (2018) Estimation of the stress strength parameter for the generalized exponential-Poisson distribution. *J Test Eval* 46(5):2184–2202
53. Muhammad I, Wang X, Li C, Yan M, Chang M (2020) Estimation of the reliability of a stress-strength system from Poisson half logistic distribution. *Entropy* 22(11):1307
54. Muhammad M, Alshanbari HM, Alanzi AR, Liu L, Sami W, Chesneau C, Jamal F (2021) A new generator of probability models: the exponentiated sine-g family for lifetime studies. *Entropy*. 23(11):1394
55. Muhammad M, Bantan RA, Liu L, Chesneau C, Tahir MH, Jamal F, Elgarhy M (2021) A new extended cosine-g distributions for lifetime studies. *Mathematics* 9(21):2758
56. Arnold BC, Balakrishnan N, Nagaraja HN (1992) A first course in order statistics. SIAM
57. Team RC (2019) R. a language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria. R Core Team
58. Okorie I, Akpanta A, Ohakwe J, Chikezie D (2017) The extended erlang-truncated exponential distribution: properties and application to rainfall data. *Heliyon* 3(6):00296
59. Merovci F, Yousof H, Hamedani G (2020) The Poisson Topp–Leone generator of distributions for lifetime data: theory, characterizations and applications. *Pakistan J Stat Oper Res* 343–355
60. Barreto-Souza W, Cribari-Neto F (2009) A generalization of the exponential-Poisson distribution. *Stat Prob Lett* 79(24):2493–2500
61. Al-Saiary ZA, Bakoban RA (2020) The Topp–Leone generalized inverted exponential distribution with real data applications. *Entropy* 22(10):1144
62. Gupta RD, Kundu D (1999) Theory and methods: generalized exponential distributions. *Aust N Z J Stat* 41(2):173–188
63. Nadarajah S, Kotz S (2006) The beta exponential distribution. *Reliab Eng Syst Safe* 91(6):689–697

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

## Authors and Affiliations

Mustapha Muhammad<sup>1</sup>  · Lixia Liu<sup>2</sup> · Badamasi Abba<sup>3,4</sup> ·  
Isyaku Muhammad<sup>5</sup> · Mouna Bouchane<sup>6</sup> · Hexin Zhang<sup>7</sup> · Sani Musa<sup>8</sup>

Mustapha Muhammad  
mmmahmoud12@sci.just.edu.jo; mmuhammad.mth@buk.edu.ng

Badamasi Abba  
babba@yumsuk.edu.ng; badamasiabba@csu.edu.cn

Isyaku Muhammad  
isyakuedu@yahoo.com

Mouna Bouchane  
mouna.bouchane@gmail.com

Hexin Zhang  
corehebtu@126.com

Sani Musa  
musasani1010@gmail.com

- <sup>1</sup> Department of Mathematics, Guangdong University of Petrochemical Technology, Maoming 525000, China
- <sup>2</sup> School of Mathematical Sciences, Hebei Normal University, Shijiazhuang 050024, People's Republic of China
- <sup>3</sup> School of Mathematics and Statistics, Central South University, Changsha, China
- <sup>4</sup> Department of Mathematics, Yusuf Maitama Sule University, Kano, Nigeria
- <sup>5</sup> School of Mechanical and Electrical Engineering, University of Electronic Science and Technology of China, Chengdu 611731, China
- <sup>6</sup> Key Laboratory of Augmented Reality College of Mathematics and Information Science, Hebei Normal University, Shijiazhuang, China
- <sup>7</sup> School of Mathematical Sciences, Hebei Normal University, Shijiazhuang 050024, People's Republic of China
- <sup>8</sup> Department of Mathematics and Computer Science, Sule Lamido University, Kafin-Hausa, Jigawa, Nigeria