



# Wnt Signalling Inhibitors Potently Drive Trans-differentiation Potential of Mesenchymal Stem Cells Towards Neuronal Lineage

Antara Banerjee<sup>1,3</sup> · Praveen Rowlo<sup>1</sup> · Ganesan Jothimani<sup>1</sup> · Asim K. Duttaroy<sup>2</sup> · Surajit Pathak<sup>1</sup>

Published online: 15 October 2022

© Taiwanese Society of Biomedical Engineering 2022

## Correction to:

**Journal of Medical and Biological Engineering**  
<https://doi.org/10.1007/s40846-022-00730-7>

In the previous version, figures legends were misplaced and arranged wrongly by the Springer Nature production department of the journal, which is corrected now.

---

The original article can be found online at <https://doi.org/10.1007/s40846-022-00730-7>.

---

✉ Antara Banerjee  
antarabanerjee@care.edu.in; antara.banerjee27@gmail.com

<sup>1</sup> Department of Medical Biotechnology, Faculty of Allied Health Sciences, Chettinad Academy of Research and Education (CARE), Chettinad Hospital and Research Institute (CHRI), Chennai, India

<sup>2</sup> Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo, Norway

<sup>3</sup> Faculty of Allied Health Sciences, Chettinad Academy of Research and Education, Kelambakkam 603103, India