CORRECTION

Correction to: Twist-minimal trace formula for holomorphic cusp forms

Kieran Child

*Correspondence:
kieran.child@bristol.ac.uk
University of Bristol, Beacon
House, Queens Road, Bristol BS8
1QU, UK
The original article can be found
online at <https://doi.org/10.1007/s40993-021-00302-9>

Correction to: Res. Number Theory

<https://doi.org/10.1007/s40993-021-00302-9>

Errata

- (1) On page 3, equation 2.5, the congruence subgroup $\Gamma_0(N)$ should be defined as the following subgroup of $SL_2(\mathbb{Z})$:

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z}) : N \mid c \right\}.$$

- (2) At the top of page 4, the second condition for a modular form should state that $f|_k\gamma(z)$ is bounded as $\Im(z) \rightarrow z$ for all $\gamma \in SL_2(\mathbb{Z})$.
- (3) On page 4, a cusp form is a modular form for which $f|_k\gamma(z) \rightarrow 0$ as $\Im(z) \rightarrow \infty$ for all $\gamma \in SL_2(\mathbb{Z})$.

The original article can be found online at <https://doi.org/10.1007/s40993-021-00302-9>.

Published online: 20 January 2022

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.