Advances in composite-based structural equation modeling

Marko Sarstedt^{1,2} · Heungsun Hwang³

Published online: 19 February 2020
© The Author(s) 2020

Structural equation modeling (SEM) has become a quasi-standard tool for analyzing complex inter-relationships between observed and latent variables. Two conceptually different approaches to SEM have been proposed: factor- vs. component-based SEM. Factor-based SEM approximates latent variables by common factors as in common factor analysis, whereas component-based SEM regards them as weighted composites of observed variables as in multivariate statistics such as canonical correlation analysis and principal component analysis. Factor-based SEM is represented by covariance structure analysis, whereas composite-based SEM includes generalized structured component analysis (GSCA; Hwang and Takane 2004), partial least squares (PLS; Lohmöller 1989), regularized generalized canonical correlation analysis (Tenenhaus and Tenenhaus 2011), and several others. Although factor-based SEM remains prevalent in practice, numerous methodological advances (e.g., Hwang et al. 2010; Suk and Hwang 2016; Schlittgen et al. 2016) and tutorial articles, which have made the methods accessible to applied researchers (e.g., Hair et al. 2019, 2020; Sarstedt et al. 2019), have contributed to component-based SEM's growing popularity in recent years.

Parallel to these developments, recent research in psychometrics calls the central tenets of the common factor model into question. For example, Rigdon (2016, p. 602) notes that “common factor proxies cannot be assumed to carry greater significance than composite proxies in regard to the existence or nature of conceptual variables.” Similarly, Rhemtulla et al. (2020) observe that “there is a growing appreciation within some areas of psychology that the latent variable model may not be the right model to capture relations between many psychological constructs and their observed indicators.” This notion has been echoed in numerous

✉ Marko Sarstedt
marko.sarstedt@ovgu.de
Heungsun Hwang
heungsun.hwang@mcgill.ca

¹ Otto-Von-Guericke-University Magdeburg, Magdeburg, Germany

² Monash University Malaysia, Subang Jaya, Malaysia

³ McGill University, Montreal, Canada

other publications in a variety of fields (e.g., Hair and Sarstedt 2019; Henseler et al. 2014; Rigdon 2012; Rigdon et al. 2017). More fundamentally, Rigdon et al. (2019) show that the indeterminacy of common factors creates a band of (metrological) uncertainty in the relationship between the factor inside the model and any variable outside the model—including the conceptual variable that the factor seeks to represent (Steiger 1979). The standard treatment of construct measures in covariance structure analysis—such as using only few indicators to measure a construct—increases factor indeterminacy and hence the degree of metrological uncertainty, hindering the replicability of behavioral science research (Rigdon et al. 2020). These results do not imply that component-based SEM techniques excel covariance structure analysis per se. However, they cast doubt on the universal applicability of the common factor model.

While there has always been a controversy between factor-based and composite-based approaches to SEM, recently, the tenor of this controversy has become more intense. Whereas some researchers strongly advocate the use of component-based SEM (Sarstedt et al. 2016), others believe that this approach should be abandoned (Rönkkö et al. 2016). The debates also led to a diversification of the composite-based SEM community, with differing viewpoints on the nature of measurement, the role of model fit, and the methods' scope of application. For example, Hwang et al. (2017) proposed GSCA with measurement errors incorporated, called $GSCA_M$, which aims to estimate the parameters of factor-based SEM via GSCA. Similarly, whereas some researchers stress the need to consider model fit metrics, others emphasize statistics for assessing a model's out-of-sample predictive accuracy (Cho et al. 2019; Hair et al. 2019; Shmueli et al. 2019).

In light of these controversies and debates, composite-based SEM is at the crossroads. The following years will show under which conditions composite-based-SEM methods will routinely be used and how sustainable their current popularity will be. With these developments in mind, this special issue of *Behaviormetrika* seeks to serve as a platform for advancing and furthering our understanding of composite-based SEM methods.

The lead article in this special issue by Hwang et al. (2020) contrasts PLS-SEM and GSCA, arguably the most prominent composite-based SEM methods in the field. After the conceptual comparison of the two approaches, the authors present the result of a concept analysis of methodological research on PLS-SEM and GSCA to identify dominant topics that characterize the joint research domain. The results illustrate the field's maturation, showing, for example, that researchers have become aware of the conceptual differences between composite and factor models and their implications for the methods' performance. Based on the results, the authors identify numerous research avenues for research on composite-based SEM methods.

Tying in with this lead article, Cho and Choi (2020) offer a comparison of PLS-SEM and GSCA on the grounds of a simulation study. For this purpose, the authors propose a new data generation approach where components are constructed to explain the variances of their indicators as well as those of endogenous components. Their simulation study considers different measurement model set-ups and PLS-SEM-based estimation modes. Their result pattern is similar to Hair et al.'s (2017), in that GSCA recovers measurement model parameters more effectively than

PLS–SEM, while both approaches perform very similarly with regard to structural model parameter recovery.

The third paper in this special issue illustrates the use of GSCA in the specific context of brain connectivity research. Using data collected during encoding of source memory in a functional magnetic resonance imaging (fMRI) study, Jung et al. (2020) demonstrate how to specify and evaluate a fully and bidirectionally connected structural model of brain connectivity using GSCA. This application to fMRI data nicely ties in with the increasing number of studies that use composite-based SEM in fields other than the social sciences to explore natural/biological phenomena (Sarstedt 2019). Based on their results, the authors discuss various implications for future extensions of the GSCA approach.

The fourth paper by Ryoo et al. (2020) offers such an extension by combining GSCA with optimal scaling and fuzzy clustering to capture unobserved class-level heterogeneity in the data. The authors test their new approach on real-world data to show that it yields the same results as maximum likelihood-based latent class analysis, while avoiding identification issues. The new approach, therefore, nicely expands the applicability and capability of latent class analysis in composite-based SEM.

The final paper in this special issue by Schamberger et al. (2020) offers a robust variant of standard PLS–SEM and consistent PLS–SEM (PLSc–SEM). Their simulation study with various population models and simulation conditions underlines the efficacy of the approach to reliably recover model estimates in the presence of outliers.

We are confident that the papers in this special issue will trigger significant interest in the field and inspire exciting follow-up research. We would like to thank Behaviormetrika's Editor-in-Chief, Maomi Ueno, for giving us the opportunity to edit this special issue. In addition, we would like to thank the numerous reviewers without whom this special issue would not have been possible—thank you!

Acknowledgements Open Access funding provided by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Cho G, Choi JY (2020) An empirical comparison of generalized structured component analysis and partial least squares path modeling under variance-based structural equation models. *Behaviormetrika*. <https://doi.org/10.1007/s41237-019-00098-0>
- Cho G, Jung K, Hwang H (2019) Out-of-bag prediction error: a cross validation index for generalized structured component analysis. *Multivar Behav Res* 54(4):505–513
- Hair JF, Sarstedt M (2019) Composites vs factors: implications for choosing the right SEM method. *Project Management Journal* 50(6):1–6
- Hair JF, Hult GTM, Ringle CM, Sarstedt M, Thiele KO (2017) Mirror, mirror on the wall: a comparative evaluation of composite-based structural equation modeling methods. *J Acad Mark Sci* 45(5):616–632
- Hair JF, Risher JJ, Sarstedt M, Ringle CM (2019) When to use and how to report the results of PLS–SEM. *Eur Bus Rev* 31(1):2–24
- Hair JF, Howard MC, Nitzl C (2020) Assessing measurement model quality in PLS–SEM using confirmatory composite analysis. *J Bus Res* 109:101–110
- Henseler J, Dijkstra TK, Sarstedt M, Ringle CM, Diamantopoulos A, Straub DW, Ketchen DJ, Hair JF, Hult GTM, Calantone RJ (2014) Common beliefs and reality about partial least squares: comments on Rönkkö & Evermann (2013). *Org Resh Methods* 17(2):182–209
- Hwang H, Takane Y (2004) Generalized structured component analysis. *Psychometrika* 69(1):81–99
- Hwang H, Ho M-HR, Lee J (2010) Generalized structured component analysis with latent interactions. *Psychometrika* 75(2):228–242
- Hwang H, Takane Y, Jung K (2017) Generalized structured component analysis with uniqueness terms for accommodating measurement error. *Front Psychol* 8:2137
- Hwang H, Sarstedt M, Cheah JH, Ringle CM (2020) A concept analysis of methodological research on composite-based structural equation modeling: bridging PLSPM and GSCA. *Behaviormetrika*. <https://doi.org/10.1007/s41237-019-00085-5>
- Jung K, Cho SS, Lee J, Kim S, Ryoo JH (2020) An illustrative application of generalized structured component analysis for brain connectivity research. *Behaviormetrika*. <https://doi.org/10.1007/s41237-019-00080-w>
- Lohmöller J-B (1989) *Latent variable path modeling with partial least squares*. Springer, Berlin
- Rhemtulla, M., van Bork, R., & Borsboom, D. (2020). Worse than measurement error: Consequences of inappropriate latent variable measurement models. *Psychol Methods* (advance online publication).
- Rigdon EE (2012) Rethinking partial least squares path modeling: in praise of simple methods. *Long Range Plan* 45(5–6):341–358
- Rigdon EE (2016) Choosing PLS path modeling as analytical method in European management research: a realist perspective. *European Management Journal* 34(6):598–605
- Rigdon, E. E., Sarstedt, M., & Ringle, C. M. (2017). On comparing results from CB-SEM and PLS–SEM Five perspectives and five recommendations. *Marketing ZFP—Journal of Research and Management*, 39(3), 4–17.
- Rigdon EE, Becker J-M, Sarstedt M (2019) Factor indeterminacy as metrological uncertainty: implications for advancing psychological measurement. *Multivar Behav Res* 54(3):429–443
- Rigdon EE, Sarstedt M, Becker J-M (2020) Quantify uncertainty in behavioral research. *Nat Hum Behav* (forthcoming)
- Rönkkö M, McIntosh CN, Antonakis J, Edwards JR (2016) Partial least squares path modeling: time for some serious second thoughts. *J Oper Manag* 47–48(November):9–27
- Ryoo JH, Park S, Kim S (2020) Categorical latent variable modeling utilizing fuzzy clustering generalized structured component analysis as an alternative to latent class analysis. *Behaviormetrika*. <https://doi.org/10.1007/s41237-019-00084-6>
- Sarstedt M (2019) Der Knacks and a silver bullet. In: Babin BJ, Sarstedt M (eds) *The great facilitator: reflections on the contributions of Joseph F Hair, Jr to marketing and business research*. Springer, Cham, Switzerland, pp 155–164
- Sarstedt M, Hair JF, Ringle CM, Thiele KO, Gudergan SP (2016) Estimation issues with PLS and CBSEM: where the bias lies! *J Bus Res* 69(10):3998–4010
- Sarstedt M, Hair JF, Cheah J-H, Becker J-M, Ringle CM (2019) How to specify, estimate, and validate higher-order models. *Australas Mark J* 27(3):197–211

- Schamberger T, Schuberth F, Henseler J, Dijkstra TK (2020) Robust partial least squares path modeling. *Behaviormetrika*. <https://doi.org/10.1007/s41237-019-00088-2>
- Schlittgen R, Ringle CM, Sarstedt M, Becker J-M (2016) Segmentation of PLS path models by iterative reweighted regressions. *J Bus Res* 69(10):4583–4592
- Shmueli G, Sarstedt M, Hair JF, Cheah J-H, Ting H, Ringle CM (2019) Predictive model assessment in PLS–SEM: Guidelines for using PLSpredict. *Eur J Mark* 53(11):2322–2347
- Steiger JH (1979) The relationship between external variables and common factors. *Psychometrika* 44(1):93–97
- Suk HW, Hwang H (2016) Functional generalized structured component analysis. *Psychometrika* 81(4):940–968
- Tenenhaus A, Tenenhaus M (2011) Regularized generalized canonical correlation analysis. *Psychometrika* 76(2):257–284

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.