Correction to: Wind of change: zooarchaeological approach to the Middle–Upper Palaeolithic transition in Cova Gran of Santa Linya (Lleida, south-eastern Pre-Pyrenees)

Sofía C. Samper Carro^{1,2,3} · Jorge Martínez-Moreno³ · Rafael Mora³

Published online: 19 September 2020
© Springer Nature Switzerland AG 2020

Correction to: Journal of Paleolithic Archaeology
<https://doi.org/10.1007/s41982-020-00066-1>

The original version of this article unfortunately contained a mistake. Table 3 is now presented here correctly.
The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s41982-020-00066-1>

✉ Sofia C. Samper Carro
sofia.samper@anu.edu.au

¹ School of Archaeology and Anthropology, College of Arts and Social Sciences, The Australian National University, Canberra 2601, Australia

² Department Archaeology & Natural History, School of Culture, History and Language, College of Asia & the Pacific, The Australian National University, Canberra 2601, Australia

³ Centre d'Estudis del Patrimoni Arqueològic (CepArQUAB), Facultat de Lletres-Edifici B, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

Table 3 Taxonomic list, including fragments classified by mammal size and unidentified fragments, for archaeological units 497D and S1B. Numbers in brackets indicate the percentage

	497D		S1B					
	NR(%)	NISP(%)	MNE(%)	MNI(%)	NR(%)	NISP(%)	MNE(%)	MNI(%)
<i>Cervus elaphus</i>	33(9%)	33(29%)	19(21%)	2(18%)	35(8%)	35(21%)	18(12%)	2(13%)
<i>Bos/Bison</i> sp.	2(1%)	2(2%)	2(2%)	1(9%)	8(2%)	8(5%)	7(5%)	1(6%)
<i>Capra pyrenaica</i>	27(8%)	27(24%)	17(19%)	2(18%)	6(1%)	6(4%)	5(3%)	1(6%)
<i>Sus scrofa</i>	-	-	-	-	1(0.2%)	1(1%)	1(1%)	1(6%)
<i>Equus ferus</i>	7(2%)	7(6%)	6(7%)	1(9%)	41(10%)	41(24%)	21(14%)	2(13%)
<i>Equus hydruntinus</i>	-	-	-	-	6(1%)	6(4%)	5(3%)	1(6%)
<i>Stephanorhinus hemitoechus</i>	26(7%)	26(23%)	9(10%)	1(9%)	6(1%)	6(4%)	3(2%)	1(6%)
<i>Vulpes vulpes</i>	3(1%)	3(3%)	3(3%)	1(9%)	-	-	-	-
<i>Oryctolagus cuniculus</i>	14(4%)	14(12%)	14(15%)	2(18%)	62(14%)	62(36%)	59(39%)	5(31%)
Aves	2(1%)	2(2%)	2(2%)	1(9%)	5(1%)	5(3%)	5(3%)	1(6%)
<i>Rana</i> sp.	-	-	-	-	1(0.2%)	1(1%)	1(1%)	1(6%)
Carnivore:herbivore ratio	0.03	0.03	0.04	0.1	-	-	-	-
Small size (<60kg)	95(27%)	-	9(9.8%)	-	24(6%)	-	6(4%)	-
Medium size (60-300kg)	91(26%)	-	5(5.4%)	-	107(25%)	-	7(5%)	-
Large size (>300kg)	23(6%)	-	3(3.3%)	-	43(10%)	-	8(5%)	-
Very large size (>700kg)	4(1%)	-	3(3.3%)	-	35(8%)	-	4(3%)	-
Unidentified mammal	29(8%)	-	-	-	50(12%)	-	-	-
Total(%NRt)	356 (6%)	114	92	11	430(6%)	171	150	16
Unidentified fragments	338(6%)	-	-	-	538(8%)	-	-	-
Unidentified fragments <2 cm	5174(88%)	-	-	-	6170(86%)	-	-	-
NRt	5868	114	92	11	7138	171	150	16