



Correction to: Measured Phenology Response of Unchanged Crop Varieties to Long-Term Historical Climate Change

Dingrong Wu¹ · Peijuan Wang¹ · Chaoyang Jiang¹ · Jianying Yang¹ · Zhiguo Huo¹ · Qiang Yu^{2,3,4}

Published online: 28 January 2019
© Springer Nature Switzerland AG 2019

Correction to: International Journal of Plant Production
<https://doi.org/10.1007/s42106-018-0033-z>

The original version of this article unfortunately contained mistakes. Table 1 was incorrect. The corrected Table 1 is given below.

Table 1 Information of sites which the planted variety kept unchanged for at least 15 years

Site	Elevation (m)	Year of planting	Number of planting years	Variety name
Winter wheat				
Changzhi	991.8	1988–2004	17	CZ648
Hancheng	458.1	1991–1994, 1996–2003, 2006–2009	15	XY6
Jincheng	752.6	1981–1986, 1989–1992, 1994–1998, 2001, 2003–2006	21	5819
Huanghua	6.6	1989–2011	23	71321
Laizhou	48.4	1990–2010	21	YN15
Fushan	53.9	1987–2011	26	YN15
Tianshui	1141.6	1991–2009	19	7464
Single rice				
Tonghua	380.1	1985–2010	26	QG
Muling	266.3	1994–1995, 1997–2011	16	SY397
Spring maize				
Jiamusi	82.0	1992–2002, 2004–2010	18	DN248
Meihekou	340.5	1986–1993, 1995–2001	15	TD4
Tonghua	380.1	1983–2002	20	JD101

The original article can be found online at <https://doi.org/10.1007/s42106-018-0033-z>.

✉ Qiang Yu
yuq@igsnr.ac.cn

¹ State Key Laboratory of Severe Weather (LASW), Chinese Academy of Meteorological Sciences, Beijing 100081, China

² State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, Northwest A&F University, Yangling 712100, China

³ Plant Functional Biology and Climate Change Cluster, University of Technology Sydney, Broadway, PO Box 123, Ultimo, NSW 2007, Australia

⁴ College of Resources and Environment, University of Chinese Academy of Science, Beijing 100049, China