



First report of powdery mildew caused by *Golovinomyces monardae* on peppermint in Italy

Angelo Garibaldi¹ · Giovanna Gilardi¹ · Sara Franco-Ortega¹ · Maria Lodovica Gullino¹

Published online: 23 March 2018
© Società Italiana di Patologia Vegetale (S.I.Pa.V.) 2018

Keywords Medicinal plants · Foliar pathogens · Disease symptoms

Peppermint (*Mentha × piperita*) is widespread in cultivation in many regions of the world as a medicinal plants. During the summer of 2017, in a garden near Biella (northern Italy), 6-month-old peppermint plants exhibited a white powdery mildew growth on the adaxial surfaces of leaves, stems and petioles at temperature range from 15 to 28 °C. Infected leaves turned yellow and dropped. Conidia were hyaline, elliptical to doliform, without fibrosin bodies, and measured 25.2 to 39.8 × 15.7 to 28.9 (31.3 × 22.6 average) µm. Chasmothecia were not observed. Based on morphology, the causal agent was determined as *Golovinomyces* spp. (Scholler et al. 2016). The internal transcribed spacer region of rDNA of the pathogen was amplified using the primers ITS1/ITS4 and sequenced (GenBank Accession MG205583). BLASTn analysis (Altschul et al. 1997) of the 436 bp obtained showed 99% similarity with *Golovinomyces monardae* (G.S. Nagy) M. Scholler, U. Brawn & Anke Schmidt (LC076842, voucher KR:M-43413). Pathogenicity was confirmed in a greenhouse at 18 to 25 °C by spraying leaves of healthy peppermint plants with the pathogen at 1 × 10⁵ conidia/ml. Twenty plants were inoculated, and the same number of non-inoculated plants served as control. Typical spots of powdery mildew developed

on inoculated plants 15 to 17 days after inoculation. Non-inoculated plants did not show symptoms. To our knowledge, this is the first report of powdery mildew caused by *Golovinomyces monardae* on peppermint in Italy. According to taxonomic and phylogenetic studies of the *Golovinomyces biocellatus* complex (Scholler et al. 2016) the same pathogen has been reported on peppermint in Germany, on *Mentha spicata* in Italy and on other hosts belonging to Lamiaceae and Verbenaceae (Garibaldi et al. 2010; Scholler et al. 2016).

References

- Altschul SF, Madden TL, Schaffer AA, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programme. *Nucleic Acids Res* 25:3389–3402
- Garibaldi A, Bertetti D, Pensa P, Gullino ML (2010) Powdery mildew caused by *Golovinomyces biocellatus* on spearmint (*Mentha spicata*) in Italy. *Plant Dis* 94:1171
- Scholler M, Schmidt A, Siahaan SAS, Takamatsu S, Braun U (2016) A taxonomic and phylogenetic study of the *Golovinomyces biocellatus* complex (*Erysiphales*, Ascomycota) using asexual state morphology and rDNA sequence data. *Mycol Prog* 15:56

✉ Giovanna Gilardi
giovanna.gilardi@unito.it

¹ Centre of Competence for the Innovation in the Agro-Environmental Sector (AGROINNOVA), University of Torino, Largo P. Braccini 2, 10095 Grugliasco, Italy