First report of *Gilbertella persicaria* causing soft rot in eggplant fruit in China

Ibatsam Khokhar^{1,2} · Irum Mukhtar³ · Junhuan Wang¹ · Yang Jia¹ · YanChun Yan¹

Received: 25 December 2018 / Accepted: 2 May 2019 / Published online: 17 May 2019
© Società Italiana di Patologia Vegetale (S.I.Pa.V.) 2019

Keywords *Solanum melongena* · *Gilbertella persicaria* · Soft rot · Fungal pathogen · Post harvest

In August 2018, eggplant fruits (*Solanum melongena*) were collected with circular, light brown and slightly sunken lesions at the apex containing little white mycelium (infection incidence was 1–2%) from a local market in the Haidian district, Beijing, China. Purified fungal isolates initially produced white colonies; later individual colonies covered the entire PDA plate surface (90 mm diam) within 3 days. Sporangiophores were smooth walled, simple to branched, erect and curved below the sporangium. Sporangia were globose to subglobose, $36.5\text{--}247.5 \times 37.4\text{--}250.5 \mu\text{m}$, columella (ovoid, pyriform or subglobose), $21.5\text{--}108.2 \times 24.5\text{--}135.6 \mu\text{m}$. Sporangiospores were globose to ellipsoid, hyaline, smooth, one-celled, $7.8\text{--}13 \times 5.2\text{--}10.4 \mu\text{m}$. Based on the morphological characteristics the fungus was identified as *Gilbertella persicaria* (Hesseltine 1960). Fungal DNA of eight isolates were amplified for partial ITS regions using ITS1/ITS4 primers and sequenced. Obtained sequences showed a 100% similarity with various ITS sequences of *G. persicaria* (KR076759, KR076758, KT213049). The

ITS sequence (MH753636) of a representative isolate (YC-1K2) was submitted in GenBank and showed 99.83% similarity with the ex-type sequence NR_111692. Koch's postulates were fulfilled by inoculating three asymptomatic eggplant fruits. A sporangiospores suspension (1×10^5 sporangiospores/ml) was sprayed on punctured sites on each fruit, kept in plastic boxes and placed in a moist chamber at 25 °C for 5 days. Sterile distilled water was sprayed on a set of three eggplant fruits for control. Inoculated fruits showed symptoms (circular, light brown spots) after 48 h of inoculation and sporangiophores evolved into rotted lesion after 5 days. The re-isolated and identified fungus from infected fruits corresponded to *G. persicaria*. Recently, *G. persicaria* has been reported to cause soft rot on dragon fruits (*Hylocereus* spp.) in China and Taiwan (Guo et al. 2012; Lin et al. 2014). To our knowledge, this is the first report of *G. persicaria* soft rot in eggplant fruit in mainland China.

Ibatsam Khokhar and Irum Mukhtar contributed equally to this work.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s42161-019-00321-7>) contains supplementary material, which is available to authorized users.

✉ Ibatsam Khokhar
ibatsamk@yahoo.com

- ¹ Graduate School of Chinese Academy of Agricultural Sciences, Beijing 100081, China
- ² Department of Biological Sciences, Forman Christian College (A Chartered University), Lahore, Pakistan
- ³ Mycological Research Center (MRC), College of Life Sciences, Fujian Agriculture and Forestry University, Fuzhou 350002, Fujian, China

References

- Guo LW, Wu YX, Mao ZC, Ho HH, He YQ (2012) Storage rot of dragon fruit caused by *Gilbertella persicaria*. Plant Dis 96:1826
- Hesseltine CW (1960) *Gilbertella* gen. nov. (Mucorales). Bull Torrey Bot Club 87:21–30
- Lin JP, Ann PJ, Tsai JN, Hsu ZH, Chang JT (2014) Flower and fruit wet rot of pitaya (*Hylocereus* spp.) caused by *Gilbertella persicaria*, a new disease record in Taiwan. Plant Pathol Bull 23:109–124

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.