First report of *Nigrospora sphaerica* causing leaf spot of date palm in Pakistan

Muhammad Waqar Alam¹ · Abdul Rehman² · Shahbaz Ahmad¹ · Mubeen Sarwar¹ · Aamir Nawaz³ · Samiya Mahmood Khan⁴ · Sajid Ali³ · Sidra Aslam⁵ · Abdul Mannan¹

Received: 3 November 2018 / Accepted: 3 July 2019 / Published online: 12 July 2019
© Società Italiana di Patologia Vegetale (S.I.Pa.V.) 2019

Keywords *Nigrospora sphaerica* · Leaf spot · Pathogenicity

Date palm (*Phoenix dactylifera* L.) is one of the most cultivated fruit trees around the world. Pakistan is the 6th largest producer of dates (Mangan et al. 2016). During November 2017, leaf spot symptoms were observed in different date palm orchards. Symptoms included yellow to brown, semicircular to irregular spots (3 mm diameter), which gradually enlarged up to 8 mm diameter and coalesced. Symptomatic tissues of infected leaves were cut into small pieces, surface sterilized in 75% ethanol for 30 s, rinsed three times in sterile distilled water, and cultured onto potato dextrose agar (PDA) for 5–7 days at 25 °C. A purified culture of the isolated pathogen (PD20181) was stored at the Fungal Culture Collection Centre, Department of Plant Pathology, University of Agriculture, Faisalabad. The fungus produced single celled, black, spherical to subspherical conidia (10 to 14 × 18 to 20 µm), borne on a hyaline vesicle at the tip of each conidiophore. Morphologically, the fungus was identified as *Nigrospora sphaerica* (Sacc.) Mason (Mason 1927; Wang et al. 2017). Identification was confirmed by amplifying and sequencing the ITS region and translation elongation factor 1- α (*TEF1- α*) using primers ITS1/ITS4 (White et al. 1990) and EF1-728F (Carbone and Kohn 1999), respectively. BLAST

analysis of GenBank Accession Nos. MH815137 and MK875828 showed 99 and 100% identity with *N. sphaerica* strains (KX986034 and KY019414) from China. Pathogenicity was verified on 10 detached, surface-sterilized, healthy leaves (cv. Aseel). Leaves were wounded and inoculated with a mycelial plug of 0.5 cm. Leaves inoculated with sterile PDA plug served as negative control. Leaves were placed in a moist chamber at 25 °C and 80% RH. Twelve days post inoculation, inoculated leaves had symptoms similar to those originally observed on affected trees; control remained asymptomatic. The morphological characteristics of the fungi reisolated from inoculated leaves were the same as of original isolates, fulfilling Koch's postulates. *Nigrospora sphaerica* has been reported to cause leaf spot of Kinnow mandarin in Pakistan (Alam et al. 2017). To our knowledge, this is the first report of leaf spot caused by *N. sphaerica* on date palm in Pakistan.

References

- Alam MW, Rehman A, Gleason ML, Riaz K, Saira M, Aslam S, Rosli R, Muhammad S (2017) First report of *Nigrospora sphaerica* causing leaf spot of Kinnow mandarin in Pakistan. J Plant Pathol 107:295
- Carbone I, Kohn LM (1999) A method for designing sets for speciation studies in filamentous ascomycetes. Mycologia 91:553–556
- Mangan T, Abdul SJ, Mehar NR, Muhammad IK, Abdul GK (2016) Profit earned by date palm farmers and contractors in district Khairpur, Sindh, Pakistan. Sci Int 28:775–779
- Mason S.C. (1927). Date culture in Egypt and the Sudan. USDA bulletin no. 1457, Washington, DC, pp. 72
- Wang M, Liu F, Crous PW, Cai L (2017) Phylogenetic reassessment of *Nigrospora*: ubiquitous endophytes, plant and human pathogens. Persoonia 39:118–142
- White TJ, Bruns TD, Taylor JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) PCR protocols: a guide to methods and applications. Academic Press, San Diego, pp 315–322

✉ Muhammad Waqar Alam
waqaralam2009@gmail.com

¹ Institute of Agricultural Sciences, University of the Punjab, Lahore, Pakistan

² Department of Plant Pathology, University of Agriculture, Faisalabad, Pakistan

³ Department of Horticulture, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Pakistan

⁴ Department of Plant Pathology, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Pakistan

⁵ Department of Ecology, College of Life Sciences, Beijing Normal University, Beijing, China