CORRECTION

Correction to: Distribution and association of agro-ecological factors influencing garlic rust (*Puccinia allii*) epidemics in Eastern Amhara, Ethiopia

Zemenu Endalew¹ · Habtamu Terefe² · Mashilla Dejene² · Alok Kumar³

Published online: 2 March 2021
© Indian Phytopathological Society 2021

Correction to: Indian Phytopathology
<https://doi.org/10.1007/s42360-020-00297-w>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original version of this article unfortunately contained a mistake. The presentation of equations was incorrect. The corrected equations are given below.

The original article has been corrected.

$$\text{Prevalence (\%)} = \frac{\text{Number of fields showing rust symptom}}{\text{Total number of fields visited}} \times 100$$

$$\text{Incidence (\%)} = \frac{\text{Number of plants infected in the quadrat}}{\text{Total number of plants in the quadrat}} \times 100$$

$$\text{PSI} = \frac{\text{Sum of numerical ratings}}{\text{Total number of plants scored} \times \text{maximum score on scale}} \times 100$$

The original article can be found online at <https://doi.org/10.1007/s42360-020-00297-w>.

✉ Habtamu Terefe
habmam21@gmail.com

¹ Sekota Dryland Agricultural Research Center, ARARI, P.O. Box 62, Sekota, Ethiopia

² Plant Protection Program, School of Plant Sciences, Haramaya University, P.O. Box 138, Dire Dawa, Ethiopia

³ Plant Breeding and Genetics Program, School of Plant Sciences, Haramaya University, P.O. Box 138, Dire Dawa, Ethiopia