Correction to: Evaluation of the tolerance and biotransformation of ferulic acid by *Klebsiella pneumoniae* TD 4.7

Maitê Bernardo Correia dos Santos¹ · Josiane Aniele Scarpassa¹ · Diego Alves Monteiro¹ · Guillermo Ladino-Orjuela¹ · Roberto Da Silva² · Mauricio Boscolo² · Eleni Gomes¹

Published online: 12 April 2021

© Sociedade Brasileira de Microbiologia 2021

Correction to: Braz J Microbiol

<https://doi.org/10.1007/s42770-021-00462-x>

In the original publication of the article, the following sentences as the last part of the “Determination of ferulic acid decarboxylase activity” were missing.

The enzymatic activity assays were measured in a reaction mixture composed of 0.05 mL of the extracts (10% v v⁻¹) and 0.45 mL of a solution composed of 0.5 g L⁻¹ of ferulic acid. The reaction mixture was incubated at 30 °C for 15 min. The activity of the ferulic acid decarboxylase enzyme was measured by the reduction of ferulic acid and the formation of 4-vinylguaiaicol (μmol). One unit of activity was defined as the production of 1 μmol of 4-vinylguaiaicol per min. For the determination of specific activity, the total enzyme unit (U) and the biomass weight (g) were considered as a reference for the total cell used in the reaction. Additionally, the total protein was quantified according to the Bradford method, using bovine serum albumin as the standard [26].

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s42770-021-00462-x>

✉ Maitê Bernardo Correia dos Santos
maite.bernardo@unesp.br

¹ Department of Biology, São Paulo State University-UNESP, Institute of Biosciences, Humanities and Exact Sciences, São José do Rio Preto, SP, Brazil

² Department of Chemistry and Environment Science, São Paulo State University-UNESP, Institute of Biosciences, Humanities and Exact Sciences, São José do Rio Preto, SP, Brazil