



Correction to: Removal of a urinary catheter before discontinuation of epidural analgesia is associated with an increased risk of postoperative urinary retention and hospital episode costs in patients undergoing surgical correction for adolescent idiopathic scoliosis

Assem A. Sultan¹ · Ryan J. Berger¹ · William A. Cantrell^{1,2} · Linsen T. Samuel¹ · Erin Ohliger¹ · Joshua Golubovsky² · Salam Bachour² · Selena Pasadyn² · Jaret M. Karnuta² · Pierre Tamer² · Phuc Le³ · Thomas E. Kuivila¹ · David P. Gurd¹ · Ryan C. Goodwin¹

Published online: 2 March 2020
© Scoliosis Research Society 2020

Correction to: Spine Deformity
<https://doi.org/10.1007/s43390-020-00039-y>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original version of this article unfortunately contained a mistake. The spelling of Dr. Erin Ohliger's name was incorrect. The correct name is Erin Ohliger.

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s43390-020-00039-y>.

✉ Assem A. Sultan
assem.sultan@gmail.com; sultana@ccf.org

¹ Department of Orthopaedic Surgery, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195, USA

² Cleveland Clinic Lerner College of Medicine, 9500 Euclid Avenue NA21, Cleveland, OH 44195, USA

³ Center for Value-Based Care Research, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195, USA