

## Correction to: Insecticidal activity, and SAR studies of semicarbazide, thiosemicarbazide, urea and thiourea derivatives against *Spodoptera littoralis* (Boisd.)

Mohamed A. Gad<sup>1</sup> · Esam A. Alqurashi<sup>2</sup> · Nawaf I. Alsenani<sup>2</sup> · Fawy M. Abd El Latif<sup>2,3</sup> · Safwat A. Aref<sup>1</sup> · N. A. Ahmed<sup>1</sup> · Antar A. Abdelhamid<sup>2,4</sup> · Ahmed M. M. El-Saghier<sup>4</sup>

Published online: 19 June 2023

© The Author(s) 2023 **OPEN**

### Correction to:

**Journal of Umm Al-Qura University for Applied Sciences**

<https://doi.org/10.1007/s43994-023-00037-6>

In this article the affiliation details for Author Nawaf I. Alsenani were incorrectly given as 'Research Institute of Plant Protection, Agricultural Research Center, Giza 12112, Egypt' but should have been 'Department of Chemistry, Faculty of Science, Albaha University, Albaha 1988, Saudi Arabia'. The original article has been corrected.

adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing,

The original article can be found online at <https://doi.org/10.1007/s43994-023-00037-6>.

✉ Esam A. Alqurashi, [ealqurashi@bu.edu.sa](mailto:ealqurashi@bu.edu.sa); Mohamed A. Gad, [samy\\_adjey@yahoo.com](mailto:samy_adjey@yahoo.com); Nawaf I. Alsenani, [n.alsenani@bu.edu.sa](mailto:n.alsenani@bu.edu.sa); Fawy M. Abd El Latif, [fabdellatif@bu.edu.sa](mailto:fabdellatif@bu.edu.sa); Safwat A. Aref, [aml.bsmet@yahoo.com](mailto:aml.bsmet@yahoo.com); N. A. Ahmed, [nabilaaliamed8@gmail.com](mailto:nabilaaliamed8@gmail.com); Antar A. Abdelhamid, [Drantar25@yahoo.com](mailto:Drantar25@yahoo.com); Ahmed M. M. El-Saghier, [el\\_saghier@yahoo.com](mailto:el_saghier@yahoo.com) | <sup>1</sup>Research Institute of Plant Protection, Agricultural Research Center, Giza 12112, Egypt. <sup>2</sup>Department of Chemistry, Faculty of Science, Albaha University, Albaha 1988, Saudi Arabia. <sup>3</sup>Chemistry Department, Faculty of Science, Aswan University, Aswan 81528, Egypt. <sup>4</sup>Department of Chemistry, Faculty of Science, Sohag University, Sohag 8252, Egypt.



J.Umm Al-Qura Univ. Appl. Sci. (2023) 9:391 | <https://doi.org/10.1007/s43994-023-00056-3>