



Even Values of Ramanujan's Tau-Function

Jennifer S. Balakrishnan¹ · Ken Ono² · Wei-Lun Tsai²

Received: 1 February 2021 / Revised: 26 May 2021 / Accepted: 8 September 2021 /

Published online: 10 December 2021

© The Author(s), under exclusive licence to Springer Science+Business Media LLC, part of Springer Nature 2021

Abstract

In the spirit of Lehmer's speculation that Ramanujan's tau-function never vanishes, it is natural to ask whether any given integer α is a value of $\tau(n)$. For odd α , Murty, Murty, and Shorey proved that $\tau(n) \neq \alpha$ for sufficiently large n . Several recent papers have identified explicit examples of odd α which are not tau-values. Here we apply these results (most notably the recent work of Bennett, Gherga, Patel, and Siksek) to offer the first examples of even integers that are not tau-values. Namely, for primes ℓ we find that

$$\tau(n) \notin \{\pm 2\ell : 3 \leq \ell < 100\} \cup \{\pm 2\ell^2 : 3 \leq \ell < 100\} \\ \cup \{\pm 2\ell^3 : 3 \leq \ell < 100 \text{ with } \ell \neq 59\}.$$

Moreover, we obtain such results for infinitely many powers of each prime $3 \leq \ell < 100$. As an example, for $\ell = 97$ we prove that

$$\tau(n) \notin \{2 \cdot 97^j : 1 \leq j \not\equiv 0 \pmod{44}\} \cup \{-2 \cdot 97^j : j \geq 1\}.$$

The first author acknowledges the support of NSF Grant (DMS-1702196), the Clare Boothe Luce Professorship (Henry Luce Foundation), a Simons Foundation Grant (Grant #550023), and a Sloan Research Fellowship. The second author thanks the Thomas Jefferson Fund and the NSF (DMS-1601306, DMS-2002265 and DMS-2055118).

✉ Ken Ono
ken.ono691@virginia.edu

Jennifer S. Balakrishnan
jbala@bu.edu

Wei-Lun Tsai
wt8zj@virginia.edu

¹ Department of Mathematics and Statistics, Boston University, Boston, MA 02215, USA

² Department of Mathematics, University of Virginia, Charlottesville, VA 22904, USA

The method of proof applies *mutatis mutandis* to newforms with residually reducible mod 2 Galois representation and is easily adapted to generic newforms with integer coefficients.

Keywords Lehmer’s conjecture · Ramanujan’s tau-function · Newforms · Modular forms

1 Introduction and Statement of Results

Ramanujan’s tau-function [7, 15], the coefficients of the unique normalized weight 12 cusp form for $SL_2(\mathbb{Z})$ (note: $q := e^{2\pi iz}$ throughout)

$$\begin{aligned}\Delta(z) &= \sum_{n=1}^{\infty} \tau(n)q^n \\ &:= q \prod_{n=1}^{\infty} (1 - q^n)^{24} = q - 24q^2 + 252q^3 - 1472q^4 + 4830q^5 - \dots,\end{aligned}\tag{1.1}$$

has been a remarkable prototype in the theory of modular forms. Despite many advances that reveal its deep properties, Lehmer’s conjecture [13] that $\tau(n)$ never vanishes remains open.

In the spirit of this conjecture, it is natural to ask whether any given integer α is a value of $\tau(n)$. Much is known for odd α thanks to the convenient fact that

$$\Delta(z) \equiv \sum_{n=0}^{\infty} q^{(2n+1)^2} \pmod{2}.\tag{1.2}$$

Murty et al. [14] proved that $\tau(n) \neq \alpha$ for sufficiently large n . Craig and the authors [4, 5] proved some effective results concerning potential odd values of $\tau(n)$ and, more generally, coefficients of newforms with residually reducible mod 2 Galois representation. Their methods have been carried further in subsequent work by Amir and Hong [2], Dembner and Jain [11], and Hanada and Madhukara [12]. For example, for $n > 1$, these papers prove that

$$\tau(n) \notin \{\pm 1, \pm 691\} \cup \{\pm \ell : 3 \leq \ell < 100 \text{ prime}\}.\tag{1.3}$$

Recently, Bennett et al. [6] proved a number of spectacular results regarding odd values of $\tau(n)$. For example, they prove (see Theorem 6 of [6]) that $|\tau(n)| \neq \ell^b$, where $3 \leq \ell < 100$ prime and b is a positive integer.

Much less is known for even α . To this end, we make use of lower bounds for the number of prime divisors of tau-values. Craig and the authors proved (see¹ Theorem

¹ Theorem 2.5 of [5] concerns the case of generic newforms with integer coefficients.

1.5 of [5]) that

$$\Omega(\tau(n)) \geq \sum_{\substack{p|n \\ \text{prime}}} (\sigma_0(\text{ord}_p(n) + 1) - 1) \geq \omega(n), \tag{1.4}$$

where $\omega(n)$ (resp. $\Omega(\tau(n))$) is the number of distinct prime factors of n (resp. $\tau(n)$) with multiplicity, and $\sigma_0(N)$ is the number of positive divisors of N . Therefore, if $\tau(n) = \pm 2$, then $n = p^m$, where p and $m + 1$ are both prime.² Similarly, if $\tau(n) = \pm 2\ell$, where ℓ is an odd prime, then this inequality implies that n has at most two distinct prime factors. Moreover, if $n = p_1^{m_1} p_2^{m_2}$, where $p_1 \neq p_2$ are prime and $m_1, m_2 \geq 1$, then $m_1 + 1$ and $m_2 + 1$ are both prime.

Combining these results with the recent work of Bennett et al. [6], we show that certain even numbers never arise as tau-values. To make this precise, we require sets of triples (ℓ, r, t) , where $3 \leq \ell < 100$ is prime and $r \pmod t$ is an arithmetic progression with modulus $t \mid 44$:

$$S^+ := \left\{ \begin{array}{l} (3, 0, 44), (5, 0, 22), (7, 0, 44), (7, 19, 44), (11, 0, 22), (13, 0, 44), (17, 0, 44), \\ (19, 0, 22), (23, 0, 4), (29, 0, 22), (31, 0, 22), (37, 0, 44), (37, 35, 44), (41, 0, 22), \\ (43, 0, 44), (43, 37, 44), (47, 0, 4), (53, 0, 44), (59, 0, 22), (61, 0, 22), (67, 0, 44), \\ (67, 43, 44), (71, 0, 22), (73, 0, 44), (79, 0, 22), (83, 0, 44), (89, 0, 22), (97, 0, 44) \end{array} \right\} \tag{1.5}$$

$$S^- := \{(3, 15, 44), (5, 11, 22), (17, 33, 44), (59, 3, 22), (83, 11, 44), (89, 11, 22)\}. \tag{1.6}$$

Then we define the set of pairs

$$N^\pm := \{(\ell, j) : 1 \leq j \not\equiv r \pmod t \text{ for all } (\ell, r, t) \in S^\pm\}. \tag{1.7}$$

These sets determine values of the form $\pm 2 \cdot \ell^j$ that we rule out as possible even tau-values.

Theorem 1.1 *If $j \geq 1$ and $3 \leq \ell < 100$ is prime, then for every n we have*

$$\tau(n) \notin \{2\ell^j : (\ell, j) \in N^+\} \cup \{-2\ell^j : (\ell, j) \in N^-\}.$$

Moreover, we have that $\tau(n) \notin \{\pm 2 \cdot 691\}$.

Example The triples $(7, r, t) \in S^+$ are $(7, 0, 44)$ and $(7, 19, 44)$. Therefore, Theorem 1.1 gives

$$\tau(n) \notin \{2 \cdot 7^j : j \not\equiv 0, 19 \pmod{44}\}.$$

² In Sect. 2 we shall show that $\tau(n) = \pm 2$ requires that n is prime.

Example Let $\Omega := \{7, 11, 13, 19, 23, 29, 31, 37, 41, 43, 47, 53, 61, 67, 71, 73, 79, 97\}$ be the set of primes $3 \leq \ell < 100$ for which there are no triples of the form $(\ell, r, t) \in S^-$. For these primes, N^- contains (ℓ, j) for every $j \geq 1$, and so Theorem 1.1 gives

$$\tau(n) \notin \{-2\ell^j : \ell \in \Omega \text{ and } j \geq 1\}.$$

As an immediate corollary, we obtain the following conclusion for primes $3 \leq \ell < 100$.

Corollary 1.2 *For every n , we have*

$$\begin{aligned} \tau(n) \notin \{ \pm 2\ell : 3 \leq \ell < 100 \} \cup \{ \pm 2\ell^2 : 3 \leq \ell < 100 \} \\ \cup \{ \pm 2\ell^3 : 3 \leq \ell < 100 \text{ with } \ell \neq 59 \}. \end{aligned}$$

Remark The first examples of $\tau(n) = \pm 2\ell$, where ℓ is prime, are

$$\tau(277) = -2 \cdot 8209466002937 \quad \text{and} \quad \tau(1297) = 2 \cdot 58734858143062873.$$

We note that 277 and 1297 are both prime. Every such value with $n \leq 200,000$ has prime n .

The proof of Theorem 1.1 is a modification of the method employed in [4,5]. These tools are based on the observation that integer sequences of the form $\{1, \tau(p), \tau(p^2), \tau(p^3), \dots\}$, where p is prime, are *Lucas sequences*. Important work of Bilu et al. [8] on primitive prime divisors of Lucas sequences applies to α -variants of Lehmer’s conjecture. Loosely speaking, their work implies that each $\tau(p^m)$ is divisible by at least one prime ℓ for which $\ell \nmid \tau(p)\tau(p^2) \cdots \tau(p^{m-1})$. In [4,5], this property is combined with the theory of newforms to obtain variants of Lehmer’s conjecture. Namely, certain odd integers α are ruled out as tau-values, as well as coefficients of newforms with residually reducible mod 2 Galois representation. Such conclusions follow from the absence of special integer points (X, Y) on specific curves, including hyperelliptic curves and curves defined by Thue equations. These special points (if any) have the property that $X = p$ or p^{2k-1} , where p is prime and $2k$ is the weight of the newform.

In Sect. 2, we recall the main tools from [5] and essential facts about newform coefficients, such as Ramanujan’s tau-function. In Sect. 3 we combine these facts with (1.3), the work of Bennett et al. (i.e. Theorem 6 of [6]), and Ramanujan’s famous tau-congruences to prove Theorem 1.1.

Remark The proof of Theorem 1.1 applies *mutatis mutandis* to integer weight newforms with integer coefficients and residually reducible mod 2 Galois representation. A minor modification holds for arbitrary integer weight newforms $f(z)$ with integer coefficients, regardless of its 2-adic properties. Indeed, suppose that $f(z) = \sum_{n=1}^{\infty} a_f(n)q^n$, and let α be any non-zero integer. We consider the “equation” $a_f(n) = \alpha$. Theorem 2.5 of [5] offers the generalization of (1.4) which constrains the

possible prime factorizations of n ; the number of distinct prime factors of n generally does not exceed $\omega(\alpha)$. By the multiplicativity of newform coefficients, for $d \mid \alpha$, we must solve the equation $a_f(p^m) = d$, where $m \geq 1$, and p is prime. To this end, one applies Theorem 3.2 of [5] which identifies the finitely many m that must be considered. As explained in [5], a solution for p , when $m \geq 2$, requires special integer points on specific curves. In many cases, there are no such points, which leads to restrictions such as those in Theorem 1.1 using the methods employed in [4–6].

2 Nuts and Bolts

Here we recall essential facts about Lucas sequences and properties of newform coefficients.

2.1 Properties of Newforms

We recall basic facts about even integer weight newforms (see [3]), along with the deep theorem of Deligne [9,10] that bounds their Fourier coefficients.

Theorem 2.1 *Suppose that $f(z) = q + \sum_{n=2}^{\infty} a_f(n)q^n \in S_{2k}(\Gamma_0(N))$ is a newform with integer coefficients. Then the following are true:*

- (1) *If $\gcd(n_1, n_2) = 1$, then $a_f(n_1n_2) = a_f(n_1)a_f(n_2)$.*
- (2) *If $p \nmid N$ is prime and $m \geq 2$, then*

$$a_f(p^m) = a_f(p)a_f(p^{m-1}) - p^{2k-1}a_f(p^{m-2}).$$

- (3) *If $p \nmid N$ is prime and α_p and β_p are roots of $F_p(x) := x^2 - a_f(p)x + p^{2k-1}$, then*

$$a_f(p^m) = \frac{\alpha_p^{m+1} - \beta_p^{m+1}}{\alpha_p - \beta_p}.$$

Moreover, we have $|a_f(p)| \leq 2p^{\frac{2k-1}{2}}$, and α_p and β_p are complex conjugates.

2.2 Implications of Properties of Lucas Sequences for Newforms

Suppose that α and β are algebraic integers for which $\alpha + \beta$ and $\alpha\beta$ are relatively prime non-zero integers, where α/β is not a root of unity. Their *Lucas numbers* $\{u_n(\alpha, \beta)\} = \{u_1 = 1, u_2 = \alpha + \beta, \dots\}$ are the integers

$$u_n(\alpha, \beta) := \frac{\alpha^n - \beta^n}{\alpha - \beta}. \tag{2.1}$$

In particular, in the notation of Theorem 2.1, for primes $p \nmid N$ and $m \geq 1$, we have

$$a_f(p^m) = u_{m+1}(\alpha_p, \beta_p) = \frac{\alpha_p^{m+1} - \beta_p^{m+1}}{\alpha_p - \beta_p}. \tag{2.2}$$

The following well-known relative divisibility property is important for the proof of Theorem 1.1.

Proposition 2.2 (Prop. 2.1 (ii) of [8]) *If $d \mid n$, then $u_d(\alpha, \beta) \mid u_n(\alpha, \beta)$.*

To prove Theorem 1.1, we employ bounds on the first occurrence of a multiple of a prime ℓ in a Lucas sequence. We let $m_\ell(\alpha, \beta)$ be the smallest $n \geq 2$ for which $\ell \mid u_n(\alpha, \beta)$. We note that $m_\ell(\alpha, \beta) = 2$ if and only if $\alpha + \beta \equiv 0 \pmod{\ell}$. The following proposition is well known.

Proposition 2.3 (Corollary³ 2.2 of [8]) *If $\ell \nmid \alpha\beta$ is an odd prime with $m_\ell(\alpha, \beta) > 2$, then the following are true.*

- (1) *If $\ell \mid (\alpha - \beta)^2$, then $m_\ell(\alpha, \beta) = \ell$.*
- (2) *If $\ell \nmid (\alpha - \beta)^2$, then $m_\ell(\alpha, \beta) \mid (\ell - 1)$ or $m_\ell(\alpha, \beta) \mid (\ell + 1)$.*

Remark If $\ell \mid \alpha\beta$, then either $\ell \mid u_n(\alpha, \beta)$ for all n , or $\ell \nmid u_n(\alpha, \beta)$ for all n .

A prime $\ell \mid u_n(\alpha, \beta)$ is a *primitive prime divisor* of $u_n(\alpha, \beta)$ if $\ell \nmid (\alpha - \beta)^2 u_1(\alpha, \beta) \cdots u_{n-1}(\alpha, \beta)$. Bilu, Hanrot, and Voutier [8] proved that every Lucas number $u_n(\alpha, \beta)$, with $n > 30$, has a primitive prime divisor. Their work is comprehensive; they have classified *defective* terms, the integers $u_n(\alpha, \beta)$, with $n > 2$, that do not have a primitive prime divisor. Their work, combined with a subsequent paper⁴ by Abouzaid [1], gives the *complete classification* of defective Lucas numbers. In [4,5], these results were applied to even weight newforms, including $\Delta(z)$. Arguing as in these papers, we obtain the following lemma.

Lemma 2.1 *Suppose $2k \geq 4$ is even, and α and β are roots of the integral polynomial*

$$F(X) = X^2 - AX + p^{2k-1} = (X - \alpha)(X - \beta), \tag{2.3}$$

where p is prime, $|A| = |\alpha + \beta| \leq 2p^{\frac{2k-1}{2}}$, and $\gcd(\alpha + \beta, p) = 1$. Then there are no defective Lucas numbers $u_n(\alpha, \beta) \in \{\pm 2\ell^i\}$, where $i \geq 0$ and ℓ is an odd prime. Also, if $u_n(\alpha, \beta) = \pm \ell$ is a defective Lucas number, then one of the following is true.

- (1) *We have $(A, \ell, n) = (\pm m, 3, 3)$, where $3 \nmid m$ and $(p, \pm m)$ satisfies $Y^2 = X^{2k-1} \pm 3$.*
- (2) *We have $(A, \ell, n) = (\pm \ell, \ell, 4)$, where $(p, \pm \ell)$ satisfies $Y^2 = 2X^{2k-1} - 1$.*

Remark Thanks to Lemma 2.1 and (1.4), if $\tau(n) = \pm 2$, then n must be prime.

³ This corollary is stated for Lehmer numbers. The conclusions hold for Lucas numbers because $\ell \nmid (\alpha + \beta)$.

⁴ This paper included a few cases that were omitted in [8].

Proof As mentioned above, [1,8] classify defective Lucas numbers. This classification includes a finite list of sporadic examples and a list of parameterized infinite families. Theorem 2.2 of [5] uses these results to describe the defective Lucas numbers that can arise as newform coefficients, i.e. sequences defined by (2.3). Tables 1 and 2 of [5] list the possible defective cases.

An inspection of Table 1 of [5], which concerns the sporadic examples, reveals that the only possible defective numbers with $2k \geq 4$ have $2k = 4$. Moreover, they are the odd numbers $u_3(\alpha, \beta) = 1$ or $u_4(\alpha, \beta) = \pm 85$.

To complete the proof, we consider the parametrized infinite families in Table 2 of [5]. If $u_n(\alpha, \beta)$ is even, then we only have to consider rows four, five, six, and seven of the table. A simple inspection reveals that $\{\pm 2\ell^i\}$ for $i \geq 0$ never arises. This then leaves $u_n(\alpha, \beta) = \pm \ell$ as the only cases to consider. However, Lemma 2.1 of [5] includes these cases, giving (1) and (2) above. \square

3 Proof of Theorem 1.1

Here we use the previous section to prove Theorem 1.1.

3.1 Ramanujan’s Congruences

Ramanujan’s classical congruences for the tau-function imply the following convenient fact involving the sets N^ε defined in (1.7).

Lemma 3.1 *If $3 \leq \ell < 100$ is prime and $(\ell, j) \in N^\varepsilon$, then for every prime p we have that*

$$\tau(p) \neq \varepsilon 2\ell^j.$$

Proof We recall the famous Ramanujan congruences (see [7,15]):

$$\tau(n) \equiv \begin{cases} n^3\sigma_1(n) & (\text{mod } 4), \\ n^2\sigma_1(n) & (\text{mod } 3), \\ n\sigma_1(n) & (\text{mod } 5), \\ n\sigma_3(n) & (\text{mod } 7). \end{cases}$$

where $\sigma_v(n) := \sum_{1 \leq d|n} d^v$. Furthermore, if $p \neq 23$ is prime, Ramanujan proved that

$$\tau(p) \equiv \begin{cases} 0 & (\text{mod } 23) & \text{if } \left(\frac{p}{23}\right) = -1, \\ \sigma_{11}(p) & (\text{mod } 23^2) & \text{if } p = a^2 + 23b^2 \text{ with } a, b \in \mathbb{Z}, \\ -1 & (\text{mod } 23) & \text{otherwise.} \end{cases}$$

If $p \neq 23$ is prime, then the collection of these congruences imply

$$\begin{aligned} \tau(p) &\equiv 0 \pmod{2}, \quad \tau(p) \equiv 0, 2 \pmod{3}, \quad \tau(p) \equiv 0, 1, 2 \pmod{5}, \\ \tau(p) &\equiv 0, 1, 2, 4 \pmod{7}, \quad \text{and} \quad \tau(p) \equiv 0, -1, 2 \pmod{23}. \end{aligned}$$

These congruences are easily reformulated in terms of N^ε . This completes the proof for $p \neq 23$. Finally, we note that $\tau(23) = 18643272 = 2^3 \cdot 3 \cdot 617 \cdot 1259$. \square

3.2 Proof of Theorem 1.1

Theorem 1.1 consists of two different types of α .

- (1) The case where $\alpha = \pm 2\ell$, where $3 \leq \ell \leq 100$ is prime or $\ell = 691$.
- (2) The case where $\alpha = \pm 2\ell^j$, where $3 \leq \ell \leq 100$ is prime and $j \geq 2$.

By Lemma 2.1 with $2k = 12$, the numbers $\{\pm 2\ell^i\}$ for $i \geq 0$ (if they arise) are never defective Lucas numbers in $\{\tau(p), \tau(p^2), \tau(p^3), \dots\}$, where p is prime. Lemma 2.1 (1) and (2) covers the cases apart from $\pm\ell$, which were ruled out by Lemma 2.1 of [4].

Case (1). Thanks to (1.4), if $\tau(n) = \pm 2\ell$, where ℓ is an odd prime, then either $n = p_1^{m_1}$, or $n = p_1^{m_1} p_2^{m_2}$, where the p_i are prime and the $m_i \geq 1$. Using Theorem 2.1 (1) and (1.3), the latter case requires $|\tau(p_1^{m_1})| = 2$ and $|\tau(p_2^{m_2})| = \ell$. Thanks to (1.3) again, this is impossible for $\ell = 691$ and primes $3 \leq \ell < 100$.

Therefore, we may assume that $\tau(p_1^{m_1}) = \pm 2\ell$. Thanks to Theorem 2.1, we have that $p_1 \neq 2$, as $4 \mid \tau(2^m)$ for every positive integer m . Therefore, (1.2) implies that m_1 is odd. Moreover, since $\tau(p_1)$ is even, it must be that $\tau(p_1^{m_1})$ is the first term in the Lucas sequence that is divisible by ℓ . Otherwise, $\pm 2\ell$ would be defective, contradicting Lemma 2.1. If $m_1 + 1$ has a non-trivial divisor other than 2, then by the relative divisibility of Lucas numbers given in Proposition 2.2, and the nondefectivity of ± 2 in Lemma 2.1, we obtain a contradiction. Hence, it boils down to considering the case when $m_1 + 1 = 4$, $\tau(p) = \pm 2$, and $\tau(p^3) = \pm 2\ell$ for some prime p . However, using the Hecke relation ((2) in Theorem 2.1), we have that $\pm 2\ell = \pm 4(p^{11} - 2)$, and there is no such p , as the left hand side is $2 \pmod{4}$ while the right hand side is $0 \pmod{4}$. Hence, it follows that $m_1 + 1$ is prime. Therefore, we have $m_1 = 1$, which in turn leads to $\tau(p_1) = \pm 2\ell$. The proof in this case is complete as Lemma 3.1 shows that $\tau(p) \neq \pm 2\ell$.

Case (2). Since $3 \leq \ell < 100$ is prime, (1.3) and Theorem 6 of [6] implies that $|\tau(n)| \neq \ell^b$ for all n and $b \geq 1$. Therefore, we may assume that $\tau(p^m) = \pm 2\ell^j$, where p is an odd prime. Here we again use the fact that $4 \mid \tau(2^m)$ for every positive integer m , and consider the degenerate case $m_1 + 1 = 4$, $\tau(p) = \pm 2$, and $\tau(p^3) = \pm 2\ell^j$ for some prime p , which gives the corresponding equation $\pm 2\ell^j = \pm 4(p^{11} - 2)$. The argument in Case (1), where the conclusion is that $m = 1$, applies *mutatis mutandis*. Therefore, the proof is complete as Lemma 3.1 shows that $\tau(p) \neq \pm 2\ell^j$.

Acknowledgements We thank Matthew Bisatt for several helpful discussions about root numbers of Jacobians of hyperelliptic curves which were part of initial discussions related to this project. We are indebted to the referees who made suggestions that substantially improved the results in this paper. Finally, we thank Kaya Lakein and Anne Larsen for comments that improved the exposition in this paper.

References

1. Abouzaid, M.: Les nombres de Lucas et Lehmer sans diviseur primitif. *J. Théor. Nombres Bordeaux* **1**(8), 299–313 (2006)
2. Amir, M., Hong, L.: On L -functions of modular elliptic curves and certain $K3$ surfaces. *Ramanujan J.* [arXiv:2007.09803](https://arxiv.org/abs/2007.09803) (accepted for publication)
3. Atkin, A.O.L., Lehner, J.: Hecke operators on $\Gamma_0(m)$. *Math. Ann.* **185**, 134–160 (1970)
4. Balakrishnan, J. S., Craig, W., Ono, K.: Variations of Lehmer’s conjecture for Ramanujan’s tau-function. *J. Numb. Theory (Prime)*. [arXiv:2005.10345](https://arxiv.org/abs/2005.10345) (accepted for publication)
5. Balakrishnan, J. S., Craig, W., Ono, K., Tsai, W.-L.: Variants of Lehmer’s speculation for newforms. [arXiv:2005.10354](https://arxiv.org/abs/2005.10354) (2020)
6. Bennett, M. A., Gherga, A., Patel, V., Siksek, S.: Odd values of the Ramanujan tau function. [arXiv:2101.02933](https://arxiv.org/abs/2101.02933). (2021)
7. Berndt, B.C., Ono, K.: Ramanujan’s unpublished manuscript on the partition and tau functions with proofs and commentary. *Sém. Lothar. Combin.* **42**, Art. B42c (1999)
8. Bilu, Y., Hanrot, G., Voutier, P.M.: Existence of primitive divisors of Lucas and Lehmer numbers. *J. Reine Angew. Math.* **539**, 75–122 (2001)
9. Deligne, P.: La conjecture de Weil. I. *Publ. Math. IHES* **4**(3), 273–307 (1974)
10. Deligne, P.: La conjecture de Weil. II. *Publ. Math. IHES* **5**(2), 137–252 (1980)
11. Dembner, S., Jain, V.: Hyperelliptic curves and newform coefficients. *J. Numb. Theory*. [arXiv:2007.08358](https://arxiv.org/abs/2007.08358) (accepted for publication)
12. Hanada, M., Madhukara, R.: Fourier coefficients of level 1 Hecke Eigenforms. *Acta Arith.* ([arXiv:2007.08683](https://arxiv.org/abs/2007.08683)) (accepted for publication)
13. Lehmer, D.H.: The vanishing of Ramanujan’s $\tau(n)$. *Duke Math. J.* **1**(4), 429–433 (1947)
14. Murty, V.K., Murty, R., Shorey, T.N.: Odd values of the Ramanujan tau function. *Bull. Soc. Math. France* **115**, 391–395 (1987)
15. Ramanujan, S.: On certain arithmetical functions. *Trans. Camb. Philos. Soc.* **22**(9), 159–184 (1916)