
A Chemical Ionization Study of Deuteron Transfer Initiated Propene Loss from Propoxyppyridines

Eric S. E. van Beelen, Tineke A. Molenaar-Langeveld,
and Steen Ingemann

Faculty of Science, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam,
The Netherlands

The mechanism of propene loss from the metastable $[M + D]^+$ ions of isomeric 2-, 3-, and 4-*n*-propoxyppyridines and the related isopropoxyppyridines has been examined by chemical ionization (CI) and tandem mass spectrometry in combination with deuterium labeling. The $[M + D]^+$ ions were generated with CD_3OD , CD_3CN , $(CD_3)_2CO$, or pyrrole- D_5 (listed in order of increasing proton affinity) as the CI reagent. The results reveal that the deuteron added in the CI process is not interchanged with the hydrogen atoms of the propyl group prior to propene loss from the metastable $[M + D]^+$ ions of the propoxyppyridines. The site selective labeling of the α -, β -, or γ -position of the propyl group indicates that the $[M + D]^+$ ions of 2-*n*-propoxyppyridine expel propene with formation of an ion-neutral complex composed of a propyl carbenium ion and 2-pyridone. By contrast, the $[M + D]^+$ ions of 3-*n*-propoxyppyridine expel propene by: (1) Formation of ion-neutral complexes, and (2) a conventional 1,5-hydride shift from the β -position of the *n*-propyl group to the ring and/or a 1,2-elimination type process. For the 4-isomer, the results suggest the occurrence of propene loss by a 1,2-elimination in addition to the intermediate formation of ion-neutral complexes. Loss of propene with one deuterium atom is the only reaction of the $[M + D]^+$ ions of the isopropoxyppyridines labeled at the α -position of the isopropyl group. The results for the isopropoxyppyridines labeled with three deuterium atoms at the β -position are consistent with: (1) The loss of propene by ion-neutral complex formation and the occurrence of a substantial isotope effect in the subsequent proton/deuteron transfer within the complex, and/or (2) the loss of propene by a 1,2-elimination type reaction. (J Am Soc Mass Spectrom 2003, 14, 1166–1176) © 2003 American Society for Mass Spectrometry

Site-specific proton affinities (PA) and the occurrence of proton transfer reactions in the absence of solvent molecules are of fundamental importance in mass spectrometry [1–6].

The site of protonation depends not only on the proton affinity of the distinct sites in a molecule, but also on the existence of possible energy barriers for the proton transfer reaction [7–10]. In particular, proton transfer to the aromatic ring can be hindered by a kinetic barrier even though this entity of a substituted benzene is often associated with a higher proton affinity than a substituent such as an OH or OR group [11–18].

Insight into the molecular properties that determine the regioselectivity of proton transfer reactions is also of importance for studies concerned with structure elucidation with the use of Chemical Ionization [1]. This is particularly pertinent as it is often observed that the

initial step in protonation of an organic compound is related directly to the dissociation reactions of the $[M + H]^+$ ions. Furthermore, polyfunctional organic species can show a complex behavior as protonation can occur at distinct sites and nonetheless lead to dissociation by a single process. For example, the metastable $[M + H]^+$ ions of phenyl *n*-propyl ether are known to dissociate exclusively by propene loss irrespective of the fact that protonation can occur at the phenyl ring as well as at the oxygen atom of the substituent [16, 17, 19–23]. As emphasized in several CI studies of aryl *n*-propyl ethers with deuterium labeled reagents, the metastable $[M + D]^+$ ions expel propene molecules that contain the deuteron transferred from the conjugate acid of the CI reagent to the ether in the ion source [16, 17, 19–22]. In previous reports, the incorporation of the deuterium atom from the CI reagent was suggested to involve initial deuteron transfer to the oxygen atom of phenyl *n*-propyl ether and fluorine or methyl substituted analogues [16, 17]. By contrast, deuteron transfer to the aromatic ring was proposed to result in the loss of a

Published online August 11, 2003

Address reprint requests to Dr. S. Ingemann, Faculty of Science, Swammerdam Institute for Life Sciences, University of Amsterdam, Nieuwe Achtergracht 166, Amsterdam 1018 WV, The Netherlands. E-mail: ingemann@science.uva.nl

propene molecule that contained only the hydrogen atoms of the original propyl group.

In order to obtain more insight into the influence of the initial protonation site on the mechanism of alkene elimination from protonated aromatic ethers, we decided to examine propene loss from a series of isomeric propoxy-pyridines. The propoxy-pyridines were chosen as model compounds because it can be anticipated that protonation at the nitrogen atom of the ring is preferred considerably over protonation at the oxygen atom of the substituent. This expectation is corroborated by the calculated proton affinity of the nitrogen atom in 2-hydroxypyridine ($PA = 931 \text{ kJ mol}^{-1}$), which is 183 kJ mol^{-1} higher than the calculated value of the oxygen atom (748 kJ mol^{-1}) [24]. In other words, the propoxy-pyridines offer the possibility to study the loss of propene from aromatic ethers that may be protonated only at the heteroatom of the ring. In addition, the selection of 2-, 3- and 4-propoxy-pyridines allows an examination of influence of the distance between the protonation site and the substituent on the mechanism of propene loss.

In the studies concerned with propene loss from $[M + D]^+$ ions of phenyl *n*-propyl ethers and their derivatives, a number of CI reagent gasses were used in order to generate the ionic species [16, 17]. In these studies it was shown that the incorporation of the deuterium atom from the CI reagent in the propene molecule is enhanced if the CI reagent gas has a relatively high proton affinity. For example, the metastable $[M + D]^+$ ions of phenyl *n*-propyl ether expel C_3H_5D and C_3H_6 in ratio of 18:82 if CD_3OD ($PA = 754 \text{ kJ mol}^{-1}$) [25] is the CI reagent, whereas the ratio between C_3H_5D and C_3H_6 loss is 27:73 with CD_3CN ($PA = 779 \text{ kJ mol}^{-1}$) [25] as the CI gas [16]. With the purpose of examining the possible influence of the dynamics of the initial deuterium transfer to the propoxy-pyridines on the ensuing loss of propene, we decided to form the metastable $[M + D]^+$ ions of the propoxy-pyridines with CD_3OD , CD_3CN , $(CD_3)_2CO$ or pyrrole- D_5 (listed in order of increasing proton affinity) as the reagent gas.

Experimental

The Chemical Ionization (CI) spectra and the mass-analyzed ion kinetic energy (MIKE) [26, 27] spectra of the (un)labeled propoxy-pyridines were recorded with the use of a Micromass (Manchester, UK) VG ZAB-HFqQ reverse geometry double focusing hybrid mass spectrometer [28, 29]. The CI reagent (CD_3OD , CD_3CN , $(CD_3)_2CO$, or pyrrole- D_5) were mixed with a propoxy-pyridine in a volume ratio of 9:1. The binary chemical sample was introduced into a combined electron ionization/chemical ionization (EI/CI) source through a heated septum inlet system (temperature $\sim 175^\circ\text{C}$) until the pressure was in the range of 10^{-5} – 10^{-4} mbar as measured by an ionization gauge placed in a side arm to the entrance of the diffusion pump situated beneath the ion source housing. A few experiments were also

performed at a different total pressure (between 10^{-6} mbar and 10^{-4} mbar) of the CI reagent and the propoxy-pyridine in the combined EI/CI source. The ion source parameters were: Electron energy 70 eV, temperature 175 – 200°C , ion repeller potential 0–2 V, and acceleration voltage 8 kV. Collision-induced Dissociation (CID) experiments were performed with helium as the collision gas and with the reduction of the main beam intensity to 50%.

Materials and Synthesis

The unlabeled 2-, 3-, and 4-hydroxypyridines, $CH_3CH_2CD_2OH$, $CD_3CD_2CD_2OD$, $CH_3CD(OH)CH_3$ and $CD_3CD(OD)CD_3$ were commercially available and used without further purification. The labeled propanols were reacted with 48% HBr in water in order to obtain the related bromopropanes. The unlabeled 2-, 3-, and 4-propoxy-pyridines and their deuterated analogues were prepared by reacting the 2-, 3-, and 4-hydroxypyridines with sodium ethanolate followed by reaction with an appropriately labeled 1- or 2-bromopropane [30, 31].

$CH_3CD_2CH_2OH$

1-propanol labeled at the 2-position was prepared by repeated hydrogen-deuterium exchange of methyl malonic acid with D_2O to afford $DOOCCD(CH_3)COOD$. Decarboxylation of one of the carboxylic groups leading to CH_3CD_2COOH was followed by reduction of the other carboxylic group by $LiAlH_4$ to give $CH_3CD_2CH_2OH$ [32].

$CD_3CH_2CH_2OH$

1-propanol labeled at the 3-position was prepared by treating $HOOCCCH_2COOH$ with NaH followed by reaction with CD_3I . Decarboxylation and reduction as described above gave $CD_3CH_2CH_2OH$.

$CD_3-CH(OH)-CH_3$

Grignard reaction of CD_3MgI with freshly distilled acetaldehyde gave the appropriately labeled 2-propanol.

All ethers were purified by preparative gas chromatography (column: Reoplex 400, temperature 125°C). The identity of the ethers as well as the labeling content ($>99\%$) was determined by 1H NMR and EI mass spectrometry.

Results

Chemical Ionization of the propoxy-pyridines with one of the selected reagents (CD_3OD , CD_3CN , $(CD_3)_2CO$ and Pyrrole- D_5) leads in the ion source to relatively abundant $[M + D]^+$ ions. The $[M + D]^+$ ions of the unlabeled propoxy-pyridines react on the μs time-scale only by the loss of C_3H_6 with the formation of $C_5H_5DNO^+$







Figure 1. Reactions of the metastable $[M + D]^+$ ions of 2-*n*-propoxy pyridine labeled with two deuterium atoms at the β -position of the propyl group.

ions (eq 1) irrespective of the CI gas. The exclusive loss of C_3H_6 from the metastable $[M + D]^+$ ions of the unlabeled ethers reveals that the deuterium atom from the reagent is not involved directly in the dissociation reaction. This is further corroborated by the sole occurrence of the loss of C_3D_6 from the metastable $[M + D]^+$ ions of propoxy pyridines with a perdeuterated propyl group (vide infra).




n-Propoxy pyridines

Site-specific deuterium labeled *n*-propoxy pyridines were examined subsequently in order to obtain more insight into the origin of the H atom transferred from the propyl group to the remaining part of the ionized species prior to the expulsion of propene. The presence of two D atoms at the α -position of the 2-*n*-propoxy pyridines is observed to lead to the losses of C_3H_5D and $C_3H_4D_2$ from the metastable $[M + D]^+$ ions (eq 2). The average ratio between the losses of C_3H_5D and $C_3H_4D_2$ is determined to be 24:76 by measuring the area of the Gaussian shaped metastable peaks (see Figure 1).




The same ratio between the losses of C_3H_5D and $C_3H_4D_2$ is observed for the $[M + D]^+$ ions of the $[\beta, \beta-D_2]$ -2-*n*-propoxy pyridines as for the $[\alpha, \alpha-D_2]$ -labeled species (see Table 1). For the $[\gamma, \gamma, \gamma-D_3]$ -labeled species, the loss of a propene molecule with one D atom less than in the propyl group in the parent species is relatively pronounced; that is, $C_3H_4D_2$ and $C_3H_3D_3$ are expelled in a ratio of 33:67 from the metastable $[M + D]^+$ ions.

The introduction of two deuterium atoms at the α -position of the 3-*n*-propoxy pyridines results in the losses of C_3H_5D and $C_3H_4D_2$ in an average ratio of 21:79 (Table 2). For the related $[\beta, \beta-D_2]$ -labeled ions (eq 3), loss of C_3H_5D is more pronounced than observed for the $[\alpha, \alpha-D_2]$ -labeled species.



The results for the loss of propene from the metastable $[M + D]^+$ ions of 4-*n*-propoxy pyridine are dependent on the position of the labeling (Table 3). For example, deuterium incorporation at the α -position results in a ratio of 22:78 between the losses of C_3H_3D and $C_3H_4D_2$ from the metastable ions, whereas the ratio between these losses is 38:62 if two D atoms are present at the β -position (eq 4).



Isopropoxy pyridines

In order to investigate the influence of the structure of the propyl group on the process of propene loss, a series of $[\beta, \beta, \beta-D_3]$ -labeled isopropoxy pyridines were studied. In contrast to the results for the *n*-propoxy pyridines, the ratio between the losses of $C_3H_4D_2$ and $C_3H_3D_3$ from the metastable $[M + D]^+$ ions of the isopropoxy pyridines appeared to be independent of the position of the substituent with re-

Table 1. Relative extent (in %) of the losses of C₃H₆, C₃H₅D, C₃H₄D₂, C₃H₃D₃ and C₃D₆ from the metastable [M + D]⁺ ions of 2-*n*-propoxypropyridine and deuterium labeled analogues

Compound M	Reactant ion	Cl reagent	Relative extent of the loss of ^a				
			C ₃ H ₆	C ₃ H ₅ D	C ₃ H ₄ D ₂	C ₃ H ₃ D ₃	C ₃ D ₆
2-(C ₅ H ₄ N)-OCH ₂ CH ₂ CH ₃ ^b	[M + D] ⁺	CD ₃ OD	100				
		CD ₃ CN	100				
		(CD ₃) ₂ C=O	100				
		Pyrrole-D ₅	100				
2-(C ₅ H ₄ N)-OCD ₂ CH ₂ CH ₃ ^c	[M + D] ⁺	CD ₃ OD		24	76		
		CD ₃ CN		23	77		
		(CD ₃) ₂ C=O		25	75		
		Pyrrole-D ₅		26	74		
2-(C ₅ H ₄ N)-OCH ₂ CD ₂ CH ₃ ^c	[M + D] ⁺	CD ₃ OD		24	76		
		CD ₃ CN		25	75		
		(CD ₃) ₂ C=O		24	76		
		Pyrrole-D ₅		23	77		
2-(C ₅ H ₄ N)-OCH ₂ CH ₂ CD ₃ ^c	[M + D] ⁺	CD ₃ OD				32	68
		CD ₃ CN				33	67
		(CD ₃) ₂ C=O				33	67
		Pyrrole-D ₅				33	67
2-(C ₅ H ₄ N)-OCD ₂ CD ₂ CD ₃ ^d	[M + D] ⁺	CD ₃ OD					100
		CD ₃ CN					100
		(CD ₃) ₂ C=O					100
		Pyrrole-D ₅					100

^aThe ratios are considered to be accurate to within a few percent.^bPropene loss from the [M + D]⁺ ions yields only C₅H₅DNO⁺ (*m/z* 97) ions.^cPropene loss from the [M + D]⁺ ions yields C₅H₅DNO⁺ (*m/z* 97) and C₅H₅D₂NO⁺ (*m/z* 98) ions.^dPropene loss from the [M + D]⁺ ions yields only C₅H₄D₂NO⁺ (*m/z* 98) ions.

spect to the nitrogen atom (see Table 4 and eq 5). In addition, deuterium incorporation at the α -position of the isopropoxy group revealed that the α -hydrogen atom is not involved in the loss of propene.

**Table 2.** Relative extent (in %) of the losses of C₃H₆, C₃H₅D, C₃H₄D₂, C₃H₃D₃ and C₃D₆ from the metastable [M + D]⁺ ions of 3-*n*-propoxypropyridine and deuterium labeled analogues

Compound M	Reactant ion	Cl reagent	Relative extent of the loss of ^a				
			C ₃ H ₆	C ₃ H ₅ D	C ₃ H ₄ D ₂	C ₃ H ₃ D ₃	C ₃ D ₆
3-(C ₅ H ₄ N)-OCH ₂ CH ₂ CH ₃ ^b	[M + D] ⁺	CD ₃ OD	100	0			
		CD ₃ CN	100	0			
		(CD ₃) ₂ C=O	100	0			
		Pyrrole-D ₅	100	0			
3-(C ₅ H ₄ N)-OCD ₂ CH ₂ CH ₃ ^c	[M + D] ⁺	CD ₃ OD		21	79		
		CD ₃ CN		20	80		
		(CD ₃) ₂ C=O		21	79		
		Pyrrole-D ₅		21	79		
3-(C ₅ H ₄ N)-OCH ₂ CD ₂ CH ₃ ^c	[M + D] ⁺	CD ₃ OD		46	54		
		CD ₃ CN		46	54		
		(CD ₃) ₂ C=O		48	52		
		Pyrrole-D ₅		46	54		
3-(C ₅ H ₄ N)-OCH ₂ CH ₂ CD ₃ ^{c,d}	[M + D] ⁺	CD ₃ OD				22	78
		CD ₃ CN				18	82
		(CD ₃) ₂ C=O				14	86
		Pyrrole-D ₅				22	78
3-(C ₅ H ₄ N)-OCD ₂ CD ₂ CD ₃ ^e	[M + D] ⁺	CD ₃ OD					100
		(CD ₃) ₂ C=O					100
		Pyrrole-D ₅					100

^aThe ratios are considered to be accurate to within a few percent.^bPropene loss from the [M + D]⁺ ions yields only C₅H₅DNO⁺ (*m/z* 97) ions.^cPropene loss from the [M + D]⁺ ions yields C₅H₅DNO⁺ (*m/z* 97) and C₅H₄D₂NO⁺ (*m/z* 98) ions.^dThe results for the [γ,γ,γ -D₃] labeled ions as based upon peak areas are considered to be less accurate than the other values in the table due to enhanced overlap of the metastable peaks.^ePropene loss from the [M + D]⁺ ions yields only C₅H₄D₂NO⁺ (*m/z* 98) ions.

Table 3. Relative extent (in %) of the losses of C₃H₆, C₃H₅D, C₃H₄D₂, C₃H₃D₃ and C₃D₆ from the metastable [M + D]⁺ ions of 4-*n*-propoxyppyridine and deuterium labeled analogues


Compound M	Reactant ion	CI reagent	Relative extent of the loss of ^a				
			C ₃ H ₆	C ₃ H ₅ D	C ₃ H ₄ D ₂	C ₃ H ₃ D ₃	C ₃ D ₆
4-(C ₅ H ₄ N)-OCH ₂ CH ₂ CH ₃ ^b	[M + D] ⁺	CD ₃ OD	100	0			
		CD ₃ CN	100	0			
		(CD ₃) ₂ C=O	100	0			
		Pyrrole-D ₅	100	0			
4-(C ₅ H ₄ N)-OCD ₂ CH ₂ CH ₃ ^c	[M + D] ⁺	CD ₃ OD		21	79		
		CD ₃ CN		21	79		
		(CD ₃) ₂ C=O		23	77		
		Pyrrole-D ₅		22	78		
4-(C ₅ H ₄ N)-OCH ₂ CD ₂ CH ₃ ^c	[M + D] ⁺	CD ₃ OD		38	62		
		CD ₃ CN		39	61		
		(CD ₃) ₂ C=O		37	63		
		Pyrrole-D ₅		34	66		
4-(C ₅ H ₄ N)-OCH ₂ CH ₂ CD ₃ ^c	[M + D] ⁺	CD ₃ OD				74	
		CD ₃ CN				73	
		(CD ₃) ₂ C=O				71	
		Pyrrole-D ₅				72	
4-(C ₅ H ₄ N)-OCD ₂ CD ₂ CD ₃ ^d	[M + D] ⁺	CD ₃ OD					100
		CD ₃ CN					100
		(CD ₃) ₂ C=O					100
		Pyrrole-D ₅					100

^aThe ratios are considered accurate to within a few percent.

^bPropene loss from the [M + D]⁺ ions yields only C₅H₅DNO⁺ (*m/z* 97) ions.

^cPropene loss from the [M + D]⁺ ions yields C₅H₅DNO⁺ (*m/z* 97) and C₅H₄D₂NO⁺ (*m/z* 98) ions.

^dPropene loss from the [M + D]⁺ ions yields only C₅H₄D₂NO⁺ (*m/z* 98) ions.



For the isopropoxyppyridines, the results for propene loss from the metastable ions are independent of the nature of the CI gas. In other words the initial transfer

of a deuteron from the Brønsted acid of the CI gas has no influence on the origin of the H or D atoms incorporated in the ionic product of propene loss from the [M + D]⁺ ions. Similar results were obtained for the *n*-propoxyppyridines (see Tables 1–3). In addition, the same ratio between the losses of C₃H₅D and C₃H₄D₂ from the metastable [M + D]⁺ ions of [β,β-D₂]-3-*n*-propoxyppyridines is obtained for a total pressure of 5 × 10⁻⁶ mbar and at a pressure of 1 × 10⁻⁴ mbar in the ion source (see also Experimental).

Table 4. Relative extent (in %) of the losses of C₃H₆, C₃H₅D, C₃H₄D₂, C₃H₃D₃ and C₃D₆ from the metastable [M + D]⁺ ions of isopropoxyppyridine and deuterium labeled analogues

Compound M	Reactant ion	Relative extent of the loss of ^a				
		C ₃ H ₆	C ₃ H ₅ D	C ₃ H ₄ D ₂	C ₃ H ₃ D ₃	C ₃ D ₆
2-(C ₅ H ₄ N)-OCH(CH ₃) ₂ ^b	[M + D] ⁺	100				
2-(C ₅ H ₄ N)-OCD(CD ₃) ₂ ^c	[M + D] ⁺					100
2-(C ₅ H ₄ N)-OCD(CH ₃) ₂ ^b	[M + D] ⁺		100			
2-(C ₅ H ₄ N)-OCH(CD ₃)(CH ₃) ^d	[M + D] ⁺			31	69	
3-(C ₅ H ₄ N)-OCH(CH ₃) ₂ ^b	[M + D] ⁺	100				
3-(C ₅ H ₄ N)-OCD(CD ₃) ₂ ^b	[M + D] ⁺					100
3-(C ₅ H ₄ N)-OCD(CH ₃) ₂ ^b	[M + D] ⁺		100			
3-(C ₅ H ₄ N)-OCH(CD ₃)(CH ₃) ^d	[M + D] ⁺			34	66	
4-(C ₅ H ₄ N)-OCH(CH ₃) ₂ ^b	[M + D] ⁺	100				
4-(C ₅ H ₄ N)-OCD(CD ₃) ₂ ^b	[M + D] ⁺					100
4-(C ₅ H ₄ N)-OCD(CH ₃) ₂ ^b	[M + D] ⁺		100			
4-(C ₅ H ₄ N)-OCH(CD ₃)(CH ₃) ^d	[M + D] ⁺			34	66	

^aThe relative abundances are independent of the CI gas used (see Results section) and accurate to within a few percent.

^bPropene loss from the [M + D]⁺ ions yields only C₅H₅DNO⁺ (*m/z* 97) ions.

^cPropene loss from the [M + D]⁺ ions yields only C₅H₄D₂NO⁺ (*m/z* 98) ions.

^dPropene loss from the [M + D]⁺ ions yields C₅H₅DNO⁺ (*m/z* 97) and C₅H₄D₂NO⁺ (*m/z* 98) ions.

Collision-Induced Dissociation

A series of CID experiments [33] were performed with the $[M + D]^+$ ions of the propoxy pyridines (see Experimental). The collisionally induced loss of propene from the $[M + D]^+$ ions of the unlabeled 2-, 3-, and 4-*n*-propoxy pyridines involves only the expulsion of C_3H_6 . For the $[M + D]^+$ ions of the $[\alpha, \alpha-D_2]$ - or $[\beta, \beta-D_2]$ -labeled 2-, 3-, and 4-*n*-propoxy pyridines, the ratio between the losses of C_3H_5D and $C_3H_4D_2$ is essentially the same in the CID experiments as observed for the metastable ions. Notably, the ratio between the losses of C_3H_5D and $C_3H_4D_2$ in the CID experiments is 75:25 for the $[M + D]^+$ ions of the $[\beta, \beta-D_2]$ -2-*n*-propoxy pyridine and about 40:60 for the $[\beta, \beta-D_2]$ -3-*n*-propoxy pyridine (see also Tables 1 and 2). Similar results are obtained for the competing losses of $C_3H_4D_2$ and $C_3H_3D_3$ from the $[M + D]^+$ ions of the $[\gamma, \gamma, \gamma-D_3]$ -labeled *n*-propoxy pyridines. It should be mentioned, however, that propene loss from the metastable $[M + D]^+$ ions contribute to the intensities of the peaks in the CID spectra and—a priori—it is not possible to separate the collisionally induced processes from the spontaneous reactions of the ions. The relative peak intensities in the CID spectra, therefore, may not necessarily provide a unambiguous measure of the relative extent of the collisionally induced losses of C_3H_5D and $C_3H_4D_2$ from the $[M + D]^+$ ions of, for example, the $[\beta, \beta-D_2]$ -3-*n*-propoxy pyridine.

The structures of the product ions of propene loss were also examined by high kinetic energy CID experiments. These experiments were performed with the product ions of propene loss as formed to a minor extent in the ion-source reactions of the $[M + D]^+$ ions of the unlabeled propoxy pyridines. The spectrum obtained for the $C_5H_5DNO^+$ ions generated by C_3H_6 loss from the $[M + D]^+$ ion of 3-*n*-propoxy pyridine is shown in Figure 2a and appears to very similar to the CID spectrum of the $[M + D]^+$ ion of 3-hydroxy pyridine (see Figure 2b). Comparable results were obtained for the other product ions of propene loss from the ionized propoxy pyridines and the related protonated hydroxy pyridines.

Discussion

The dissociation reactions of ions in the gas phase often involve the intermediate formation of ion-neutral complexes held together by electrostatic interactions and/or hydrogen bonds [21, 34–40]. The involvement of ion-neutral complexes is generally manifested in the energetics of the processes and the loss of the positional identity of, for example, hydrogen atoms prior to dissociation. For example, the metastable ions formed by proton transfer to *n*-propyl benzene under CI conditions are reported to isomerize irreversibly to protonated isopropylbenzene prior to propene loss [36, 41, 42]. The suggested pathway involves cleavage of the bond to the *n*-propyl group concomitant with a 1,2-hydride shift in the incipient carbenium ion. The ion-neutral





Figure 2. (a) The Collision-induced Dissociation mass spectrum of the $C_5H_5DNO^+$ ion formed by C_3H_6 loss from the metastable $[M + D]^+$ ions of 3-*n*-propoxy pyridine. (b) The CID spectrum of the $[M + D]^+$ ion of 3-hydroxy pyridine as generated with CD_3COCD_3 as the CI reagent gas.

complex of a benzene molecule and a secondary propyl carbenium ion reacts reversibly by proton transfer thus leading to partial incorporation of the hydrogen atom from the CI reagent into the propene molecule expelled from the metastable ions. Comparable studies have been reported for the loss of propene from the metastable protonated aryl *n*-propyl ethers [16, 17, 20]. For these compounds, protonation under CI conditions may occur either on the aromatic ring or on the oxygen atom of the ether function, thus leading to isomeric ions that can expel propene by different pathways. In previous studies it was concluded that only the transfer of a deuterium to the oxygen atom results in interchange between the propyl hydrogen atoms and the deuterium atom of the labeled CI reagent [16, 17]. The extent of incorporation of the deuterium atom from the CI reagent into the propene molecule was observed to depend on the presence and position of a methyl group or a fluorine atom on the ring. In particular, the ratio



Scheme 1. Proposed mechanism of propene loss from the metastable $[M + D]^+$ ions of 2-*n*-propoxy-pyridine with intermediate formation of ion-neutral complexes and complete loss of the positional identity of the hydrogen/deuterium atoms of the propyl carbenium ion.


between the losses of C_3H_5D and C_3H_6 from the metastable $[M + D]^+$ ions of *n*-propyl-3-methylphenyl ether was reported to be 13:87, whereas this ratio becomes 27:73 if the methyl group is situated at the 4-position of the ring [16]. In terms of the initial protonation site, the presence of a methyl at the 4-position may disfavor proton transfer to the ring [43] and—as a result—protonation at the oxygen atom becomes relatively important under CI conditions.

For the present series of aromatic ethers, the results reveal that the loss of propene from $[M + D]^+$ ions occurs without interchange between the deuterium atom from the CI reagents and the hydrogen atoms of the propyl group. Even though the site-specific proton affinities of the pyridine ring of the ethers are unknown, deuterium transfer to the nitrogen atom is expected to be associated with strong thermodynamic driving force. The existence of a strong thermodynamic preference for transfer to the nitrogen atom is substantiated by the calculated proton affinity of 946 kJ mol^{-1} for this site in 3-hydroxypyridine, whereas the value for the oxygen atom is suggested to be much lower (726 kJ mol^{-1}) [44, 45]. For the propoxy-pyridines, it may be anticipated that the proton affinity of the oxygen atom may be somewhat higher as compared to the same position in the hydroxypyridines due to the stabilization of the positive charge by the alkyl group in the oxygen protonated species. This could imply that some endothermic deuterium transfer to the oxygen atom of the propoxy-pyridines may occur in the experiments with CD_3OD (PA = 754 kJ mol^{-1}) and CD_3CN (PA = 779 kJ mol^{-1}), whereas the reagents, $(CD_3)_2CO$ and pyrrole- D_5 , may be too strong bases (PA = 812 and 875 kJ mol^{-1} , respectively) [25] to allow the occurrence of deuterium transfer to this site in the ethers under the present CI conditions. Notwithstanding that it cannot be excluded that some deuterium transfer occurs to other sites than the nitrogen atom in the present experiments,

the combined results indicate that loss of propene involves predominantly $[M + D]^+$ ions formed by deuteriation at the nitrogen atom.

In spite of the conclusion that the results are in line with formation of nitrogen deuterated species, the results suggest the occurrence of propene loss by a number of pathways; that is, propene loss from the metastable $[M + D]^+$ ions of the propoxy-pyridines can involve: (1) Ion-neutral complex formation and loss of the positional identity of all hydrogen atoms of the propyl carbenium ion part of the complex, (2) ion-neutral complex formation by cleavage of the ether bond associated with an irreversible 1,2-hydride shift in the carbenium ion if a *n*-propyl group is present in the parent compound or with no loss of positional identity of the hydrogen atoms of the carbenium ion if the ether contains an isopropyl group, (3) a 1,5-hydride shift from the β -position of the propyl group to the pyridine ring concomitant with dissociation, and (4) a 1,2-elimination type process.

The formation of ion-neutral complexes is suggested to be the predominant or even the exclusive pathway involved in the loss of propene from the $[M + D]^+$ ions of the 2-*n*-propoxy-pyridine. For this particular species, the experimental ratio between the losses of C_3H_5D and $C_3H_4D_2$ (24:76) from the $[D_2]$ -labeled propoxy-pyridines is slightly different from the ratio (29:71) calculated on the basis of formation of an ion-neutral complex with a propyl carbenium ion that may react by reversible 1,2-hydride and deuteride shifts prior to proton or deuterium transfer (Scheme 1). This is further confirmed by the ratios of the loss of propene from the $[\gamma,\gamma,\gamma-D_3]$ -2-*n*-propoxy-pyridines, which expel $C_3H_4D_2$ and $C_3H_3D_3$ in a ratio (33:67) that is also different from the statistical value (43:57). This statistical analysis is based, however, upon a neglect of isotope effects on the ensuing acid-base reaction between the partners of the complex. Possibly, the initial cleavage of the bond between the



Scheme 2. Proposed mechanisms of propene loss from the metastable $[M + D]^+$ ions of 3-*n*-propoxypropyridine. The first pathway (I) involves intermediate formation of ion-neutral complexes, the second pathway (II) is a 1,2-elimination type process and the third reaction (III) involves a 1,5-H shift.


oxygen atom and the α -carbon atom of the propyl group is followed by a proton/deuteron transfer step with a small but discernible isotope effect (vide infra).

Even though the precise structure of the C_5H_4NDO molecule in the complexes shown in Scheme 1 is uncertain, it may be anticipated that cleavage of the ether bond leads to 2-pyridone. Upon the assumption that 2-pyridone is generated, proton transfer from the propyl carbenium ion to the oxygen atom is estimated to be exothermic with $\sim 180 \text{ kJ mol}^{-1}$ as based on the experimental proton affinity of propene ($PA = 751 \text{ kJ mol}^{-1}$) [25] and the calculated value of the oxygen atom in 2-pyridone (933 kJ mol^{-1}) [24]. A large difference in proton affinity between propene and the 2-pyridone molecule within the ion-neutral complex shown in Scheme 1 will cause proton transfer to be essentially irreversible in line with the absence of incorporation of the deuterium atom from the CI reagent into the propene molecules.

The loss of propene from the metastable $[M + D]^+$ ions of the 3-*n*-propoxypropyridine is likely to involve one of the other pathways in addition to the formation of ion-neutral complexes as indicated by the different experimental ratios between the losses of C_3H_5D and $C_3H_4D_2$ from the $[\alpha, \alpha\text{-D}_2]$ - and the $[\beta, \beta\text{-D}_2]$ -labeled species (Table 2). For the $[\alpha, \alpha\text{-D}_2]$ -labeled ions, the average ratio of 21:79 suggests that transfer of a deuterium atom from this position is less important than predicted for formation of an ion-neutral complex and complete loss of the positional identity of the hydrogen and deuterium atoms of the propyl group (29% C_3H_5D loss and 71% $C_3H_4D_2$ loss). In addition, the loss of $C_3H_4D_2$ from the $[\beta, \beta\text{-D}_2]$ -labeled ions is significantly

less pronounced than expected for the ion-neutral complex pathway shown in Scheme 1, for the 2-isomer (on average 54% instead of the predicted 71%). Evidently, part of the observed 46% C_3H_5D loss (Table 2) arises by a pathway that involves the specific transfer of a deuterium atom from the β -position of the propyl group. A possible reaction would be a 1,5-hydride shift to the 2-position of the ring concomitant with propene loss and formation of the keto-species shown in Scheme 2. It should be emphasized, that the CID spectrum of the product ions of propene loss is similar to the spectrum obtained of the deuterated 3-hydroxypropyridine (see also Results and Figure 2). A competing reaction such as the 1,2-elimination with direct formation of the 3-hydroxypropyridine (deuterated at the nitrogen atom), therefore, cannot be excluded.

The occurrence of propene loss by another pathway than ion-neutral complex formation can reflect that the initial cleavage of the ether bond will lead to the zwitterion form of 3-hydroxypropyridine (see Scheme 2). The relative stability of this form of 3-hydroxypropyridine is uncertain but it is expected to be significantly less stable than the hydroxy form [46, 47]. As a result, the pathway involving ion-neutral complexes may be associated with a relatively high critical energy and allow other processes to be able to compete with this reaction pathway. Estimation of the relative extent of ion-neutral complex formation and, for example, the 1,5-hydride shift initiated propene loss is hampered, however, by the likely occurrence of kinetic isotope effects. For example, a statistical analysis (see Appendix) of propene loss from the metastable $[M + D]^+$ ions of the $[\beta, \beta\text{-D}_2]$ -labeled 3-*n*-propoxypropyridine suggest that



Scheme 3. Proposed mechanism of propene loss from the metastable $[M + D]^+$ ions of the 2-, 3-, and 4-*n*-isopropoxy-pyridines with intermediate formation of ion-neutral complexes and no loss of the positional identity of the hydrogen/deuterium atoms of the isopropyl carbenium ion.

about 24% of the ions dissociate by the pathway involving a 1,5-hydride shift (see also reference [48] for a review on composite atom interchange in gaseous ions). A similar analysis for the $[\alpha, \alpha\text{-D}_2]$ -labeled ions, indicates that slightly more ions (about 30%) dissociate by such a mechanism, whereas a significantly larger part of the $[\gamma, \gamma, \gamma\text{-D}_3]$ -labeled ions (45–50%) appears to react by the 1,5-hydride initiated propene loss as based upon the simplified statistical analysis. For the $[\alpha, \alpha\text{-D}_2]$ - and $[\beta, \beta\text{-D}_2]$ -labeled ions, the occurrence of an isotope effect in the 1,5-hydride shift could be held responsible for the suggestion that a lower fraction of the ions reacts by this pathway if the label is at the β -position. On the other hand, the apparent pronounced loss of propene by the 1,5-hydride shift from the $[\gamma, \gamma, \gamma\text{-D}_3]$ -labeled species would tend to suggest the occurrence of a significant isotope effect in the ion-neutral complex pathway thus favouring the other reaction channel(s). Despite the limitations of the statistical analysis (as based on the absence of isotope effects), these considerations lead to the conclusion that more than one pathway is involved in propene loss from the 3-*n*-propoxy-pyridine.

The results for the metastable $[M + D]^+$ ions of the 4-*n*-propoxy-pyridine appear to some extent to be intermediate to the findings for the 2-*n*- and 3-*n*-propoxy-pyridine isomers. In mechanistic terms, the results reveal a slight preference for the involvement of the β -position of the propyl group as exemplified by the observation that the average ratio between the losses of C_3H_5D and $C_3H_4D_2$ (38:72) differs from the ratio (29:71) predicted for the sole occurrence of ion-neutral complex formation and randomization of the hydrogen and deuterium atoms within the propyl carbenium ion. For the 4-isomer, it is proposed that the competing process may be a 1,2-elimination in line with the expectation that the charge distribution within the $[M + D]^+$ ions will disfavour a 1,5-hydride shift to the aromatic ring.

Even though the precise nature of the competing process is unknown, it is clear that it should involve a specific transfer of a hydrogen or deuterium atom from the β -position prior to propene loss.

For the isopropoxy-pyridines, the absence of the loss of C_3H_5D from the $[M + D]^+$ ions of 2-, 3- and 4-(C_5H_4N)- $OCD(CH_3)_2$ indicates that the positional identity of the hydrogen atom at the α -position of the isopropyl group is retained during propene loss (Scheme 3). In the context of ion-neutral complex formation this observation suggests that the initial cleavage of the bond to the isopropyl group leads to an isopropyl carbenium ion with insufficient internal energy to undergo reversible 1,2-hydride shifts (see also reference [49]).

For the metastable $[M + D]^+$ ions of the $[\beta, \beta, \beta\text{-D}_3]$ -isopropoxy-pyridines the competing losses of $C_3H_4D_2$ and $C_3H_3D_3$ should occur to an equal extent in the absence of an isotope effect. By contrast, the observed average ratio is 33:67 from all the isomeric ions (Table 4). The experimental results indicate clearly the occurrence of a significant kinetic isotope effect of ~ 2 in propene loss from the ionized and labeled isopropoxy-pyridines. A similar kinetic isotope effect has been reported for the loss of propene from the metastable $[M + H]^+$ ions of $C_6H_5CH(CH_3)CD_3$ and ascribed to the existence of a significant energy barrier between the $[C_6H_6 \text{ iso-}C_3H_7^+]$ and $[C_6H_7^+ C_3H_6]$ complexes involved in the process [42]. For the present systems, it should be mentioned that the results can be interpreted also in terms of the occurrence of propene loss by a 1,2-elimination as shown in Scheme 2 for the $[M + D]^+$ ions of 3-*n*-propoxy-pyridine. Such a process may well be associated with a significant isotope effect. In other words, a distinction between the different possible pathways in propene loss from the $[M + D]^+$ ions of the

isopropoxy pyridines cannot be reached on the basis of the present results.

Conclusions

Chemical Ionization of isomeric propoxy pyridines with deuterium labeled reagents (CD_3OD , CD_3CN , $(\text{CD}_3)_2\text{CO}$ and pyrrole- D_5) yields abundant $[\text{M} + \text{D}]^+$ ions that expel only propene on the μs time-scale. The main mechanism of propene loss is proposed to be the formation of ion-neutral complexes composed of a propyl carbenium ion and a hydroxypyridine in the keto form if the propoxy group is situated at the 2- or 4-position, whereas a zwitterion is formed if the propoxy group is at the 3-position in the parent compound. The subsequent proton/deuteron transfer is associated with an isotope effect that is particularly pronounced for the loss of propene from the $[\text{M} + \text{D}]^+$ of the isopropoxy pyridines. In addition to the pathway involving ion-neutral complex formation, propene loss can involve: (1) A 1,5-hydride shift as indicated by the results of the 3-*n*-propoxy pyridine and/or (2) a 1,2-elimination type process as suggested for the 4-*n*-propoxy pyridine.

Acknowledgments

The authors thank the Netherlands Organization for Scientific Research (SON/NWO) for continuing financial support. They acknowledge Mr. F. Pinkse for technical support and Mrs A. M. van der Burg for assistance during the synthesis of some of the labeled compounds.

Appendix

The relative extent of the losses of the partially labeled propene molecules from the metastable $[\text{M} + \text{D}]^+$ ions of 3-*n*-propoxy pyridines is analyzed by considering the complete loss of the positional identity of the hydrogen and deuterium atoms in combination with the occurrence of a specific transfer of a H/D atom from the β -position. For the $[\alpha, \alpha\text{-D}_2]$ -labeled 3-*n*-propoxy pyridine, the statistical ratio is 2/7 (29%) $\text{C}_3\text{H}_5\text{D}$ loss and 5/7 (71%) $\text{C}_3\text{H}_4\text{D}_2$ loss, whereas the average experimental ratio between the losses of $\text{C}_3\text{H}_5\text{D}$ and $\text{C}_3\text{H}_4\text{D}_2$ is about 20:80. Upon the assumption that the 20% loss of $\text{C}_3\text{H}_5\text{D}$ only involves the formation of a $[\text{C}_5\text{H}_4\text{DNO} \text{C}_3\text{H}_5\text{D}_2]^+$ ion-neutral complex, the statistical ratio is matched if 50% of the total loss of propene (in the form of $\text{C}_3\text{H}_4\text{D}_2$) arises from this channel. In conclusion, the remaining part of the $\text{C}_3\text{H}_4\text{D}_2$ molecules (30% of the total loss of propene) is formed by the specific transfer of a hydrogen atom from the β -position.

For the $[\text{M} + \text{D}]^+$ ions of the $[\beta, \beta\text{-D}_2]$ -labeled 3-*n*-propoxy pyridine the experimental ratio between the losses of $\text{C}_3\text{H}_5\text{D}$ and $\text{C}_3\text{H}_4\text{D}_2$ is approximately 46:54. For these $[\text{M} + \text{D}]^+$ ions, the 54% loss of $\text{C}_3\text{H}_4\text{D}_2$ is assumed to arise only by formation of ion-neutral complexes. As a result, the statistical ratio of 2/7 $\text{C}_3\text{H}_5\text{D}$ loss and 5/7 $\text{C}_3\text{H}_4\text{D}_2$ loss is obtained if 22% of the total propene loss

in the form of $\text{C}_3\text{H}_5\text{D}$ is by this pathway. The remaining 24% of the total loss of propene (as $\text{C}_3\text{H}_5\text{D}$) originates from a specific transfer of a D from the β -position. Similarly, the loss of $\text{C}_3\text{H}_4\text{D}_2$ from the $[\text{M} + \text{D}]^+$ ions of the $[\gamma, \gamma, \gamma\text{-D}_3]$ -*n*-propoxy pyridines is assumed to originate only from ion-neutral complexes mediated processes as based on the experimental ratio 22:78 between the losses of $\text{C}_3\text{H}_4\text{D}_2$ and $\text{C}_3\text{H}_3\text{D}_3$. As a result, it is estimated that 49% of the total propene loss (in the form of $\text{C}_3\text{H}_3\text{D}_3$) is expelled by the pathway involving a specific H-transfer from the β -position.

References

- Harrison, A. G. *Chemical Ionization Mass Spectrometry*, 2nd ed.; CRC Press: Boca Raton, FL, 1992, pp 1–208.
- Kebarle, P. Gas-phase Ion Thermochemistry Based on Ion-equilibria from the Ionosphere to the Reactive Centers of Enzymes. *Int. J. Mass Spectrom.* **2000**, *200*, 313–330.
- McMahon, T. B. Thermochemical Ladders: Scaling the Ramparts of Gaseous Ion Energetics. *Int. J. Mass Spectrom.* **2000**, *200*, 187–199.
- Taft, R. W. Protonic Acidities and Basicities in the Gas Phase and in Solution: Substituent and Solvent Effects. *Prog. Phys. Org. Chem.* **1983**, *14*, 247.
- Szulejko, J. E.; McMahon, T. B. Progress Toward an Absolute Gas-phase Proton Affinity Scale. *J. Am. Chem. Soc.* **1993**, *115*, 7839–7848.
- Kuck, D. Mass Spectrometry of Alkylbenzenes and Related Compounds 2. Gas-phase Ion Chemistry of Protonated Alkylbenzenes (Alkylbenzenium ions). *Mass Spectrom. Rev.* **1990**, *9*, 583–630.
- Meot-Ner, M.; Sieck, L. W. Proton Affinity Ladders from Variable-Temperature Equilibrium Measurements. 1. A Re-evaluation of the Upper Proton Affinity Range. *J. Am. Chem. Soc.* **1991**, *113*, 4448–4460.
- Jasinski, J. M.; Brauman, J. I. Rates of Proton Transfer Between Pyridine Bases in the Gas Phase. Steric and Isotope Effects. *J. Am. Chem. Soc.* **1980**, *102*, 2906–2913.
- Büker, H.-H.; Grützmacher, H.-F. A Fourier Transform-Ion Cyclotron Resonance Study of Steric Effects on Proton Transfer Reactions of Polyalkyl Benzenes. *Int. J. Mass Spectrom. Ion Processes* **1991**, *109*, 95–104.
- Bouchoux, G.; Salpin, J. Y.; Leblanc, D. A Relationship Between the Kinetics and Thermochemistry of Proton Transfer Reactions in the Gas Phase. *Int. J. Mass Spectrom. Ion Processes* **1996**, *153*, 37–48.
- Hunt, D. F.; Sethi, S. K. Gas-Phase Ion/Molecule Isotope-Exchange Reactions: Methodology for Counting Hydrogen Atoms in Specific Organic Structural Environments by Chemical Ionization Mass Spectrometry. *J. Am. Chem. Soc.* **1980**, *102*, 6953–6963.
- Lau, Y. K.; Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes. *J. Am. Chem. Soc.* **1976**, *98*, 7452–7453.
- Defrees, D. J.; McIver, R. T., Jr.; Hehre, W. J. The Proton Affinities of Phenol. *J. Am. Chem. Soc.* **1977**, *99*, 3853–3854.
- Vulpis, T.; Hammerum, S.; Houriet, R. Protonation of Anisole and Halobenzenes. *Adv. Mass Spectrom.* **1989**, *11*, 578–579.
- Audier, H.; Berthomieu, D.; Leblanc, D.; Morton, T. H. Multiple Protonation Sites in Aryl Ethers. *Int. J. Mass Spectrom.* **1998**, *175*, 133–147.
- Bogdanov, B.; Matimba, H. E. K.; Ingemann, S.; Nibbering, N. M. M. Chemical Ionization of Phenyl *n*-Propyl Ether and Methyl Substituted Analogs: Propene Loss Initiated by Com-

- peting Proton Transfer to the Oxygen Atom and the Aromatic Ring. *J. Am. Soc. Mass Spectrom.* **1996**, *7*, 636–652.
17. Bogdanov, B.; Matimba, H. E. K.; Ingemann, S.; Nibbering, N. M. M. Chemical Ionization of Fluorophenyl *n*-Propyl Ethers: Loss of Propene from the Metastable $[M + D]^+$ Ions. *J. Am. Soc. Mass Spectrom.* **1998**, *9*, 121–129.
 18. Bogdanov, B.; van Duijn, D.; Ingemann, S.; Hammerum, S. Protonation of Fluorophenols and Fluoroanisoles in the Gas Phase: Experiment and Theory. *Phys. Chem., Chem. Phys.* **2002**, *4*, 2904–2910.
 19. Benoit, F. M.; Harrison, G. Hydrogen Migrations in Mass Spectrometry. *Org. Mass Spectrom.* **1976**, *11*, 599–608.
 20. Kondrat, R. W.; Morton, T. H. Ion Neutral Complexes from Protonated Propyl Phenyl Ethers. Gas-phase Solvolysis versus Elimination. *Org. Mass Spectrom.* **1991**, *26*, 410–415.
 21. Kondrat, R. W.; Morton, T. H. Gas-phase Analogs of Solvolysis—The Conjugate Acid of Normal-Propyl Phenyl Ether. *J. Org. Chem.* **1991**, *56*, 952–957.
 22. Jacquet, J. P.; Morton, T. H. Chemical Ionization of *n*-Propyl Phenyl Ether with D_2O Reagent Gas. A Test of Statistical Models. *J. Mass Spectrom.* **1997**, *32*, 251–252.
 23. Audier, H. E.; Morton, T. H. Ring Versus Oxygen Protonation in Metastable Ion Decompositions of Protonated Isopropyl Phenyl Ether. *Int. J. Mass Spectrom.* **1999**, *187*, 393–399.
 24. Wolken, J. K.; Tureček, F. Modeling Nucleobase Radicals in the Gas Phase. Experimental and Computational Study of 2-Hydroxypyridinium and 2-(1H)-Pyridone Radicals. *J. Phys. Chem. A* **1999**, *103*, 6268–6281.
 25. Hunter, E. P. L.; Lias, S. G. Evaluated Gas Phase Basicities and Proton affinities of Molecules: An Update. *J. Phys. Chem. Ref. Data.* **1996**, *27*, 413–656.
 26. Cooks, R. G.; Beynon, J. H.; Caprioli, R. M.; Lester, G. R. *Metastable Ions*; Elsevier: Amsterdam, The Netherlands, 1976, pp 1–296.
 27. Uggerud, E. Account: The 1995 Maccoll Lecture: Fifty Years of Metastable Ions. Experiment and Theory in the Study of Mass Spectrometric Fragmentation Reactions. *Eur. J. Mass Spectrom.* **1996**, *2*, 205–211.
 28. Harrison, A. G.; Mercer, R. S.; Reiner, E. J.; Young, A. B.; Boyd, R. K.; March, R. E.; Porter, C. J. A Hybrid BEQQ Mass Spectrometer for Studies in Gaseous Ion Chemistry. *Int. J. Mass Spectrom. Ion Processes* **1986**, *74*, 13–31.
 29. Zappey, H. W.; Ingemann, S.; Nibbering, N. M. M. Gas-Phase Ion-Molecule Reactions of the CH_3O^+ Cation. *J. Am. Soc. Mass Spectrom.* **1992**, *3*, 515–517.
 30. Molenaar-Langeveld, T. A.; Gremmen, C.; Ingemann, S.; Nibbering, N. M. M. Isomeric Dependence of the Formation of Ion/Neutral Complexes in Dissociation Reactions of Ionized Propoxyppyridines. *Int. J. Mass Spectrom.* **2000**, *199*, 1–16.
 31. Schmid, G. H.; Wolkoff, A. W. The Formation of Alkenes by the Thermal Elimination Reaction of *N*-Methyl-4-Alkoxyppyridinium Iodides. *Can. J. Chem.* **1972**, *50*, 1181–1187.
 32. Wilson, C. L. The Preparation of Organic Compounds Containing Deuterium, Dideuteromalic Deuteroacid and Tri-deuteroacetic Deuteroacid. *J. Chem. Soc.* **1935**, 492–494.
 33. Bush, K. L.; Glish, G. L.; McLuckey, S. A. *Mass Spectrometry/Mass Spectrometry, Techniques and Application of Tandem Mass Spectrometry*; VCH Publishers: New York, NY, 1988, pp 1–333.
 34. Morton, T. H. Gas Phase Analogues of Solvolysis Reactions. *Tetrahedron* **1982**, *38*, 3195–3243.
 35. McAdoo, D. J. Ion Neutral Complexes in Unimolecular Decompositions. *Mass Spectrom. Rev.* **1988**, *7*, 363–393.
 36. Harrison, A. G.; Wang, J.-Y. Fragmentation of Deuteronated Aromatic Derivatives: The Role of Ion-Neutral Complexes. *Int. J. Mass Spectrom.* **1997**, *160*, 157–166.
 37. McAdoo, D. J.; Morton, T. H. Gas-Phase Analogs of Cage Effects. *Acc. Chem. Res.* **1993**, *26*, 295–302.
 38. Longevialle, P. Ion-Neutral Complexes in the Unimolecular Reactivity of Organic Cations in the Gas-Phase. *Mass Spectrom. Rev.* **1992**, *11*, 157–192.
 39. Morton, T. H. The Reorientation Criterion and Positive-Ion Neutral Complexes. *Org. Mass Spectrom.* **1992**, *27*, 353–368.
 40. Matimba, H. E. K.; Ingemann, S.; Nibbering, N. M. M. Competing Reactions Leading to Propene Loss from the Molecular Ions of Aryl *n*-Propyl Ethers. *J. Mass Spectrom.* **1996**, *31*, 609–622.
 41. Herman, J. A.; Harrison, A. G. Effect of Protonation Exothermicity on the Chemical Ionization Mass Spectra of Some Alkylbenzenes. *Org. Mass Spectrom.* **1981**, *16*, 423–427.
 42. Berthomieu, D.; Audier, H.; Denhez, J.-P.; Monteiro, C.; Mourgues, P. $[C_6H_6, C_3H_7^+]$ and $[C_6H_7^+, C_3H_6]$ Ion-Neutral Complexes Intermediates in the Reactions of Protonated Propylbenzenes. *Org. Mass Spectrom.* **1991**, *26*, 271–275.
 43. Koblenz, T.; van Beelen, E. S. E.; Ingemann, S.; Hammerum, S.; in preparation.
 44. Wolken, J. K.; Tureček, F. Heterocyclic Radicals in the Gas Phase. An Experimental and Computational Study of 3-Hydroxypyridinium Radicals and Cations. *J. Am. Chem. Soc.* **1999**, *121*, 6010–6018.
 45. Tureček, F. Modeling Nucleobase Radicals in the Mass Spectrometer. *J. Mass Spectrom.* **1998**, *33*, 779–795.
 46. Trikoupis, M. A.; Gerbaux, P.; Lavorato, D. L.; Flammang, R.; Terlouw, J. K. Hydrogen Shift Isomers of Ionic and Neutral Hydroxypyridines: A Combined Experimental and Computational Investigation. *Int. J. Mass Spectrom.* **2002**, *217*, 1–22.
 47. Huq, F.; Yu, J. Q. Molecular Modeling Analysis: “Why is 2-Hydroxypyridine Soluble in Water but not 3-Hydroxypyridine?” *J. Mol. Model.* **2002**, *8*, 81–86.
 48. Kuck, D. Half a Century of Scrambling in Organic Ions: Complete, Incomplete, Progressive, and Composite Atom Interchange. *Int. J. Mass Spectrom.* **2002**, *213*, 101–144.
 49. Koch, W.; Liu, B.; Schleyer, P. v. R. Definitive Characterization of the $C_2H_7^+$ Potential Energy Surface. *J. Am. Chem. Soc.* **1989**, *111*, 3479–3480.