

- A
- A₁ ions, enhancement for peptide sequencing and protein identification, 20:1214
- α₃ Ions, observation frequency, MS/MS, 19:1764
- a₃⁺ ions, study of formation, protonated tetraglycine, 17:1654
- Ab initio and modified neglect of differential overlap bond orders and modified neglect of differential overlap energy partitioning, application of: the effect of protonation site on bond strengths in simple peptides, 5:704
- Ab initio calculations on 1,2-hydrogen shifts in the benzene radical cation and on carbon scrambling via isomerization to the fulvene structure, 6:513
- Ab initio calculations on isomerization of 1,5-hexadiyne radical cation, 8:599
- of the dimethylene cyclobutene and 1,2,4,5-hexatetraene radical cations to the benzene structure, 8:594
- Ab initio calculations on the isomerization of alkene radical cations, 10:577
- Ab initio calculations on the isomerization of the 1,3-hexadien-5-ynyl radical cation to the benzene structure and to non-classical ion structures, 7:731
- Ab initio characterization of ring-opening H-transfer in ionized cyclopentanone, similarity to ion-neutral complex-mediated alkane eliminations, 7:1251
- Ab initio study of intramolecular interactions in cations and hypervalent ammonium radicals, 8:1111
- Ab initio transition state, elimination of methane from protonated acetaldehyde, 5:1102
- Ablation
- irradiation effects in, 2–5-dihydroxybenzoic acid monocrystals in, 14:893
- resonant laser, as a selective metal ion source for gas-phase ion molecule reactions, 7:664
- Abundance plot, in detection of isobaric interferences in DEMALDI-TOFMS of complex polymer mixtures, 12:911
- Abundance sensitivity, enhanced, by resonant ejection of atomic ions, plasma source ion trap mass spectrometry, 7:1161
- Acceleration, mass correlated, in reflectron MALDI-TOF mass spectrometer, 13:135
- Acceleration-deceleration mass spectrometer, tandem quadrupole, novel, for neutralization-reionization studies, 3:493
- Accuracy in tandem mass spectrometry measurements, a kinetics-based protocol for instrument-independent measurements of collision-activated dissociation in RF-only quadrupoles, 1:272
- Accurate base composition of double-strand DNA by mass spectrometry, 7:1266
- Accurate mass capability, radical ion species, MALDI-TOF-MS, 17:672
- Accurate mass measurement of low molecular weight compounds by MALDI-TOFMS, 11:458
- Accurate masses for structure confirmation, 5:57
- Acetaldehyde, protonated, elimination of methane from, the ab initio transition state, 5:1102
- Acetals, cyclic, fast-atom bombardment of, evidence indicating the predominant involvement of condensed-phase processes in ionization, 4:493
- Acetate-anion complex, monoprotonated, intrinsic hydration of, 15:777
- Acetate dimers, proton-bound, characterization by IMS, 19:1361
- Acetone
- copper ion solvation by, 13:453
- ionized and ethylated, MS/MS and FTICR of, 8:225
- isotope labeling, RABA, quantitative proteomics, 20:1366
- protonated peptide derivatization in gas phase with, 11:244
- tandem mass spectrometry studies of, 2:113
- Acetone cation, collision-activated dissociation of, kinetics and mechanism of, 3:427
- Acetone enolate anion, gas-phase base-catalyzed Claisen-Schmidt reactions of, with various para-substituted benzaldehydes, 7:82
- Acetone/water cluster ions, tandem mass spectrometry studies of, 2:113
- Acetonitrile
- chemical ionization tandem mass spectrometry, of polyunsaturated fatty acid methyl esters, 10:1253
- gas phase reactions of, 8:1124
- Ni(II) complex ion reactions with, in gas phase, 15:1128
- propionitrile identification in, APPI MS, 19:1579
- Acetophenones, *para*-substituted, chemical mass shifts of, correlation with Brown's σ_p^+ constants, 13:623
- N-Acetyl neuraminic acid oligomers, associated with tetanus toxin C fragment, ESI-MS, 17:967
- Acetylation
- high mobility group 1 proteins, LC MS/MS, 18:1569
- peptide, detection and location of acetylated lysine residues, 18:1423
- Acetylcholine
- isomerization of linear C₃H₃⁺ in its reaction with, in a quadrupole ion trap mass spectrometer, 3:631
- mechanism of trimethylamine loss, density functional theory study, 20:238
- Acetylcholinesterase, inhibitor screening by MALDI/FTMS, 19:1850
- Acetaldehyde and methanol, protonated, gas-phase reaction between, 1:481
- N-Acetylheparosan, hyaluronic acid distinguished from, by MS/MS, 19:82
- Acid-denatured myoglobin, stepwise refolding of, evidence from electrospray mass spectrometry, 4:638
- Acid hydrolysis
- microwave-assisted, of protein, LC MALDI MS/MS with, 16:471
- proteins, MALDI, 20:2106
- Acidic heteroxylans, uronosyl substitution patterns in, ESI-MS/MS of, 15:43
- Acidic phospholipids, unsaturated, negative ESI-MS structural characterization of, 16:46
- Acidic residues, in gas phase H/D exchange of peptides and peptide dimers, 14:1477

Acidities

- apparent gas-phase, of multiply protonated peptide ions, ubiquitin, insulin B, and renin substrate, 7:1211
- gas-phase
of diols, 4:723
of germane and methylgermane, Fourier transform-ion cyclotron resonance study of, bond dissociation energy of germane, 4:54
of methylthioacetic acid, determining with Cooks' kinetic method, 16:535
and organic molecule responsiveness to negative ion ESI MS, 16:446
- Acidity of fluorobenzene and 1,4-difluorobenzene on the, gas-phase reactions between O⁻ and C₆H₅F, 4:73
- Acidity of uracil and uracil analogs in gas phase, four surprisingly acidic sites and biological implications, 13:985
- Acid(s), bitter, in hop extract, LC-APCI-MS of, 15:180
- Acids and esters, gas-phase methylation and dissociation of, effects of functional group interactions on, 7:565
- Acoustic desorption, laser-induced, charged phenyl radicals evaporated by, gas-phase reactions of, 13:192
- Acrolein, protonated thioether conjugates of, fragmentation of, 8:727
- Acrylonitrile cations, long-lived excited electronic states of, 14:110
- Activated carbon substrate surface for MALDI-MS, 11:644
- Activated ion electron capture dissociation, optimization for proteins, 20:763
- Activation energies, estimation using fragmentation yield:CID of iron(II)phenanthroline, ESI-MS, 17:962
- Activation energy, for unimolecular dissociation of noncovalent gas phase peptide, 14:1282
- Activation of C₂H₆, C₃H₈, HC(CH₃)₃, and c-C₃H₆ by gas-phase Ru⁺ and the thermochemistry of ru-ligand complexes, 10:821
- Active site location, from high-resolution tandem mass spectrometry, thiaminase I (42 kDa heterogeneity) and sequence refinement, 6:981
- Acute myeloid leukemia, histone modifications in, HPLC and MS of, 15:77
- Acyclic analogs, orders of relative alkali metal ion affinities of, by the kinetic method, 3:543
- Acyclic distonic acylium ions, dual free radical and acylium ion reactivity in single molecule, 11:697
- Acyl analogues of platelet-activating factor, heptafluorobutyryl derivatives of, mass spectral fragmentation, 2:476
- Acyl carrier protein, conformation by ESI-MS, 18:1525
- Acyl coenzyme A thioesters, long-chain fatty, by negative ion fast-atom bombardment mass spectrometry and tandem mass spectrometry, 5:416
- N-Acylanilines, ortho effect in ESI-MS, 19:1114
- Acylcarnitine, acyl groups of, studies on picolinyl ester fragmentation mechanisms, 17:1620
- Acylium ions
gaseous, ketalization of, 12:150
iso(thio)cyanate interactions with, 16:1602
- Acylphosphatidylglycerols, from salmonella typhimurium, ES-IMS/MS of, 15:1
- Acylpolyamides, from spider venom, MS/MS of, 15:1636
- ADAMS. *See* algorithm for discriminant analysis of mass spectra
- Additives for stabilization of double-stranded DNA in UVMALDI-MS, 13:1129
- Additives to the liquid chromatographic eluent, ion formation of N-methyl carbamate pesticides in thermospray mass spectrometry, 5:913
- Adduct formation, regioselectivity of I, fast atom bombardment-induced condensation of glycerol with ammonium surfactants, 3:318
- Adduction, noncovalent, in electrospray mass spectrometry, mass and charge state assignment for proteins and peptide mixtures via, 6:1175
- á-Adducts, anionic, in gas phase, using electrospray ion source as chemical reactor, 15:927
- Adducts
copper, of protonated ubiquitin, electrospray ionization mass spectrometry of, 6:521
deoxynucleoside, polycyclic aromatic hydrocarbon dihydrodiol-epoxides, low energy tandem mass spectrometry of, 6:248
fragmentation mechanism of, ion-molecule reaction of alkanenitriles and transition-metal ions in the gas phase, 2:120
- Adducts (molecular mass=98 μ) attached to peptide and protein ions in electrospray ionization mass spectra, origin and removal of, 1:382
- Adenine, protonated, tautomers, solvated clusters, and dissociation mechanisms, 16:1713
- Adenine-thymine binary-base oligonucleotide dissociations, positive-ion MALDI-TOF-MS of, 13:1052
- Adenosine 5'-monophosphate, multiply charged negative clusters of, in gas phase, 16:1840
- Adenosine-5'-phosphosulfate reductase, *M. tuberculosis*, noncovalent complexes analyzed using ESI-FTICR MS, 18:167
- Adenosine monophosphate, gas phase H/D exchange of, conformational dependence of, 12:780
- Adenosine triphosphate
binding to growth factors detected by MALDI-TOF and ESI-MS, 19:91
release from PP/ATP films, EC/ESMS, 18:919
- Adenosylated peptides acting as bisubstrate analogs of protein kinases, mass spectrometric characterization of a series of, 5:443
- Adenovirus proteins, extraction by SDS-PAGE, MALDI-MS of, 11:356
- Adenylate kinase, association constants of, with noncovalent inhibitors, 14:442
- Adenylyl cyclase 8, CaM and, isotope-labeled cross-linkers and FTIR MS analysis of, 17:1100
- Adipocytes, free fatty acid transport in, multi imaging MS of, 15:1572
- Adipose tissue, human, highly chlorinated dimethanofluorenes in, 1:99
- ADME (absorption, distribution, metabolism, and elimination) assays, parallel LC-MS in, 13:155
- Adsorption/desorption in ²⁵²Cf plasma desorption mass spectrometry, substrates for, small molecules as, 2:29
- Advanced glycation end products/peptides, enzymatic digestion and MS studies of, 15:496
- Advantages of *Thermococcus kodakaraensis* (KOD) DNA polymerase for PCR-mass spectrometry based analyses, 14:601
- Aerodynamic drag model for protein ions, 5:17
- Aerosol matrix-assisted laser desorption/ionization, vitamin B12 fragmentation in, 8:140
- Aerosol matrix-assisted laser desorption ionization mass spectrometry, 5:1
- Aerosol TOFMS, L2MS with, for analysis of individual organic particles, 9:1068
- Aerosols, mass spectra of, chemically assigned classification of, 13:826
- Affinity chromatography, in phosphopeptide quantification, 15:363
- Affinity probes, for peptides and proteins, carbon nanotubes as, in MALDI MS, 15:1629

- Affinity purified cross-linked peptides, MS detection of, 15:832
- Aflatoxin B1-guanine adducts, identification and positional mapping of, 9:676
- Afterglow technique, flowing, for ion-molecule reaction studies, early development of, 3:479
- Afterglow-guided ion beam, in MS/MS, 8:688
- Afterpeak time regime of pulsed glow discharge plasmas, ion formation processes in, 4:727
- Ag⁺-phenylalanine complex, gas phase fragmentation of, 13:408
- AGG. *See* Alanylglucylglycine
- Aglycones, flavonoid, LC-ESI-MS/MS of, 12:707
- Aglycons, glycoalkaloids, comparison of high- and low-energy collision-induced dissociation tandem mass spectrometry in, 7:173
- Agostic effects in the gas-phase, SiC13⁺ and SiC1⁺ affinities for pyridines determined by using the kinetic method with multiple stage mass spectrometry, 7:198
- Agricultural matrices, ferric chelates in, as fertilizers, LC-ESI-MS, 18:37
- Air
- electrospraying water in, at atmospheric pressure, 15:253
 - laboratory, soil surfaces exposed to, cyclohexamine on, static secondary ionization mass spectrometry detection of, 7:168
 - urban, selected organic compounds in, automated high-speed analysis of, by on-line isotopic dilution cryofocusing gas chromatography/mass spectrometry, 5:1001
 - volatile organic compounds in, PIT-MS detection and identification of, 16:1316
- Air amplifier, mechanistic studies on, LTQ-FT-ICR-MS, 18:1909
- Air analysis using tenax collection with jet-separator enrichment and ion trap mass spectrometric analysis, 7:1172
- Air and helium, on the performance of the ion trap detector when used for real-time monitoring of some atmospheric pollutants, 4:774
- Airborne particles, on-line analysis of, 10:648
- Airborne particulate matter and soil, nitro-substituted polynuclear aromatic hydrocarbons in, sub-parts-per-billion determination of, by electron capture-tandem mass spectrometry, 7:1255
- Akabori reaction, microwave-enhanced, for peptide analysis, 13:839
- Akt, inhibitor interaction with, analyzed by cross-linking and MS, 20:1504
- Alanine, gas-phase acidity of, theoretical calculations of, 10:318
- Alanylglucylglycine, protonated, loss of CO and ammonia, MS/MS, 18:1291
- Albumin
- human serum
 - carbon isotope variability in, direct analysis of, by liquid flow-injection isotope ratio mass spectrometry, 7:605
 - ESI-MS of, 8:1247
 - specific protein carbonylation sites, 17:1172
- Alcohol ammonia and water-alcohol protonated clusters, on the structure of, 4:507
- Alcohol complexation, properties of mono- and diphosphate bridged cavitands, ESI-FTICR MS study, 17:213
- Alcohol consumption, ethyl sulfate as marker of, by LC-ESI-MSMS, 15:1644
- Alcohol derivatization reagent, anthraquinone-2-carbonyl chloride as, for negative ion chemical ionization mass spectrometry, 6:148
- Alcohol-water and ammonia-alcohol protonated clusters, on the structure of, 4:507
- Alcohols
- amino, site-selective gas-phase reaction of, with dimethyl ether ions, 4:230
 - detection of, by electron capture negative ion chemical ionization mass spectrometry, pentafluorobenzyl chloroformate derivatization for, 6:525
 - meta-nitrobenzyl, and dipyrrocyanine dyes, beam-induced reaction between, in liquid-secondary-ion mass spectrometry, 5:1017
 - multifunctional, including triacylglycerols, fragmentation of conjugate bases of esters derived from, 6:498
 - reactions of Sc⁺, Y⁺ and Lu⁺ with, effects of the class and chain length of alcohols on the nature of primary products, 7:1157
- Aldehyde dehydrogenases, structure of, enzyme inhibitors and, 12:97
- Aldehydes, short chain, protein modification by, ESI-MS/MS of, 215
- Aldehyde-ketose isomerization, gas-phase, study using tandem mass spectrometry and theoretical calculations, 4:943
- Aldehyde reductase inhibitors, binding of, crystallographic and MS studies of, 10:635
- Alginate, oligosaccharides derived from, sequence analysis by negative-ion ES MS/MS, 17:621
- Algorithm for discriminant analysis of mass spectra (ADAMS), aerosol MS chemical classification by, 13:826
- Algorithms
- identification of PRISM, 18:563
 - mass spectral library search, for compound identification, optimization and testing of, 5:859
 - for peptide mass fingerprint measurement, 16:13
- ALiPHAT method, evaluation of PC-IDMS and LOD correlation, 20:2006
- Aliphatic amino acids, [M + Ni]⁺ adducts of, fragmentation of, 8:749
- Aliphatic compounds, 1:10.5-eV photoionization mass spectrometry of, 1:419
- Aliphatic disulfides, bond-selective photodissociation of, 6:872
- Aliphatic ethers, ionized, ion-neutral complex-mediated alkane eliminations from, size effects in, 2:261
- Aliphatic peptidyl hydroperoxides, as source of secondary oxidation, hydroxyl radical footprinting of proteins, 20:1123
- Aliphatic phosphopeptides, dissociation of, ESI-FTICR-MS and IRMPD of, 15:121
- Alkali/alkaline earth metal cation selectivity, of lariat ether amides, ESI-MS evaluation of, 14:1215
- Alkali cation attachment to derivatized fullerene, MALDI of, 13:1448
- Alkali-cationed polysaccharides, MS/MS of, 8:987
- Alkali-cationized peptides, dissociation pathways of, 12:497
- Alkali chloride cluster formation in liquid secondary ion mass spectrometry, ion-ion recombination for, 4:410
- Alkali chloride cluster ion dissociation, examined by kinetic method, heterolytic bond dissociation energies, effective temperatures, and entropic effects of, 13:1388
- Alkali ions, poly(ethylene terephthalate) oligomers cationized by, 10:883
- Alkali metal binding selectives, of caged aza-crown ether ligands, MESI-MS of, 11:884
- Alkali-metal-carboxylate radical anions, in fast-atom bombardment, 6:1252
- Alkali metal cation-valinomycin complexes, gas-phase studies of, attachment rates and cation affinities, 5:632
- Alkali metal cationized oligosaccharides, collision-induced dissociation of, influence of the collision energy and of the collision gas for the assignment of linkage position, 4:197
- Alkali metal complexes, of polyether ionophore antibiotics, gas phase stabilities of, by CID, 16:779
- Alkali metal ions
- cleavage reactions of, positive ion ESI-MS/MS of, 15:689

- formation of, in ESI-MS of peptides and proteins, 11:967
poly(ethylene glycol) doubly and singly cationized by, 13:177
- Alkali metal ion affinities, relative, of crown ethers and acyclic analogs, by the kinetic method, 3:543
- Alkali metal ions and peptides, gas-phase anionic complexes of, structure and collision activated decompositions, 5:137
- Alkali metal-mediated resorcarene capsules, ESI-FTICR-MS study on gas phase structure and cation binding of tetraethyl resorcarene and its per-methylated derivative, 13:851
- Alkaline earth complexes, gas phase reactions of, 8:1124
- Alkaline earth metal ions
fatty acids cationized with, charge remote fragmentation of, 1:320
reactions with benzene, 12:250
- Alkaloids, structure determination of, by MS, 13:1254
- Alkane eliminations
from ionized alkanes, 9:21
ion-neutral complex-mediated from ionized aliphatic ethers, size effects in, 2:261
similarity to, *Ab initio* characterization of ring-opening H-transfer in ionized cyclopentanone, 7:1251
- Alkanenitriles, in the gas phase, ion-molecule reaction of, study on fragmentation mechanism of the adducts, 2:120
- Alkanes
air plasma APCI-MS study, 20:697
atmospheric chemical ionization of, 5:177
cyclopentadienylcobalt ion reactions with, 14:51
erratum, 5:1026
long chain, atomic transition metal ion reactions with, 12:367
in supersonic molecular beams, electron impact mass spectrometry of, 6:120
- Alkanols, charge-remote fragmentation and the two-step elimination of, from fast atom bombardment-desorbed ($M + H$)⁺, ($M + Cat$)⁺, and ($M-H$)⁻ ions of aromatic beta-hydroxyoximes, 4:819
- Alkenamides, fragmentation of, by collisional activation in triple quadrupole mass spectrometer, 15:1462
- Alkene radical ions, isomerization of, 10:577
- Alkenes, atmospheric chemical ionization of, 5:177, 5:1026
- Alkenones
double bond locations using GC-MS analysis, 17:710
polyunsaturated long chain, double bond location in, GC-MS in derivatization for, 15:1161
- Alkyl and aryl esters of benzenedicarboxylic acids, gas-phase reactions of O₂⁻ with, 5:990
- Alkyl anilines, isomeric, resolution by MS, 20:269
- S-Alkyl cysteine sulfoxide, peptide ions containing, gas-phase fragmentation reactions, 18:1690
- Alkyl derivatives, in glycosidic bonds, CID of, 14:63
- Alkyl dipeptides, protonated, in gas phase, bimolecular hydrogen-deuterium exchange behavior of, 6:466
- Alkyl diselenide, mass spectra of, 15:1325
- Alkyl ethoxysulfate, APCI-MS of, 10:529
- Alkyl methylphosphonic acids, response enhancement in negative ESI LC-MS/MS, 18:1821
- Alkyl phosphates, from an industrial effluent, identification of, multispectral, 2:419
- Alkyl side chains, fragmentation of deprotonated peptides containing, 12:1
- Alkylamine thiohydantoin amino acid derivatives, electrospray ionization study of, 6:1211
- Alkylammonium ions, gas phase hydrates of, magic numbers in, 8:1175
- Alkylaniline guanine adducts, C8-substituted, differentiation by ESI and quadrupole ion trap MS/MS, 14:1488
- Alkylation
halo, gas-phase, of H₆₀-fullerene by ion-molecule reaction under chemical ionization, 4:894
reductive, isotope labeling, RABA, quantitative proteomics, 20:1366
- Alkylisocyanate polymers, structural studies on, by thermal degradation tandem mass spectrometry, 2:130
- Alkylnicotinium halides, *N*⁻, a class of cationic matrix additives for enhancing the sensitivity in negative ion fast-atom bombardment mass spectrometry of poly anionic analytes, 5:935
- Allophycocyanine, localization in MALDI-preparations by CLSM, 17:1599
- Allosteric inhibition, of protein-protein interaction, MS of, 14:8
- Allyl bromide molecular ion, isomers of, kinetic and spectroscopic characterization of, 2:372
- ALPI MS. *See* Atmospheric pressure laser ionization mass spectrometry
- Alternating RF/DC isolations for quantitation with coeluting internal standards in gas chromatography/ion trap mass spectrometry, 3:85
- Alternative approaches to infrared multiphoton dissociation in an external ion reservoir, 14:1413
- Alumina, porous silica and, peptide fingerprinting on, LDI MS, 19:632
- Aluminosilicate matrix
a particulate isotopic standard of uranium and plutonium in, 5:852
plutonium in, particulate standard of, 2:81
- Alzheimer patients, treated, quantitative plasma profiling of arecoline in, an automatic reaction control chemical ionization technique in ion trap detector for, 2:168
- AMBER*, in ion mobility spectroscopy of charge-solvated vs salt bridge structures of protonated trimers, 16:1009
- Ambient air, BTEX analysis using APCI-MS/MS, 20:829
- Ambient desorption/ionization mass spectrometry, direct current atmospheric pressure discharges for, 20:837
- Ambient ionization techniques, lessons from, 20:1947
- Ambient pressure ion mobility spectrometry, carbohydrate isomer resolution by ESI-APIMS-TOF-MS, 18:1163
- Ambient pressure pyroelectric ion source, applications to chemical nerve agent detection, 20:2093
- American Society for Mass Spectrometry, 45th conference of, course descriptions and authors of, 8:387
- Amide anions
in solution, ESI-MS of, 11:1061
in solution by ESI-MS, 11:1061
- Amide bond cleavage mechanism, of protonated oligopeptides, MS/MS of, 15:103
- Amide hydrogen, migration of, during ESI, 12:410
- Amide hydrogen/deuterium exchange, modeling data from, in protein-ligand interactions, 15:388
- Amides, protonated, fragmentation of, 9:454
- Amides and amines, proton-bound homodimers and heterodimers of, in gas phase, FTICR-MS of, 13:1273
- Amine-glycerol solutions, solution chemistry and secondary ion emission from, 2:33
- Amine/polyether complexes, protonated, in quadrupole ion trap, threshold dissociation energies of, 14:383
- Amine-trifluoroacetic acid self-assembly, 15:356
- Amines
ES-generated copper(II) complex ions of, CID of, in formation of molecular radical cations of enkephalin derivatives, 12:1114
isomeric arylglycoside gas phase reactivity toward, 15:244
macrocyclic, C_{*n*}H_{*n-1*}NH₂ fragmentation of, electron ionization induced, evidence for the rearrangement of aminocarbene

- radical cations and a comparison with long-chain esters, 1:238
- protonated, 9:1146
- 9-Amino-9-deoxyepiquinine, Michael-Michael-retro Michael addition catalyzed by, ESI-MS, 18:2074
- Amino acid-cyclodextrin complexes, gas phase, chiral recognition in, 12:278
- Amino acid derivatives, alkylamine thiohydantion, electrospray ionization study of, 6:1211
- Amino acid esters, fragmentation induced by triphenylcorrole-(CuIII) complexes, MALDI MS, 18:791
- Amino acid residue, C-terminal, loss of, rearrangement involving, metastable decomposition of peptide $[M + H]^+$ ions via, 1:249
- Amino acid sequence prerequisites for the formation of C_n ions, 4:874
- Amino acid sequences, known, predicting mass spectrometric information from, a computer program (COMPOST) for, 2:174
- Amino acid sequences in a protein database, an approach to correlate tandem mass spectral data of peptides with, 5:976
- Amino acid side chain losses, in electron capture dissociation, 13:241
- Amino acids. *See also specific amino acid*
- α -AA, revising the proton affinity scale of, 17:1275
- analysis by MALDI TOF/TOF MS/MS, 18:279
- aromatic, complexation with Na^+ and K^+ in gas phase, cation- π effects in, 11:1037
- carbon isotope ratio determination in, ICP-MS for, 295
- cationized heterodimers of, decompositions of, in relation to charge location in peptide ions, 4:461
- charged clusters, chirality in formation and stability of, 17:275
- common, encapsulation of by cyclodextrins, electrospray ionization mass spectrometry of, 6:866
- D- and L-, effect of iodine substituents on chiral distribution, 18:1516
- deprotonated, investigation of hydration mechanism, 20:632
- deprotonated, hydration energies from gas-phase equilibria measurements, 19:1091
- determination of unique substitutions, peptide mass mapping, FT-ICR MS, 17:508
- gas-phase enantioselectivity and zwitterionic formation, 17:442
- influence of composition on CID of model peptides, MDS, 18:1625
- intrinsic size parameters of, in predicting mobilities of peptide ions, 12:885
- isotopically labeled, proteome analysis with, 11:78
- ozone oxidation products of, identification by ESI-MS and MS/MS, 11:526
- peptide composition and role in charge inversion, gas-phase ion/ion reactions, 20:180
- polarities of, ECD product ion abundances correlated with, 20:2273
- potassiated, hydration in gas phase, 18:2083
- in probing structure of cytochrome c by ESI-MS, 15:1612
- protonated, hydrogen/deuterium exchange of, in gas phase, 16:1846
- the reactions of ground Cu^+ and Fe^+ with, 7:722
- residues, Cu-binding proteins, identified using MCO reactions and MS, 17:1552
- silver binding energies of, 9:760
- single, distinguished in protein and tertiary structure probed by ECD, 15:1087
- sodiated, gas phase H/D exchange, 17:29
- substitution in the hemoglobin "core," an improved method to determine, electrospray ionization-tandem mass spectrometry analysis of peptides derived by enzymatic digestion of oxidized globin subunits, 7:1040
- substitutions mapped by nano-electrospray MS, avian influenza virus (H5N1) matrix protein 1, 20:312
- underivatized, HPLC/AP-MIPI-MS of, 12:117
- unusual, in a semisynthetic polypeptide, use of high- and low-energy collision-induced dissociation tandem mass spectrometry in the identification of, 7:1034
- Amino acid(s), calixarene non-covalent complexes with, formed by MALDI-MS, 13:964
- α -Amino acids, reevaluation of computed proton affinities for, 20:2116
- β -Amino acids
- cyclic and bicyclic, diastereochemical differentiation, 20:34
- diastereochemical differentiation, FT-ICR-MS, 18:1038
- enantiomers of, separation by MS, 20:1235
- Amino acids or alcohols, detection of, by electron capture negative ion chemical ionization mass spectrometry, pentafluorobenzyl chloroformate derivatization for, 6:525
- Amino alcohols, site-selective gas-phase reaction of, with dimethyl ether ions, 4:230
- Aminocarbene radical cations, evidence for the rearrangement of, electron ionization induced fragmentation of macrocyclic amines $C_nH_{2n-1}NH_2$, 1:238
- Aminocyclitol-aminoglycoside antibiotics, CAD of, 11:200
- Aminoethyl-triphenylphosphonium derivatives of fatty acids, FAB-CAD-B/E linked-scan mass spectrometry of, charge-remote fragmentation during, 3:769
- 9-Aminofluorene-labeled oligosaccharides, MALDI-FTMS characterization of, 12:1254
- Aminoglycosides, silver-coated gold nanoparticles as a matrix in SALDI MS for analysis of, 20:1925
- 2-Aminoimidazolone, oligodeoxynucleotide fragmentation with, 13:1190
- Aminonitropyrene, LC-MS/MS analysis of, 15:1853
- Aminophenylboronic acid, immobilization on magnetic beads, by MALDI, 16:1456
- Ammonia
- complexes with ketones, mobility resolution and mass analysis of ions from, 18:940
- elimination in condensation reactions in protonated dipeptides, 15:1025
- loss of CO and, protonated AGG, MS/MS, 18:1291
- Ammonia-alcohol and water-alcohol protonated clusters, on the structure of, 4:507
- Ammonia laser desorption/chemical ionization with ammonium bromide, Fourier transform ion cyclotron resonance mass spectrometry of aromatic hydrocarbons, 2:299
- Ammonia radical cation, water-solvated, reactions with carbonyl compounds, 15:966
- Ammonium acetate, thermospray reagent plasma from, controversial ions, 5:186
- Ammonium bromide, ammonia laser desorption/chemical ionization with, Fourier transform ion cyclotron resonance mass spectrometry of aromatic hydrocarbons, 2:299
- Ammonium-counter ion clusters, studied by ESI MS, 18:285
- Ammonium ion, oligonucleotide charge state distribution and, in ESI-MS, 8:155
- Ammonium ion affinities and gas-phase basicities of polyethers, by kinetic method and ligand exchange technique, 4:718
- Ammonium ion-mediated resorcarene capsules, ESI-FTICR-MS study on gas phase structure and ammonium ion affinity of tetraethyl resorcarene and its per-methylated derivative, 143
- Ammonium ion/polyether complexes, gas-phase, hydrogen-bonding interactions in, 5:260

- Ammonium ions, quaternary, CSR fragmentation reactions of, 12:571
- Ammonium radical, hypervalent, intramolecular interactions in, 8:1111
- Ammonium salts, diquaternary, observed charge states of, and ion pairing, in electrospray ionization mass spectrometry, effects of solvent and counterion on, 7:1050
- Ammonium sulfate in-source decay MALDI-TOFMS, peptide sequence by, 11:1000
- Ammonium surfactants
- condensation of glycerol with, fast-atom bombardment-induced, II, time dependence of mass spectra and tandem mass spectra, 5:92
 - fast atom bombardment-induced condensation of glycerol with, I, regioselectivity of the adduct formation, 3:318
- Amoxicillin degradation products, ion trap MS confirmation of, ESI-TOF mass determination in, 16:1670
- AMPPNP, ESI-TOF-MS of, 15:1392
- AMUSE. *See* Array of Micromachined UltraSonic Electrospray
- Amyloid- β
- peptide analysis via mini-ball mill and MALDI MS, 18:1533
 - production and clearance rates, quantified using SILT, 18:997
- Amyloid- β -peptide
- interaction with oleuropein, ESI-MS, 17:568
 - localization of noncovalent binding site between oleuropein and, ESI-FT-ICR MS, 19:1078
- Amyloid precursor proteins, potential cleavage sites of, LCMS/MS of, 15:1780
- A_n cross-ring cleavage, in glycoconjugate mixtures, 15:1863
- Anabolic steroid glucuronides, ESI and APCI-MS/MS of, 11:722
- Anabolic steroids, fragmentation pathways, ESI-MS/MS, 17:477
- Analog-to-digital converters, two, increasing the dynamic range of a transient recorder by using, 7:107
- Analysis and estimation of systematic errors originating from Coulomb mutual interaction and image charge in Fourier transform ion cyclotron resonance precise mass difference measurements, 4:855
- Analysis by fast-atom bombardment tandem mass spectrometry of phosphatidylcholine isolated from heart mitochondrial fractions, evidence of incorporation of monohydroxylated fatty acyl moieties, 7:50
- Analysis of an Egyptian mummy resin by mass spectrometry, 3:582
- Analysis of coumarins by micro high-performance liquid chromatography-mass spectrometry with a particle beam interface, 6:132
- Analysis of diacyl peroxides by Ag⁺ CIS-MS/MS: free radical pathways of complex decomposition, 12:449
- Analysis of hydrocarbon dendrimers by laser desorption time-of-flight and Fourier transform mass spectrometry, 5:731
- Analysis of hydrocarbon polymers by MALDI-FTMS, 8:225
- Analysis of hydroxy fatty acids as pentafluorobenzyl ester, trimethylsilyl ether derivatives by electron ionization gas chromatography/mass spectrometry, 6:40
- Analysis of lipid hydroperoxides and long-chain conjugated keto acids by negative ion electrospray mass spectrometry, 6:1190
- Analysis of long-chain fatty acyl coenzyme A thioesters by negative ion fast-atom bombardment mass spectrometry and tandem mass spectrometry, 5:416
- Analysis of oligodeoxynucleotides by negative-ion matrix-assisted laser desorption mass spectrometry, 4:955
- Analysis of peptides, proteins, protein digests, and whole human blood by CE/ESI-MS using an in-capillary electrode sheathless interface, 9:1081
- Analysis of peptides and proteins containing nitrotyrosine by MALDI-MS, 12:438
- Analysis of polycyclic aromatic hydrocarbons by ion trap MS/MS, 8:183
- Analysis of quaternary protein ensembles by MALDI-MS, 8:1046
- Analysis of salen complexes by FAB-MS and APCI-MS, 8:996
- Analysis of serum protein precipitated with antiserum by matrix-assisted laser desorption ionization/time-of-flight and electrospray ionization mass spectrometry as a clinical laboratory, 6:854
- Analysis of solution residues by glow discharge mass spectrometry, 4:47
- Analysis of the accuracy of determining average molecular weights of narrow dispersity polymers by MALDI-TOFMS, 9:275
- Analysis of triacylglycerols and whole oils by MALDI/TOFMS, 10:983
- Analysis of unsaturated compounds by Ag⁺ coordination ion-spray MS: studies of formation of Ag⁺/lipid complex, 872
- Analyte
- characteristics of, and organic molecule responsiveness to negative ion ESI MS, 16:446
 - concentration of, effect on peptide ion internal energies produced by ESI and AP-MALDI, 16:743
- Analyte charge state, signal intensity, and stability in negative ion electrospray mass spectrometry, implications for the mechanism of negative ion formation, 4:546
- Analyte electrolysis, in electrospray emitter, 11:951
- Analyte ions, reaction of, with neutral chemical ionization gas, 3:549
- Analyte response
- laser ablation ICPMS, 17:641
 - salt concentration and, ESI-MS of, 10:625
- Analytes
- ionization of, effect of solvent and surface effects on, DIOS MS, 19:8
 - neutral radical induced transformations in APPI and near-VUV APLI, 20:1868
 - polyanionic, negative ion fast-atom bombardment mass spectrometry of, enhancing sensitivity in, a class of cationic matrix additives for, N-alkylnicotinium halides, 5:935
 - with two basic functional groups, identification by FTICR, 20:1251
- Analytes from aqueous solutions, mass spectrometry of, a heated electrospray source for, 5:791
- Analytical approach for description of ion motion in quadrupole mass spectrometer, 14:818
- Analytical chemistry, mass spectrometry and, 6:993
- Analytical information from mass spectrometry, past and future, 1:1
- Analytical mass spectrometry
- environmental applications of, 13:898
 - principles of collisional activation in, 3:599
- Analytical performance of a high-pressure radio frequency-only quadrupole collision cell with an axial field applied by using conical rods, 9:775
- Androstadienolones, MS of, 16:1660
- Androstanolones, MS of, 16:1660
- Anesthetic, local, cyclodextrin non-covalent complexes with, comparison using CE and ESI-MS, 13:928
- Angiotensin-converting enzyme (ACE) inhibitors, MS fragmentation of, 15:1300
- Angiotensin II, doubly protonated, formed by electrospray ionization, charge-separation reactions in tandem mass spectrometry of, experimental considerations and structural implications, 7:30
- Angiotensin II (M + 2H)²⁺ ions, location of charge in, 11:40

- Angiotensin peptides and zinc metal ions, interaction of, by electrospray ionization mass spectrometry, 5:959
- Angle-resolved neutralization-reionization mass spectrometry, 2:492
- Angle-resolved translational energy spectra of survivor ions, high-energy collisional activation studied via, 4:835
- Aniline oligomers, surface-assisted reduction of, in APCI and APPI, 13:109
- Anilino, membrane introduction MS of, 9:234
- 1-Anilino-8-naphthalene-sulfonate, protein binding by, ESI-MS of, 12:428
- Animal feed, multiresidue screening in food and feed matrices, Oribtrap MS, 20:1464
- Anion
 distonic radical, sulfur-containing, CH₂S-CH-CN, formation and characterization of, in gas phase, 6:71
 electrospray-generated, ion/ion reactions in, 13:614
- Anion adduct generation, ionization and, during MALDI, 13:1218
- Anion detection, dicationic reagents in, using positive ion mode ESI-MS, 19:261
- Anion- π interactions, ESI-MS study of, 20:257
- Anionic α -adducts, in gas phase, using electrospray ion source as chemical reactor, 15:927
- Anionic adducts
 APCI-MS identification of, 16:1377
 of carbohydrates, 16:622
- Anionic complexes of alkali metal ions and peptides, gas-phase, structure and collision activated decompositions, 5:137
- Anionic nucleophiles
 fluorotrimethylsilane affinities of, 16:697
 gas phase reactions of chloramine with, 12:139
- Anionic peptides, hydrogen atom scrambling, collisional activation by MALDI TOF/TOF MS, 19:1719
- Anions
 alkali-metal-carboxylate radical, in fast-atom bombardment, 6:1252
 copper phthalocyanine, heterolytic dissociation of negative ions into counter ions, 19:375
 electron, from deoxy-polyadenylates, decomposition of, 8:148
 oligonucleotide, multiply charged, proton transfer reactions of protonated pyridine with, gas phase, ion-ion reactions in, 6:529
 tauroconjugated bile acid, fast-atom bombardment mass spectrometry and low energy collision-induced tandem mass spectrometry of, 6:588
- Anodic stripping voltammetry, with ICP-MS, NCBFE and NCMFE for, 17:945
- Anomeric configuration of glycosyl esters of nucleoside pyrophosphates and polyisoprenyl phosphates, by fast-atom bombardment tandem mass spectrometry, 7:541
- ANRORC reaction, ionic products from, isomer differentiation by charge inversion tandem mass spectrometry, 3:467
- Anthracycline aminodisaccharides, collision-induced dissociation mass spectra of, internal monosaccharide losses in, 9:710
- Anthraquinone-2-carbonyl chloride, as an alcohol derivatization reagent for negative ion chemical ionization mass spectrometry, 6:148
- Antibiotic-containing mixtures from fermentation broths, rapid analysis of, by using liquid chromatography-electrospray ionization-mass spectrometry matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry, 7:1227
- Antibiotics
 complex oligosaccharide, quadrupole ion trap multiple-stage MS analysis of, 13:1313
 linked to peptides and proteins released by ECD-FTICR-MS, 14:302, 14:682
- macrolide
 application of nanoscale packed capillary liquid chromatography (75 m id) and capillary zone electrophoresis/electrospray ionization mass spectrometry to, 3:563
 fast-atom bombardment and tandem mass spectrometry of, 5:151
 polyether ionophore, gas phase stabilities of, by CID, 16:779
 in vancomycin group, bacterial cell wall peptide precursor interaction with, hydrogen/deuterium exchange in, 9:1255
- Antibodies
 IgG, disulfide linkages characterized by reversed phase HPLC with negative ESI MS, 17:1590
 improving mass accuracy of HPLC-EI-TOF-MS, 17:867
 molecular mass analysis by on-line SEC-MS, 19:502
 mouse switch variant, by matrix-assisted laser desorption ionization mass spectrometry and electrospray ionization mass spectrometry, 7:707
- Antibody beads, peptides affinity-bound to, direct MALDIMS/MS of, 14:1076
- Anticancer drugs, low-molecular weight, and their analogues, electrospray ionization mass spectrometry for, 4:588
- Antifungal lipopeptide iturin A₂, sodium stabilization of, during ESI-MS, 12:505
- Antigens, mycobacterial glycopeptidolipid, fast atom bombardment mass spectrometry of, structural characterization by charge remote fragmentation, 3:656
- Antimalarials, solid artesunate, quantization of active ingredients by DESI MS, 19:380
- Antimony potassium tartrate, bis(L-alanine)ethylenediamine cobalt(III) association with, chiral recognition in, 11:804
- Antimony(III)-D, L-tartrates, enantioselective binding in solution and gas phase, ESI-MS, 20:2100
- Antioxidants, predicting activity of, 15:848
- Antisense therapeutic, sequencing of, by CID, 12:805
- Antiserum, serum protein precipitated with by matrix-assisted laser desorption ionization/time-of-flight and electrospray ionization mass spectrometry as a clinical laborator, 6:854
- Antithrombotic, nonpeptide, in dog plasma, by microbore height-performance liquid chromatography-tandem mass spectrometry utilizing pneumatically assisted electrospray ionization, 7:1238
- Antiviral agents, pyrimidinal, CID mechanisms of ion formation, ESI-MS/MS, 18:1477
- AP. *See* Atmospheric pressure; high performance liquid chromatography/atmospheric pressure microwave-induced plasma ionization MS
- AP MALDI. *See* Atmospheric pressure matrix-assisted laser desorption/ionization
- AP-MIPI. *See* Atmospheric pressure microwave-induced plasma ionization
- APCI. *See* Atmospheric pressure chemical ionization
- APCI (atmospheric pressure chemical ionization) mass spectrometry, salen complex analysis by, 8:996
- APCI low energy collision-induced dissociation fragmentation of protonated ortho silicates, McLafferty or ion-neutral complex rearrangements, 5:544
- APCI-MS. *See* Atmospheric pressure chemical ionization mass spectrometry
- APCI-MS/MS *See* Atmospheric pressure chemical ionization tandem mass spectrometry
- API. *See* Atmospheric pressure ionization
- APIMS. *See* Ambient pressure ion mobility spectrometry
- ApoE* gene, genotyping of SNPs and indels by MS, 17:3

- Apomyoglobin, conformation of gas-phase ions, 19:1907
- Apoprotein B-100, human oxidized, cysteic acid residue in, tandem mass spectrometric characterization of, 6:242
- Apparent gas-phase acidities of multiply protonated peptide ions, ubiquitin, insulin B, and renin substrate, 7:1211
- Apparent proton affinities of highly charged peptide ions, 10:483
- APPI. *See* Atmospheric pressure photoionization mass spectrometry
- APPI-MS. *See* Atmospheric pressure photoionization mass spectrometry
- APPIS. *See* Ambient pressure pyroelectric ion source
- Application of electrospray mass spectrometry and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry for molecular weight assignment of peptides in complex mixtures, 4:670
- Application of electrospray mass spectrometry in probing protein-protein and protein-ligand noncovalent interactions, 4:624
- Application of ESI-FAIMS-MS to analysis of tryptic peptides, 13:1282
- Application of integrated MALDI-TOFMS, ESI-MS, MS/MS, to characterize complex polyol mixtures, 12:55
- Application of ion-impact energy measurement to ESI-MS of proteins and protein mixtures, 12:826
- Application of microelectrospray liquid chromatography techniques to FTICR-MS to enable high-sensitivity biological analysis, 9:333
- Application of nanoscale packed capillary liquid chromatography (75 m id) and capillary zone electrophoresis/electrospray ionization mass spectrometry to the analysis of marcolide antibiotics, 3:563
- Application of nonpolar matrices for analysis of low molecular weight nonpolar synthetic polymers by MALDI-TOFMS, 11:731
- Application of nonresonance excitation to ion trap tandem mass spectrometry and selected ejection chemical ionization, 7:668
- Application of normal phase liquid chromatography/mass spectrometry by using coaxial continuous flow fast atom bombardment, 3:575
- Application of parallel LC-MS for high throughput microsomal stability screening of compound libraries, 13:155
- Application of self-ionization for enhancing stereochemical and positional effects from arylglycosides under electron ionization conditions in an ion trap mass spectrometer, 14:851
- Application of simultaneous excitation/detection to generate real-time excitation profiles in FTICR-MS, 11:1009
- Application of sustained off-resonance irradiation, the beat frequency measurement and radial separation of mass-selected ions, 8:86
- Applications of collision dynamics in quadrupole mass spectrometry, 9:101
- Applications of electrospray ionization mass spectrometry to neutral organic molecules including fullerenes, 4:596
- Applications of external customized waveforms to a commercial quadrupole ion trap, 10:355
- Applied voltages and reaction times, negative chemical ionization in quadrupole ion trap mass spectrometry, 5:757
- Applying charge discrimination with electrospray ionization-mass spectrometry to protein analyses, 6:1098
- Approach to correlate tandem mass spectral data of peptides with amino acid sequences of a protein database, 5:976
- Aquatic environments
lake water testing of chemistry with portable and AUV-based MS, 19:1403
- membrane inlet MS for online and in situ analysis of methane, 19:1395
- Aqueous sample derivatizing agents, comparing highly fluorinated chloroformates as, 16:803
- Aqueous samples, pesticides in, by solid-phase microextraction in-line coupled to gas chromatography-mass spectrometry, 6:1119
- Aqueous solution chemistry, are the electrospray mass spectra of proteins related to?, 3:216
- Aqueous solutions
analytes from, mass spectrometry of, a heated electrospray source for, 5:791
desorption/ionization from, 13:354
- Aqueous solutions of metal salts, electrospray mass spectrometry, 3:281
- Arabidopsis thaliana*, phosphatidylglycerol, analysis by ESI QIT and MS/MS, 18:783
- Arachidonate containing glycerophosphocholines, molecular species analysis of, by tandem mass spectrometry, 2:45
- Arachidonic acid
chemically reactive metabolites of, MA analysis of, 13:1227
free-radical oxidation of, four regioisomers of F2-isoprostanes formed by, mass spectrometric analysis of, 7:490
- Archiving of data, electronic, networking mass spectrometer data systems for, 4:742
- ARE. *See* Axial resonant excitation
- Are the electrospray mass spectra of proteins related to their aqueous solution chemistry?, 3:216
- Arecoline, plasma profiling of, in treated Alzheimer patients, an automatic reaction control chemical ionization technique in ion trap detector, 2:168
- Argentinated peptides, CID of, product from, 12:163
- Arginine
chiral resolution of, by ESI-ion trap MS/MS, 16:825
methylated, quantified with nanoflow LS-ETD-Orbitrap MS, 20:507
molecular recognition by anionic salt, 15:616
- Arginine-aldehyde adducts, in cytochrome c modification by 4-hydroxy-2-nonenal, 15:1136
- Arginine-containing peptide ions formed by electrospray ionization, modeling the maximum charge state of, 7:972
- Arginine dipeptide isomers, discrimination of, ESI-ion trap MS and kinetic method for, 15:840
- Arginine residues, selective titration by sulfonates, 12:810
- Argon
collision cross sections of myoglobin and cytochrome c with, 8:681
effect on fluorescence yield of trapped gas-phase ions, 17:855
quantitative analysis of, by quadrupole ion trap MS, 13:1120
reaction with CO₂ and SO₂, rate constants for as functions of temperature, 10:815
- Argon and helium, as collision gases in the high energy collision-induced decomposition of MH⁺ ions of peptides, 3:502
- Argon-helium inductively coupled plasma mass spectrometry, matrix-induced interferences in, 2:355
- Argon/hydrogen mixture, high-resolution glow discharge mass spectrometer operated with, relative ion yields measured with, 4:278
- Aristolochic acids, DNA adducts, LC-MS/MS analysis, 18:642
- Aromatic amines, protonated, with ND₃, H/D exchange reactions, reactive collisions in quadrupole cells and, 6:19
- Aromatic beta-hydroxyoximes, (M + H)⁺, (M + Cat)⁺, and (M-H)⁻ ions of, charge-remote fragmentation and the two-step elimination of alkanols from, 4:819
- Aromatic carbonyl compounds, protonated, remote fragmenta-

- tions of, via internal reactions in intermediary ion-neutral complexes, 3:417
- Aromatic compounds
 polycyclic, analysis with APCI-MS, charge exchange and proton transfer in, 15:301
 simple, at atmospheric and at reduced pressure conditions, a comparison of the reactions of the oxide radical anion (O^-) with, 5:367
- Aromatic diazoamino compounds, thermolysis of, 9:234
- Aromatic hydrocarbon diolepoxide adducted peptides, aromatic, characteristics of high energy collision-induced dissociation tandem mass spectra of, 5:649
- Aromatic hydrocarbon polycyclic. *See* Polycyclic aromatic hydrocarbon
- Aromatic hydrocarbons
 Fourier transform ion cyclotron resonance mass spectrometry of, ammonia laser desorption/chemical ionization with ammonium bromide, 2:299
 polycyclic
 in ancient rocks, μ L2MS vs GC/MS of, 12:989
 C_{10} to C_{160} , flame-generated, by atmospheric-pressure chemical ionization mass spectrometry with liquid introduction via heated nebulizer interface, 7:276
 by high-performance liquid chromatography-particle beam-mass spectrometry, 4:898
 in solid matrices, 14:562
 polynuclear, nitro-substituted, in airborne particulate matter and soil, sub-parts-per-billion determination by electron capture-tandem mass spectrometry, 7:1255
- Aromatic nucleophilic substitutions, reactions of halonitrobenzenes and dinitrobenzene with carbanions, gas phase, 18:1351
- Aromatic phosphopeptides, dissociation of, ESI-FTICR-MS and IRMPD of, 15:121
- Aromatic rings in metal chelation, peptide side chain influences on metal ion binding site in metal ion-cationized peptides and, 6:1079
- Aromatics, nitrogen-containing, speciation by APPI or FT-ICR MS, 18:1265
- ArPd(PPh₃)₂I complexes, gas phase fragmentation reactions, ESI-FTICR-MS, 17:1582
- Array detector, improved detection limits for electrospray ionization on a magnetic sector mass spectrometer by using, 5:194
- Array of Micromachined UltraSonic Electrospray, ion source for MS, charge separation in, 20:1684
- Array of micromachined ultrasonic electrosprays, Venturi-assisted, comparison of internal energy deposition with Venturi-assisted ESI, 19:1320
- Arrhenius activation parameters, for loss of neutral nucleobases, 15:55
- Arsenic-glutathione complexes, analysis by HPLC/ES MS/MS, 19:1559
- Artemisinin-heme noncovalent complexes, characterization by ESI-MS and CID-MS, 15:1181
- Artesunate antimalarials, solid, quantization of active ingredients by DESI MS, 19:380
- Artifact peak intensity, in linked scans for dissociations occurring in the first field-free region of sector spectrometers, 8:554
- Artifact peaks, in linked scans, 8:554
- Artificial neural networks, using high-energy collision-induced dissociation spectra, peptide sequence determination from, 6:947
- Aryl and alkyl esters of benzenedicarboxylic acids, gas-phase reactions of O_2^- with, 5:990
- Aryl radicals, membrane introduction MS of, 9:234
- Arylamine DNA adducts, isotope dilution LC-MS/MS detection method for, 14:1057
- Arylazido groups, photolabile, peptides modified with, MALDI-MS analysis of, 15:1156
- Aryldiazonium, membrane introduction MS of, 9:234
- Arylglycosides
 isomeric, chemical ionization of, 15:244
 self-ionization of, 14:851
- Arylnitrenium ions, generation of, by nitro-reduction and gas phase N-heterocycle synthesis, 15:1679
- Arylnitrile radical cations, nitroaromatic compound denitration by, 11:33
- ASAP. *See* Atmospheric solids analysis probe
- ASGDI. *See* Atmospheric sampling glow discharge ionization
- Asilomar Conference on Mass Spectrometry, report of the twenty-third meeting, R19:1 (July)
- Asilomar conference on mass spectrometry
 the 11th, highlights of, 7:505
 the 12th, elemental mass spectrometry, 7:613, 7:693, 7:760, 7:987
- ASMS conference on mass spectrometry and allied topics, June 2000 in Long Beach, CA, review of, 11:921
- Asparagine, protonated, thermodynamics and mechanism of decomposition, GIB-MS/MS, 20:852
- Aspartic acid
 isoaspartic acid and, differentiated by electron transfer dissociation, 17:15
 residues of proline and, effect on peptide dissociation, 19:344
 side chain effects, experimental and theoretical insights, 18:121
- Aspartic acid peptides, of fixed charge derivative, hydrogen/deuterium exchange of, 16:1067
- Association constants, of adenylate kinase, with noncovalent inhibitors, 14:442
- ASV. *See* Anodic stripping voltammetry
- Asymmetric waveform ion mobility spectrometry (FAIMS), tandem, coupled to TOFMS, ion trapping in, 12:1320
- "Ate" class, distonic ions of, 10:896
- Atmospheric pressure photoionization (APPI) mass spectrometry, ionization efficiency in, solvent flow rate and, 16:1399
- Atmospheric chemistry, of hydroxymethylpropanal, ITMS in, 13:530
- Atmospheric conditions, comparison of the reactions of the oxide radical anion (O^-) with simple aromatic compounds at, 5:367
- Atmospheric gases, glow discharge in, structures in NO_3^- formed via, 1:217
- Atmospheric pollutants, real time monitoring of, ion trap detector used for, the effects of air and helium on, 4:774
- Atmospheric pressure
 electrospraying water in air at, 15:253
 environmental, and ionization pathways of polyfluorinated chemicals, 14:1177
 in HPLC of underivatized amino acids, 12:117
 ion trapping in tandem FAIMS-FAIMS coupled to a TOFMS, studies with electrospray-generated gramicidin S ions, 12:1320
 negative, chemical ionization liquid chromatography-mass spectrometry, effect of high-performance liquid chromatography mobile phase components on, 7:1059
- Atmospheric pressure (AP) liquid chromatography (LC) mass spectrometry, GC/MS ion source combined with, advantages of, 16:1730
- Atmospheric pressure (AP) matrix-assisted laser desorption/ionization (MALDI)
 ionization mechanism, analyte concentrations, and ion cool time effects on peptide ion internal energies produced by, 16:743
 on QqLIT instrument, 16:176

- Atmospheric pressure (AP) matrix-assisted laser desorption/ionization (MALDI) Fourier transform (FT) mass spectrometry, of normal and chemically modified RNA, 16:199
- Atmospheric pressure chemical ionization (APCI). *See also* liquid chromatography-atmospheric pressure chemical ionization.
- of alkanes, alkenes and cycloalkanes, 5:177; Erratum, 5:1026
 - archaeal tetraether lipids analyzed by LC-MS/MS, 20:51
 - of fluorinated penols in APCI-MS, MS/MS, and IMS, 10:1157
 - for drug analysis, 13:204
 - ionization mechanisms related to negative ion APPI, DART and, 20:1518
 - liquid chromatography mass spectrometry interface, 7:69
 - in LC-MS/MS quantification of lanosterol and FF-MAS in vitro, 15:1216
 - of mutagenic nitropyrenes, 15:1853
 - reduction in, 13:109
 - for studying non-polar isomeric hydrocarbons using IMS and MS with different ionization techniques, 13:1078
- Atmospheric pressure chemical ionization (APCI) mass spectrometry, in anionic adduct identification, 16:1377
- Atmospheric pressure chemical ionization Fourier transform ion cyclotron resonance magnetic resonance spectrometry permanent magnet, 18:1552
- Atmospheric pressure chemical ionization liquid chromatography mass spectrometry interface, optimization of, 7:69
- Atmospheric pressure chemical ionization mass spectrometry
- air plasma, study of alkanes, 20:697
 - hydrolysis products of sulfur mustards analyzed by, 18:1364
 - positive ion mechanisms, 19:1927
 - for routine sample analysis, 6:387
 - tetracycline degradation by ozone in aqueous medium, 18:679
- Atmospheric pressure chemical ionization-mass spectrometry
- high-performance liquid chromatography, determination of the renin inhibitor CP-80,794 in human serum, 2:164
 - salen complex analysis by, 8:996
- Atmospheric pressure chemical ionization-mass spectrometry alkyl ethoxysulfate mixture analysis by, 10:529
- Atmospheric pressure chemical ionization mass spectrometry (APCI-MS)
- charge exchange and proton transfer in, in analyzing polycyclic aromatic compounds, 15:301
 - gas and liquid reactions in, 15:311
- Atmospheric pressure chemical ionization mass spectrometry with liquid introduction via heated nebulizer interface, characterization of flame-generated C₁₀ to 160 polycyclic aromatic hydrocarbons by, 7:276
- Atmospheric pressure chemical ionization-mass spectrometry-alkyl ethoxysulfate mixture analysis by, of nitrocatechol glucuronides, 10:537
- Atmospheric pressure chemical ionization tandem mass spectrometry, ambient air BTEX analysis using, 20:829
- Atmospheric pressure chemical ionization tandem mass spectrometry (APCI-MS/MS), of eight anabolic steroid glucuronides, 11:722
- Atmospheric pressure corona discharge ionization, nitric oxide-assisted, for the analysis of automobile hydrocarbon emission species, 5:1107
- Atmospheric pressure interfaces, in LC-MS/MS, in residue determination of compounds, 16:1619
- Atmospheric pressure interference, high-performance liquid chromatography-mass spectrometry of porphyrins by using, 7:965
- Atmospheric pressure ion lens, improving nebulizer-assisted electrospray ion sources with, 13:906
- Atmospheric pressure ion mobility (APIM), in separation of so-diated isobaric saccharides, 16:660
- Atmospheric pressure ionization
- ESI source with, 9:1196
 - LC/MS and, 9:187
 - techniques for the analysis of N-methyl carbamate pesticides by liquid chromatography mass spectrometry, 3:378
- Atmospheric pressure ionization mass spectrometry
- chemical results obtained by, internal standard correction, 2:427
 - detection mass bias in, 2:232
 - detection of thermally unstable intermediate in Wittig reaction by, 9:1168
- Atmospheric pressure laser ionization, near-VUV, neutral radical induced transformations of analyte ions in APPI and, 20:1868
- Atmospheric pressure laser ionization mass spectrometry, spatially resolved DIA in, 19:400
- Atmospheric pressure MALDI, influence of matrix on analyte fragmentation, 17:1005
- Atmospheric pressure matrix-assisted laser desorption/ionization
- infrared
 - of aqueous solutions, 13:354
 - in study of noncovalent sugar-sugar complexes, 11:58
- phosphopeptide fragmentation by, 13:274
 - time scale of internal energy relaxation, by nano-ESI and, in QITMS, 20:1801
- Atmospheric pressure microwave-induced plasma ionization source for molecular mass spectrometry, 9:42
- Atmospheric pressure photo ionization, neutral radical induced transformations of analyte ions in near-VUV APLI and, 20:1868
- Atmospheric pressure photo ionization mass spectrometry, dopant-assisted, of polyisobutylene derivatives, 20:2342
- Atmospheric pressure photoionization (APPI)
- ionization mechanisms related to negative ion APCI, DART and, 20:1518
 - in LC-MS/MS quantification of lanosterol and FF-MAS in vitro, 15:1216
 - of mutagenic nitropyrenes, 15:1853
 - negative ion, EC, DEC, proton transfer and anion attachment, 18:1789
 - proton transfer investigated by FT-ICR, 18:1682
 - reduction in, 13:109
- Atmospheric pressure photoionization mass spectrometry
- dopant for charge-exchange ionization investigated by, 19:955
 - mechanism of [M+H]⁺ formation in, 19:1579
 - mobile phases and VUV lamps, detection of PAH compounds, 18:589
 - of polyisobutylene derivatives, 19:656
 - speciation of nitrogen-containing aromatics FT-ICR MS or, 18:1265
- Atmospheric pressure photoionization mass spectrometry (APPI-MS), negative ion, 15:203
- Atmospheric pressure thermal dissociation, sulfopeptides and phosphopeptides, 19:1898
- Atmospheric sampling glow discharge ionization, analytical performance of, factors influencing, 14:1315
- Atmospheric solids analysis probe, analysis of ergosterol pathway inhibition in fungi, 18:1274
- Atomic fragment losses, slower than polyatomic fragment losses, 14:136
- Atomic ionization, volumetric intensity dependence on, in laser focus, 16:82
- Atomic ions, resonance ejection of, enhanced abundance sensi-

- tivity by, plasma source ion trap mass spectrometry, 7:1161
- Atomic transition metal ions, reactions with long chain alkanes, 12:367
- Atomic transition metals, ion mobility source for guided ion beam experiments, 18:1196
- Atomic weight standard, $12\text{H}_6\text{O}$ as, unsuitability of, 6:1243
- Atorvastatin and active metabolites, in plasma, HPLC-MS/MS of, 10:55
- ATPase supramolecular protein complex, H/D exchange on, MALDI-MS analysis of, 14:471
- Attachment sites to peptidoglycan, Braun's lipoprotein, characterization of, by ^{242}Cf -PD and MALDI time-of-flight mass spectrometry, 6:892
- Attomole/zeptomole per microliter sensitivity for peptides, microelectrospray, 5:867
- Aurum Dataset, MALDI-TOF/TOF MS for protein standards, 18:850
- Automated assignment of charge states from resolved isotopic peaks for multiply charged ions, 6:52
- Automated charge state determination, of isotope-resolved MS by PTFT, 19:46
- Automated data interpretation based on concept of negative signature mass \ddot{o} for mass-mapping disulfide structures of cystinyl proteins, 14:1032
- Automated digestion and droplet deposition microfluidic chip, for MALDI/TOF MS, 19:964
- Automated high-speed analysis of selected organic compounds in urban air by on-line isotopic dilution cryofocusing gas chromatography/mass spectrometry, 5:1001
- Automated nanospray, chip-based emitters in, in pharmaceutical compound quantitative analysis, 16:363, 16:798 (erratum)
- Automated orthogonal control system, for ESI, 15:1201
- Automated reduction and interpretation of high resolution ESI-MS of large molecules, 11:320
- Automatic analysis of H/D exchange mass spectra of peptides and proteins using calculations of isotopic distributions, 12:1153
- Automatic gain control, in MS, using jet disrupter electrode, 16:244
- Automatic reaction control chemical ionization technique in ion trap detector for quantitative plasma profiling of arecoline in treated Alzheimer patients, 2:168
- Automation
 data-controlled, of liquid chromatography/tandem mass spectrometry analysis of peptide mixtures, 7:532
 of data massaging, interpretation, and e-mailing modules for high throughput open access MS, 10:1174
 of FT-ICR MS for acquisition, e-mailing of high-resolution exact-mass ESI MS data, 10:1166
 of high resolution MS for the synthetic chemistry, 10:546
 of MS imaging with a MALDI-TOF instrument, 10:67
- Automobile hydrocarbon emission species, analysis, nitric oxide-assisted atmospheric pressure corona discharge ionization for, 5:1107
- Autonomous underwater vehicles, lake water testing of chemistry with MS based in, 19:1403
- Avian influenza virus (H5N1), matrix protein 1, AA substitutions mapped by nanoelectrospray MS, 20:312
- Avidin tetramer, observation of noncovalent complexes to, by electrospray ionization mass spectrometry, 5:201
- Axial energy, cyclotron frequency and, 19:467
- Axial ion ejection, from linear quadrupole ion trap, 14:1130
- Axial resonant excitation, linear ion trap mass selective ejection by, 17:685
- Axialization, selective parent ion, for improved efficiency of collision-induced dissociation in laser desorption/ionization Fourier transform ion cyclotron resonance mass spectrometry, 5:900
- Aza-18-crown-6-ether, proton affinity of, 13:493
- 4-Aza-3-ene-1,6-diyne, adduct between DNA and, ESI-MS, 17:1342
- 4-Azasteroids, fragmentation of, 12:385
- 3-Azidopropionitrile, transition metal complexation by, ESI-MS, 18:453
- Azo dyes, fast-atom bombardment mass spectra of, effect of tautomerization on, 5:407
- Azobeneze group, cyclodextrin bearing by, identification by ESI-MS, 14:1116
- B
- b^* , γ cleavages, polyamidoamine dendrimer ions, observation of, ECD-MS, 17:536
- β -Diketones, m/z 69 ion from, 9:29
- B-DNA helix, stability in a solvent-free environment, 18:1188
- b_2 ion, proline-containing, gas phase ion structure of, 17:20
- b ions
 cyclic oxazolone structure of, 16:1941
 in ECD, 16:1932
- b_2 ions
 MS analysis of, 9:1002
 in peptide mass spectra, structure and fragmentation of, 11:427
- b_5 ions, peptide sequence scrambling through cyclization of, CID, 19:1776
- b -type ions
 b_9 ion cyclization, 20:2248
 b_{2+} ion formation, fragmentation of doubly-protonated tripeptides, ESI MS, 20:2135
 b_2 ion structure, generated from doubly protonated tryptic peptides, 20:618
 influence of peptide size on sequence scrambling during CID, of, 20:2174
- Bacillus licheniformis*, Glu enantiomers analyzed by ESI/FTICR MS, 17:442
- Backbone-modified peptides, structural determination of, tandem mass spectrometry for, 1:174
- Background ions, produced by fast-atom bombardment of glycerol, 5:72
 origins and structures of, 5:72
- Background reduction, in orthogonal TOF-MS, 8:125
- Background reduction in ^{252}Cf -plasma desorption mass spectrometry, pulsed ion deflection system for, 3:706
- Bacteria
 MALDI-MS of, internal control for, 10:1131
 quality and reproducibility of, 14:342
 whole, MS profiling of, electron and chemical ionization mode in, 10:747
 whole cell
 FTMS of, 14:1306
 MALDI-TOFMS of, 14:1306
- Bacterial analysis, by MALDI-TOF MS, inter-laboratory comparison of, 16:456
- Bacterial cell proteins, surfactant-treated, MALDI-MS of, 16:1422
- Bacterial cell wall precursor peptides, vancomycin antibiotic interaction with, hydrogen/deuterium exchange in, 9:1255
- Bacterial classification, neural network models for, 13:10
- Bacterial fatty acids, picolinyl esters of, 14:58
- Bacterial identification
 flow injection ESI-MS in, 13:118
 by ICM-MS, limits of, 12:49
- Bacterial lipooligosaccharides, LDI TOF-MS of, 8:645
- Bacterial spores, MALDI-TOF-MS of, 15:1444
- Bacteriotoxin, E9:Im9 complex, analysis using lysine acetylation and nano-LC-MALDI-MS/MS, 17:983

- Bacteriovorax stolpii*, UKi2, identification of SPNL molecular species, qTOF-II MS, 18:394
- Baculovirus expressed mouse interleukin-3, application to, liquid secondary ion mass spectrometry analysis of permethylated, n-hexylamine derivatized oligosaccharides, 3:345
- Baeyer-Villiger solid-state organic reactions, monitoring kinetics of, DESI, 20:2087
- Ballpoint dyes, photofading on paper analyzed by LDI and MALDI MS, 17:297
- Base composition of double-strand DNA, accurate, by mass spectrometry, 7:1266
- Bases, strong, reactions with vinyl fluoride formation and characterization of 1-fluorovinyl anion and the fluoride-acetylene hydrogen-bonded complex, 3:91
- Basic fibroblast growth factor, ATP binding to, detected by MALDI-TOF and ESI-MS, 19:91
- Basicities, gas-phase
and ammonium ion affinities of polyethers, by kinetic method and ligand exchange technique, 4:718
of deprotonated monomeric and dimeric ferulic and sinapic acids, 10:1111
of histidine and lysine and their selected di- and tripeptides, 7:1203
of serine and dipeptides of serine and glycine, 5:718
- Basicity, role in determining selectivity in ESI MS, 19:719
- Beam deflection for temporal encoding in time-of-flight mass spectrometry, 1:440
- Beam-induced dehalogenation of organic compounds, liquid secondary ionization mass spectrometry, secondary electron capture in, evidence for a mechanism that involves, 7:1109
- Beam-induced reaction between meta-nitrobenzyl alcohol and dipridocyanine dyes in liquid-secondary-ion mass spectrometry, 5:1017
- Beam-induced reactions, occurring in liquid secondary ion mass spectrometry/fast-atom bombardment by tandem mass spectrometry, 4:312
- Beam tandem mass spectrometer, new flowing afterglow-guided ion, applications to the thermochemistry of polyiodide ions, 8:688
- Behavior of oxygen, as CID target gas, 12:23
- Bench marking survey, on supporting chemical synthesis and structural elucidation in pharmaceutical industry, 16:599
- Benezenes, chlorinated, [m–1] ions of, and D₂O or ND₃, hydrogen/deuterium exchange between, 5:282
- Benzaldehydes, para-substituted, gas-phase base-catalyzed Claisen-Schmidt reactions of the acetone enolate anion with various, 7:82
- Benzamides, proton affinity of, and proton-bound complexes, dissociation of, 5:826
- Benzannulated perylene diimide ligands, perylene diimide and, binding to DNA, ESI-MS, 17:593
- Benzen structure, 1,3-hexadien-5-yne radical cation to, and to nonclassical ion structures, Ab initio calculations of, 7:731
- Benzene
alkaline earth metal ion reactions with, 12:250
loss from trityl cations, mechanistic study, 17:730
metal halide ion reactivity and binding energy with, 13:477
per(onio)-substituted, electrospray of, 9:15
toluene, ethylbenzene and xylenes, ambient air, analysis using APCI-MS/MS, 20:829
- Benzene radical cation, Ab initio calculations on, and carbon scrambling via isomerization to fulvene structure, 1,2-hydrogen shifts in, 6:513
- Benzenedicarboxylic acids, alkyl and aryl esters of, gas-phase reactions of O₂[–] with, 5:990
- Benzenedicarboxylic esters, molecular ions of, collision with oxygen, in quadrupole MS, 10:1248
- Benzidine in environmental samples, FT-ICR MS of, 10:1152
- Benzodiazepin-4-one,-6-one, and -4,6-dione, pyrrolo[1,2-a], mass spectrometry and molecular orbital calculations on, characterization and differentiation of heterocyclic isomers, 7:653
- Benzo[ghi]fluoranthene tetrols, PSD studies of, 12:376
- Benzoic acid matrix 4:2-(4-hydroxyphenylazo), matrix-assisted laser desorption ionization mass spectrometry with, 4:399
- Benzo[a]pyrene, conjugated, collision-activated dissociation of, 8:50
- Benzonitrile hydrogen interchange in the protonated species, protonation of, 1:225
- Benzophenones, para-substituted, chemical mass shifts of, correlation with Brown's σ_p^+ constants, 13:623
- Benzoxazole, DNA duplex with, ESI-MS of, 15:1593
- Benzoxazole ligands, metal-mediated DNA binding of, evaluation by ESI, 19:209
- Benzoylcegonine, in urine, HPLC and MS/MS in confirmation/quantitation of, 11:257
- Benzoylnitrene radical anion, new reagent for M-2H anion generation, 18:2014
- BEqQ hybrid mass spectrometer, determination of high-energy fragmentation of protonated peptides using, 4:470
- Beta-cyclodextrin inclusion complexes, ternary gas phase, ES-MS of, 14:449
- Beta-lactoglobulin, in methanol, ESI-MS of, 12:317
- Betabellin 15D, metal binding and proteolysis of, 10:969
- Beyer-Swinehart algorithm, modification for calculation of RRKM unimolecular reaction rate constants for biopolymers, 18:1063
- Bezimidazole, DNA duplex with, ESI-MS of, 15:1593
- Bi³⁺ ion, binding to transferrin, indirect detection with ESI-MS, 15:1658
- Biemann Medal
awarded to Julia Laskin, editorial, i (June) 2005, Gary J. Van Berkel, 17:887
2006, David E. Clemmer, 18:1161
2007, recipient Roman Zubarev, 19:751
- Bifidobacterium lactis, stress-related fingerprint identification from, by MALDI-TOF-MS, 15:1222
- Bile acid anions, tauroconjugated, fast-atom bombardment mass spectrometry and low energy collision-induced tandem mass spectrometry of, 6:588
- Billionfold data increase from mass spectrometry instrumentation, 8:1
- Bimetallic oxovanadium complexes, ESI-QTOF-MS/MS analysis, 19:1247
- Bimetallic silver-gold clusters, MALDI of, 15:1455
- Bimolecular elimination reactions, microsolvated gas-phase, deuterium KIEs, 18:1046
- Bimolecular hydrogen-deuterium exchange, behavior of protonated alkyl dipeptides in the gas phase, 6:466
- Bimolecular quadruplexes, DNA, transitions to higher order molecular structures, ESI-FTICR-MS, 18:1467
- Bimolecular reactions of diols, effects of functional group interactions on, 3:535
- Binding affinities, protein-mutant DNA complexes, laser spray MS, 17:611
- Binding of aldose reductase inhibitors, correlation of crystallographic and MS studies, 10:635
- BINOLate ligands, iron II and III gaseous complexes with, 19:121
- Bioaerosol detection, pyrolysis MS in, 10:502
- Bioaerosol mass spectrometry (BAMS)
efficient ion generation in, 19:315

- of ionization of biomarkers from micrometer particles, 15:900
TOF, sensitivity and mass range of, 16:1866
- Bioimaging, metabolite imaging using ME-SALDI-MS, 20:80
- Biological mass spectrometry
asymmetric charge partitioning in protein homodimers, 15:1408
charge competition and linear dynamic detection range in ESI-MS, 15:1416
ESI-FTICR-MS of GlcNAc-6-O-sulfotransferase, 15:1400
ESI mechanism features that affect protein complexes, 15:1424
ESI-TOF-MS of protein kinase G, cGMP, and AMPPNP, 15:1392
monomeric structures and dimer formation in α -synuclein, 15:1435
MS of, 12:125
positive and negative ion ESI-MS of drug-DNA complexes, 15:1382
- Biological materials, degradation of, correlation with CID of Gln-Gly cleavage, 12:337
- Biological molecules
gas-phase conformations of, 10:9
gaseous Fe(III)-heme cation binding to, 16:589
secondary ion mass spectrometry for resolution and imaging of, 17:1195
- Biological nucleophiles, model, reactions of polycyclic aromatic hydrocarbon radical cations with, 7:628
- Biological samples, solvent-free MALDI-MS of, via mini-ball mill approach, 16:542
- Biological soft tissues, molecular secondary ion mass spectrometry studies of, a model matrix for, doped gelatin films as, 4:419
- Biological tissues and fluids, 3-nitrosine in, quantification of, 11:578
- Biomarkers, human urine, modified nucleosides and ribosylated metabolites, ESI-FTICR MS and ESI-IT-MS, 19:1500
- Biomedical applications, of MALDI-TOF-MS, 13:1015
- Biomolecules
analysis using IR-MALDESI coupled to FT-ICR, 20:667
anhydrous, conformational landscape of, 20:1775
ESI-MS characterization of, 11:1
gold nanoparticles as assisted matrices for, MALDI-TOF-MS, 20:875
high-resolution accurate mass measurements of, using a new electrospray ionization ion cyclotron resonance mass spectrometer, 4:566
large
Fourier-transform ion-cyclotron resonance mass spectrometry of, 4:190
IR laser post-ionization MS of, 15:934
isotope depletion of, 9:149
MALDI of, matrix solutions for, 9:166
sputtering, shock wave model for, using massive cluster impacts, 3:311
top-down identification and characterization, MS, 19:1045
- Biophysics, MS in, 16:2064
- Biopolymer ions, generated by MALDI, photodissociation of, tandem TOF MS for, 15:1248
- Biopolymer sequencing, by triple quadrupole mass spectrometer in ESI nozzle-skimmer/precursor ion MS/MS mode, 12:846
- Biopolymers, RRKM unimolecular reaction rate constants for, BS algorithm modification for calculation of, 18:1063
- Bioreactive probes, transthyretin variant detection using, 11:172
- Biotin binding to streptavidin, electrospray ionization mass spectrometry of, 6:912
- Biotin-tags, biotinylated probe for selective enrichment of phosphopeptides, 19:741
- Biphenyls, polychlorinated, correlation between electron capture negative chemical ionization mass spectrometric fragmentation and calculated internal energies for, 5:44
- BIRD. *See* Blackbody infrared radiative dissociation
- α -Bis(4-hydroxybutyl)poly(dimethylsiloxane), GPC-MALDITOF-MS analysis of, 13:170
- Bis-intercalators, threading, binding to DNA, screening by ESI-MS/MS, 18:311
- Bis(imidazole-1-carboxylate) end-functionalized polymers, MALDI-MS study of, 14:117
- Bis(L-alaninate)ethylenediamine cobalt(III), antimony potassium tartrate association with, chiral recognition in, 11:804
- Bismuth cluster ion source, to improve biological TOF-secondary ion MS imaging, 16:1608
- Bisphenol A diglycidyl methacrylate, adsorbed and covalently bound, LDPI-MS of, 18:1097
- Bis-phosphate crown ethers, anionic salt bridge formation with, arginine molecular recognition by, 15:616
- Bisubstrate analogs of protein kinases, series of adenosylated peptides acting as, mass spectrometric characterization of, 5:443
- Blackbody infrared radiative dissociation
of heme-globin complexes, 8:519
of leucine enkephalin, 8:771
to measure deoxyribose nucleotide dimer anion dissociation energies, 10:1095
of nonspecific protein-carbohydrate complexes produced by nanoESI, 16:1583
of oligonucleotide anions, 9:1117
- Blood, human, measuring volatile organic compounds in, by using purge-and-trap gas chromatography/mass spectrometry, importance of enhanced mass resolution in, 3:831
- β_2 M. *See* $\beta_{20:2}$ -Microglobulin
- Bn (n greater than or equal to 3) ions in peptide spectra, structure and fragmentation of, 7:233
- Bombardment ionization method, model pulsed, study of time-dependent desorption using, improved detection limits for fast atom bombardment mass spectrometry, 3:637
- Bond cleavage reactions, competing, entropy of activation for, 16:2039
- Bond dissociation energy, heterolytic, in alkali chloride cluster ion dissociation, 13:1388
- Bond dissociation energy of germane, Fourier transform-ion cyclotron resonance study of gas-phase acidities of germane and methylgermane, 4:54
- Bond orders, differential overlap, modified neglect of, and ab initio and modified neglect of differential overlap energy partitioning, effect of protonation site on bond strengths in simple peptides, 5:704
- Bond-selective photodissociation of aliphatic disulfides, 6:872
- Bond strengths, gas-phase, of CO₂ and N₂O, with the halide ions, 4:58
- Bond strengths in simple peptides, effect of protonation site on, application of ab initio and modified neglect of differential overlap bond orders and modified neglect of differential overlap energy partitioning, 5:704
- Book Reviews. *See also* Conference reviews
Advances in Gas Phase Ion Chemistry, Vols. 1 and 2, 7:1178
Advances in Mass Spectrometry, Volume 12, Proceedings of the 12th International Mass Spectrometry Conference Held in Amsterdam, August 26–30, 5:53
Analysis of Carbohydrate by GLC and MS, 4:975
Analytical Advances for Hydrocarbon Research, 14:1327
Analytical Mass Spectrometry: Strategies for Environmental and Related Applications, 13:898
Analytical Methods in Forensic Chemistry, 4:362

- Applications of Inorganic Mass Spectrometry*, 14:1204
Applications of Mass Spectrometry to Organic Stereochemistry, 6:152
Applications of Plasma Source Mass Spectrometry, 4:428
Applications of Plasma Source Mass Spectrometry II, 6:610
Assigning Structures to Ions in Mass Spectrometry, 18:1145
A Beginner's Guide to Mass Spectral Interpretation, Introduction to Mass Spectrometry, 9:852
Chemical Ionization Mass Spectrometry, Second Edition, 4:286
Comprehensive Analytical Chemistry Volume 52, Protein Mass Spectrometry, R20:29 (August)
Computational Methods for Mass Spectrometry Proteomics, R19:3 (November)
Continuous-Flow Fast Atom Bombardment Mass Spectrometry, 3:867
Current Practice of Gas Chromatography-Mass Spectrometry, 12:1348
Electrospray Ionization Mass Spectrometry: Fundamentals, Instrumentation, and Applications, 1191
Encyclopedia of Mass Spectrometry. Volume 1: Theory and Ion Chemistry, 15:762
Encyclopedia of Spectroscopy and Spectrometry, 12:1064
Expanding Role of Mass Spectrometry in Biotechnology, 15:625
Experimental Mass Spectrometry, 6:277
First North American Fourier Transform Ion Cyclotron Resonance Mass Spectrometry Conference, 8:813
Focus on Analytical Spectrometry: A Compendium of Applied Spectroscopy Focal Point Articles (1994–1997), 10:783
Forensic Analysis on the Cutting Edge-New Methods for Trace Evidence Analysis, RI (March)
Fundamentals of Contemporary Mass Spectrometry, R18:1 (November)
Fundamentals of Gas-Phase Ion Chemistry, 4:832
Gas Chromatography and Mass Spectrometry: A Practical Guide, 8:294
GC/MS: A Practical User's Guide, 10:364
GC/MS: A practical User's Guide 2nd Edition, R19:1 (August)
GC/MS in Clinical Chemistry, 12:127
Global View of LC/MS, 10:78
Handbook of Mass Spectra of Environmental Contaminants, Second Edition, 5:598
HPLC for Pharmaceutical Scientists, 18:1344
HPLC of Acyl Lipids, 17:885
Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy, 8:671
Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectrometry, 16:1902
Identification of Microorganisms by Mass Spectrometry, 17:1037
Important Peak Index of the Registry of Mass Spectral Data, 4:82
Inductively Coupled Plasmas in Analytical Atomic Spectrometry, Second Edition, 4:756
Inductively Coupled Plasma-Mass Spectrometry: Practices and Techniques, 12:1226
Inductively Coupled Plasma Mass Spectrometry Handbook, 18:1345
Inorganic Mass Spectrometry: Fundamentals and Applications, 11:822
Inorganic Mass Spectrometry: Principles and Applications, R19:1 (November)
Instrumentation for Trace Organic Monitoring, 4:755
Instrumentation in Analytical Chemistry 1988–1991, 4:286
Integrated Strategies for Drug Discovery Using Mass Spectrometry, 17:108
Interpretation of Mass Spectra, Fourth Edition, 5:949
Interpreting Protein Mass Spectra: A Comprehensive Resource, 13:107
Introduction to Mass Spectrometry, An ACS Video Course, 7:298
Introduction to Mass Spectrometry: Instrumentation, Applications, and Strategies of Data Interpretation, Fourth Edition, R19:5 (July)
Introduction to Proteomics: Tools for the New Biology, 13:1149
Ion Mobility Spectrometry, 6:152
Isotope Tracers in Metabolic Research: Principles and Practice of Kinetic Analysis, 2nd Edition, 16:977
LC/MS: A Practical User's Guide, 17:1193
Liquid Chromatography-Mass Spectrometry, 6:222
Liquid Chromatography-Mass Spectrometry, 18:1135
Liquid Chromatography/Mass Spectrometry Applications in Agricultural, Pharmaceutical, and Environmental Chemistry, 4:831
Mass Spectral and GC Data of Drugs, Poisons, Pesticides, Pollutants and Their Metabolites, Volume 1: Methods and Tables; Volume 2: Mass Spectra, Third Revised and Enlarged Edition, R1 (January)
Mass Spectrometry, 4:82
Mass Spectrometry: A Textbook, 16:793
Mass Spectrometry (Analytical Chemistry by Open Learning), 4:831
Mass Spectrometry and Hyphenated Techniques in Neuropeptide Research, 14:81, 14:287
Mass Spectrometry Basics, 14:924
Mass Spectrometry for Biotechnology, 7:1179
Mass Spectrometry for the Characterization of Microorganisms, 6:1262
Mass Spectrometry for Chemists and Biochemists, 2nd Edition, 649
Mass Spectrometry, Clinical and Biomedical Applications, 6:453
Mass Spectrometry Desk Reference, Second Edition, R18:1 (September)
Mass Spectrometry in Biomolecular Sciences, 7:1273
Mass Spectrometry in Biophysics: Conformation and Dynamics of Biomolecules, 16:2064
Mass Spectrometry in Cancer Research, 7:1273
Mass Spectrometry in Drug Discovery, 13:1356
Mass Spectrometry in Drug Metabolism and Pharmacokinetics, R20:19 (July)
Mass Spectrometry in Medicinal Chemistry: Applications in Drug Discovery (Methods and Principles in Medicinal Chemistry), Volume 36, R19:1 (February)
Mass Spectrometry in the Biological Sciences, 7:692
Mass Spectrometry in the Biological Sciences, A Tutorial, NATO Advanced Science Institutes Series C, Mathematical and Physical Sciences 353, 3:867
Mass Spectrometry of Biological Materials, 12:125
Mass Spectrometry of Lipids, 5:124
Mass Spectrometry of Peptides, 4:519
Mass Spectrometry of Polymers, 13:1250
Mass Spectrometry of Protein Interactions, R19:1 (April)
Mass Spectrometry of Proteins and Peptides, 12:126
Mass Spectrometry of Soils, 8:200
Mass Spectrometry PittCon2 2001, 12:607
Mass Spectrometry PittCon 2002, 13:749
Mass Spectrometry PittConTM 2000, 11:472
Mass Spectrometry PittCon® 2007, 18:1145
Mass Spectrometry PITTCon'96, 7:500
Mass Spectrometry: Principles and Applications, 8:1193
MassFinder 3, 18:1137
Metabolome Analysis: An Introduction, 18:1R (August)
Methods and Mechanisms for Producing Ions from Large Molecules, 3:780
Methods in Enzymology, Volume 193, Mass Spectrometry, 3:779
New and Emerging Proteomic Techniques, 18:162
Plasma Source Mass Spectrometry, 4:428
Plasma Source Mass Spectrometry: The New Millennium, 14:171
Practical Aspects of Ion Trap Mass Spectrometry, 10:80
Practical Spectroscopy Series, Volume 32, Applied Electrospray Mass Spectrometry, 13:1013

- Preview of Personal Computer Literature Abstract Sources for Mass Spectrometry*, 7:299
- Principles of Mass Spectrometry Applied to Biomolecules*, 18:802
- Protein and Peptide Analysis by Mass Spectrometry*, 8:296
- Proteomics Today: Protein Assessment and Biomarkers Using Mass Spectrometry, 2D Electrophoresis, and Microarray Technology*, 17:111
- Quadrupole Ion Trap Mass Spectrometry, Second Edition*, 17:746
- Quantitative Proteomics by Mass Spectrometry*, R18:3 (September)
- Radioactive and Stable Isotope Tracers in Biomedicine*, 4:193
- Redox Proteomics: From Protein Modification to Cellular Dysfunction and Diseases*, 18:588
- Report of 2008 ASMS Fall Workshop: Protein-Protein and Protein-Ligand Interactions*, R20:1 (May)
- Report of the 24th Asilomar Conference on Mass Spectrometry*, R20:1 (January)
- A Review of Electronic Mass Spectral Databases*, R20:22 (July)
- Review of Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry, 4th Edition*, 18:803
- Review of the 19th Asilomar Conference on Mass Spectrometry: Bimolecular Interactions: Identification and Characterization of Protein Complexes*, 15:759
- Secondary Ion Mass Spectrometry SIMS VIII*, 4:755
- Spectral Techniques in Proteomics*, R18:3 (November)
- Techniques in Protein Chemistry*, 4:519
- Techniques in Protein Chemistry II*, 4:519
- Techniques in Protein Chemistry III*, 4:519
- Time-of-Flight Mass Spectrometry*, 7:123
- Time-of-Flight Mass Spectrometry and Its Applications*, 5:949
- Time-of-Flight Mass Spectrometry Instrumentation and Applications in Biological Research*, 9:1104
- Trace Quantitative Analysis by Mass Spectrometry*, R20:4 (January)
- Twelfth Sanibel Conference on Mass Spectrometry: Field-portable and Miniature Mass Spectrometry*, 11:468
- Understanding Mass Spectra: A Basic Approach*, 11:664
- Understanding Mass Spectra: A Basic Approach*, 16:792
- Using Mass Spectrometry for Drug Metabolism Studies*, 17:1188
- Vacuum Ultraviolet Photoionization and Photodissociation of Molecules and Clusters*, 4:974
- ZEKE Spectroscopy*, 10:456
- Borane salts, electrospray mass spectrometry of, electrospray needle as an electrochemical cell, 6:860
- Borate, gas-phase ion-molecule chemistry of, 5:169
- Boron derivatization, gas phase, in phosphopeptide fragmentation pattern manipulation, 16:1905
- Boronate esters, gas-phase ion-molecule chemistry of, 5:169
- Borrelia burgdorferi*, P13 membrane protein of, MS of, 13:295
- Boscalid, pesticide metabolism simulation using EC/LC/MS, 20:138
- Boundary activation, to effect non-resonant CISE, 13:650
- Boundary ejection, co-occurrence of resonance ejection and, ion traps, multiplexed multilinear MS, 17:56
- Bovine alpha-lactalbumin, characterization of, by electrospray ionization and liquid secondary ion mass spectrometry, 5:377
- Bovine pancreatic trypsin inhibitor, hydroiodic acid attachment to, 10:552
- Bovine serum albumin
ion trapping using large diameter frequency-adjusted linear quadrupole ion traps, 19:1943
three-dimensional structure of, chemical cross-linking and MS of, 15:1237
- Bowers, Michael T., personal retrospective, 16:983
- Bradbury-Nielsen gates, template-based method for production, 18:1901
- Bradykinin
CID of, in interface region of ESI, 12:772
fragmentation analysis of, by ²⁵²Cf-plasma desorption mass spectrometry, 2:379
- Bradykinin analogues, comparison of 193-nm photodissociation and TOF-TOFMS analysis, 17:721
- Bradykinin fragment 1–5 ions, in gas phase, IM-MS of, 16:893
- Bradykinin ions, gas phase, H/D exchange of, 14:85
- Brain
crustacean, neuropeptide and lipid mapping by 3D MSI, 20:1068
developing, secondary ion images of, 15:1116
HT_{2a} antagonist and metabolite in, LC-MS/MS of, 8:371
phosphatidylcholines in, MALDI-MS/MS of, 16:2052
phospholipids in, MALDI-TOF and MALDI-IM-TOF MS of, 16:133
protein expression in, MALDI-MS of, 16:1093
rat, peptide imaging in, MALDI-FTICR MS, 18:145
secondary ion images of, 11:191
in situ characterization of glycerophospholipids and sulfatides using MALDI-MS/MS, 18:17
- Brain-derived neurotrophic factor, ATP binding to, detected by MALDI-TOF and ESI-MS, 19:91
- Branched-chain fatty acids, novel method for, low-energy tandem mass spectrometry of the molecular ion derived from fatty acid methyl esters, 4:223
- Branching pattern and sequence analysis of underivatized oligosaccharides by combined MS/MS of singly and doubly charged molecular ions in negative ion ESI MS, 13:671
- Brassinosteroid analogs, FAB-MS of, 8:270
- Braun's lipoprotein and determination of its attachment sites to peptidoglycan by ²⁴²Cf-PD and MALDI time-of-flight mass spectrometry, characterization of, 6:892
- Breast, tumor and healthy tissue in, detecting estrogen DNA adducts in, nano-LC combined with nano-ES MS/MS in, 482
- Breast cancer, serine phosphorylation site in NTD of ER α , from MCF7 cells, 19:729
- Broad energy range focusing reflectrons, design of, method for, 7:1002
- Broad mass range in the quadrupole ion trap, selective ion isolation/rejection over, 2:11
- Bromide molecular ion, allyl, isomers of, kinetic and spectroscopic characterization of, 2:372
- Bromoacetic acid
negative ESI of, 11:809
TOFMS mass measurement of, 11:809
- Bromoacetonitrile, proton affinities of, using kinetic method with full entropy analysis, 11:601
- 4-Bromomethyl-6-methoxycoumarin, derivatization of 5-fluorouracil with, for LC/MS determination, 9:971
- Bromotrichloromethane and carbon microsomal metabolism, tandem mass spectrometry study of C-phenyl-N-tert-butyl nitrene spin adducts, in vitro rat liver and, 6:847
- Broths, fermentation, rapid analysis of antibiotic-containing mixtures from, by using liquid chromatography-electrospray ionization-mass spectrometry matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry, 7:1227
- Brown's constant, correlation of *para*-substituted acetophenone and benzophenone chemical mass shifts with, 13:623
- BS algorithm. *See* Beyer-Swinehart algorithm
- Buffer gas, pulsed introduction of, in quadrupole ion trap, 8:532
- Buffer gases, effect on fluorescence yield of trapped gas-phase ions, 17:855
- Buffer tolerance, in MALDI-MS, 9:805
- Building mass spectrometers and a philosophy of research,

- 119C₇F₁₄, molecular anions of, decomposition reactions of, cluster-assisted, 1:372
- Butene isomer mixtures, simultaneous analysis of, by process MS, 11:1079
- Butyrylcholinesterase, inhibition by OP and CB pesticides, MALDI-TOF-MS, 18:698
- C
- C²⁺60, addition of 1-butene to, charge separation and “ball and chain” propagation in, collision-induced dissociation evidence for, 7:261
- C8- and N²-deoxyguanosine adducts of 2-acetylaminofluorene, isomeric, by fast-atom bombardment and tandem mass spectrometry, 5:58
- C₃₇-C₄₀ alkenones, unsaturated, double bond location in, GC-MS in derivatization for, 15:1161
- C-C bond cleavage, in collision-activated dissociation of polyatomic ions at low energy, fundamentals of tandem mass spectrometry, 1:6
- C-C bond formation and H-transfer between partners, and channeling of energy into dissociation, photoionization study of the ion-neutral complexes [CH₃CH⁺3, CH₂CH₃] and [CH₃CH₂CH⁺CH₃, CH₃] in the gas phase, 7:73
- C₆₀-fullerene, gas-phase alkylation of, by ion-molecule reaction under chemical ionization, 4:894
- ¹³C-Oleate transport, in adipocytes, multi imaging MS of, 15:1572
- C-terminal amino acid residue, loss of, rearrangement involving, metastable decomposition of peptide [M + H]⁺ ions via, 1:249
- C-terminal derivatization, in de novo sequencing of proteolytic peptides, 11:673
- C₁₀ to C₁₆₀ polycyclic aromatic hydrocarbons, flame-generated, by atmospheric-pressure chemical ionization mass spectrometry with liquid introduction via heated nebulizer interface, 7:276
- CAD. *See* Collision-activated dissociation; Collisionally activated decomposition; Collisionally activated dissociation; Collisionally activated dissociation entries
- Cadmium, isotopically enriched, isotope dilution ICP-MS applied to, 16:708
- Cadmium thiolate, conjugates, characterization by ESI and triple quadrupole MS MS, 17:1442
- Caffeic acid oxidation, products of, analysis by on-line EC-ESI-MS, 15:1228
- Caged aza-crown ether ligands, alkali metal binding selectivities of, MESI-MS of, 11:884
- Calcitonins, folding analysis of, by ESI-MS with H/D exchange, 15:1556
- Calcium
- calmodulin interaction with, H/D exchange in, 10:711
 - small conformational changes induced by, in cardiac troponin C, 10:703
- Calcium affinity, of peptides, 11:770
- Calcium binding, to EF-hand motifs probed by ESI-MS, 12:1296
- Calcium binding sites, in peptides, MS/MS determination of, 9:1020
- Calcium-induced noncovalently linked tetramers of MRP8 and MRP14 are confirmed by ESI-MS analysis, 11:780
- Calcium-induced noncovalently linked tetramers of MRP8 and MRP14 detected by ultraviolet MALDI-MS, 10:1124
- Calcote mechanism for soot formation, initial reaction of, 7:559
- Calculated internal energies for polychlorinated biphenyls, correlation between electron capture negative chemical ionization mass spectrometric fragmentation, 5:44
- Calcyclin, isolated from granulocytes, nonspecific cleavage of, 12:1180
- CALDI-MS. *See* Charge assisted laser desorption/ionization mass spectrometry
- Calibrant ions, delivery for MS, 18:991
- Calibration
- in liquid chromatography particle beam mass spectrometry, 3:662
 - of mass ranges up to m/z 10,000 in electrospray mass spectrometers, 10:273
 - mass scale, factors in, TOF-SIMS, 17:514
 - point for electron ionization MS/MS spectra measured with multiquadrupole mass spectrometers, 5:576
 - stepwise-external, FT-ICR MS, sub part-per-million mass accuracy, 17:1681
- Calibration experiments, of trilinolenoylglycerol, chemometric approaches by LC-ion trap MA, 16:388, 16:801 (erratum)
- Calibration standards, for ESI-MS, tris (perfluoroalkylethyl)silyl alkyl amines as, 12:1050
- Californium-252 plasma desorption, high resolution and tandem Fourier-transform mass spectrometry with, 1:427
- Calixarene, amino acid non-covalent complexes with, formed by MALDI-MS, 13:964
- Calmodulin
- calcium-induced association of, with melittin, 8:809
 - calcium interaction with, ESI-MS and H/D exchange of, 10:711
 - divalent cation binding analyzed by ESI-MS, 20:1159
 - methionine oxidation in, quantification of, 16:1470
 - peptide interactions with, investigated by MALDI-MS, 20:576
 - α-synuclein and, metal interactions monitored by SNAPP-MS, 19:1668
- Calmodulin-melittin complex interface, hydroxyl radical probe of, by ESI MS, 16:225
- Calutron, history of, 8:943
- CaM, AC8 and, isotope-labeled cross-linkers and FTIR MS analysis of, 17:1100
- Camera-based detection system, acquisition and quantification of ion images with, 1:37
- Camphene, α-pinene and, bimolecular and unimolecular contributions to SCI of isomers, 18:2026
- Cancer
- human urine biomarkers, modified nucleosides and ribosylated metabolites, ESI-FTICR MS and ESI-IT-MS, 19:1500
 - N-linked oligosaccharides with uncommon glycoform, MALDI MS, 19:671
- Cancer research, MS in, 14:79
- Candida rugosa*, lipase, immobilized on PDMS membrane, use in MIMS, 18:973
- Capacitive coupling gated trapping, in MALDI-FTICR, 8:195
- Capacitively coupled open cell, ion cyclotron, high-resolution ion isolation with, 6:533
- Capillaries
- packed, LC-RAM-MS of, 13:82
 - viscous gas flow through, ion transport by, 5:873
- Capillary dissociation, of cyclodextrin complexes, 8:244
- Capillary electrochromatography-microelectrospray MS, nucleoside adduct separation and identification by, 9:823
- Capillary electrophoresis
- enhanced sample loading by, 8:15
 - Fourier transform ion cyclotron resonance mass spectrometry with sustained off-resonance irradiation for the characterization of protein and peptide mixtures, 5:894
 - of local anesthetic-cyclodextrin non-covalent complexes, 13:928
 - of peptides, 8:495
 - protein modification by, 8
 - sheathless ESI-MS interface with, 8:561
- Capillary electrophoresis-atmospheric pressure chemical ioniza-

- tion-mass spectrometry, setup and parameters using orthogonal interface, 20:1311
- Capillary electrophoresis/electrospray ionization-mass spectrometry, analysis of peptides and protein mixtures by, 9:1081
- Capillary electrophoresis-electrospray ionization-time-of-flight mass spectrometry, of single intact cell hemoglobin, 10:184
- Capillary electrophoresis-electrospray mass spectra of herbicides paraquat and diquat, 7:981
- Capillary electrophoresis-mass spectrometry
detection of noncovalent FKBP-FK506 and FKBP-rapamycin complexes by, 6:85
electroosmotic flow in, electrospray needle voltage and, 10:261
- Capillary electrophoresis/mass spectrometry, with ion trap/reflection TOFMS detector, applications of myelin basic protein, 9:701
- Capillary electrophoresis-nanoelectrospray mass spectrometry, electrochemical properties of, 10:309
- Capillary electrophoresis tandem mass spectrometry (CE-MS/MS)
detection of noncovalent FKBP-FK506 and FKBP-rapamycin complexes by, 6:85
in structural analysis of chromophore-labeled disaccharides, 11:292
- Capillary electrospray source, heated, charge-dependent/independent ion energy distribution from, selective generation of, 5:221
- Capillary gas chromatography/ion trap mass spectrometry, quantitative methods, combined, using labeled or unlabeled internal standards, 1:265
- Capillary interface, in leveling response factors in ESI, 16:772
- Capillary isoelectric focusing-mass spectrometry, on-line desalting in, 8:743
- Capillary LC-TOFMS, proteome analyses with, 14:980
- Capillary liquid chromatography, nanoscale packed (75 m id), application of, to the analysis of macrolide antibiotics, 3:563
- Capillary liquid chromatography tandem mass spectrometry, on IT/reTOFMS using sequence tag database search approach for peptide sequencing and protein identification, 11:127
- Capillary-scale particle-beam LC/MS interface, can electron ionization sustain the competition?, 9:993
- Capillary separation-tandem mass spectrometry, for protein digest analysis, 8:1237
- Capillary tubes, flared inlet, incorporation in FT-ICR, 17:772
- Capillary zone electrophoresis, continuous-flow fast atom bombardment mass spectrometry, an integral probe for, 3:198
- Capillary zone electrophoresis-continuous flow-fast atom bombardment mass spectrometry, identification of deoxynucleoside-polyaromatic hydrocarbon adducts by, 3:757
- Capillary zone electrophoresis/electrospray ionization, parameters for the mass spectrometry and tandem mass spectrometry analysis of peptides, 3:289
- Capillary zone electrophoresis/electrospray ionization mass spectrometry, application of, to the analysis of macrolide antibiotics, 3:563
- Capillary zone electrophoresis/electrospray ionization parameters, for the mass spectrometry and tandem mass spectrometry analysis of peptides, 3:289
- Capillary zone electrophoresis-mass spectrometry, sheathless, applied to peptide analysis, 10:1271
- Carbamate pesticides
N-methyl, ion formation of, in thermospray mass spectrometry, the effects of additives to the liquid chromatographic eluent and of the vaporizer temperature, 5:913
thermospray interfaces with, comparative study of different, influence of the ion source geometry, 6:656
- Carbamates, thermally-labile, in supersonic molecular beams, fast, very fast and ultra-fast gas chromatography-mass spectrometry of, 7:737
- Carbanions
gaseous, hydrogen/chlorine exchange reactions of, 16:2045
nucleophilic substitution reactions of halonitrobenzenes and dinitrobenzene with, gas phase, 18:1351
on the stabilization of, by adjacent phenyl, cyano, methoxycarbonyl, and nitro groups in the gas phase, 5:159
- Carbohydrate isomers, resolution by ESI-APIMS-TOF-MS, 18:1163
- Carbohydrates
fragmentation of negative ions from
high mannose *N*-linked glycans in, 16:631
hybrid and complex *N*-linked glycans in, 16:645
nitrate and other anionic adducts in, 16:622
glycoproteins and, distinguishing phosphorylation and sulfation, ion-pairing and MS, 17:1282
ILMs for peptides, peptides and, analysis by MALDI MS, 20:1790
ionization and fragmentation in MS, labeling group and, 16:683
LC-MS quantitative analysis of, 16:1805
N-linked derivatized, fragmentation of, 11:900
from ovalbumin
composition of, 11:564
postsources decay fragmentation of, 11:572
phosphorylated, isomeric differences of, probing with ion/molecule reactions and FTICR-MS, 14:323
phytochemicals and, profiling in tulip tissues and bulbs, PESI MS, 20:2304
plant single-cell extracts, pressure probe and UV-MALDI-TOF MS, 19:1842
- Carbon
Meisenheimer complexes bonded at, 15:998
polymeric electrodes filled with, as ion optical elements for TOF-MS, 13:1170
surface-induced dissociation and chemical reactions on stainless steel, diamonds and, 20:927
- Carbon-13 labeling, for improved organic material profiling, secondary ion MS, 17:1142
- Carbon and nitrogen isotope ratios, in cells, 15:478
- Carbon-carbon double bonds, location of, in H_6-C_{23} linear alkenes from carbon addition reactions in a quadrupole ion trap equipped with a pulsed sample-inlet system, 2:305
- Carbon dioxide
electronic sputtering produced by fission fragments, condensed CO and, 17:1120
isotope ratios, in expired air from mice and men following the administration of ^{13}C -methyl methionine, the quadrupole-based mass spectrometric evaluation of, 3:742
- Carbon ions, energetic, silver surface bombardment by, 16:1677
- Carbon isotope ratio determination in amino acids, proteins, and oligosaccharides by ICP-MS, 14:295
- Carbon isotope variability in albumins, direct analysis of, by liquid flow-injection isotope ratio mass spectrometry, 7:605
- Carbon microsomal metabolism, tandem mass spectrometry study of *C*-phenyl-*N*-tert-butyl nitron spin adducts, in vitro rat liver and, 6:847
- Carbon monoxide
electronic sputtering produced by fission fragments, condensed CO_2 and, 17:1120

- elimination in condensation reactions in protonated dipeptides, 15:1025
- loss of ammonia and, protonated AGG, MS/MS, 18:1291
- Carbon nanotubes
- as adsorbent of solid phase extraction, in MALDI MS, 16:263
 - 2,5-dihydroxybenzoyl hydrazine derivative, use in trace peptide analysis, MALDI, 17:1023
 - immobilized, for MALDI-TOF MS, 16:333
 - oxidized, for MALDI-TOF MS analysis, 16:883
 - oxidized carbon, monitoring enzyme reactions and screening inhibitors, MALDI-TOF-MS with, 17:1616
 - as peptide and protein affinity probes, in MALDI MS, 15:1629
- Carbon scrambling via isomerization to fulvene structure 1,2-hydrogen shifts in, benzene radical cation, Ab initio calculations on, 6:513
- Carbon stable isotope analysis, organic matter and nitrocellulose, *Sphagnum* and *Polytrichum* mosses, 18:1453
- Carbon thin films, fluorinated amorphous, elemental distribution in, FIB-SIMS of, 16:126
- Carbonyl anions, metal, insertion reactions of, with methyl formate in the gas phase as revealed by ¹³C- and D-labeling, 5:525
- Carbonyl compounds
- aromatic, remote fragmentations of, via internal reactions in intermediary ion-neutral complexes, 3:417
 - O⁻ chemical ionization of, 2:292
 - water-solvated ammonia radical cation reactions with, 15:966
- Carbonylation, specific protein sites for, human serum albumin, 17:1172
- Carboxonium, gaseous, carbosulfonium transacetalization with, 12:14
- Carboxylate anions, low-energy collision-activated dissociation from structurally similar prostaglandins, E₂, F_{2α}, D₂, and DHKF_{2α}, 1:389
- Carboxylates, nonylphenol, in water and sludge, LC-ES-MS/MS of, 14:516
- Carboxylic acids
- interaction of cesium cations with, 20:1912
 - isoquinolines gas phase reaction to, after ESI and CID, 19:151
- Carboxylic esters, long-chain, with CH₃OBOCH₃⁺ in a small Fourier-transform ion cyclotron resonance mass spectrometer with, 7:1138
- Carcinogen-nucleoside adducts, identification of, fast atom bombardment mass spectral methods for, 3:360
- Carcinogen-peptide adducts, nano-electrospray MS/MS of, 9:202
- Carcinogenic potential, of environmental contaminants, estimation of, by ion-molecule reactions and tandem mass spectrometry, 1:110
- Cardiac glycosides, fast atom bombardment of, ions formed by, high resolution mass spectrometry and tandem mass spectrometry analysis of, mechanistic considerations of the protonation and fragmentation of highly functionalized molecules in fast atom bombardment, 1:455
- Cardiolipin
- characterization as sodiated ions, positive ion ESI with q-ion trap MS, 17:1146
 - E. coli*, characterization by ESI with multiple stage QIT MS, 17:420
 - MALDI-MS analysis from rat organ sections, 18:567
 - tandem quadrupole ion trap MS of, 16:491
- Carrageenans, sulfated oligosaccharides, structural analysis by positive mode nano-ESI-FTICR-MS and MS/MS by SORICID, 17:96
- Carrier process, enhanced ion abundances from, investigation of, in high-performance liquid chromatography particle beam mass spectrometry, 1:92
- Cassette dosing practices, in pharmaceutical industry, survey results on, 15:1374
- Catalysts, V-P-O, ion association by time-of-flight mass spectrometry, 1:129
- Catechins, green tea, isomeric differentiation using gas-phase H/D exchange reactions, 18:1749
- Cation, electrospray-generated, ion/ion reactions in, 13:614
- Cation adduction, mass and charge assignment for electrospray ions by, 4:828
- Cation affinities, gas-phase studies of valinomycin-alkali metal cation complexes, 5:632
- Cation attachment, in MALDI-MS of polymers, 8:132
- Cation exchange μ -HPLC-ESI-MS/MS, on-line strong, 14:719
- Cation-(π) effects in the complexation of NaI and KI with Phe, Tyr, and Trp in the gas phase, 11:1037
- Cationic matrix additives, class for enhancing the sensitivity in negative ion fast-atom bombardment mass spectrometry of polyanionic analytes, N-alkylnicotinium halides, 5:935
- Cationization, gas-phase, and protonation of neutrals generated by matrix-assisted laser desorption, 4:393
- Cationized heterodimers, of amino acids, decompositions of, in relation to charge location in peptide ions, 4:461
- Cations
- aminocarbene radical, rearrangement of, electron ionization induced fragmentation of macrocyclic amines C_nH_{2n}-1NH₂, 1:238
 - binary mixtures of, multiple charging of polypropylene glycol by, in electrospray, 16:152
 - distonic radical ⁺CH₂-O-CH₂, ionized methylene transfer from heterocyclic compounds, pentaquadrupole mass spectrometric study, 6:554
 - divalent, calmodulin binding of, analyzed by ESI-MS, 20:1159
 - in the gas phase, 1,2-eliminations, 19:1491
 - intramolecular interactions in, 8:1111
 - isomeric, C₃H₆Br⁺, gas phase bimolecular chemistry of, 6:11
 - isomerization of, 8:594, 8:599
 - radical, methyl propionate, 7:482
 - rhodamine 575 fluorescence and photodissociation of, quadrupole ion trap, 20:96
- Cavitands, mono- and diphosphonate bridged, alcohol complexation properties, ESI-FTICR MS study, 17:213
- CD4/lc-copurifying proteins, identification with PAGE and MS, 15:558
- CE. *See* Capillary electrophoresis
- CE-MS/MS. *See* Capillary electrophoresis tandem mass spectrometry
- CEC- μ ESI/MS. *See* Capillary electrochromatography-microelectrospray MS
- Cell culture media, pyrolysis MS of, 10:502
- Cells
- carbon and nitrogen stable isotope ratios in, 15:478
 - laser microdissected, MALD-MS analysis of, 13:1292
- Cellulomonas fimi*, binding of xylobiose inhibitors to β -1,4 exoglucanase from, MS/MS, 18:64
- Central ring electrode, for trapping and excitation/detection, in FTICR-MS, 12:30
- Centrin-2, metal-binding properties determined by μ ESI-MS and UV spectroscopy, 17:1158
- CEPH family 1362 STR database, as online resource for characterization of PCR products using ESI-MS, 13:87
- Ceramide ions, high-energy collision-induced dissociation of, from permethylated glycosphingolipids, 5:558
- Ceramides
- characterization CID-MS/MS with negative ion ESI, 13:558
 - lithiated, low energy CAD MS/MS of, with ESI, 13:680
 - structure determination of, by collisional activation of [M + Li]⁺ ions, 3:260

- Cerebellum, gangliosides from, automated nanoelectrospray MS/MS of, 15:1649
- Cesium cations, interaction with carboxylic acids, 20:1912
- $C_7F_7^+$ ions, gas phase, structures of, 4:813
- CFCs substitutes recommended by EPA, ion chemistry in air plasma and reactions with atmospheric ions, 16:1081
- C_2H_6 , C_3H_8 , $HC(CH_3)_3$, and $c-C_3H_6$, activation by gas-phase Ru^+ , 10:821
- CH_3^- anion, photoelectron spectrum of, 10:810
- $C_6H_5^+$ and $C_6H_6^+$ cations, structural probe for, charge inversion as, 10:878
- $C_4H_4^+$ isomers
 case study in the use of the MS3 capabilities of a pentaquadrupole mass spectrometer, 3:518
 ion-molecule reactions and collision-activated dissociation of, case study in the use of the MS3 capabilities of a pentaquadrupole mass spectrometer, 3:518
- $C_2H_4/N_2O/Ar$ flame, study of, triple quadrupole mass spectrometry as applied to flame diagnostics, 4:152
- $CH_2-S-CH-CN$, sulfur-containing distonic radical anion, formation and characterization of, in gas phase, 6:71
- CH_2^+ transfer to pyridine nucleophiles, means of producing alpha-distonic ions, 4:117
- Characteristics of high energy collision-induced dissociation tandem mass spectra of polycyclic aromatic hydrocarbon diolepoxide adducted peptides, 5:649
- Characterization and differentiation of heterocyclic isomers mass spectrometry and molecular orbital calculations on pyrrolo[1,2-a][1.4]benzodiazepin-4-one,-6-one, and -4,6-dione, 7:653
 tandem mass spectrometry and molecular orbital calculations on 3-methylisooxazolo- and 2-methylloxazolopyridines, 6:962
- Characterization and relative ionization efficiencies, of end-functionalized polystyrenes by MALDI-MS, 14:648
- Characterization of a cross-linked DNA-endonuclease VIII repair complex by ESI-MS, 11:503
- Characterization of a glow discharge ion source for mass spectrometric analysis of organic compounds, 5:305
- Characterization of a high-pressure quadrupole collision cell for low-energy collision-induced dissociation, 5:1042
- Characterization of a single crystal cubic Prussian blue $Co_8(tacn)_8(CN)_{12}$ cluster by ion trap and FT-ICR MS with microelectrospray ionization, 10:352
- Characterization of a specific cysteic acid residue in oxidized human apoprotein B-100, tandem mass spectrometric, 6:242
- Characterization of allosteric inhibition of a protein-protein interaction by MS, 14:8
- Characterization of amino acid side chain losses in electron capture dissociation, 13:241
- Characterization of an improved thermal ionization cavity source for MS, 10:1008
- Characterization of anthraquinone-2-carbonyl chloride as an alcohol derivatization reagent for negative ion chemical ionization mass spectrometry, 6:148
- Characterization of bacterial lipooligosaccharides by delayed extraction matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, 8:645
- Characterization of beam-induced reactions occurring in liquid secondary ion mass spectrometry/fast-atom bombardment by tandem mass spectrometry, 4:312
- Characterization of bis(alkylamine)mercury cations from mercury nitrate surfaces by using an ion trap second ion mass spectrometry, 9:35
- Characterization of Braun's lipoprotein and determination of its attachment sites to peptidoglycan by ^{242}Cf -PD and MALDI time-of-flight mass spectrometry, 6:892
- Characterization of ceramides by low energy CID-MS/MS with negative ion ESI, 13:558
- Characterization of diacylglycerolphosphocholine molecular species by FAB-CAD-MS/MS, general method not sensitive to the nature of the fatty acyl groups, 3:71
- Characterization of disulfide linkages and disulfide bond scrambling in recombinant human macrophage colony stimulating factor by fast-atom bombardment mass spectrometry of enzymatic digests, 6:638
- Characterization of erythromycin analogs by collisional activated dissociation and infrared multiphoton dissociation in a quadrupole ion trap, 13:630
- Characterization of femtomole levels of proteins in solution using rapid proteolysis and nanoelectrospray ionization mass spectrometry, 8:483
- Characterization of flame-generated C_{10} to C_{160} polycyclic aromatic hydrocarbons by atmospheric-pressure chemical ionization mass spectrometry with liquid introduction via heated nebulizer interface, 7:276
- Characterization of labeled oligonucleotides using enzymatic digestion and tandem mass spectrometry, 9:660
- Characterization of large heterogeneous proteins by electrospray ionization-mass spectrometry, 7:1219
- Characterization of lipooligosaccharides from *Haemophilus ducreyi* containing poly lactosamine repeats, 13:724
- Characterization of long-chain carboxylic esters with $CH_3OBOCH_3^+$ in a small Fourier-transform ion cyclotron resonance mass spectrometer, 7:1138
- Characterization of low molecular weight alkoxylated polymers using long column SFC-MS and image analysis-based quantitation approach, 13:1195
- Characterization of mouse switch variant antibodies by matrix-assisted laser desorption ionization mass spectrometry and electrospray ionization mass spectrometry, 7:707
- Characterization of MSAD, implications for ESI-MS characterization of iomolecules, 11:1
- Characterization of noncovalent complexes formed between minor groove-binding molecules and duplex DNA by electrospray ionization-mass spectrometry, 6:1154
- Characterization of phosphatidylinositol, phosphatidylinositol 4-phosphate, and phosphatidylinositol 4,5-bisphosphate by ESI-MS/MS, 11:986
- Characterization of poly(ethylene glycol) esters using low energy CID in ESI-MS, 13:888
- Characterization of polysiloxanes with different functional groups by TOFMS, 9:292
- Characterization of protein N-glycosylation by reversed-phase microbore liquid chromatography/electrospray mass spectrometry, complementary mobile phases, and sequential exoglycosidase digestion, 5:350
- Characterization of the $C_3H_6O^+$ ion from 2-methoxyethanol, mixture analysis by dissociation and neutralization-reionization, 6:1030
- Characterization of the dissociation behavior of gas-phase protonated and methylated lactones, 3:47
- Characterization of the elusive disulfide bridge forming human Hb Ta-Li $\beta 83$ (EF7)Gly \rightarrow Cys variant by ES MS/MS, 13:187
- Characterization of the opioid peptide-containing protein and of bovine alpha-lactalbumin by electrospray ionization and liquid secondary ion mass spectrometry, 5:377
- Characterization of the P13 membrane protein of *Borrelia burgdorferi* by MS, 13:295
- Characterization of the primary thermal degradation processes

- of peptides using the mass spectrometric technique
- K^+ IDS, K^+ ionization of desorbed species, 5:564
- Characterization of the product ions from the CID of argentinated peptides, 12:163
- Characterization of the reactive and dissociative behavior of transition metal oxide cluster ions in the gas phase, 2:212
- Characterization of VX on concrete using IT-SIMS, 11:69
- Charge and mass assignment for electrospray ions by cation adduction, 4:828
- Charge assisted laser desorption/ionization mass spectrometry, of droplets, 19:833
- Charge compensation for imaging large insulating samples by using secondary ion tandem mass spectrometry, 5:37
- Charge competition, in ESI-MS, 15:1416
- Charge-dependent/independent ion energy distribution from a heated capillary electrospray source, selective generation of, 5:221
- Charge derivatization
 - in fragmentation of multiply protonated peptides, 13:47
 - methionine-containing peptide analysis by, 16:1131
- Charge-derivatized peptides, fragmentation of, 12:399
- Charge detection mass spectrometry, of megadalton-DNA electrospray ions, 9:305
- Charge discrimination with electrospray ionization-mass spectrometry, applying to protein analyses, 6:1098
- Charge-driven fragmentation processes, in diacyl glycerophosphatidic acids upon low-energy collisional activation, a mechanistic proposal, 11:797
- Charge effects for differentiation of oligodeoxynucleotide isomers containing 8-oxo-dG residues, 13:195
- Charge exchange
 - in APCI-MS, in analyzing polycyclic aromatic compounds, 15:301
 - doubly and singly charged tungsten hex acarbonyl ions generated by, internal energy distributions deposited in, 1:16
- Charge exchange ionization
 - in collision cells as a method to detect the presence of long lived excited electron states of polyatomic ions, 12:1120
 - of long-lived excited electronic states of dichloroethylene cation isomers, 15:1266
 - of PAHs, comparison of dopants for, RP-LC-APPI-MS, 20:73
- Charge injection device, two-dimensional, direct generation of ion beam images with, 6:140
- Charge inversion as a structural probe for $C_6H_5^+$ and $C_6H_6^+$ cations, 10:878
- Charge inversion tandem mass spectrometry, isomer differentiation by, an investigation into the structure of the ionic products from an Sn(ANRORC) reaction, 3:467
- Charge localization of and charge state, the effect on the collision-induced dissociation of peptide ions, 5:966
- Charge location in peptide ions, decompositions of cationized heterodimers of amino acids in relation to, 4:461
- Charge partitioning
 - asymmetric, in protein homodimers, 15:1408
 - dipeptide dications, between *c* and *z* fragments after ECD, 20:1881
- Charge/radical site, initiation vs Coulombic repulsion for cleavage of multiply charged ions, charged solvation in poly(alkene glycol) ions, 12:565
- Charge reduction, peptide ions, soft landed surface effects on desorption kinetics and, 20:901
- Charge reduction electrospray ionization mass spectrometry, Critical Insight, 19:629
- Charge-remote and charge-driven fragmentation processes in diacyl glycerophosphoethanolamine upon low-energy collisional activation, a mechanistic proposal, 11:892
- Charge remote fragmentation
 - to determine oxofatty acid structure, 9:216
 - energy dependence of, 9:840
 - during FAB-CAD-B/E linked-scan mass spectrometry of aminoethyl-triphenyl phosphonium derivatives of fatty acids, 3:769
 - of fatty acids cationized with alkaline earth metal ions, 1:320
 - in a hybrid (BEqQ) mass spectrometer to determine isotopic purity in selectively polydeuterated surfactants, 1:85
 - of peptides, MALDI of, 8:501
 - of peptides flowing attachment of a fixed positive charge, a matrix-assisted laser desorption/ionization postsource decay study, 8:501
 - some examples of, fast atom bombardment and collision-induced dissociation of prostaglandins and thromboxanes, 1:325
 - structural characterization by, fast atom bombardment mass spectrometry of mycobacterial glycopeptidolipid antigens, 3:656
 - under MALDI-PSD, 10:217
 - on TOF-TOF MS, lithiated fatty acids, 18:1967
 - and the two-step elimination of alkanols from atom bombardment-desorbed $(M + H)^+$, $(M + Cat)^+$, and $(M-H)^-$ ions of aromatic beta-hydroxyoximes, 4:819
- Charge separation
 - AMUSE ion source for MS, 20:1684
 - and “ball-and-chain” propagation in the addition of 1-butene to $C_2^+/60$, collision-induced dissociation evidence for, 7:261
 - effect of peptide chain length, 20:1890
 - reactions in tandem mass spectrometry of doubly protonated angiotensin II formed by electrospray ionization, experimental considerations and structural implications, 7:30
- Charge site in “charge-remote” fragmentations, implications of, 3:776
- Charge-site-remote fragmentation reactions, ionization mode and, 12:571
- Charge sites, primary and secondary locations, in angiotensin II $(M + 2H)^{2+}$ ions formed by ESI, 11:40
- Charge-space effects in plasma mass spectrometry, time-resolved measurements of individual ion cloud signals to investigate, 7:362
- Charge state
 - distribution shifting of protein ions observed in matrix-assisted laser desorption ionization mass spectrometry, 6:1183
 - and the localization of charge, the effect on the collision-induced dissociation of peptide ions, 5:966
 - maximum, of arginine-containing peptide ions formed by electrospray ionization, 7:972
 - of protein ions produced by ESI-MS, solvent effects on, 11:976
 - from resolved isotopic peaks for multiply charged ions, automated assignment of, 6:52
- Charge state assignment, of native ES spectra of protein complexes, 20:435
- Charge state distribution shifts, of protein, in nanoESI triple quadrupole MS, 16:409
- Charge state distributions
 - in ESI-TOF-MS, IgG1 κ mAb characterization using, 20:1030
 - macromolecular structure, ESI MS, 19:1239
 - proteins, in ESI MS, 20:1617
- Charge-state reduction with improved signal intensity of oligonucleotides in electrospray ionization mass spectrometry, 7:697
- Charge stripping, doubly and singly charged tungsten hex acarbonyl ions generated by, internal energy distributions deposited in, 1:16

- Charge tagging, in fragmentation pathway studies of oligonucleotides in MALDI-MS, 13:1407
- Charge-to-mass ratio scale, 5:120
- Charge transfer interactions, in stability of lithium atom complexes with formaldehyde and formalimine, 12:1229
- Charged derivatives for high energy collision-induced dissociation tandem mass spectrometry, comparison of, 6:428
- Charged droplets, ion formation from, roles of geometry, energy, and time, 4:524
- Charged phenyl radicals, gas-phase reactions of, 13:192
- Charged state distribution, proteins, effect of stabilization on, ESI MS, 18:1605
- Charged states, observed, of diquatery ammonium salts, and ion pairing, in electrospray ionization mass spectrometry, effects of solvent and counterion on, 7:1050
- CHARM, in ion mobility spectroscopy of charge-solvated vs salt bridge structures of protonated trimers, 16:1009
- $C_3H_6Br^+$ isomeric cations, gas phase bimolecular chemistry of, 6:11
- CHCA. *See* α -Cyano-18:4-hydroxycinnamic acid
- CH_3CH_3 , formation from $C_3H_6O^+$ isomers, 13:1235
- $CH_2CH_2CH^+OH$, dicationic ion, keto ion $CH_3CH_2CH_2O^+$, enol ion $CH_3CH=CHOH^+$, and related $C_3H_6O^+$ radical cations, stabilities and isomerization proclivities studied by dissociation and neutralization-reionization, 7:573
- $CH_3CH_2NH=CH_2^+$, ethene elimination from, 9:138
- $[CH_2CHOH^+, CH_3CHO]$ solvated enol radical cation, 11:705
- Chelation
effect on ESI-MS of diketone/metal ion complexes, 10:402
metal, of dinucleotide analogs in the gas phase by fast-atom bombardment mass spectrometry, 7:42
- ChemApplex program, for peptide mass fingerprinting, scoring methods in, 13:22
- Chemical analysis, direct, of UV laser ablation products of organic polymers by using selective ion monitoring mode in gas chromatography/mass spectrometry, 5:106
- Chemical analysis of UV laser ablation products of organic polymers by using, 5:106
- Chemical behavior, electrospray ionization-mass spectrometry as tool to predict, 6:207
- Chemical cross-linking, of bovine serum albumin three-dimensional structure, 15:1237
- Chemical electron-transfer reactions, gas-phase molecular dications formed from neutral organics in solution via, by using electrospray ionization mass spectrometry, 5:689
- Chemical equilibria, shrinking droplets and, ESI, 18:385
- Chemical ionization
atmospheric pressure, of alkanes, alkenes, and cycloalkanes, 5:177, 5:1026
automatic reaction control, in ion trap detector for quantitative plasma profiling of arecoline in treated Alzheimer patients, 2:168
cluster, and deuterium exchange mass spectrometry in supersonic molecular beams, 7:550
of deuteron transfer initiated propene loss from propoxypyridines, 11:66
early days of, 1:277
the early days of, a reminiscence, 1:277
ejection, selected, and ion trap tandem mass spectrometry, application of nonresonance excitation to, 7:668
of fluorophenyl n-propyl ethers: loss of propene from the metastable $[M+D]$ ions, 9:121
ion-molecule reaction under, gas-phase halo alkylation of H_6O -fullerene by, 4:894
negative
with oxygen, dibenzothiophene subjected to, oxidation reactions of, 4:949
in quadrupole ion trap mass spectrometry, effects of applied voltages and reaction times, 5:757
negative ion, polarizable stationary phase, and mass-resolved chromatography, high-temperature gas chromatography/mass spectrometry with, high-resolution triglycerol mixture analysis using, 7:350
 O_2^+ , of carbonyl compounds, 2:292
 O_2^+ and OH^+ , of some fatty acid methyl esters and triacylglycerols, 5:553
of phenyl n-propyl ether and methyl substituted analogs, propene loss initiated by competing proton transfer to the oxygen atom and the aromatic ring, 7:639
of polyfluorinated chemicals, connection to environmental atmospheric pressure, 14:1177
selective, of nitrogen and sulfur heterocycles in petroleum fractions by ion trap mass spectrometry, 4:322
silver nitrate, for analysis of hydrocarbon polymers by laser desorption Fourier transform mass spectrometry, 4:453
- Chemical ionization/ammonia laser desorption with ammonium bromide, Fourier transform ion cyclotron resonance mass spectrometry of aromatic hydrocarbons, 2:299
- Chemical ionization-collision-activated dissociation, disulfur monoxide-water charged complexes, isotope exchanges, FT-ICR, TQ and, 18:1664
- Chemical ionization gas, neutral, reaction of analyte ions with, 3:549
- Chemical ionization liquid chromatography, negative atmospheric pressure, mass spectrometry, effect of high-performance liquid chromatography mobile phase components on, 7:1059
- Chemical ionization liquid chromatography mass spectrometry interface, atmospheric pressure, optimization of, 7:69
- Chemical ionization mass spectrometric fragmentation, electron capture negative, and calculated internal energies for polychlorinated biphenyls, correlation between, 5:44
- Chemical ionization mass spectrometry
atmospheric pressure
high-performance liquid, quantitative analysis by, the determination of the renin inhibitor CP-80,794 in human serum, 2:164
results obtained by, internal standard correction of, 2:427
with liquid introduction via heated nebulizer interface, characterization of flame-generated C_{10} to C_{160} polycyclic aromatic hydrocarbons by, 7:276
of gas phase reactivity of isomeric arylglycoside toward amines, 15:244
utilizing an isotopically labeled reagent gas, 5:596
- Chemical ionization-MS/MS, internal energy deposition in, 14:102
- Chemical ionization of peptides, Fe^+ , 4:106
- Chemical ionization reagents, oxygenated, with vincamine, ion-molecule reactions of, 7:250
- Chemical ionization tandem mass spectrometry, desorption, of polyprenyl and dolichyl phosphates, 7:958
- Chemical modification of fluorinated self-assembled monolayer surfaces by low energy reactive ion bombardment, 6:187
- Chemical noise
in MALDI mass spectra, 13:129
in MALDI-MS, 13:129
- Chemical reaction interface mass spectrometry
implementation on a Hewlett-Packard mass-selective detector, 5:765
to measure hydrogen and deuterium, 11:400
nitrogen trifluoride in, for detection of phosphorus, deuterium, chlorine, and sulfur, 6:421
- Chemical reaction interface mass spectrometry/gas chromatog-

- raphy, selective detection of sulfur-containing compounds by, 2:250
- Chemical substructure identification by mass spectral library searching, 6:644
- Chemical vapor deposition, tetramethyl- and tetraethyl-distibines, low temperature thermal lysis behavior, 19:1336
- Chemical warfare agents
degradation products detected by negative mode IM(tof) MS, 17:241
detection by ion/molecule reactions, 182
- Chemical weapons-related compounds, detection of, with postcolumn derivatization LC-MS, 10:440
- Chemically assigned classification of aerosol mass spectra, 13:826
- Chemically oxidized dexamethasone in human plasma, detection of, electron capture negative ionization in, comparison of tandem and conventional mass spectrometry using, 1:341
- Chemokine ligands
analysis of noncovalent complexes with polyanionic molecules, ESI FT-ICR MS, 17:524
CCL2, effects of sulfate position on heparin octasaccharide binding to, MS/MS, 17:1114
- Chemometric approach to evaluate the parameters affecting the determination of reaction rate constants by ion trap MS, 9:938
- Chemoprevention agents, alkylation sites by, Keap-1, 18:2226
Chenopodium quinoa, triterpene saponin mixture, nLC-ESI-MS/MS, 17:795
- C_6H_5F and O_2^- , gas-phase reactions between, on the acidity of fluorobenzene and 1,4-difluorobenzene, 4:73
- $C_5H_5Fe^+$, gas phase reactions of, 9:314
- $CH_3 + HCH_2CH_2$ to $CH_3CH_2CH_2OH$, isomerization of, five-membered ring shift in by ab initio theory, preference for ion-neutral complex-mediated pathway over, 6:1037
- Chip-based emitters, automated nanospray using, in pharmaceutical compound quantitative analysis, 16:363, 16:798 (erratum)
- Chip-based nanoelectrospray, evaluation of, parameters for protein identification, 18:1714
- Chiral enrichment of serine, via ion soft-landing, 15:1360
- Chiral recognition characteristics of metal and proton complexes of di-*o*-benzoyl-tartaric acid dibutyl ester and L-tryptophan in gas phase, 14:571
- Chiral recognition in association between antimony potassium tartrate and bis(L-alaninate)ethylenediamine cobalt(III) complexes using ESI-MS, 11:804
- Chiral recognition in gas phase
cyclodextrin-amino acid complexes, is the three point interaction still valid in the gas phase, 12:278
mass spectrometric studies of diastereomeric cobalt complexes, 5:452
- Chiral resolution of arginine, by ESI-ion trap MS/MS, 16:825
- Chirality
iodine substituents and, D- and L- amino acids, 18:1516
in microcystins, 5:724
role in formation and stability of charged amino acid clusters, 17:275
- Chloramine, gas phase reactions of, with anionic nucleophiles, 12:139
- Chlordane, technical, highly chlorinated dimethanofluorenes in, 1:99
- Chloride adduct ions, formation and decomposition of, in negative ion ESI-MS, 11:933
- Chloride affinities, of monosaccharides, ranking of, 12:1193
- Chlorinated benzenes, $[m + 1]^+$ ions of, and D_2O or ND_3 , hydrogen/deuterium exchange between, mechanistic study of, 5:282
- Chlorinated dimethanofluorenes, highly, in technical chlordane and in human adipose tissue, 1:99
- Chlorinated phenoxyacid herbicides, mass spectra of, obtained by particle beam liquid chromatography/mass spectrometry, evidence for thermal decomposition contributions to, 3:823
- Chlorine, chemical reaction interface mass spectrometry in detection of, nitrogen trifluoride in, 6:421
- Chlorine isotope ratios, by inductively coupled plasma mass spectrometry, alleviation of polyatomic ion interferences for, 1:284
- Chlorine isotopic abundances, tropylium, monomeric metaphosphate anion, and Conestoga wagon theory, 4:761
- 4-Chloro- α -cyanocinnamic acid, PE chloramines detected using MALDI-TOF-MS and, 20:867
- Chloroacetic acid
negative ESI of, 11:809
TOFMS mass measurement of, 11:809
- Chloroalkyl phosphates, from an industrial effluent, identification of, multispectral, 2:419
- Chloroaromatic compounds, oxygen addition to, collision-induced, 7:1144
- 3-Chlorobenzidine in environmental samples, FT-ICR MS of, 10:1152
- Chloroformates, highly fluorinated, comparison of, 16:803
- Chlorophylls, modified structure and properties of, by using fast atom bombardment combined with tandem mass spectrometry, 1:72
- $CH_3NH_3^+$, formation of, from the molecular ions of octopamine and synephrine, the role of distonic ions in, 1:104
- CH_3NO_2 isomers, nitromethane and methyl nitrate, $[M - H]$ anions of, 1:295
- CH_3O^+ cation, gas-phase ion-molecule reactions of, 3:515
- $CH_3O-C(=O)-C(=O)-OCH_3^+$, breaking of, loss of CH_3 , CO_2 versus $CH_3O-C=O$ in, 10:869
- $C_3H_9O^+$ formation from ionized phytyl methyl ether, structure and mechanisms of, 7:205
- $C_3H_7O^+$ ions, collisional studies of, 1:301
- $CH_3OBOCH_3^+$, characterization of long-chain carboxylic esters in a small Fourier-transform ion cyclotron resonance mass spectrometer with, 7:1138
- $CH_3OCH_2^+$, reactions with hydrocarbons, 13:1015
- C_2H_5OD , H/D exchange reactions of emolates ions with 2, reactive collisions in quadrupole cells, 3:853
- CH_3OD , H/D exchange reactions of emolates ions with, reactive collisions in quadrupole cells, 3:853
- Cholesterol oxidation products, trimethylsilyl ethers of, mass spectra of, trimethylsilyl group migration in, product ion characterization by linked-scan tandem mass spectrometry, 4:327
- Cholesterol ozonolysis, hydroperoxy bishemiacetal product derived from, ESI-MS/MS of, 15:194
- Cholesteryl neoglycolipids, analyzed using ESI and CID-MS/MS, 18:294
- Cholesteryl polyethoxy neoglycolipids, synthetic amphiphilic, ESI MS/MS of, 16:565
- Chondroitin sulfate
bovine aorta, analysis of oversulfation using nano-ESI-Q-TOF-MS/MS and FTIRCR MS, 18:179
sulfated disaccharides in, ESI-MS/MS of, 11:916
- Chondroitin sulfate oligosaccharides, MS/MS of, 14:1270
- $C_3H_6O + II$ on from 2-methoxyethanol, characterization of, mixture analysis by dissociation and neutralization-reionization, 6:1030

- C_4H_8O+II potential energy surface, fragmentation of 1- and 3-methoxypropene ions, 3:99
- Chorismate mutase inhibitor, noncovalent, binding evaluation by ESI-MS, 14:1470
- $C_5H_5RFe^+$ (R=F, Cl, Br, Oh, OCH_3 , H_6H_5) ions, structure of in the gas phase and the generation of their neutral counterparts by neutralization-reionization mass spectrometry, 6:1143
- Chromatin-associated factors, proteomic analysis of, 14:687
- Chromatium vinosum, iron-sulfur metalloprotein from, 3+ oxidation state for, 12:817
- Chromatographic pipette tips, nano-ESI-MS/MS from, 15:1333
- Chromatography
 capillary gas/ion trap mass spectrometry quantitative methods, using labeled unlabeled internal standards, 1:265
 electrospray liquid/mass spectrometry, selective detection of phosphopeptides in complex mixtures by, 4:710
 gas
 -mass spectrometry, fast, very fast, and ultra-fast, of thermally labile steroids, carbamates, and drugs in supersonic molecular beams, 7:737
 electron ionization mass spectrometry, relative sensitivities in, correlations, with molecular parameters, 1:397
 ion trap mass spectrometry, alternating RF/DC isolations for quantitation with coeluting internal standards in, 3:85
 isotope dilution/mass spectrometry, for platinum determination in urine, 2:85
 isotope-ratio-monitoring/mass spectrometry, nitrogen isotope analyses by, 5:387
 mass spectrometry components, overlapping, spectral deconvolution for, 3:558
 on-line isotopic dilution cryofocusing/mass spectrometry, on-line, automated high-speed analysis of selected organic compounds in urban air by, 5:1001
 purge-and-trap/mass spectrometry, measuring volatile organic compounds in human blood by, importance of enhanced mass resolution in, 3:831
 suspended trapping/Fourier transform mass spectrometry for analysis of complex organic mixtures, 1:431
 gas/mass spectrometry, selective ion monitoring mode in, direct chemical analysis of UV laser ablation products of organic polymers by using, 5:106
 gas/mass spectrometry and particle beam liquid chromatography/mass spectrometry, poly(amic acid) and polyimide characterization using, 5:299
 liquid
 -electrospray ionization-mass spectrometry matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry, rapid analysis of antibiotic-containing mixtures from fermentation broths by using, 7:1227
 atmospheric pressure chemical ionization mass spectrometry interface, optimization of, 7:69
 electrospray, -tandem mass spectrometry, quantification of SR 27417 in human plasma using, a study of ion suppression, 7:1099
 high performance, particle beam mass spectrometry, investigation of enhanced ion abundances from a carrier process in, 1:92
 high-performance-particle beam-mass spectrometry, determination of polycyclic aromatic hydrocarbons by, 4:898
 liquid-mass spectrometry, high performance, of porphyrins by using an atmospheric pressure interface, 7:965
 liquid/tandem mass spectrometry analysis of peptide mixtures, data-controlled automation of, 7:532
 mass-resolved, negative ion chemical ionization, polarizable stationary phase, and mass-resolved chromatography, high-temperature gas chromatography/mass spectrometry with, high-resolution triglycerol mixture analysis using, 7:350
 mass spectrometry
 for the analysis of N-methyl carbamate pesticides, an evaluation of atmospheric pressure ionization techniques for, 3:378
 with collision-induced dissociation, the determination of glycopeptides by, 3:804
 mass transport and calibration in, 3:662
 normal phase, by using coaxial continuous flow fast atom bombardment, 3:575
 particle beam mass spectra of chlorinated phenoxyacid herbicides obtained by, evidence for thermal decomposition contributions to, 3:823
 microbore high-performance-tandem mass spectrometry utilizing pneumatically assisted electrospray ionization, quantitative determination of a nonpeptide antithrombotic in dog plasma by, 7:1238
 nanoscale packed capillary (75 μ m id), application of, to the analysis of macrolide antibiotics, 3:563
 on-line immobilized metal-ion affinity, phosphopeptides isolated by, electrospray ionization mass spectrometry of, 4:662
 particle beam liquid/mass spectrometry and gas chromatography/mass spectrometry, poly(amic acid) and polyimide characterization using, 5:299
 particle beam/mass spectrometry interface, performance of, effect of a dc glow discharge on, 5:908
 particle beam-mass spectrometry with massive cluster impact, 7:293
 reversed phase-microbore/electrospray mass spectrometry, complementary mobile phases, and sequential exoglycosidase digestion, characterization of protein N-glycosylation by, 5:350
 Chromatography efficiency, evaluation of, 16:1757
 Chromatography-liquid, mass spectrometry, mass spectrometric analysis of nitrogen- and phosphorus-containing pesticides by, 5:655
 Chromium oxyanions, gas phase, ion-molecule reactions of, 14:1067
 Chromophore-labeled disaccharides, CE-MS/MS of, 11:292
 Chromophore labeling, closed-ring, in determining linkages of linear and branched oligosaccharides, 13:1322
 Chronic lymphocytic leukemia, histone modifications in, HPLC and MS of, 15:77
 Chronopotentiometry experiments, electrospray emitter analyte electrolysis in, 11:951
 $C_4H_{12}Si^+$ isomers, metastable and collision-induced fragmentation studies of, a systematic study of structure-reactivity relations, 5:537
 $C_6H_5SSH_6H_5Cr^+$ ions and gas phase $RSH_6H_5Cr^+L$ (R=CCH₅ L⁺ arene), tandem mass spectrometry study of, 5:928
 $C_2H_5X^+$ ions (X=Cl, Br, I), tandem mass spectrometric investigation of, involvement of classical and nonclassical ethyl structures therein, 5:814
 CID. *See* Collision-induced dissociation.
 CID (collision-induced dissociation), of poly(alkylmethacrylate), 8:1206
 CID-MS. *See* Collision-induced dissociation mass spectrometry
 CINP+NCl, reactivity toward cyclic organic ethers, 10:379
 Cirrannulene and coronene, H⁰, electron affinities of, by using the kinetic method, 7:619
 CIS-MS. *See* Coordination ionspray mass spectrometry
 CIS-MS/MS *See* Coordination ionspray tandem MS.
cis-[Ru(CN)₂(bpy)₂], under ion bombardment, electron and proton transfers observed for, 13:1209
 CISE. *See* Collision-induced signal enhancement

- Cisplatin
determination of cytochrome *c* binding site for, MS, 20:1141
effect on gas phase dissociation of oligonucleotides, ESI-MS/
MS, 20:792
- Citrullination, protein, isocyanic acid as diagnostic marker, CID,
20:723
- Claisen-Schmidt reactions, gas-phase base-catalyzed, of the ace-
tone enolate anion with various para-substituted benzal-
dehydes, 7:82
- Classical and nonclassical ethyl structures, the involvement
therein, tandem mass spectrometric investigation of
(C₂H₅)X⁺ ions (X=Cl, Br, I), 5:814
- Classical quantification algorithms, acquisition and quantifica-
tion of ion images with, 1:37
- Classification of lactate dehydrogenase of different origin by
LC-MS and multivariate analysis, 14:236
- Cleavage mechanisms in multiply charge ions, 12:565
- Cleavage reactions, of complex ions, positive ion ESI-MS/MS of,
15:689
- Clemmer, David E., Biemann Medal, 2006, 18:1161
- Clinical chemistry, GC/MS in, 12:127
- Clogged probes, two thermospray probe insert designs that al-
low to be rapidly replaced, 4:182
- Closed ring labeling, N-linked oligosaccharide linkage and
branched determination using ion trap MS, 18:248
- Cluster-assisted decomposition reactions of the molecular anions
of SF₆ and C₇F₁₄, 1:372
- Cluster chemical ionization and deuterium exchange mass spec-
trometry in supersonic molecular beams, 7:550
- Cluster division models, electrospray process, LSI analysis, 17:
151
- Cluster impact, massive, liquid chromatography particle beam-
mass spectrometry with, 7:293
- Cluster impacts, massive, shock wave model for sputtering
biomolecules using, 3:311
- Cluster ions
ammonium-counter ion clusters studied by ESI MS, 18:285
ejected from an Li-Mg alloy liquid metal ion source, observa-
tion of g₂²⁺ and Mg₃²⁺, 2:76
of inorganic transition metal compounds, LA-MS, 19:488
on a VG analytical ZAB-2FQ, dissociation studies of, high and
low energy collision-induced, a compact high pressure ion
source for, 3:33
- Clusters
matrix to analyte proton transfer in, 12:726
silver, interferences in MALDI-MS of nonpolar polymers, 12:
732
- Cn ions, formation of, amino acid sequence prerequisites for,
4:874
- CNC and CNH bond activation in the fragmentation of the
[M + Ni]⁺ adducts of aliphatic amino acids, 8:749
- CNF bond cleavage, in perfluorohexane, 8:749
- CnH_{2n} + 1NH₂ macrocyclic amines, fragmentation of, electron
ionization induced, evidence for the rearrangement of
aminocarbene radical cations and a comparison with
long-chain esters, 1:238
- CNTs. *See* Carbon nanotubes
- Co, Al, Fe, Cu, ScO, M(acetone)₂⁺, collision-induced and infra-
red multiphoton dissociation studies on, in the gas phase,
4:135
- CO₂ and N₂O, gas-phase bond strengths of, comparative study
of, with the halide ions, 4:58
- Coaxial continuous flow fast atom bombardment, the applica-
tion of normal phase liquid chromatography/mass spec-
trometry by using, 3:575
- Coaxial matrix delivery, precolumn *versus*, continuous flow fast
atom bombardment with packed microcolumns, 1:312
- Cobalt, diastereomeric, glucosyl-glucose disaccharide complexes,
tandem mass spectrometry of, differentiation of stereo-
chemistry of glycosidic bond configuration, 7:953
- Cobalt complexes, diastereomeric, mass spectrometric studies of,
chiral recognition in the gas phase, 5:452
- Cobalt-coordinated hexosamine monosaccharides, stereoselective
dissociations of, 12:517
- Cobalt-coordinated pentasaccharides, linkage position determi-
nation in, ion trap-MS for, 9:1125
- Cobalt ions, comparison of gold and cobalt ion cluster projec-
tiles, molecular depth profiling, SIMS, 18:406
- Cobalt(III)-hexammine, ESI-MS and, RNA hairpins probed us-
ing, 17:1376
- Cocaine
metabolites and, screening in human urine by TOF-MS, 20:891
in urine, HPLC and MS/MS in confirmation/quantitation of,
11:257
- Coeluting internal standards in gas chromatography/ion trap
mass spectrometry, quantitation with, alternating RF/DC
isolations for, 3:85
- Coenzyme A thioesters, long-chain fatty acyl, by negative ion
fast-atom bombardment mass spectrometry and tandem
mass spectrometry, 5:416
- Cold spray ionization mass spectrometry, acid-induced protein
conformational changes and noncovalent complexes, 20:
845
- Collision
high-energy, glutathione conjugates activated by, fragmenta-
tion characteristics of, 3:815
reactive, in quadrupole cells. 2. H/D exchange reactions of
emolate ions with CH₃OD and C₂H₅OD, 3:853
- Collision-activated decomposition of protonated peptides, influ-
ence of cysteine to cysteic acid oxidation on, evidence for
intraionic interactions, 3:337
- Collision-activated decompositions, gas-phase anionic complexes
of alkali metal ions and peptides, 5:137
- Collision-activated dissociation
of conjugated benzo[*a*]pyrene, 8:50
of cyclodextrin complexes, 8:244
erythromycin analog characterization by, 13:630
of flavonoid transition metal complex threshold dissociation,
16:139
of gas phase salt complexes, stereoselectivity in, 15:1509
of ions containing S- bonds, 8:605
low energy
d-type ions from, 8:25
of lithiated ceramides, 13:680
neutral and sialylated oligosaccharides, electron detachment
dissociation, 18:2162
of oligonucleotide/metal complex fragmentation, 16:28
in quadrupole ion trap, cyclic protonated peptide and metal
complex characterization by, 15:1039
tryptic peptide dications, bifurcating fragmentation behavior,
19:1755
- Collision-activated dissociation-B/E linked scan-mass spectrom-
etry-fast atom bombardment, of aminoethyl-triphenyl-
phosphonium derivatives of amino acids, charge remote
fragmentation during, 3:769
- Collision-activated dissociation in RF-only quadrupoles, instru-
ment-independent measurements of, kinetics-based proto-
col for, precision and accuracy in tandem mass spectrom-
etry measurements, 1:272
- Collision-activated dissociation of C₄H₄⁺ isomers, a case study
in the use of the MS3 capabilities of a pentaquadrupole
mass spectrometer, 3:518
- Collision-activated dissociation of large multiply charged

- polypeptides and proteins produced by electrospray ionization, 1:53
- Collision-activated dissociation of polyatomic ions at low energy, simple C-C bond cleavage in, a dynamic study of, fundamentals of tandem mass spectrometry, 1:6
- Collision-activated dissociation of the acetone cation, kinetics and mechanism of, 3:427
- Collision-activated fragmentation of proferrioxamines, evidence for a succinimide-mediated mechanism, 6:836
- Collision cell
- charge exchange ionization in, to detect excited electron states of polyatomic ions, 12:1120
 - influence of the length and position of, on resolving power attainable from collisionally activated processes in sector instruments, 6:212
 - ion mobility cell combined with, TOF/MS and, 17:691
 - quadrupole, for low-energy collision-induced dissociation, 5:1042
- Collision cross-section database, singly-charged peptide ions, 18:1232
- Collision cross sections
- of myoglobin and cytochrome c ions with Ne, Ar, and Kr, 8:681
 - for protein ions, 4:616
 - quadrupole, segmented radiofrequency-only for, 8:697
- Collision dynamics, in quadrupole mass spectrometry, 9:101
- Collision energies, low, enhancement of ion transmission at, via modifications to the interface region of a four-sector tandem mass spectrometer, 5:460
- Collision energies and high resolution, high accuracy fragment ion selection to improve glycopeptide detection by precursor ion scanning, 14:777
- Collision energy
- arbitrary distribution of, 14:1348
 - effect on N-protonated diglycine surface-induced dissociation, 14:02
 - influence of, for the assignment of linkage position, collision-induced dissociation of alkali metal cationized and permethylated oligosaccharides, 4:197
- Collision experiments, high energy, electron loss from multiply protonated lysozyme ions in, 12:889
- Collision gas, influence of, for the assignment of linkage position, collision-induced dissociation of alkali metal cationized and permethylated oligosaccharides, 4:197
- Collision gases, helium and argon as, in the high energy collision-induced decomposition of H⁺ ions of peptides, 3:502
- Collision-induced and infrared multiphoton dissociation studies on (acetone)₂⁺ (M=Al, Fe, Co, Cu, ScO) in the gas phase, 4:135
- Collision-induced decomposition
- high-energy, of MH⁺ ions of peptides, comparison of helium and argon as collision gases in, 3:502
 - of leukotriene A₄, 13:1227
- Collision induced decomposition of peptides, choice of collision parameters, 7:677
- Collision-induced dissociation (CID). *See also* Electrospray ionization collision-induced dissociation-tandem mass spectrometry; electrospray ionization-collision-induced dissociation; matrix-assisted laser desorption/ionization-collision-induced dissociation
- α - and β -alanine-containing peptides, 19:1743
 - analysis of LTC₄ isomers using 157nm photofragmentation and, 19:14
 - of argentinated peptides, product ions from, 12:163
 - arginine-containing protonated dipeptides, fragmentation reactions, 18:1959
 - B. anthracis* petrobactin, 18:842
 - of *b*-type ions, influence of peptide size on sequence scrambling during, 20:2174
 - of bradykinin ions, in interface region of ESI, 12:772
 - branched permethylated oligosaccharides analyzed by ECD and, 19:138
 - ceramide characterization by, 13:558
 - of charge-reduced species, peptide analysis by decoupling ETD and, 20:567
 - in chiral recognition of zinc(II) ion complexes, 15:795
 - chirality in formation and stability of charged amino acid clusters, 17:275
 - comparison with SID for large system structural studies, 19:903
 - of copper ion solvation by acetone, 13:453
 - of derivatized N-linked glycans, 12:926
 - of duplex DNA, multistep mechanism in, 13:91
 - dynamic
 - in QIF MS with two-frequency excitation waveform, 18:2017
 - in a quadrupole ion trap, 18:749
 - E. coli* microcin C7-C51 characterization by, 19:1187
 - ECD, EDD and, structural analysis of polyamidoamine dendrimers, 19:1312
 - EDC and, metal ion complexes with polyamidoamine dendrimers, 20:674
 - effects of target gas in, using a double quadrupole mass spectrometer and radiofrequency glow discharge, 5:845
 - electrospray low energy, of protonated cross-linked peptides, 13:709
 - energy quantification for small molecule characterization, 20:1759
 - of ES-generated copper(II) complex ions, in formation of molecular radical cations of enkephalin derivatives, 12:1114
 - in ESI/MALDI-MS, free-radical-induced site-specific peptide cleavage in gas phase, 18:807
 - ESI-MS with, in qualitative assessment of ionic terpolymer sample monomer ratios, 13:921
 - ETD and, in linear ion traps, protonated peptides, 20:1672
 - evidence for charge separation and “ball-and-chain” propagation in the addition of 1-butene to C²⁺+60, 7:261
 - fast excitation, in quadrupole ion trap MS, 14:785
 - of fatty acids, fragment ions in, charge-associated mechanism in, 16:280
 - fragmentation:CID, activation energy of iron(II)phenanthroline, ESI-MS, 17:962
 - gas-phase peptide conformation probed by, 17:786
 - gas phase stabilities of polyether ionophore antibiotics by, 16:779
 - of Gln-Gly cleavage, biological degradation correlation with, 12:337
 - of guanosine H₂O adducts, 16:1291
 - HASTE, low mass product ion detection, quadrupole ion trap MS, 17:81
 - high-energy
 - of ceramide ions from permethylated glycosphingolipids, 5:558
 - structural elucidation of O-linked glycopeptides by, 7:319
 - in tandem mass spectrometry, low-mass ions produced from peptides by, 4:882
 - high measured mass accuracy of peptide fragments, 19:891
 - higher energy, phosphopeptide analysis using Orbitrap HCD, 20:1425
 - improved, on MALDI-TOF, 15:522
 - in-source, guanosine, 17:1050
 - incorporated with LC-IMS-TOFMS techniques, 14:1424
 - influence of amino acid composition on, model peptides, MDS, 18:1625
 - intramolecular cross-linked peptides, 20:557

- ion trap
 cations from hemoglobin α chain, 17:923
 multiply deprotonated RNA, 19:1832
- IRMPD and CID of peptides in ion traps, 20:349
- isoquinolines gas phase reaction to carboxylic acids ESI and, 19:151
- in laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry, selective parent ion axialization for, 5:900
- LIAD with, in FT-ICR-MS, peptide sequencing using, 18:525
- liquid chromatography/mass spectrometry with, determination of glycopeptides by, 3:804
- of long-range glycosyl transfer reactions in gas phase, 13:325
- low energy, of poly(ethylene glycol) esters, 13:888
- low-energy
 characterization of a high-pressure quadrupole collision cell for, 5:1042
 peptide disulfide bond cleavage via salt bridge mechanisms, 18:1109
 peptide radical dications via, 16:1795
- mechanisms of ion formation, ESI-MS/MS, 18:1477
- of methionine oxidation in peptides, 14:605
- of metal cationized peptide ions, fragmentation pathway for, 16:1305
- multiple-resonance, leucine enkephalin low-energy sequential decomposition pathways established in, 15:1794
- of multiply charged polypeptides, high-energy, produced by electrospray, 4:652
- negative ion, mobile protons in oligosaccharides, 18:952
- neutral loss of isocyanic acid, peptides, 20:723
- in nozzle-skimmer region, in biopolymer sequencing, 12:846
- oxygen as target of, 12:23
- of peptide ions, effect of charge state and the localization of charge on, 5:966
- peptide mono-oxidized thio-ether bond evaluation by ECD and, 18:493
- peptide sequence scrambling through cyclization of b_5 ions, 19:1590, 19:1776
- of peralkylated disaccharides, 14:63
- in pesticide studies, 12:590
- of poly(styrene), 9:269
- possible combination with ECD, 19:753
- of protein mixtures, 15:1341
- protein ubiquitination analyzed by ETD-based MS fragmentation or, 20:1652
- in quadrupole TOF MS, generating protein sequence tags by, 15:1478
- in quantifying methionine oxidation in calmodulin, 16:1470
- single series peptide fragment ion spectra from, in triple quadrupole, 9:606
- SORI, structural analysis of sulfated oligosaccharides by positive mode nano-ESI-FTICR-MS and MS/MS, 17:96
- SORI-, study of isomeric diglycosyl flavonoids, ESI FTICR MS, 18:2127
- SORI induced, precalculating parameters for, 13:318
- spectra of the immonium ions of two isomeric hexaglycosylceramides, electron ionization-tandem mass spectrometry of glycosphingolipids, 3:353
- sustained off-resonance irradiation, mono-deprotonated peptide nucleic acid sequencing by, 12:805
- of tantalum oxide ions, measurement in ion trap, 11:1072
- and theoretical studies of Cu-dimethoxyethane complexes, 12:480
- threshold, with guided ion beam mass spectrometer, ESI source for, 18:1124
- triacylglycerols, CID MALDI-TOF/RTOF, 20:1037
- in triple quadrupole, single series peptide fragment generated by, 9:606
- of triply protonated tryptic peptide, histidine residue position and, 12:1262
- uranyl cation in, 14:1205
- Collision-induced dissociation mass spectra
 of anthracycline aminodisaccharides, internal monosaccharide losses in, 9:710
 tandem high energy, peptide sequencing from, pattern-based algorithm for, 3:326
- Collision-induced dissociation mass spectrometry
 high-energy, of synthetic mannose-6-phosphate oligosaccharides, 7:182
 of nonionic surfactants following direct supercritical fluid injection, 3:150
 sequencing of proteolytic peptides by, 11:673
- Collision-induced dissociation of alkali metal cationized and permethylated oligosaccharides, influence of the collision energy and of the collision gas for the assignment of linkage position, 4:197
- Collision-induced dissociation of prostaglandins and thromboxanes, some examples of charge remote fragmentation, 1:325
- Collision-induced dissociation of uracil and its derivatives, 5:339
- Collision-induced dissociation spectra of peptides
 new fragment ion type in, identification of, formation of α -2–16 ions, 4:424
 uninterpreted high-energy, with sequence databases, 7:1089
- Collision-induced dissociation studies, high and low energy, of cluster ions on a VG analytical ZAB-2FQ, a compact high-pressure ion source for, 3:33
- Collision-induced dissociation tandem mass spectra, high energy, of polycyclic aromatic hydrocarbon diolepoxide adducted peptides, 5:649
- Collision-induced dissociation tandem mass spectrometry
 cholesteryl neoglycolipids analyzed using ESI and, 18:294
 high- and low-energy, in the analysis of glycoalkaloids and their aglycons, 7:173
 low- and high-energy, identification of an unusual amino acid in a semisynthetic polypeptide, 7:1034
- Collision-induced dissociation tandem mass spectrometry., with 157nm laser-induced photofragmentation, isomeric N-glycan identification by, 19:1027
- Collision-induced dissociation tandem mass spectrometry (CID MS/MS), low energy ESI, of polyglycols, lithium and transition metal ions in, 12:832
- Collision-induced fragmentation studies of all $C_4H_{12}Si^+$ isomers, a systematic study of structure-reactivity relations, 5:537
- Collision-induced ion ejection, in FTICR cell, 16:422
- Collision-induced oxygen addition to chloroaromatic compounds, 7:1144
- Collision-induced photon emissions, study of keV $CO_2^+ \bullet / He$ collisions by, 19:1551
- Collision-induced reactions of double charged porphyrins, 3:235
- Collision-induced signal enhancement (CISE), use of boundary activation to effect non-resonant CISE, 13:650
- Collision of molecular anions, of benzenedicarboxylic esters with oxygen in a triple quadrupole mass spectrometer, 10:1248
- Collision parameters, choice of, collision induced decomposition of peptides, 7:677
- Collision rate constants for polarizable ions, 14:501
- Collisional activation
 in analytical mass spectrometry, principles of, 3:599
 β -substituted meso-tetraphenylporphyrin fragmentation induced by, 13:1427

- diacyl glycerophosphatidic acid fragmentation in, 11:797, 11:892
- high-energy
 delayed dissociation spectra of survivor ions from, 4:845
 studied via angle-resolved translational energy spectra of survivor ions, 4:835
- intermolecular migration during, monitored by H/D exchange and FTICR MS/MS, 15:639
- of ion, ESI of, 9:15
- ion trap
 of deprotonated deoxymononucleoside and deoxydinucleoside monophosphates, 6:102
 relative dissociation energy measurements using, 5:250
- in ion trap, of double-stranded oligonucleotide complexes with DNA-binding drugs, 11:450
- of octaethylporphyrin decomposition, 9:767
- of oxofatty acids, 9:620
- principles of, in analytical mass spectrometry, 3:599
- in radio frequency ion traps and guides, 15:1616
- of a series of homoconjugated octadecadienoic acids with fast atom bombardment and tandem mass spectrometry, 1:183
- in triple quadrupole mass spectrometer, alkenamide fragmentation by, 15:1462
- Collisional activation and collision-activated dissociation of large multiply charged polypeptides and proteins produced by electrospray ionization, 1:53
- Collisional cooling, within the nitric oxide synthase oxygenase domain, 15:629
- Collisional damping interface for an ESI-TOFMS, 9:569
- Collisional energy, ion fragmentation and, peptides, 20:469
- Collisional focusing effects in radio frequency quadrupoles, 3:398
- Collisional relaxation of metastable electronic states of Fe^+ , 6:543
- Collisional stabilization of the $[\text{C}_5\text{H}_5^+]$ collision complex in a quadrupole ion trap mass spectrometer, 3:631
- Collisional studies of some $\text{C}_3\text{H}_7\text{O}^+$ ions, 1:301
- Collisionally-activated decomposition, peptides, cyclization reaction of fragment ions, 18:663
- Collisionally activated decomposition (CAD)
 of aminocyclitol-aminoglycoside antibiotics and their application in the identification of a new compound in tobramycin samples, 11:200
 in structural determination of sphingomyelin by MS/MS, 11:437
- Collisionally activated dichlorocarbene dications studied by tandem mass spectrometry, reactivity of, 6:1019
- Collisionally activated dissociation
 ESI-MS using of Li^+ adducts of unsaturated fatty acids, 10:600
 in evaluation of noncovalent interactions of polyether compounds with peptides, 14:1148
 of metal complexes, flavonoid glycoside characterization by, 12:537
 of poly(ethylene glycol), 12:565
 in structural characterization of triacylglycerols, 10:587
 of substituted indole, by quadrupole TOFMS, 12:1145
- Collisionally-activated dissociation
 analysis of trypanosome flagellar proteome using ETD/ETD/CAD, 20:167
 local protein hydrogen exchange data from HDX CAD MS using deconvolution, 17:1543
 of peptide ions in Fourier transform mass spectrometry, efficiency of, 1:288
 protonated 2'-deoxycytidine and 2'-deoxyuridine, 17:1335
- Collisionally activated dissociation-MS, of doubly charged hyperbranched polyesteramides, 14:332
- Collisionally activated dissociation-quadrupole MS/MS, on phosphatidylcholines, fragmentation processes in, 14:352
- Collisionally activated dissociation tandem mass spectrometry (CAD MS/MS)
 of glycosphingolipids as lithiated adducts, 12:61
 of noncovalent protein assemblies, 12:329
- Collisions of slow polyatomic ions with surfaces, scattering method and results, 14:1360
- Color bodies, on polymer surface, LC-MS/MS of, 16:295
- Column switching, DNA-phosphate adduct analysis with, 15:593
- Combination of two matrices results in improved performance of MALDI-MS for peptide mass mapping and protein analysis, 14:992
- Combinatorial libraries, synthetic failures in, hybrid gas phase separation of, 11:352
- Combined capillary gas chromatography/ion trap mass spectrometry quantitative methods using labeled or unlabeled internal standards, 1:265
- Combined linear ion trap for mass spectrometry, 8:962
- Combined mass spectrometry methods, for the characterization of a new variant of human hemoglobin, the double mutant hemoglobin Villeparisis $\beta 77(\text{EF1}) \text{His} \rightarrow \text{Tyr}$, $\beta 80(\text{EF}_4) \text{Asn} \rightarrow \text{Ser}$, 7:163
- Compact high pressure ion source for high and low energy collision-induced dissociation studies of cluster ions on a VG analytical ZAB-2FQ, A 2:438:441, 3:33
- Comparative ESI-MS study of hemocyanins from deep-sea and shore crabs, from protein oligomeric state to biotope, 14:419
- Comparative evaluations of mass spectral data bases
- Comparative studies of poly(dimethyl siloxanes) using automated GPC-MALDI-TOFMS and on-line GPC-ESITOFMS, 195
- Comparative study of different thermospray interfaces with carbamate pesticides, influence of the ion source geometry, 6:656
- Comparative study of the gas-phase bond strengths of CO_2 and N_2O with the halide ions, 4:58
- Comparing similar spectra, from similarity index to spectral contrast angle, 13:85
- Comparison of laser desorption and MALDI for ruthenium and osmium trisbipyridine complexes using FTMS, 14:393
- Comparison of algorithms and databases for matching unknown mass spectra, 9:92
- Comparison of charged derivatives for high energy collision-induced dissociation tandem mass spectrometry, 6:428
- Comparison of different tests for identification of a compound from its mass spectrum, 14:262, 14:264
- Comparison of equilibrium ion density distribution and trapping force in Penning, Paul and combined ion traps, 9:473
- Comparison of helium and argon as collision gases in the high energy collision-induced decomposition of MH^+ ions of peptides, 3:502
- Comparison of high- and low-energy collision-induced dissociation tandem mass spectrometry in the analysis of glycoalkaloids and their aglycons, 7:173
- Comparison of local anesthetic-cyclodextrin non-covalent complexes using CE and ESI-MS, 13:928
- Comparison of mass spectrometric techniques for generating molecular weight information on a class of ethoxylated oligomers, 9:282, 9:651
- Comparison of microprobe two-step laser desorption/laser ionization MS and GC/MS of polycyclic aromatic hydrocarbons in ancient terrestrial rocks, 12:989
- Comparison of negative and positive ESI-MS/MS, for LC MS/MS of oxidized deoxynucleosides, 12:80

- Comparison of SSI with APCI as interface of HPLC-MS for analysis of a drug and its metabolites, 13:204
- Comparison of tandem and conventional mass spectrometry using electron capture negative ionization in the detection of chemically oxidized dexamethasone in human plasma, 1:341
- Comparison of the CID of duplex DNA at different collision regimes, multistep dissociation mechanism in, 13:91
- Comparison of the orders of gas-phase basicities and ammonium ion affinities of polyethers by kinetic method and ligand exchange technique, 4:718
- Comparison of the reactions of the oxide radical anion (79 II) with simple aromatic compounds at atmospheric and at reduced pressure conditions, 5:367
- Compensation voltage peak shapes, using domed FAIMS, 16:948
- Competition between resonance ejection and ion dissociation during resonant excitation in a quadrupole ion trap, 5:1031
- Competitive binding to the oligopeptide-binding protein OppA, in-trap cleanup in FTICR-MS, 11:1023
- Competitive ionization of tetraphenylporphyrin in a laser generated metal ion plasma, 4:159
- Complement, Efb complex with C3d domain of, resolution by lysine acetylation and MS, 19:55
- Complementary mobile phases
reversed-phase microbore liquid chromatography/electrospray mass spectrometry, and sequential exoglycosidase digestion, characterization of protein N-glycosylation by, 5:350
and sequential exoglycosidase digestion, characterization of protein N-glycosylation by, 5:350
- Complete large-molecule high-resolution mass spectra from 50-femtomole microvolume injection, 6:220
- Complete sequence confirmation of large peptides by high energy collisional activation of multiply protonated ions, 6:234
- Complete sequencing of mono-deprotonated peptide nucleic acids by sustained off-resonance irradiation CID, 12:805
- Complex I, mitochondrial, FMN-binding subunit, MS profile of intramolecular disulfides, 19:1876
- Complex-mediated reactions, effect of the polarity of the neutral partner on, isomeric ion-neutral complexes generated from ionized 2-methylpropanol and n-butanol, 3:409
- Complex mixtures
low abundance species in, nano-ES-IQTOF-MS fucosylation study of, 12:964
peptides in, molecular weight assignment of, application of electrospray mass spectrometry and matrix-assisted laser desorption ionization time-of-flight mass spectrometry for, 4:670
phosphopeptides in, selective detection of, by electrospray liquid chromatography/mass spectrometry, 4:710
- Complex organic mixtures, analysis of, suspended trapping gas chromatography/Fourier transform mass spectrometry, 1:431
- Complexes of iron(II) with cysteine-containing peptides in the gas phase, 7:977
- Complexes of Li atoms with formaldehyde (LiOCH₂) and charge transfer interactions, 12:1229
- Complexes of silver(I) with peptides and proteins as produced in electrospray mass spectrometry, 8:781
- Composition-based sequencing, mass determination by FTI-CRMS, 15:703
- Composition of N-linked carbohydrates from ovalbumin and co-purified glycoproteins, 11:564
- COMPOST, computer program for predicting mass spectrometric information from known amino acid sequences, 2:174
- Compound identification, mass spectral library search algorithms for, optimization and testing of, 5:859
- Compound libraries, microstability screening of, parallel LC-MS in, 13:155
- Comprehensive on-line RPLC-CE-MS, of peptides, 8:495
- Computational analysis, of metal-induced conformational mutations, 12:1272
- Computational chemistry, of ionic and neutral P(OH)₂+/- in gas phase, 13:250
- Computational study of double hydrogen migration in molecular ions of endo tricyclo[5.2.1.0^{2,6}] decene, 13:1186
- Computer-assisted interpretation of copolymer mass spectra, 6:1112
- Computer program (COMPOST) for predicting mass spectrometric information from amino acid sequences, 2:174
- Computer programs, MoWeD as, to determine component molecular weights, 12:599
- Computer simulation of single-ion trajectories in Paul-type ion traps, 4:687
- Computer simulations
enhancement of dipolar resonance excitation, linear ion trap study, 18:578
for high focusing and mass filtering TRIPOLE, 18:413
- COMSPARI (comparison of spectral retention information), in GC-MS and LC-MS of identifying minor compounds in complex mixtures, 15:580
- Concentric tube vacuum chamber for high magnetic field, high-pressure ionization in a Fourier transform ion cyclotron resonance mass spectrometer, 4:168
- Condensation of glycerol with ammonium surfactants, fast-atom bombardment-induced, II, time dependence of mass spectra and tandem mass spectra, 5:92
- Condensation reactions, in protonated dipeptides, 15:1025
- Condensed phase, of oxidized peptides containing cysteine, fragmentation of, 12:228
- Condensed-phase processes in ionization, evidence indicating, fast-atom bombardment of the cyclic acetals, 4:493
- Conductive carbon-filled polymeric electrodes, novel ion optical elements for TOF-MS, 13:1170
- Conestoga wagon theory, tropylium, chlorine isotopic abundances, monomeric metaphosphate anion and, 4:761
- Conference reviews
Asilomar Conference on Mass Spectrometry, 8:673
ASMS Fall workshop on polymer MS 2004, 16:291
Eighth Lake Louise Workshop on Tandem Mass Spectrometry, 7:611
First North American FTICR Conference, 8:813
Mass Spectrometry and Allied Topics, June 2000 in Long Beach, CA, 11:921
Mass Spectrometry Pitt (Con) '97, 8:573
Mass Spectrometry PittCon[®] 2008, R19:1 (June)
Mass Spectrometry PittCon '99, 10:457
Mass Spectrometry PittCon 2004, 15:942
Mass Spectrometry PittConn 2005, 16:967
MS PittCon '98, 9:548
PDMS'93, Hirschegg, Kleinwalsertal, Austria, June 19–26, 4:971, 1993
Sanibel Conference on Mass Spectrometry: Quadrupole Ion Traps, 8:569
The Second European Meeting on Tandem Mass Spectrometry, Institute of Mass Spectrometry, University of Warwick, Coventry, UK, July 19–22, 4:196, 1992
13th Asilomar on MS, 9:545
14th International MS, Tampere, 9:178
10th Sanibel, on MS, TOFMS in, 9:849
13th Sanibel Conference on Mass Spectrometry: Informatics and Mass Spectrometry, 12:475

- 15th Sanibel conference on mass spectrometry: ion activation in chemistry and biochemistry, 680
- Configuration assignment, absolute, of isomers, by MS, 16:431
- Confocal laser scan microscopy, allophycocyanine localization in MALDI-preparations by, 17:1599
- Conformers, elongated, of ubiquitin, ESI-FAIMS-MS of, 12:894
- Conformational dependence, of gas phase H/D exchange of AMP and dAMP, 12:780
- Conformational studies of Zn-ligand-hexose diastereomers using ion mobility measurements and density functional theory calculations, 13:284
- Conformer-dependent proton transfer reactions of ubiquitin ions, 8:954
- Conformer selection of protein ions by ion mobility in a triple quadrupole mass spectrometer, 5:127
- Conjugate bases of esters derived from multifunctional alcohols including triacylglycerols, fragmentation of, 6:498
- Conotoxins, *Conus virgo*, pyroglutamic acid identification by MALDI-MS/MS, 18:1396
- Constant neutral loss, oligopeptide fragmentation viewed as, 1:336
- Constant neutral loss scanning for characterization of glycerol phosphatidylcholine phospholipids, 9:1189
- Constant ratio of B/E, scans at, for studying decompositions of peptide metal(II)-ion complexes formed by electrospray ionization, 6:608
- Construction and characterization of a small-bore electrospray ionization source, 4:964
- Contamination
cyclic polyamide oligomers from nylon 66 membrane filter disks, LC/MS, 17:652
dimethyldioctadecylammonium ions as source of, ESI MS, 19:666
- Continuous-flow fast atom bombardment mass spectrometry, an internal probe for, 3:198
- Continuous flow-fast atom bombardment mass spectrometry capillary zone-electrophoresis, identification of deoxy-nucleoside-polyaromatic hydrocarbon adducts by, 3:757
- Continuous-flow fast atom bombardment mass spectrometry of oligonucleotides, 2:157
- Continuous flow fast atom bombardment with packed microcolumns, a comparison of precolumn *versus* coaxial matrix delivery, 1:312
- Continuous-flow ligand-exchange reactors, metal-complex formation in, ESI/MS, 18:707
- Continuum background reduction in orthogonal-acceleration time-of-flight mass spectrometry with continuous ion sources, 8:125
- Contrast angle, in spectral comparison, 13:85
- Controlled-current electrolytic process inherent to electrospray, observation of gas-phase molecular dications formed from neutral organics in solution via, 7:157
- Controlled ion fragmentation in a 2-D quadrupole ion trap for external ion accumulation in ESI FTICR-MS, 12:1312
- Controversial ions (*m/z* 59, 60, 77, and 119) in the thermospray reagent plasma from ammonium acetate, on the origin of some, 5:186
- Controversial ions (*m/z* 59, 60, 77 and 119), in the thermospray reagent plasma from ammonium acetate, On the, 5:186
- Contryphan, oxidized glutathione and, fragmentation of peptide disulfides from negative ion mass spectrometry, 19:358
- Conus virgo*, conotoxins, pyroglutamic acid identification by MALDI-MS/MS, 18:1396
- Conventional mass spectrometry and tandem mass spectrometry, comparison of, using electron capture negative ionization in the detection of chemically oxidized dexamethasone in human plasma, 1:341
- Converters, analog-to-digital, two, increasing the dynamic range of a transient recorder by using, 7:107
- Cooks, R. Graham, 2006 distinguished contribution in MS awardee, 19:159
- Cooks' kinetic method
in determining gas phase acidity of methylthioacetic acid, 16:535
extended, cysteine-polyalanine peptide gas-phase acidities determined by, 18:188
- Cooling rates, determination of, quadrupole ion traps, 17:932
- Coordination ionspray-MS, Ag⁺, in analysis of unsaturated compounds, 872
- Coordination ionspray tandem mass spectrometry (CIS-MS/MS), diacyl peroxide analysis by, 12:449
- Copolymer mass spectra, computer-assisted interpretation of, 6:1112
- Copolymers
analysis of, by laser desorption Fourier transform mass spectrometry, 1:66
with compositional drifts or sequence constraints, mass spectra of, 15:374
isobutylene/ μ -methylstyrene, compositional analysis by MALDI-MS, 15:621
- Copper
ground state and Fe⁺, the reactions of, with the 20 common amino acids, 7:722
new matrix for MALDI/MS using, 20:1263
peptide complexes with, electrogeneration in microspray MS, 19:560
reduction in MALDI-MS, 14:42
ternary copper (II) complexes with, macrocyclic ligand effects on electron and proton transfers, 20:972
- Copper adducts of protonated ubiquitin, electrospray ionization mass spectrometry of, 6:521
- Copper-dimethoxyethane complexes, CID and theoretical studies of, 12:480
- Copper-glycine solutions, electrospray ionization of, 7:25
- Copper ions, solvation by acetone, 13:453
- Copper-octaethylporphyrin, collisionally activated decomposition of, 9:767
- Copper phthalocyanine, heterolytic dissociation of negative ions into counter ions, 19:375
- Copper(II), 2-diarylcyclopropane oxidation by, 9:1, 9:966
- Copper(II) complexes
ES-generated, CID of, in formation of molecular radical cations of enkephalin derivatives, 12:1114
low energy dissociation of, 12:490
reduction of, in ESI source, 9:1217
- Corona discharge ionization, atmospheric pressure, nitric oxide-assisted, for the analysis of automobile hydrocarbon emission species, 5:1107
- Corona discharges, electrochemical electrospray ionization initiated by, 20:1988
- Coronene and cirrannulene, H₂O, electron affinities of, by using the kinetic method, 7:619
- Correct identification, estimating probabilities of, from results of mass spectral library searches, 5:316
- Correlation between electron capture negative chemical ionization mass spectrometric fragmentation and calculated internal energies for polychlorinated diphenyls, 5:44
- Correlation of stereoselectivity and ion response in electrospray mass-spectrometry, electrospray ionization-mass spectrometry as a tool to predict chemical behavior?, 6:207
- Correlations of chemical mass shifts of *para*-substituted aceto-

- phenones and benzophenones with Brown's σp^+ constants, 13:623
- Correlations of relative sensitivities in gas chromatography electron ionization mass spectrometry with molecular parameters, 1:397
- Corticosteroids, perfluoroacyl derivatives of, and related substances for potential use in quantitative mass chromatography mass spectrometry, 6:264
- Costa Rica Airborne Research and Technology Applications, 3D gas concentration mapping using portable MS, 19:1411
- Coulomb-induced frequency modulation, two-dimensional, in Fourier transform ion cyclotron resonance, a mechanism for line broadening at high mass and for large ion populations, 4:909
- Coulomb mutual interaction and image charge in Fourier transform ion cyclotron resonance precise mass difference measurements, systematic errors originating from, analysis and estimation of, 4:855
- Coulombic repulsion, for cleavage of multiply charged ions, 12:565
- Coulombic repulsion effects, on dissociation pathways of protein complexes in gas phase, 18:617
- Coumarins, analysis of by micro high-performance liquid chromatography-mass spectrometry with a particle beam interface, 6:132
- Counter ions, copper phthalocyanine, heterolytic dissociation of negative ions into, 19:375
- Counterion and solvent, effects of, on ion pairing and observed charge states of diquatery ammonium salts in electrospray ionization mass spectrometry, 7:1050
- Counterions, varying effect of, on the sensitivity of electrohydrodynamic and fast atom bombardment mass spectrometry, 1:233
- Coupling a particle beam interface directly to a quadrupole ion trap mass spectrometer, 4:38
- Coupling two-step laser desorption/ionization with aerosol TOFMS for the analysis of individual organic particles, 9:1068
- Covalent labels, strategies for
proteins, 20:1193
proteins (reprint, critical insight), 20:1193 in July 20:2009
- CP-80,794, in human serum, quantitative analysis by high-performance liquid chromatography atmospheric pressure chemical ionization mass spectrometry, 2:164
- CpHA, monitoring zinc affinity by automated nano-ESI-MS, 17:180
- Cr(CO)₅, gas-phase, relaxation of, 10:950
- Creatine kinase, complexed with ADP or ATP, deconvolution for specific *vs.* nonspecific interactions, ESI-FT-ICR MS, 17:1239
- CRIMS. *See* Chemical reaction interface mass spectrometry
- Critical energies, estimation of, an empirical approach to, by using a quadrupole ion trap, 7:1116
- Critical evaluation of a comprehensive mass spectral library, 10:287, 10:565
- Cross-linking, probability-based shotgun cross-linking site analysis, 20:1896
- Cross-ring cleavage, in glycoconjugate mixtures, 15:1863
- Cross sections
collision, for protein ions, 4:616
of peptide ions, data base of, 10:1188
- Crosslinked peptides, Lys-Lys, influence of cross-linker identity on gas-phase dissociation, MS/MS, 17:395
- Crosslinker, for MS analysis of protein complex quaternary structure, 12:222
- 18-Crown-6, protonated, FT-ICR hydration of, 10:862
- 18-Crown-6 attachment, site-specific, ESI-MS probing protein structure using, 17:1209
- Crown ether complex self-assembly, ESI-MS detection of, 16:1162
- Crown ether-metal ion complexes, direct desorption of, in liquid secondary ionization mass spectrometry, evidence for, 5:638
- Crown ethers, orders of relative alkali metal ion affinities of, determination of, by the kinetic method, 3:543
- Cryogenically cooled sample holder for polymer sample analysis by radiofrequency glow discharge mass spectrometry, 8:1214
- α -Crystallin, GlcNAcylation sites of, identification by Q-TOFMS, 12:1106
- Crystallography, of binding of aldose reductase inhibitors, 10:635
- CSD. *See* Charge state distributions
- Cu, Al, Fe, Co, ScO, M(acetone)₂⁺, collision-induced and infrared multiphoton dissociation studies on, in the gas phase, 4:135
- Cu⁺. *See* Copper
- Cu⁺ chemical ionization for analysis of hydrocarbons by gas chromatography-mass spectrometry, 9:731, 9:981
- Cubic trap, ion coherence, induced signal, and radial or z ejection in, effect of ion initial axial position on, simulated ion trajectory and induced signal in ion cyclotron resonance ion traps, 5:807
- Cucurbit[8]uril, dication complexes with, unimolecular gas-phase reactivity, 18:1863
- Cu(II) tripeptide complexes, tripeptide radical cation generation in, via intramolecular electron transfer, 16:763
- Cumulative distribution functions to characterize MS of intact proteins, 13:40
- Curie-point pyrolysis, of bacterial fatty picolinyl esters, 14:58
- Curie-point pyrolysis tandem mass spectrometry, an investigation of dipeptides containing polar and nonpolar side groups by, 3:750
- Current density profiler, electrospray, 16:312
- Current pulse, fast direct, using as pump and laser probe, determination of ion frequencies in a quadrupole ion trap by, 5:29
- Curve-crossing pictures, study of keV CO₂⁺/He collisions by collision-induced photon emissions, 19:1551
- Cuticular hydrocarbons, from wasp, GC-MS of, 9:320
- Cyanide ions, carbon-13 and nitrogen-14 labeled polyglycine, CN⁻ secondary ion formation, 17:1181
- α -Cyano-4-hydroxycinnamic acid, effect of diammonium citrate to matrix of, in detection of phosphorylated peptides, MALDI-TOF MS, 18:1925
- α -Cyano-4-hydroxycinnamic acid, peptide cocrystallization with, promotion within levitated droplets, 16:254
- α -Cyano-4-hydroxycinnamic acid clusters, MALDI-PSD and LIFT-TOF/TOF of, 15:336
- Cyano groups, adjacent, in the gas phase, on the stabilization of carbonions by, 5:159
- Cyanuric acid determination, stable association complex ESI for, 12:1085
- Cyclic acetals, fast-atom bombardment of, evidence indicating the predominant involvement of condensed-phase processes in ionization, 4:493
- Cyclic and linear polyglycols, [M + X]⁺ ions (X=H, Li, Na, K) from, dissociation characteristics of, 5:1081
- Cyclic GMP, ESI-TOF-MS of, 15:1392
- Cyclic peptide fragments, nomenclature for labeling of, 10:360
- Cyclic peptides
IM-MS of, 15:870
protonated, MSn characterization of, 15:1039

- Cyclic polyamide oligomers, contamination from nylon 66 membrane filter disks, LC/MS, 17:652
- Cyclic sudden sampling introduction mode, in MIMS, 15:823
- Cyclization, intramolecular, and electron impact ionization mass spectrometry, in 2-substituted pyrimidin-4(3H) ones, 5:113
- Cyclization
and electron-impact-induced 3,3-sigmatropic rearrangement in phenyl allenylmethyl ethers, 6:195
- peptide b_n ions, 20:2248
- Cyclization of ortho-cyclopropylphenyl benzamides in gas and liquid phases, 12:956
- Cycloalkylidene malonic acids, polycyano push-pull pentadienes from, EI-MS of, 14:189
- Cyclodepsipeptides, identification using negative ion mass spectrometry, 20:2221
- β -Cyclodextrin
complexes with polyamide acids, ESI-MS, 17:9
- use in determining MWs of low-molecular weight explosives, MALDI-TOF MS, 17:189
- Cyclodextrin-amino acid complexes, gas phase, chiral recognition in, 12:278
- Cyclodextrin bearing, by azobenzene group, identification by ESI-MS, 14:1116
- Cyclodextrin complex binding constants, ESI-MS evaluation of, 11:23
- Cyclodextrin complexes
gas phase, zwitterion formation in, 16:166
- with local anesthetic, comparison using CE and ESI-MS, 13:928
- specificity of, electrospray MS of, 13:946
- Cyclodextrin inclusion complexes, detection of by electrospray mass spectrometry, false positives and, 6:437
- Cyclodextrin-peptide complexes, dissociation of, ESI-FTMS of, 8:244
- Cyclodextrins, encapsulation of amino acids by, electrospray ionization mass spectrometry of, 6:866
- Cyclodextrin(s), unsubstituted and disubstituted, ESI-MS of, 14:124
- Cyclohexamine on soil surfaces exposed to laboratory air, static secondary ionization mass spectrometry detection of, 7:168
- Cyclohexene-fused 2-N-phenyliminoperhydro-3,1-oxazines and related thiazines, fragmentations of, effects of N-substitution on, 2:125
- 1,3,5,7-Cyclooctatetrane, electron affinity of, by kinetic method, 9:1141
- Cyclopentadienylcobalt ion, reactions with alkanes, 14:51
- Cyclopentadienylmanganese cations, substituted, in the gas phase, metastable ion study of, 4:216
- Cyclopentanone, ionized, ring-opening γ -transfer in, Ab initio characterization of, similarity to ion-neutral complex-mediated alkane eliminations, 7:1251
- Cyclopentylimino derivatization, for double bond location in polyunsaturated long chain alkenones, GC-MS in, 15:1161
- Cyclo(Pro-Gly)₃ complexes, ESI-MS of, 16:1536
- Cyclopropylphenyl benzamides, cyclization in gas and liquid phases, 12:956
- Cyclopropyltetrahydropyridine, fragmentation of, 8:724
- Cyclotrons, reduction of axial kinetic energy-induced perturbations on observed frequency, 19:467
- Cycloalkanes, atmospheric chemical ionization of, 5:177, 5:1026
- Cycloaddition, polar [4+2+] Diels-Alder, to nitrilium and immonium ions in the gas stage, applications of multiple stage mass spectrometry in a pentaquadrupole, 6:1
- Cylindrical geometry FAIMS, of differing electrode diameters, 16:349
- Cylindrical ion trap source for time-of-flight mass spectrometry, a segmented ring, 7:1009
- Cysteic acid, cysteine oxidation to, influence of, on the collision-activated decomposition of protonated peptides, evidence for intraionic interactions, 3:337
- Cysteic acid-containing peptides, low energy collision-activated dissociation of, from d-type ion, 8:25
- Cysteic acid groups, effects on gas-phase reactivity and dissociation of [M + 4H]⁴⁺ ions from insulin chain B, 10:928
- Cysteic acid residue, in oxidized human apoprotein B-100, tandem mass spectrometric characterization of, 6:242
- Cysteine
alkylated, side-chain fragmentation in ECD MS, 17:1271
- characterization of oxidation products of, in PTP1B active site, 20:1540
- to cysteine acid oxidation, on the collision-activated decomposition of protonated peptides, evidence for intraionic interactions, 3:337
- gas phase fragmentation of long-lived radical cations, 20:985
- oxidized peptides containing, fragmentation of, 12:228
- protonated, fragmentation of, 9:1275
- Cysteine and cystine in peptides and proteins, quantitative assessment of, following organomercurial derivatization and analysis by matrix-assisted laser desorption ionization mass spectrometry, 5:359
- Cysteine peptides
complexes of iron(II) with, in the gas phase, 7:977
- dehydration of, 9:945
- Fe²⁺ interaction with, gas phase studies of, 9:1285
- iron(II) complexes with, 8:199
- protonated ions of, fragmentation of, 15:697
- Cysteine-polyalanine peptides, gas-phase acidities determined by extended Cook's kinetic method, 18:188
- Cysteine reactivity, of estrogen receptor zinc fingers, 16:2017
- Cysteine sulfenic acid, in MMP-7, MS/MS, 18:1544
- Cysteinyl eicosanoids, ESI and MS/MS of, 12:763
- Cystine in peptides and proteins, quantitative assessment of, following organomercurial derivatization and analysis by matrix-assisted laser desorption ionization mass spectrometry, 5:359
- Cystinyl proteins, mass mapping disulfide structures of, negative signature mass algorithm in, 14:1032
- Cytochrome *c*
collision cross sections of, with Ne, Ar, and Kr, 8:681
- conformation of gas-phase ions, 19:1907
- determination of cisplatin binding site by MS, 20:1141
- in methanol, ESI-MS of, 12:317
- modification by 4-hydroxy-2-nonenal, 15:1136
- myoglobin and, heme binding in, by ion spray mass spectrometry, 4:631
- oxidized and reduced, regional stability of, 8:1039
- probing structure of, by ESI-MS, 15:1612
- thermal denaturation analyzed by ESI-MS, 20:819
- Cytochrome *c*₂, *R. capsulatus*, local stability probed by solution-phase H/D exchange and MS, 17:1518
- Cytochrome *c*-cytochrome b5 complex, MS/MS of, 13:59
- Cytochrome *c* ions
in gas phase, in Paul trap, 16:1493
- gas phase studies of, temperature-dependent H/D exchange in, 13:506
- multiply charged, gas-phase proton transfer reactions involving, under thermal conditions, 3:624
- Cytosine nucleoside, self-assembling into dimers, evaluation with ESI-MS/MS, X-ray diffractometry, and theoretical calculations, 15:268, 15:436
- CZE. *See* Capillary electrophoresis; Capillary zone electrophoresis

D

D. *See* Deuterium

D-labeling, insertion in the gas phase as revealed by, 5:525

D-type ions

promotion of

during the low energy collision-induced dissociation of some cysteic acid-containing peptides, 8:25

from cysteic-acid containing peptides, during low energy collisional-activated dissociation, 8:25

DA-APPI. *See* Dopant-assisted atmospheric pressure photoionization

DAD. *See* Diode array detection

DART. *See* Direct analysis in real time

DASER-MMF. *See* Data self-calibration mixture mass fingerprint searching

Data, electronic archiving of, networking mass spectrometer data systems for, 4:742

Data acquisition, simultaneous vs single-channel scanning, for ICP sector-field mass spectrometers, 14:227

Data base

mass spectra, electron-ionization mass spectra, 2:432

mass spectral, comparative evaluations of, 2:438, 2:441

Data-controlled automation of liquid chromatography/tandem mass spectrometry analysis of peptide mixtures, 7:532

Data increase, instrumentation and, 8:1

Data processing

of hydrogen exchange mass spectra using *HX-Express*, 17:1700

on-line laser MS of inorganic, organic, or biological airborne particles, 10:648

parallel processing of large data sets from nanoLC-FTICR-MS, 18:152

Data self-calibration mixture mass fingerprint searching, protein identification in complex mixtures, 19:1915

Data systems, mass spectrometer, networking, for improved productivity and electronic archiving of data, 4:742

Database search, screening MS/MS spectra for, 11:422

Databases

generation of N-glycan structure databases, 20:1739

660 peptide ion cross sections, use of intrinsic size parameters for bona fide predictions of cross sections, 10:1188

protein, amino acid sequences in, an approach to correlate tandem mass spectral data of peptides with, 5:976

sequence, with uninterpreted high-energy collision-induced dissociation spectra of peptides, 7:1089

Daughter ion formulas, by multiple stages of mass spectrometry, 1:166

DDT metabolites and PCBs, oxygen addition-induced dechlorination of, in electron capture mass spectrometry, regioselectivity of, 7:66

De novo sequencing

of peptides using MALDI-TOF-TOF, 13:784

of proteolytic peptides by a combination of C-terminal derivatization and nano-ES/CID-MS, 11:673

Deamination

EDC monitoring of, during protein and peptide sample processing, 19:855

protonated amines to yield protonated imines, 9:1146

protonated nucleobases, 18:2040

Decarboxylation, of metastable methyl benzoate, 17:807

Decay fragmentation, postsorce, of N-linked carbohydrates from ovalbumin and related glycoproteins, 11:572

Dechlorination of PCBs and DDT metabolites in electron capture mass spectrometry, oxygen-addition induced, regioselectivity of, 7:66

Declustering and fragmentation of protein ions from an ESI source, 8:1053

Decomposition, thermal, contributions to the mass spectra of

chlorinated phenoxyacid herbicides obtained by particle beam liquid chromatography/mass spectrometry, 3:823

Decomposition of peptides, collision induced, choice of collision parameters, 7:677

Decomposition pathways, low-energy sequential, of leucine enkephalin, using multiple-resonance CID in quadrupolar ion guide, 15:1794

Decomposition reactions, cluster-assisted, of the molecular anions of SF₆ and C₇F₁₄, 1:372

Decompositions

cationized heterodimers of amino acids in relation to charge location in peptide ions, 4:461

low energy, of gas-phase peptide, ions, role of the site of protonation in, 7:522

N-(substituted benzalamino)phthalimide radical cations embody ion-neutral complexes and Stevenson's rule, 5:837

odd- and even-electron anions derived from deoxy-polyadenylates, 8:148

peptide metal(II)-ion complexes formed by electrospray ionization, use of scans at a constant ratio of B/E for studying, 6:608

structure and collision activated, gas-phase anionic complexes of alkali metal ions and peptides, 5:137

Deconvolution

by an entropy-based algorithm, electrospray ionization mass spectrometry, 3:207

of natural isotopic abundance, 8:659

spectral, for overlapping GC/MS components, 3:558

Deconvolution method

CK complexed with ADP or ATP, specific vs nonspecific interactions, 17:1239

local protein data from HDX CAD MS using, 17:1543

Defect formation on surfaces bombarded by energetic multiply charged proteins, implications for the conformation of gas-phase electrosprayed ions, 7:329

Defining and using microbial spectral databases, 13:875

Deflected drift-trajectories in orthogonal acceleration time-of-flight mass spectrometry, 5:588

Deflection of laser-produced ions in laser-induced thermal-desorption/Fourier-transform mass spectrometry for surface analysis, 5:583

Degradation processes, primary thermal, of peptides, using the mass spectrometric technique K⁺IDS, K⁺ ionization of desorbed species, 5:564

Dehalogenation of organic compounds, liquid secondary ionization mass spectrometry beam-induced, secondary electron capture in, evidence for a mechanism that involves, 7:1109

Dehydration of peptide [M + H]⁺ ions in the gas phase, 4:477

Dehydroalanine, selenocysteine conversion to, 14:593

Delayed dissociation spectra of survivor ions from high-energy collisional activation, 4:845

Delayed extraction matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (DE-MALDI-TOFMS), of complex polymer mixtures, isobaric interferences in, 12:911

Denaturation, thermal

of cytochrome *c*, analyzed by ESI-MS, 20:819

of ribonuclease S, by electrospray ionization mass spectrometry, 5:614

Dendrimers, hydrocarbon, by laser desorption time-of-flight and Fourier transform mass spectrometry, 5:731

Denitration of nitroaromatic compounds by aryl nitrile radical cations, 11:33

Density functional theory

calculations, a_n and a_n^* fragments, isotope labeling, MS/MS and, 19:1788

- hydrolysis of PtenCl₂ investigated by ESI MS and, 20:1015
 mechanism of trimethylamine loss from acetylcholine, 20:238
 study of peptide ions generated by MALDI, 20:1327
- Density functional theory calculations, of Zn-ligand-hexose diastereomers, 13:284
- Deoxy-polyadenylates, odd and even electron anions from, decomposition of, 8:148
- Deoxyadenosine monophosphate, gas phase H/D exchange of, conformational dependence of, 12:780
- 2'-Deoxycytidine, protonated 2'-deoxyuridine and, collisionally-activated dissociation, 17:1335
- Deoxydinucleotides, self-complementary, complex ions derived from, cleavage reactions of, positive ion ESI-MS/MS of, 15:689
- Deoxyguanosine
 in cluster with sinapinic acid, electron transfer vs proton transfer in, 12:176
 linkage to deuterated 2-hydroxyestradiol, LC/ESI-ion trap MS of, 14:364
 N², and C8 adducts of 2-acetylaminofluorene, isomeric, differentiation of, by fast-atom bombardment and tandem mass spectrometry, 5:58
- Deoxyguanosine clusters, G-quadruplexes in, ion mobility MS of, 16:989
- Deoxymononucleoside and deoxydinucleoside monophosphates, deprotonated, ion trap collisional activation of, 6:102
- Deoxynucleoside adducts of polycyclic aromatic hydrocarbon dihydrodiol-epoxides, low energy tandem mass spectrometry of, 6:248
- Deoxynucleoside-polyaromatic hydrocarbon adducts, identification of, by capillary zone electrophoresis continuous flow-fast atom bombardment mass spectrometry, 3:757
- Deoxynucleosides
 adducts with heterocyclic aromatic amines, analysis by ion trap MS, 18:2107
 oxidized, LC of, ESI-MS/MS for, 12:80
- Deoxyoligonucleotides
 protonated, sodium controlled reactivity in gas phase, MALDI, 20:689
 thiolated, characterization and sequence variation, HPLC-MS, 17:1396
- Deoxyribose nucleotide dimer anions, dissociation energies of, 10:1095
- Deoxyribose radicals, modeling by neutralization-reionization MS, 15:1055, 15:1068
- 2'-Deoxyuridine, protonated 2'-Deoxycytidine and, collisionally-activated dissociation, 17:1335
- Deprotonated deoxymononucleoside and deoxydinucleoside monophosphates, ion trap collisional activation of, 6:102
- Deprotonated peptides, fragmentation of, 12:1
 phenylalanine and, 13:1242
- Deprotonated phenol fatty acid esters, ion-dipole complex formation from, evidenced by using gas-phase labeling combined with tandem mass spectrometry, 4:343
- Deprotonated trinucleosides, gas phase conformations of, 14:161
- Deprotonation reactions
 on globular and denatured proteins at atmospheric pressure, ESSI-MS, 19:455
 protonation, gas phase reactions and, fluorinated 1,2,4-triazines, ESI MS, 19:686
- Derivatization, organomercurial, quantitative assessment of cysteine and cystine in peptides and proteins following, by matrix-assisted laser desorption ionization mass spectrometry, 5:359
- Derivatization of 5-fluorouracil with 4-bromomethyl-7-methoxycoumarin for determination by liquid chromatography-mass spectrometry, 9:970
- Derivatization of protonated peptides via gas phase ion-molecule reactions with acetone, 11:244
- Derivatization, mass spectrometric detection of LTC₄, LTD₄, and PTE₄ by, enhancement of, 5:292
- Dermatan sulfate oligosaccharides, EDD of, 19:294
- Desalting, of fluids, capillary isoelectric focusing-MS in, 8:743
- Design and calibration of an electrostatic energy analyzer-time-of-flight mass spectrometer for measurement of laser desorbed ion kinetic energies, 6:619
- Design and performance of a high resolution electrospray ion source for a magnetic sector mass spectrometer with a heated capillary inlet, 6:1256
- Design and performance of a novel electrospray interface, 3:18
- Design and performance of a plasma-source mass spectrograph, 8:307
- Design and performance of an in-line surface-induced dissociation device in a four-sector mass spectrometer, 6:257
- Desorbed species, K⁺IDS, K⁺ ionization of, mass spectrometric technique of, characterization of the primary thermal degradation processes of peptides using, 5:554
- Desorption
 direct, of crown ether-metal ion complexes in liquid secondary ionization mass spectrometry, 5:638
 laser
 direct *versus* matrix-assisted, laser microprobe mass spectrometry of quaternary phosphonium salts, 4:798
 matrix-assisted, neutrals generated by, gas-phase cationization and protonation of, 4:393
 observed fast fragmentation of peptides in, factors that influence, 7:225
 laser ionization mass spectrometry, matrix-assisted, with 2-(4-hydroxyphenylazo)benzoic acid matrix, 4:399
 time-dependent, using a model pulsed bombardment ionization method, improved detection limits for fast atom bombardment mass spectrometry, 3:637
- Desorption/absorption in ²⁵²Cf plasma desorption mass spectrometry, substrates for, small molecules as, 2:29
- Desorption chemical ionization tandem mass spectrometry of polyprenyl and dolichyl phosphates, 7:958
- Desorption electrospray ionization
 enhanced signal using oxidizing electrosprays, 17:1091
 salt tolerance of, 18:2218
- Desorption electrospray ionization mass spectrometry
 direct analysis of liquids by, 20:10
 fluorescence spectroscopy and, attomole amounts of rhodamine 6G detected by, 18:1093
 high-throughput quantitative analysis by, 20:321
 implementation on fieldable MS, 19:1419
 internal energy distributions in, 19:420
 monitoring kinetics of Baeyer-Villiger solid-state organic reactions, 20:2087
 quantization of active ingredients in solid artesunate antimalarials, 19:380
 sphingolipid and phospholipids, MS/MS and, 19:531
 SVOC aerosols, analysis by, 19:450
- Desorption ionization
 laser, matrix-assisted, of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, high resolution end group determination of low molecular weight polymers by, 7:449
 in macromolecular mass spectrometry, editorial, 19:1041
- Desorption/ionization of biomolecules from aqueous solutions at atmospheric pressure using an infrared laser at 3(μ)m, 13:354
- Desorption ionization on silicon mass spectrometry, effect of solvent and surface effects on analyte ionization, 19:8

- Desorption kinetics, peptide ions, soft landed surface effects on charge reduction and, 20:901
- Desorption matrix-assisted. *See* Matrix-assisted laser desorption/ionization
- Detailed analysis of α - ω -bis(4-hydroxybutyl)poly(dimethylsiloxane) using GPC-MALDI-TOF-MS, 13:170
- Detailed study of the quadrupole mass analyzer operating within the first, second, and third (intermediate) stability regions
and third (intermediate) stability regions, analytical approach to, 9:50
transmission and resolution of, 9:70
- Detection and characterization of hydroxyl radical adducts by MS, 12:1214
- Detection and characterization of methionine oxidation in peptides by CID and ECD, 14:605
- Detection and identification of carcinogen-peptide adducts by nano-electrospray MS/MS, 9:202
- Detection and quantification of the sulfated disaccharides in chondroitin sulfate by ESI-MS/MS, 11:916
- Detection and sequencing of phosphopeptides affinity bound to immobilized metal ion beads by MALDI-MS, 11:273
- Detection limits, improved, for electrospray ionization on a magnetic sector mass spectrometer by using an array detector, 5:194
- Detection mass bias in atmospheric pressure ionization mass spectrometry, 2:232
- Detection of 25,000 molecules of substance P by MALDI-TOFMS and investigations into fundamental limits of detection in MALDI, 12:1055
- Detection of a thermally unstable intermediate in the Wittig reaction using low-temperature liquid secondary ion and API-MS, 9:1168
- Detection of amino acids or alcohols by electron capture negative ion chemical ionization mass spectrometry, pentafluorobenzyl chloroformate derivatization for, 6:525
- Detection of estrogen DNA-adducts in breast tumor tissue and healthy tissue by combined nano-LC and nano-ES MS/MS, 14:482
- Detection of femtomole and sub-femtomole levels of peptides by tandem magnetic sector/reflection time-of-flight mass spectrometry and matrix-assisted laser desorption ionization, 2:91
- Detection of intact double-stranded DNA by MALDI, 6:972
- Detection of noncovalent FKBP-FK506 and FKBP-rapamycin complexes by capillary electrophoresis-mass spectrometry and capillary electrophoresis-tandem mass spectrometry, 6:85
- Detection of pharmaceutical compounds in tissue by MALDI and laser desorption/chemical ionization MS/MS with a quadrupole ion trap, 10:1315
- Detector performance and ion trajectories, inductively coupled plasma mass spectrometry with an enlarged sampling orifice and offset ion lens, 4:16
- Detector pulse width and jitter in a linear orthogonal-acceleration time-of-flight mass spectrometer, resolution limitations from, 5:772
- Determination of binding selectivities in host-guest complexation by electrospray/quadrupole ion trap mass spectrometry, 9:1049
- Determination of calcium binding sites in gas phase small peptides by MS/MS, 9:1020
- Determination of 3,3'-dichlorobenzidine and its degradation products in environmental samples with a small low-field FT-ICR mass spectrometer, 10:1152
- Determination of diquat and paraquat in water by LC-(ESI)MS, 9:830
- Determination of flavone, flavonol, and flavanone aglycones by LC-ESI-MS/MS, 12:707
- Determination of glycopeptides by liquid chromatography/mass spectrometry with collision-induced dissociation, 3:804
- Determination of high-energy fragmentation of protonated peptides using a BEqQ hybrid mass spectrometer, 4:470
- Determination of ion frequencies in a quadrupole ion trap by using a fast direct current pulse as pump and a laser probe, 5:29
- Determination of isomeric dibenzo[*a,l*]pyrene-adenine adducts by six different tandem mass spectrometric experiments, 8:977
- Determination of linkages of linear and branched oligosaccharides using closed-ring chromophore labeling and negative ion trap MS, 13:1322
- Determination of monoisotopic masses and ion populations for large biomolecules from resolved isotopic distributions, 6:229
- Determination of oligonucleotide composition from mass spectrometrically measured molecular weight, 4:204
- Determination of orders of relative alkali metal ion affinities of crown ethers and acyclic analogs by the kinetic method, 3:543
- Determination of pesticides in aqueous samples by solid-phase microextraction in-line coupled to gas chromatography-mass spectrometry, 6:1119
- Determination of photomodified oligodeoxynucleotides by exonuclease digestion, MALDI, and PSDMS, 12:1127
- Determination of plutonium 240/239 ratios in Lake Ontario sediments, 2:240
- Determination of polycyclic aromatic hydrocarbons by high-performance liquid chromatography-particle beam-mass spectrometry, 4:898
- Determination of positions, velocities, and kinetic energies of resonantly elicited ions in the quadrupole ion trap mass spectrometer by laser photodissociation, 4:792
- Determination of steroids by LC-MS, 8:1010
- Determination of the activation energy for unimolecular dissociation of a noncovalent gas phase peptide, substance complex by infrared multiphoton dissociation FTICR MS, 12:82
- Determination of the anomeric configuration of glycosyl esters of nucleoside pyrophosphates by fast-atom bombardment tandem mass spectrometry, 6:516
- Determination of the anomeric configuration of glycosyl esters of nucleoside pyrophosphates and polyisoprenyl phosphates by fast-atom bombardment tandem mass spectrometry, 7:541
- Determination of the CAD of substituted indole by orthogonal acceleration Q-TOFMS, 12:1145
- Determination of the dissociation kinetics of a transient intermediate, 10:119
- Determination of the proton affinities of bromo- and iodoacetonitrile using the kinetic method with full entropy analysis, 11:601
- Determination of thermal energy and its distribution in peptides, 10:323
- Determining anomericity of the glycosidic bond in Zn(II)-diethylenetriamine-disaccharide complexes using MSn in a quadrupole ion trap, 10:269
- Determining synthetic failures in combinatorial libraries by hybrid gas phase separation methods, 11:352
- Deuterium. *See also* Hydrogen/deuterium exchange
chemical reaction interface mass spectrometry in detection of, nitrogen trifluoride in, 6:421
KIEs in microsolvated gas-phase E2 reactions, 18:1046

- kinetic isotope effects, microsolvated gas-phase E2 reactions, 19:1296
 measurement of, CRIMS in, 11:400
- Deuterium exchange and cluster chemical ionization mass spectrometry, in supersonic molecular beams, 7:550
- Deuterium exchange behavior, of ERK2, pepsin mapping of, 10:685
- Deuterium/hydrogen exchange
 gas-phase
 as a molecular probe for the interaction of methanol and protonated peptides, 5:526
 in electrospray ionization mass spectrometry as a practical tool for structure elucidation, 5:434
 methyl guanine isomer distinction by, using a Fourier transform mass spectrometer, 4:296
- Deuterium/hydrogen exchange between $[m-1]^+$ ions of chlorinated benzenes and D_2O or ND_3 , mechanistic study of, 5:282
- Deuterium/hydrogen exchange reactions, of emolate ions with CH_3OD and C_2H_5OD , reactive collisions in quadrupole cells, 3:853
- Deuterium labeling, regiospecificity in, MS of, 15:1366
- Deuteron transfer, propene loss from propylpyridines initiated by, 14:1166
- Development and application of packed-volume supercritical fluid chromatography/pneumatically assisted electrospray MS, 9:498
- Development and evaluation of on-line nanoliter flow analysis of protein digests by pneumatic-splitter electrospray chromatography mass spectrometry, 8:703
- Development and ion characterization of an ESI ion trap/linear time-of-flight mass spectrometer, 8:1085
- Development and validation of a HPLC MS/MS assay for atorvastatin, ortho-hydroxy atorvastatin, and para-hydroxy atorvastatin in human, dog, and rat plasma, 10:55
- Development of a high-resolution mass spectrometer, 2:447
- Development of a three-dimensional topographic map display for CE/MS with an ion trap/reflection TOFMS detector, applications to tryptic digests of isoforms of myelin basic protein, 9:701
- Development of an isotope dilution LC-MS/MS detection method for DNA adducts of selected aromatic amines, 10:57
- Development of fast atom bombardment mass spectral methods for identification of carcinogen-nucleoside adducts, 3:360
- Development of generic LC-MS using experimental design, 13:2
- Development of LC-IMS-CID-TOFMS techniques, analysis of a 256-component tetrapeptide combinatorial library, 14:1424
- Dexamethasone, chemically oxidized, in human plasma, electron capture negative ionization in, comparison of tandem and conventional mass spectrometry using, 1:341
- Dexamethasone transformation products, on light-activated TiO_2 surface, ion trap MS/MS of, 12:1286
- DHA. *See* Docosahexaenoic acid
- Di-o-benzoyl-tartaric acid dibutyl ester, metal and proton complexes of, in gas phase, chiral recognition characteristics of, 14:571
- Diacylene radical cation, ethylene reactions with, 15:114
- Diacyl glycerophosphatidic acids
 fragmentation processes in charge-driven, 11:797, 11:892
 fragmentation processes in charge-remote, 11:892
- Diacyl peroxides, CIS-MS/MS analysis of, 12:449
- Diacylglycerophosphocholine
 divalent metal ions complexed with, ECD, 19:978
 saturated, ESI MS, artifacts in spectra of, 17:384
- Diacylglycerolphosphocholine molecular species, by FAB-CAD-MS/MS, a general method not sensitive to the nature of the fatty acyl groups, 3:71
- Diagnosis of clinical disease, MS in, 10:561
- Diammonium citrate, effect on CHCA matrix for detection of phosphorylated peptides, MALDI-TOF MS, 18:1925
- Diamonds, surface-induced dissociation and chemical reactions on stainless steel, carbon and, 20:927
- 1,2-Diarylcyclopropanes, Cu(II) oxidation of, 9:966
- Diastereomeric benzo[ghi]fluoranthene tetrols, product ion studies of, by MALDI-TOFMS and PSD, 12:376
- Diastereomeric cobalt complexes, mass spectrometric studies of, chiral recognition in the gas phase, 5:452
- Diastereomeric cobalt-glucosyl-glucose disaccharide complexes, tandem mass spectrometry of, differentiation of stereochemistry of glycosidic bond configuration, 7:953
- Diastereomeric *N*-acetylhexosamine, quantification of, MS3 spectra for, 11:1086
- Diazinon photocatalytic degradation products, 14:803
- Diazoamino compounds, aromatic, thermolysis of, 9:234
- Diazoxon, fragmentation and reactions of, 16:515
- Dibenzothiophene subjected to negative chemical ionization with oxygen, oxidation reactions of, 4:949
- Dicarboxylic acid
 ESI-MS of, 10:557
 polyammonium macrotricyclic ligand complexes with, ESI-MS of, quantum calculations and, 12:304
- Dicationic reagents, in anion detection using positive ion mode ESI-MS, 19:261
- Dications
 dichlorocarbene, single-electron capture by, 7:266
 dichlorocarbene dications studied by tandem mass spectrometry, reactivity of collisionally activated, 6:1019
 molecular, gas-phase, formed from neutral organics in solution, via the controlled-current electrolytic process inherent to electrospray, 7:157
 oligoodeoxynucleotide, electron capture dissociation and infrared multiphoton dissociation of, 14:23
 in stabilization of gas phase ternary β -cyclodextrin inclusion complexes observed by ES-MS, 14:449
 tungsten hexacarbonyl, ground and excited states of, double-ionization energies to, 2:108
 unimolecular gas-phase reactivity, complexes with cucurbit[8]uril, 18:1863
- 3,3'-Dichlorobenzidine in environmental samples, FT-ICR MS of, 10:1152
- Dichlorocarbene dication
 single-electron capture by, mechanisms of, 7:266
 studied by tandem mass spectrometry, reactivity of collisionally activated, 6:1019
- Dichloroethylene cation isomers, long-lived excited electronic states of, charge exchange ionization of, 15:1266
- Dichloro(ethylenediamine)palladium(II), complex formation in aqueous solutions, ESMS, 18:769
- Dichloro(ethylenediamine)platinum(II), hydrolysis investigated by electrospray MS and DFT, 20:1015
- 3,5-Didehydrophenyl cation, FTICR-MS study of, 12:258
- Dielectric barrier discharge ion source, for ambient mass spectrometry, 18:1859
- Diethylene glycol monobutyl ether, as contaminant in ion trap mass spectrometer, 14:1247
- Diexo- and diendo-fused isomers, of thiouracil and thiazinopyrimidine derivatives, MS of, 12:1011
- Differential electrochemical mass spectrometry, 15:1693
- Differential electrospray emitter, potential build-up and gradient mapping within, 15:1737
- Differential mass spectrometry, analysis of complex peptide mixtures using FTMS and, 18:226

- Differential mobility analysis, parallel, nanoES GEMMA and, nanobioparticle analysis using, 19:1062
- Differential mobility spectrometry, oligosaccharides analysis using nanoESI-DMS-MS, 18:502
- Differential overlap bond orders, modified neglect of, and ab initio and modified neglect of differential overlap energy partitioning, effect of protonation site on bond strengths in simple peptides, 5:704
- Differential overlap energy partitioning, modified neglect of, and ab initio and modified neglect of differential overlap bond orders, effect of protonation site on bond strengths in simple peptides, 5:704
- Differentiation and quantitation of isomeric dipeptides by low energy dissociation of copper(II)-bound complexes, 12:490
- Differentiation of flavonoid glycoside isomers using metal complexation and ESI-MS, 14:1437
- Differentiation of isomeric C8- and N²-deoxyguanosine adducts of 2-acetylaminofluorene by fast-atom bombardment and tandem mass spectrometry, 5:58
- Differentiation of isomeric C8-substituted alkyraniline adducts of guanine by ESI and quadrupole ion trap MS/MS, 14:1488
- Differentiation of isomeric photomodified oligodeoxynucleotides by fragmentation of ions produced by MALDI and ESI, 10:329
- Differentiation of *O*-acetyl and *O*-carbamoyl esters of *N*-acetylglucosamine by decomposition of their oxonium ions, application to the structure of the nonreducing terminal residue of Nod factors, 11:301
- Differentiation of positional isomers of nitro meso-tetraphenylporphyrins by MS/MS, 12:381
- Differentiation of stereochemistry of glycosidic bond configuration, tandem mass spectrometry of diastereomeric cobalt-glucosyl-glucose disaccharide complexes, 7:953
- Differentiation of stereoisomeric steroids by reactions with phosphonium ions, 13:362
- Diffusion, ionic, in SIFT-MS, 12:863
- Diffusion measurements by ESI-MS for studying solution-phase noncovalent interactions, 14:430
- Difluorobenzene, ionized *o*-,*m*-,*p*-, a TPEPICO study, 20:20
- Digestion, of surface-bound proteins, 14:203
- Digital ion trap mass spectrometry, manipulation of rectangular waveform in, 19:620
- Diglycine, N-protonated surface-induced dissociation, collision energy effect on, 14:1402
- Diglycosyl flavonoids, isomeric, SORI CID study of, ESI FTICR MS, 18:2127
- Digoxin, fast atom bombardment of, ions formed by, high resolution mass spectrometry and tandem mass spectrometry analysis of, mechanistic considerations of the protonation and fragmentation of highly functionalized molecules in fast atom bombardment, 1:455
- 3,4-Dihydro-2,4-dioxo-2H-1,3,5-oxadiazinium ions, synthesis in gas phase, 16:1602
- Dihydrodiol-epoxides, polycyclic aromatic hydrocarbon (PAH), deoxynucleoside adducts, low energy tandem mass spectrometry of, 6:248
- 1,2-Dihydropyridine derivatives, electron-induced MS of, 8:1255
- Dihydroxybenzoic acid
gas phase acidities/basicities of, 11:544
noncovalent association with cyclic and linear oligosaccharides, MALDI-TOFMS and X-ray crystallographic study of, 11:228
in UV-MALDI, of oligonucleotides, 8:218
- 2,5-Dihydroxybenzoic acid, tripeptide VPL in cluster with, matrix to analyte proton transfer in, 12:726
- 2,5-Dihydroxybenzoic acid monocrystals, and irradiation effects on MALDI, ablation, ion production, and surface modifications, 893
- 2,5-Dihydroxybenzoyl hydrazine derivative, CNTs in trace peptide analysis, MALDI, 17:1023
- Diketone/metal ion complexes, ESI-MS of, 10:402
- Dimer formation
electrospray ionization mass spectrometry of oligonucleotides, specific and nonspecific, 6:159
in α -synuclein, pH dependence of, monomeric structures and, 15:1435
- Dimeric G-quadruplex inhibitors, of HIV-1 integrase, investigation by ESI MS, 19:550
- Dimers, proton-bound, reactions of, 5:695
- Dimethanofluorenes, highly chlorinated, in technical chlordane and in human adipose tissue, 1:99
- 2,3-Dimethyl-2-butanol, photoionization of, 15:989
- Dimethyl and monomethyl ethers, ethylene glycol, functional group-selective ion-molecule reactions of, 4:97
- Dimethyl diselenide, distinguishing conventional and distonic radical cations by using, 7:1245
- Dimethyl ether
ion/molecule reaction with, selective detection of the tolyl cation among other [C₇H₇]⁺ isomers by, 2:270
selectivity of, methylene substitution reactions in a quadrupole ion trap, 3:39
- Dimethyl ether ions
competitive reactions of in quadrupole ion trap, steric and substituent effects in phenols evaluated by, 6:627
gas-phase reactions of amino alcohols with, site-selective, an investigation of, 4:230
reactivities of, 8:610
selective ion-molecule reactions of, with lactams, 4:353
- Dimethyl methylphosphonate, detection by ion/molecule reactions, 182
- Dimethylation, N-terminal, peptides with imine/methylol adduction and, MS/MS, 17:859
- Dimethyldioctadecylammonium ions, contamination source in ESI MS, 19:666
- Dimethylene cyclobutene, isomerization of, 8:594
- Dimethylene systems, ion-molecule reactions in, 2:220
- Dimethylphenols, determination of gas-phase acidities, 19:1595
- Dimethylpyrroles, dissociation of, reactions of isomeric parent ions in, 7:930
- 2-(4,6-Dimethoxy-pyrimidin-2-ylsulfanyl)-N-phenylbenzamide, gas-phase Smiles rearrangement reactions, ESI-MS, 18:1813
- Dinitrobenzene, nucleophilic substitution reactions of halonitrobenzenes and, with carbanions, gas phase, 18:1351
- Dinitrofluorobenzene, product ion assignment with, 10:448
- Dinitrophenols, 2:2-alkyl-4,6-mass spectra of novel [M-35]⁺ fragment ions in, sequential ortho effects, 2:69
- Dinitropyrene, LC-MS/MS analysis of, 15:1853
- Dinucleotide analogs in the gas phase, metal chelation of, by fast-atom bombardment mass spectrometry, 7:42
- Dinucleotide phosphodiester, molecular dynamics and MS/MS of, 8:1257
- Dinucleotides, negative ions, gas phase H/D exchange, 18:1827
- Diode array detection. *See also* High performance liquid chromatography-diode array detection
- Diode array detection (DAD), of *Ophiopogon japonicus* homoisoflavonoids, 16:234
- Diolepoxides adducted peptides, polycyclic aromatic hydrocarbon, characteristics of high energy collision-induced dissociation tandem mass spectra of, 5:649
- Diols
bimolecular and dissociation reactions of, effects of functional group interactions on, 3:535
gas-phase acidities of, 4:723

- Diophantine mass spectrometric structure analysis, 10:231
- DIOS MS. *See* Desorption ionization on silicon mass spectrometry
- 1,4-Dioxane, phosphonium ion ketalization by, 14:182
- Dioxins, monobromopolychlorodibenzo-*p*, trace, in environmental samples, application of gas chromatography high-resolution mass spectrometry to, 3:248
- Dipeptide dications, charge partitioning between *c* and *z* fragments after ECD, 20:1881
- Dipeptides
 arginine-containing protonated, fragmentation reactions, CID, 18:1959
 containing polar and nonpolar side groups, by Curie-point pyrolysis tandem mass spectrometry, 3:750
 protonated, condensation reactions in, 15:1025
 selected, gas-phase basicities of histidine and lysine and their, 7:1203
 of serine and glycine, gas-phase basicities of, 5:718
 study of $b_1 + H_2O$ and b_1 -ions in product ion spectra, 18:1414
- Dipolar resonance excitation, enhancement of, studied by linear ion trap, 18:578
- Dipole-ion complex formation from deprotonated phenol fatty acid esters evidenced by using gas-phase labeling combined with tandem mass spectrometry, 4:343
- Dipyridocyanine dyes, and meta-nitrobenzyl alcohol, beam-induced reaction between, in liquid-secondary-ion mass spectrometry, 5:1017
- Diquat
 capillary electrophoresis-mass spectra of, 7:981
 in water, LC-(ESI)MS of, 9:830
- Diquaternary ammonium salts, observed charge states of, and ion pairing, in electrospray ionization mass spectrometry, effects of solvent and counterion on, 7:1050
- Direct analysis in real time
 implementation on fieldable MS, 19:1419
 ionization mechanisms related to negative ion APPI, APCI and, 20:1518
 TOF MS with, Chinese and American drywall sulfur materials analyzed by, 20:2082
- Direct analysis of carbon isotope variability in albumins by liquid flow-injection isotope ratio mass spectrometry, 7:605
- Direct analysis of laser capture microdissected cells by MALDI-MS, 13:1292
- Direct association equilibrium in the FT-ICR ion trap, hydration equilibrium of protonated 18-crown-6, 10:862
- Direct chemical analysis of UV laser ablation products of organic polymers by using selective ion monitoring mode in gas chromatography/mass spectrometry, 5:106
- Direct comparison of IR and UV wavelength MALDI-MS of proteins, 9:1
- Direct current pulse excitation, and multiparticle simulation of ion motion in ion trap mass spectrometer, 6:57
- Direct current quadrupole lens system, inductively coupled plasma-time-of-flight mass spectrometer for elemental analysis and, 6:400
- Direct detection and quantitation of He@C₆₀ by ultrahigh resolution FTICR-MS, 13:1349
- Direct determination of polycyclic aromatic hydrocarbons in solid matrices using laser desorption/laser photoionization ion trap MS, 14:462
- Direct dynamics study of N-protonated diglycine surface-induced dissociation: influence of collision energy, 14:1402
- Direct generation of ion beam images with a two-dimensional charge injection device, 6:140
- Direct matrix-assisted laser desorption ionization mass spectrometric analysis of proteins immobilized on nylon-based membranes, 5:230
- Direct monitoring of protein-chemical reactions utilizing nano-electrospray MS, 10:112
- Direct observation of trapping motion in elongated Fourier-transform mass spectrometry trapped ion cells, 1:351
- Direct sampling ion trap mass spectrometry (DSITMS), in U.S. space program, 12:656
- Direct *versus* matrix-assisted laser desorption, laser microprobe mass spectrometry of quaternary phosphonium salts, 4:798
- Disaccharide, glucose, -glucosyl complexes-diautomeric cobalt, tandem mass spectrometry of, differentiation of stereochemistry of glycosidic bond configuration, 7:953
- Disaccharides
 chromophore-labeled, CE-MS/MS of, 11:292
 differentiating anomers, linkage, and position isomers of, FAIMS for, 14:265
 glycosaminoglycan-derived, sulfate and linkage position in, negative ion fast-atom bombardment tandem mass spectrometry to determine, 3:797
 heparin-derived, wavelength-tunable UVPD in LIT, 20:1645
 peralkylated, low energy CID of, 14:63
 sodiated isobaric, separation using ESI-APIM-TOF MS, 16:660
 sulfated, in chondroitin sulfate, ESI-MS/MS of, 11:916
- Discovery of long-lived excited electronic states of vinylchloride, vinylbromide, vinyliodide, and acrylonitrile cations, 14:110
- Discriminator interface, for nanoflow ESI-MS, 14:1236
- Diselenide, dimethyl, distinguishing conventional and distonic radical cations by using, 7:1245
- Dissociation
 channeling of energy into, photoionization study of the ion-neutral complexes [CH₃CH⁺IIB CH₂CH₃] and [CH₃CH₂CH⁺CH₃IICH₃] in the gas phase, 7:73
 collision-activated, in RF-only quadrupoles, instrument-independent measurements of, a kinetics-based protocol for, precision and accuracy in tandem mass spectrometry measurements, 1:272
 collision-induced. *See* Collision-induced dissociation.
 effects of target gas in, using a double quadrupole mass spectrometer and radiofrequency glow discharge, 5:845
 evidence for charge separation and "ball-and-chain" propagation in the addition of 1-butene to C₂+60, 7:261
 high-energy, in tandem mass spectrometry, low-mass ions produced from peptides by, 4:882
 laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry, improved efficiency of, selective parent ion axialization for, 5:900
 liquid chromatography/mass spectrometry with, the determination of glycopeptides by, 3:804
 low-energy, characterization of a high-pressure quadrupole collision cell for, 5:1042
 of peptide ions, efficiency of, in Fourier transform mass spectrometry, 1:288
 of prostaglandins and thromboxanes, some examples of charge remote fragmentation, 1:325
 electron-induced, energy deposition during, 4:145
 of heparin disaccharides, ion trap MS/MS of, 15:1274
 high-energy collision-induced
 of ceramide ions from permethylated glycosphingolipids, 5:558
 structural elucidation of O-linked glycopeptides by, 7:319
 using artificial neural networks, peptide sequence determination from, 6:947
 metastable, kinetic energy orifice as probe of, in FTICR-MS, 11:1118
 mixture analysis by neutralization-reionization and, characterization of C₃H₆O⁺ on from 2-methoxyethanol and, 6:1030

- multipole storage assisted, 11:1
 peptide ion, energetics of, by laser desorption chemical ionization Fourier transform mass spectrometry, 6:1069
 photon-induced, with a four-sector tandem mass spectrometry, 1:107
 rate of, the effect of ion size on, RRKM calculations on model large polypeptide ions, 4:11
 on a rhenium oxide surface, high-efficiency surface-induced, 4:869
 surface-induced
 in ICR-MS/MS, 11:1107
 from a liquid surface, 4:769
 in tandem quadrupole mass spectrometers, a comparison of three designs, 3:27
 of tetrameric ions of noncovalent streptavidin complexes formed by electrospray ionization, 6:459
 Dissociation and gas-phase methylation of acids and esters, effects of functional group interactions on, 7:565
 Dissociation and neutralization-reionization, stabilities and isomerization proclivities studies, distonic ion $\text{CH}_2\text{CH}_2\text{CH}^+\text{OH}$, keto ion $\text{CH}_3\text{CH}_2\text{CH}_2\text{O}^+$ radical cations, 7:573
 Dissociation behavior, of gas-phase protonated and methylated lactones, characterizations of the, 3:47
 Dissociation characteristics, of $[\text{M} + \text{X}]^+$ ions ($\text{X}=\text{H}, \text{Li}, \text{Na}, \text{K}$) from linear and cyclic polyglycols, 5:1081
 Dissociation constants, determination for protein-ligand interactions, comparison of ESI, nanoESI and ESSI, 19:332
 Dissociation device, in-line surface-induced, in a four-sector mass spectrometer, design and performance of, 6:257
 Dissociation dynamics, of phenetole cations by photoelectron photoion coincidence, 2:464
 Dissociation energetics, of peptide ions, 9:175
 Dissociation energetics and mechanisms of leucine enkephalin $(\text{M} + \text{H})^+$ and $(2\text{M} + \text{X})^+$ ions ($\text{X} = \text{H}, \text{Li}, \text{Na}, \text{K},$ and Rb) measured by blackbody infrared radiative dissociation, 8:771
 Dissociation energy
 bond, of germane, Fourier transform-ion cyclotron resonance study of gas-phase acidities of germane and methylgermane, 4:54
 deoxyribose nucleotide dimer anions measured using blackbody infrared radiative dissociation, 10:1095
 relative, measurements using ion trap collisional activation, 5:250
 threshold, of protonated amine/polyether complexes in quadrupole ion trap, 14:383
 Dissociation fragmentation, APCI low energy collision-induced, of protonated orthosilicates, McLafferty or ion-neutral complex rearrangements, 5:544
 Dissociation kinetics, of a transient intermediate, 10:119
 Dissociation mass spectrometry, collision-induced
 high-energy, of synthetic mannose-6-phosphate oligosaccharides, 7:182
 of nonionic surfactants following direct supercritical fluid injection, 3:150
 Dissociation of different conformations of ubiquitin ions, 13:719
 Dissociation of doubly charged transition metal/polyether/pyridyl ligand complexes in a quadrupole ion trap MS, 10:126
 Dissociation of heme-globin complexes by blackbody infrared radiative dissociation: molecular specificity in the gas phase?, 8:519
 Dissociation of individual isotopic peaks, predicting isotopic distributions of product ions in MSn, 14:310
 Dissociation of multiply charged polypeptides, high-energy collision-induced, produced by electrospray, 4:652
 Dissociation of peptide ions, collision-induced, the effect of charge state and the localization of charge on, 5:966
 Dissociation of polyether-transition metal ion dimer complexes in a quadrupole ion trap, 8:620
 Dissociation of proton-bound complexes and proton affinity of benzamides, 5:826
 Dissociation of uracil and its derivatives, collision-induced, 5:339
 Dissociation pathways of alkali-cationized peptides, opportunities for C-terminal peptide sequencing, 12:497
 Dissociation reactions of diols, effects of functional group interactions on, 3:535
 Dissociation spectra, delayed, of survivor ions from high-energy collisional activation, 4:845
 Dissociation spectra of peptides, uninterpreted high-energy collision-induced, search of, with sequence databases, 7:1089
 Dissociation studies, collision-induced, high and low energy, of cluster ions on a VG analytical ZAB-2FQ, a compact high pressure ion source for, 3:33
 Dissociation tandem mass spectra, high energy collision-induced, high-energy, of polycyclic aromatic hydrocarbon diolepoxide adducted peptides, characteristics of, 5:649
 Dissociation tandem mass spectrometry
 collision induced, high- and low-energy, use of, in the identification of an unusual amino acid in a semisynthetic polypeptide, 7:1034
 high- and low-energy, in the analysis of glycoalkaloids and their aglycons, comparison of, 7:173
 Dissociative behavior, of transition metal oxide cluster ions in the gas phase, characterization of, 2:212
 Distibines, tetramethyl- and tetraethyl-, low temperature thermal lysis behavior, 19:1336
 Distinction among isomeric unsaturated fatty acids as lithiated adducts by ESI-MS using low energy CAD on a triple stage quadrupole instrument, 10:600
 Distinguishing conventional and distonic radical cations by using dimethyl diselenide, 7:1245
 Distonic and conventional radical cations, distinguishing, by using dimethyl diselenide, 7:1245
 Distonic ion $\text{CH}_2\text{CH}_2\text{CH}^+\text{OH}$, keto ion $\text{CH}_3\text{CH}_2\text{CH}_2\text{O}^+$, enol ion $\text{CH}_3\text{CH}=\text{CHOH}^+\text{II}$, and related $\text{C}_3\text{H}_6\text{O}+\text{II}$ radical cations, stabilities and isomerization proclivities studied by dissociation and neutralization-reionization, 7:573
 Distonic ions
 of the "ate" class, 10:896
 dual reactivity of, 11:697
 the role of, in the formation of CH_3NH_3^+ and $(\text{CH}_3)_2\text{NH}_2^+$ from the molecular ions of octopamine and synephrine, 1:104
 Distonic radical cation $+\text{CH}_2\text{-O-CH}_2$, ionized methylene transfer from to heterocyclic compounds, pentaquadrupole mass spectrometric study, 6:554
 Distonoid ions, characterization, 17:1014
 Distribution of ion acceptance, spatially resolved, in APLI MS, 19:400
 Distribution shifting, charge state, of protein ions observed in matrix-assisted laser desorption ionization mass spectrometry, 6:1183
 N,N'-Disubstituted imidazolium ionic liquids, ESMS fragmentation study, 17:85
 Disulfide, aliphatic, bond-selective photodissociation of, 6:872
 Disulfide bonds
 FMN-binding complex I subunit, MS profile of intramolecular disulfides, 19:1876
 gas-phase scrambling during MALDI MS, 20:1603
 insulin, mapping with Route 66 method, 20:157
 peptide, cleavage via salt bridge mechanisms, low energy CID, 18:1109

- proteins containing, MALDI MS, 20:2284
 ranid skin secreted peptides, oxidation *versus* carboxamidomethylation of, MALDI-MS, 19:479
 sulfhydryl bonds and, counting using mercurial ions, peptides and proteins, 19:1108
- Disulfide linkages and disulfide bond scrambling, in recombinant human macrophage colony stimulating factor by fast-atom bombardment mass spectrometry of enzymatic digests, 6:638
- Disulfide-linked peptides, SO₂-•electron transfer ion/ion reactions with, 16:1020
- Disulfur monoxide, water charged complexes with, isotope exchanges, FT-ICR, CI-CAD and TQ, 18:1664
- 3,3'-Dithiobis(sulfosuccinimidyl propionate), unexpected reactivity with peptides, 15:736
- Diuretics, fragmentation of, in negative ESI-MS, 14:658
- DMA. *See* Differential mobility analysis
- DMMP. *See* Dimethyl methylphosphonate
- DMPO adducts, of oxygen-centered radicals, separation and identification of, 14:862
- DMS. *See* Differential mobility spectrometry
- DNA
 adduct between 4-aza-3-ene-1,6-diyne and, ESI-MS, 17:1342
 automated de novo sequencing of, by LC-MS/MS, 15:32
 bimolecular quadruplexes and transitions to higher order molecular structures, ESI-FTICR-MS, 18:1467
 chromophore-bound, electron photodetachment dissociation, 18:1990
 double-stranded
 accurate base composition of, by mass spectrometry, 7:1266
 oxidative damage to, gas chromatographic-mass spectrometric method for the assessment of, by quantitation of thymine glycol residues, 4:336
 stabilizing in ultraviolet MALDI-MS, 13:1129
 duplex, CID of, multistep mechanism in, 13:91
 duplex and triplex, ligand binding mode assessed by ESI-MS/MS and molecular modeling, 18:1052
 evaluation of flavonoid-DNA complexes ESI-MS, 19:914
 fragmentation of isomeric intrastrand cross-link lesions, ion-trap MS, 20:611
 G-quadruplex, gas-phase stability using ESI-MS and MDS, 18:1760
 genomic, analysis by PCR-LC-ESI-MS, 17:124
 megadalton, mass determination of, using charge detection mass spectrometry, 9:305
 metal-mediated binding of benzoxazole ligands, evaluation by ESI, 19:209
 MS/MS sequencing of, 15:510
 mutant, protein complexes binding affinities with, laser spray MS, 17:611
 noncovalent complexes
 with basic polypeptides or polyamines, MALDI-TOF MS, 18:1977
 with polybasic compounds, studied by nanoES-MS, 18:346
 noncovalent interaction with peptide, MALDI-TOF detection of, 15:28
 perylene diimide and benzannulated perylene diimide ligand binding to, ESI-MS, 17:593
 polyamide binding to, evaluation by ESI-MS, 17:1742
 purine-rich triple-helix, bivalent cations and formation of, ESI-FT-MS, 20:1281
 secondary structure of, MALDI of, 10:521
 single stranded, complexes between guanidinium derivatives and, MALDI characterization of, 17:283
 stability in gas *versus* solution phases, 17:1383
 threading bis-intercalator binding to, screening by ESI-MS/MS, 18:311
- DNA-bezoxazole or DNA-benzimidazole complex, ESI-MS of, 15:11593
- DNA-binding domain, of vitamin D receptor, zinc-induced changes in, 9:8
- DNA-binding drugs, double-stranded oligonucleotide complexes with, 11:450
- DNA cross-links, determination by MS/MS, 15:1565
- DNA decay, under UV MALDI, 9:866
- DNA-drug complexes, positive and negative ion ESI-MS of, 15:1382
- DNA-endonuclease VIII repair complex, ESI-MS characterization of, 11:505
- DNA fragment ions, high resolution characterization of, 10:423
- DNA-PAH adducts, sensitive determination of, matrix design for matrix-assisted laser desorption ionization, 5:1021
- DNA-phosphate adducts, analysis with miniaturized LC-ESI-MS/MS and column switching, 15:593
- DNA-protein binding affinity, evaluation by ESI-MS, 16:116
- Do charge-remote fragmentations occur under MALDI post-source decompositions and MALDI collisionally activated decompositions?, 10:217
- D₂O or ND₃, hydrogen/deuterium exchange between [m-1]⁻ ions of chlorinated benzenes and, mechanistic study of, 5:282
- Docosahexaenoic acid, resolvin D1, protectin D1, and related derived products, analysis via ES-MS/MS, 18:128
- Dodecapeptides, triply protonated, gas phase reactivity and molecular modeling studies of, 9:716
- Dog plasma, nonpeptide antithrombotic in, quantitative determination of, by microbore high-performance liquid chromatography-tandem mass spectrometry utilizing pneumatically assisted electrospray ionization, 7:1238
- Dolichyl and polyprenyl phosphates, desorption chemical ionization tandem mass spectrometry of, 7:958
- Dominulin, A and B, in cuticle and venom of social paper wasp analyzed by MALDI-TOF, MALDI-TOF/TOF and ESI-ion trap, 17:376
- Dopant-assisted atmospheric pressure photoionization
 for charge exchange ionization, investigation by APPI, 19:955
 comparison of, for charge exchange ionization of PAHs, RP-LC-APPI-MS, 20:73
 effects of solvent flow, dopant flow, and lamp current on, 16:1275
 factors affecting primary ionization, 17:130
 of polyisobutylene derivatives, 20:2342
- Doped gelatin films as a model matrix for molecular secondary ion mass spectrometry studies of biological soft tissue, 4:419
- Double bonds
 alkenone, locations using GC-MS analysis, 17:710
 migration during EI MS, monounsaturated fatty acids, 20:1997
- Double-focusing mass spectrometer, hybrid, high pressure drift reaction cell to study thermal energy reactions of mass-selected ions, 1:197
- Double hydrogen migration, in endo tricyclo[5.2.1.0_{2,6}] decene, 13:1186
- Double-ionization energies
 of fluorinated benzene molecules, 4:513
 to ground and excited states of the tungsten hexacarbonyl dication, 2:108
- Double mutant hemoglobin Villeparisis β77(EF1) His → Tyr, β80 (EF₄) Asn → Ser, use of combined mass spectrometry methods for the characterization of a new variant of human hemoglobin, 7:163
- Double quadrupole mass spectrometer, effects of target gas in collision-induced dissociation using, 5:845
- Double-stranded DNA
 intact, detection of by MALDI, 6:972

- oxidative damage to, gas chromatographic-mass spectrometric method for the assessment of, by quantitation of thymine glycol residues, 4:336
- Doubly protonated angiotensin II formed by electrospray ionization, charge-separation reactions in tandem mass spectrometry of, the importance of, experimental considerations and structural implications, 7:30
- Drag model, for protein ions, an aerodynamic, 5:17
- DREAMS. *See* Dynamic range enhancement applied to mass spectrometry.
- Drift gas pressures, elevated, IMS using electrodynamic ion funnels and, 18:1176
- Drift gas selectivity, high resolution ESI-IMS of, 13:300
- Drift reaction cell, high pressure, to study thermal energy reactions of mass-selected ions, a hybrid double-focusing mass spectrometer, 1:197
- Drift-trajectories, spontaneous and deflected, in orthogonal acceleration time-of-flight mass spectrometry, 5:588
- Droplet surface partitioning coefficient, in ESI, 12:1002
- Droplets
 - charge assisted laser desorption/ionization mass spectrometry, 19:833
 - in electrospray
 - break-up of, physical/chemical separations in, 12:343
 - charge gradients in, ion response and, 12:206
 - protein fluorescence measurements within, 12:716
 - isolated, as ion source, 11:393
 - levitated, α -cyano-4-hydroxycinnamic acid cocrystallization with peptides in, 16:254
 - single on-line, deposition for MALDI-MS, 15:1471
- Drug analysis, SSI vs APCI for, 13:204
- Drug discovery recovery libraries, sub-ppm mass accuracy on ESI-TOF-MS, 18:1612
- Drug metabolites
 - enhanced sample loading of, 8:15
 - structural elucidation of, tandem quadrupole orthogonal TOFMS for, 10:1305
- Drug residue analysis, veterinary, hormone residue analysis, UPLC-Orbitrap MS, 20:451
- Drugs
 - illicit, LC-MS/MS bioanalysis of, matrix effect in, 14:1290
 - in a single hair, fast analysis of, 9:1311
 - thermally-labile, in supersonic molecular beams, fast, very fast and ultra-fast gas chromatography-mass spectrometry of, 7:737
- Drywall, Chinese and American, sulfur materials analyzed by DART with TOF MS, 20:2082
- DSI-MS. *See* Desorption electrospray ionization-mass spectrometry
- DTSSP. *See* 3,3'-Dithiobis(sulfosuccinimidyl propionate)
- Dual electrospray ionization source combined with hexapole accumulation to achieve high mass accuracy of biopolymers in FTICR-MS, 11:876
- Dual purpose collision cell, interface for a four-sector mass spectrometer with, 6:175
- Dueling ESI, instrumentation to study ion/ion reactions of electrospray-generated cations and anions, 13:614
- Duplex DNA and minor groove-binding molecules, noncovalent complexes formed between, by electrospray ionization-mass spectrometry, characterization of, 6:1154
- Duty cycle enhancement, axially-resonant excitation, linear ion trap, TOF-MS, 17:1669
- Dyes
 - azo, fast-atom bombardment mass spectra of, effect of tautomerization on, 5:407
 - ballpoint, photofading on paper analyzed by LDI and MALDI MS, 17:297
 - dipyridocyanine, and meta-nitrobenzyl alcohol, beam-induced reaction between, in liquid-secondary-ion mass spectrometry, 5:1017
 - pigments and, MALDI TOF MS identification, 18:2001
- Dynamic combinatorial libraries, direct screening using FTICR MS and SORI-CAD, 17:1074
- Dynamic ion cooling technique for FTICR-MS, 12:1169
- Dynamic range enhancement applied to mass spectrometry (DREAMS), increased proteome coverage with, in peptide abundance measurements, 13:954
- Dynamic range of a transient recorder, increasing the, by using two analog-to-digital converters, 7:107
- Dynamic reaction cell for inductively coupled plasma mass spectrometry (ICP-DRC-MS), reduction of interferences produced within the cell, 10:1083
- Dynorphin A, biotransformation products of, matrix-assisted laser desorption mass spectrometry of, in vitro, 5:10
- E
- E2 reactions. *See* Bimolecular elimination reactions
- Earth metal cations, alkali/alkaline, lariat ether amide selectivity of, ESI-MS evaluation of, 14:1215
- ECD. *See* Electron capture dissociation
- ECD-MS. *See* Electron capture dissociation mass spectrometry
- ECD of peptides, optimization of experimental parameters for, in Fourier transform mass spectrometer, 13:1396
- Ecotin, thermal dissociation of, in gas phase, 13:1432
- EDD. *See* Electron detachment dissociation
- Editorials, 12:243, 12:479, 12:617, 13:1, 13:417, 13:587, 13:759, 13:1253, 13:1359, 15:951, 15:1379, 15:1691, 16:1, 16:983, 16:1205, 16:1727
 - Alexander Makarov, award for distinguished contribution in mass spectrometry, i (August)
 - focus on electrospray, 11:931
 - focus on H/D exchange of proteins in solution, 10:671
 - focus on proteomics in honor of Ruedi Aebersold, 2002 Biemann awardee, 14:683
 - focus on proteomics in honor the 2002 contribution in mass spectrometry award to W. J. Henzel, J. T. Stults, and C. Watanabe, 14:930
 - focus on the 2003 Sanibel conference—ion activation in chemistry and biochemistry: mechanisms, dynamics, and applications, 13:29
 - on the Friser legacy, 10:917
 - Growth of the Journal of the American Society for Mass Spectrometry, 3:267
 - on the importance of titles in references, 9:451
 - JASMS to 12 issues means faster turnaround, 4:1
 - on MALDI, 9:865
 - New Policies and Old Issues for JASMS, 17:1
 - noncovalent interactions, 14:417
 - on oligonucleotides, 9:660
 - on physical aspects of gas-phase ion chemistry, 10:797
 - Ron Hites Award, i (Nov)
 - special issues on electrospray ionization, 4:4:615, 4:523
 - on synthetic polymers, 9:267
- EESI-MS. *See* Extractive electrospray ionization mass spectrometry
- EF-hand motifs, calcium binding to, probed by ESI-MS, 12:1296
- Efb, complex with C3d domain, resolution by lysine acetylation and MS, 19:55
- Effect of charge state and the localization of charge on the collision-induced dissociation of peptide ions, 5:966
- Effect of different solution flow rates on analyte ion signals in nano-ESI MS, 14:492
- Effect of electrospray needle voltage on electroosmotic flow in capillary electrophoresis-mass spectrometry, 10:261

- Effect of frequency sweep direction on motion of excited ions in Fourier transform ion cyclotron resonance cells, 3:847
- Effect of high-performance liquid chromatography mobile phase components on sensitivity in negative atmospheric pressure chemical ionization liquid chromatography-mass spectrometry, 7:1059
- Effect of ion-molecule collisions in the vacuum chamber of an electrospray time-of-flight mass spectrometer on mass spectra of proteins, 7:342
- Effect of ion size on rate of dissociation, RRKM calculations on model large polypeptide ions, 4:11
- Effect of location of hydrogen abstraction on the fragmentation of diuretics in negative ESI-MS, 14:658
- Effect of protonation site on bond strengths in simple peptides, application of Ab initio and modified neglect of differential overlap bond orders and modified neglect of differential overlap energy partitioning, 5:704
- Effect of tautomerization on the fast-atom bombardment tandem mass spectra of azo dyes, 5:407
- Effect of the long-range potential on ion mobility measurements, 8:275
- Effects of air and helium on the performance of the ion trap detector when used for real-time monitoring of some atmospheric pollutants, 4:774
- Effects of cysteic acid groups on the gas-phase reactivity and dissociation of $[M + 4H]^{4+}$ ions from insulin chain B, 10:928
- Effects of electron and chemical ionization modes on the MS profiling of whole bacteria, 10:747
- Effects of functional group interactions on the bimolecular and dissociation reactions of diols, 3:535
- Effects of functional group interactions on the gas-phase methylation and dissociation of acids and esters, 7:565
- Effects of heavy-atom substituents on matrices used for matrix-assisted laser desorption-ionization mass spectrometry, 5:800
- Effects of heavy gases on the tandem mass spectra of peptide ions in the quadrupole ion trap, 7:1194
- Effects of ionization mode on charge-site-remote and related fragmentation reactions of long chain quaternary ammonium ions, 12:571
- Effects of liquid phase composition on salt cluster formation in positive ion mode ES-MS: implications for clustering mechanisms in electrospray, 12:1077
- Effects of modulation defects on Hadamard transform-TOFMS, 278
- Effects of molecular entanglements during electrospray of high molecular weight polymers, 9:299
- Effects of pH on the kinetic reaction mechanism of myoglobin unfolding studied by time-resolved ESI-MS, 11:312
- Effects of position of internal histidine residues on collision-induced fragmentation of triply protonated tryptic peptides, 12:1262
- Effects of pulsed introduction of buffer gas on ion storage and detection efficiencies in a quadrupole ion trap, 8:532
- Effects of salt concentration on an analyte response using ESI-MS, 10:625
- Effects of solvent and counterion on ion pairing and observed charge states of diquaternary ammonium salts in electrospray ionization mass spectrometry, 7:1050
- Effects of solvent on maximum charge state and charge state distribution of protein ions produced by ESI, 11:976
- Effects of target gas in collision-induced dissociation using a double quadrupole mass spectrometer and radiofrequency glow discharge, 5:845
- Efficiency, improved, of collision-induced dissociation in laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry, selective parent ion azialization for, 5:900
- Efficiency of collisionally-activated dissociation and 193-nm photodissociation of peptide ions in Fourier transform mass spectrometry, 1:288
- Efficient determination and evaluation of model cyclodextrin complex binding constants by ESI-MS, 14:1123
- Egyptian mummy resin, analysis of, by mass spectrometry, 3:582
- EI. *See* Electron ionization entries
- EI/Q/oa TOF mass spectrometer, miniaturized, 12:641
- Eicosanoids
characterization by ESI-FTICR, 19:569
cysteinyl, ESI and MS/MS of, 12:763
- EIMS. *See* Electron ionization mass spectrometry.
- Ejection chemical ionization, selected, and ion trap tandem mass spectrometry, application of nonresonance excitation to, 7:668
- Electric ion resonance cell, for FTMS, 10:1000
- Electrically floating sampling interface, inductively coupled plasma mass spectrometry with, 4:733
- Electrochemical cell, electrospray needle as, 6:860
- Electrochemical electrospray ionization, initiated by a corona discharge, 20:1988
- Electrochemical electrospray ionization mass spectrometry
adenosine triphosphate release from PP/ATP films, 18:919
hemoglobin subunit analysis by, 18:1279
- Electrochemical electrospray ionization tandem mass spectrometry, duplex and triplex DNA, ligand binding mode assessed by molecular modeling and, 18:1052
- Electrochemical electrospray ionization time-of-flight mass spectrometry, carbohydrate isomer resolution by ESI-APIMS-TOF-MS, 18:1163
- Electrochemical mass spectrometry (EC-MS)
differential, 15:1693
potential build-up and gradient mapping within emitter, 15:1737
ferrocenoyl piperazide using, 15:1727
mass tags by, for protein MS, 15:1767
metabolomic applications of, 15:1717
porous flow-through electrode emitter in, 15:1755
in protein oxidation and cleavage, 15:1707
signal intensities in, ground loop current effects on, 15:1748
- Electrochemical processes in wire-in-a-capillary bulk-loaded nano-electrospray emitter, 12:853
- Electrochemical properties of capillary electrophoresis-nano-electrospray mass spectrometry, 10:309
- Electrochemistry, m-Nitrobenzyl alcohol, in fast atom bombardment mass spectrometry, 3:113
- Electrochemistry/electrospray ionization mass spectrometry (EC-ESI-MS), on-line, caffeic acid oxidation product analysis by, 15:1228
- Electrochemistry liquid chromatography mass spectrometry
evaluation of tricationic ion-pairing reagents in, 20:529
pesticide metabolism simulation using, 20:138
- Electrode diameters, in cylindrical geometry FAIMS, 16:349
- Electrodeposited polythiophene films provide reproducible samples for laser desorption ion trap mass spectrometry studies, 8:1203
- Electrodes
configurations with arbitrary geometries, use in ion trajectory simulation program, 17:1216
planar resistive, miniaturized mass analyzers, novel ion traps using, 19:1435
- Electrodynamic ion funnel
for FTICR-MS, 11:19

- jet disrupter electrode in, automatic gain control in MS using, 16:244
- Electrodynamic ion funnels, IMS using elevated drift gas pressures and, 18:1176
- Electrohydrodynamic mass spectrometry, sensitivity of, effect of varying counterions on, 1:233
- Electrolytic process, controlled current, inherent to electrospray, observation of gas-phase molecular dications formed from neutral organics in solution via, 7:157
- Electron affinities of H₂O, cirrannulene and coronene, estimation of, by using the kinetic method, 7:619
- Electron affinity, of 1,3,5,7-cyclooctatetraene determined by kinetic method, 9:1141
- Electron anions, odd and even, decomposition of, 8:148
- Electron autodetachment dynamics, of trapped gas phase oligonucleotides, 1330
- Electron capture
secondary, in the liquid secondary ionization mass spectrometry beam-induced dehalogenation of organic compounds, evidence for a mechanism that involves, 7:1109
single, by the dichlorocarbene dication, mechanisms of, 7:266
survival of peptide cations after, 17:1675
- Electron capture dissociation (ECD)
activated-ion, optimization for proteins, 20:763
amino acid side chain losses in, 13:241
b ions in, 16:1932
B. anthracis petrobactin, 18:842
branched permethylated oligosaccharides analyzed by CID and, 19:138
and CAD-MS of doubly charge hyperbranched polyester-amides, 332
characterization of phosphate-containing metabolites by, 19:799
charge location directed, peptide dications, 17:1704
charge-tagged peptides, 18:2146
CID and, metal ion complexes with polyamidoamine dendrimers, 20:674
complexes of diacylglycerophosphocholine and divalent metal ions, 19:978
dipeptide dications, charge partitioning between *c* and *z* fragments after, 20:1881
DR, protein fragment intermediate lifetimes measured by, 17:1605
EDD, CID and, structural analysis of polyamidoamine dendrimers, 19:1312
effect of fixed charge modifications on, 19:1514
effect of phosphopeptides on, 19:1263
effect of radical trap moieties on spectra of substance P, 17:1428
effects of electron kinetic energy and ion-electron inelastic collisions, 19:772
efficiency in FTICR-MS, 16:1060
electron impact conditions, MS of O,O-dialkyl N,N-dialkyl phosphoramides under, 15:547
- Fourier transform ion cyclotron resonance tandem mass spectrometry, MARCKS peptide phosphorylation sites identified by, 18:2137
- FTICR-MS, peptide- and protein-linked antibiotic released by, 14:302, 14:682
- of ganglioside GM1, 16:752
- of gaseous multiply charged ions by FTICR-MS, 12:245
- general mechanism of, 16:208
- hemoglobin Hb askew variant analysis by, 20:1707
- hydrogen atom host-guest transfer induced by, 20:639
- hydrogen rearrangements, peptides, 18:113
- in identification and localization of fatty acid modification in ghrelin, 13:1443
- implementation progress in FT-ICR MS, 19:762
- induction of protoporphyrin IX ions by, 19:809
- and infrared multiphoton dissociation of oligodeoxynucleotides dications, 14:23
- IR activated DR, folding kinetics of peptide ions measured by, 19:780
- isoaspartyl residue quantification in deaminated proteins by, 18:48
- isoquinolines gas phase reaction to carboxylic acids CID and, 19:151
- long-lived ions from, radical migration via hydrogen abstraction, 17:576
- at low temperature, 15:1869
- of methionine oxidation in peptides, 14:605
- monitoring deamidation during protein and peptide sample processing, 19:855
- of *N*-acylated glucagon-like peptide-1 derivative, 16:548
- noncovalent complexes, electrostatic interaction site identified by ECD/ETD, 20:176
- nonergodicity in, hydrated ion nanocalorimetry, 18:1217
- O*-glycosylated and *O*-phosphorylated peptide sequencing by fixed charge derivitization and, 18:1405
- oligoribonucleotide fragmentation from EDD and, 17:1369
- peptide mono-oxidized thio-ether bond evaluation by CID and, 18:493
- of peptides
amino acid side chain loss in, 16:1523
b-type ions produced in, 18:2007
in Fourier transform MS, 13:1396
peptides metalated with alkaline-earth metal ions, 17:757
periodic sequence distribution in, amphipathic peptides, 20:1182
of poly(ethylene glycol), 12:565
of polypeptides, in three-dimensional quadrupole ion trap, 16:22
possible combination with CID, 19:753
product ion abundances correlated with amino acid polarities, 20:2273
role of ubiquitin conformation on, 17:1469
single d-amino acid distinguished in protein and tertiary structure probed by, 15:1087
spin-trap capped peptides, 18:432
study of oligonucleotides, 20:213
sulfopeptide fragmentation in ETD and, 18:1617
sustained off resonance irradiation (SORI) collision-induced dissociation (CID), of lactacin, 16:481, 16:1595
of trications, divalent metal ion interactions with peptides probed by, 17:1731
- Electron capture dissociation mass spectrometry
observation of *b*^{*}, *y* cleavages, polyamidoamine dendrimer ions, 17:536
side-chain fragmentation of alkylated cysteine in, 17:1271
- Electron capture dissociation-MS, of secreted proteins from *Mycobacterium tuberculosis*, 14:253
- Electron capture ionization of explosives with a microflow rate particle beam interface, 7:753
- Electron capture mass spectrometry, oxygen addition-induced dechlorination of PCBs and DDT metabolites in, regioselectivity of, 7:66
- Electron capture negative chemical ionization mass spectrometric fragmentation and calculated internal energies for polychlorinated diphenyls, correlation between, 5:44
- Electron capture negative ion chemical ionization mass spectrometry, amino acid/alcohol detection via, pentafluorobenzyl chloroformate derivatization for, enhancement of, 6:525

- Electron capture negative ion mass spectra, not reproducible, an ion source problem, 4:270
- Electron capture negative ionization, in the detection of chemically oxidized dexamethasone in human plasma, comparison of tandem and conventional mass spectrometry using, 1:341
- Electron capture negative ionization quadrupole mass spectrometry, pLVI-GC with, H-POPs analyzed by, 18:1375
- Electron capture-tandem mass spectrometry, sub-parts-per-billion determination of nitro-substituted polynuclear aromatic hydrocarbons in airborne particulate matter and soil by, 7:1255
- Electron detachment dissociation
 dermatan sulfate oligosaccharides, 19:294
 ECD, CID and, structural analysis of polyamidoamine dendrimers, 19:1312
 of GAG tetrasaccharides, 18:234
 of ganglioside GM1, 16:752
 glycosaminoglycans, effect of charged state and sodium cationization on IRMRD and, 19:790
 neutral and sialylated oligosaccharides, 18:2162
 oligoribonucleotide fragmentation from ECD and, 17:1369
 tandem mass spectrometry, peptide sequencing using, 19:1156
- Electron impact
 doubly and singly charged tungsten hexacarbonyl ions generated by, internal energy distributions deposited in, 1:16
 ionization mass spectrometry and intramolecular cyclization in 2-substituted pyrimidin-4(3H)-ones, 5:113
 mass spectrometry, of alkanes in supersonic molecular beams, 6:120
 organosulfur compound fragmentation and, 8:801
 3,3-sigmatropic rearrangement and cyclization in phenyl allylmethyl ethers, 6:195
- Electron-induced dissociation
 energy deposition during, 4:145
 glycosaminoglycans tetrasaccharides, 19:1449
- Electron-induced mass spectrometry, of 1,2-dihydropyridine derivatives, 8:1255
- Electron ionization gas chromatography/mass spectrometry, analysis of hydroxy fatty acids as pentafluorobenzyl ester, trimethylsilyl ether derivatives by, 6:40
- Electron ionization induced fragmentation of macrocyclic amines $C_nH_{2n-1}NH_2$, evidence for the rearrangement of aminocarbene radical cations and a comparison with long-chain esters, 1:238
- Electron ionization-induced loss of methyl radical from 1-cyclopropyl-4-substituted 1,2,3,6-tetrahydropyridine derivatives, 8:724
- Electron ionization-induced metastable decompositions of isomeric trimethylsilyl-methanol, $(CH_3)_3SiCH_2OH$, and methoxytrimethylsilane, $(CH_3)_3SiOCH_3$, 9:1029
- Electron-ionization mass spectra
 data base of, an enlarged, 2:432
 of exo-endo double bond isomers of polycyano push-pull pentadienes derived from cycloalkylidene malonic acids, 14:189
- Electron ionization mass spectral library search results, evaluating, 7:313
- Electron ionization mass spectrometry
 coupling to liquid chromatography, 13:265
 double bond migration during, monounsaturated fatty acids, 20:1997
 fragmentation of trimethylsilyl derivatives of hydroxy- and oxo-carboxylic acid, 19:66
 gas chromatography, relative sensitivities in, correlations of, molecular parameters, 1:397
- Electron ionization MS/MS spectra, calibration point for measured with multiquadrupole mass spectrometers, 5:576
- Electron-ionization-tandem mass spectrometry of glycosphingolipids, identification of compound-specific sequence ions in the collision-induced dissociation spectra of the ammonium ions of two isomeric hexaglycosylcer amides, 3:353
- Electron loss from multiply protonated lysozyme ions in high energy collisions with molecular oxygen, 12:889
- Electron paramagnetic resonance spectroscopy, glutathione conjugates, ES-MS/MS and, 20:2013
- Electron photodetachment dissociation, chromophore-bound DNA, 18:1990
- Electron promoted ion coherence, effect on electric fields at axial positions, 19:467
- Electron spin resonance, in separation and identification of oxygen-centered radical DMPO adducts, 14:862
- Electron states, excited, of polyatomic ions, detection by charge exchange ionization in collision cells, 12:1120
- Electron transfer
 gas phase dissociative, perturbation molecular orbital theory of, 8:510
 observed in mass spectra, 13:1209
 ternary copper (II) complexes with histidine, macrocyclic ligand effects on proton transfers and, 20:972
 vs proton transfer within radical-cation clusters of guanosine and deoxyguanosine with substituted naphthalenes and sinapinic acid, 12:176
- Electron transfer dissociation
 analysis of trypanosome flagellar proteome using ETD/ETD/CAD, 20:167
 aspartic and isoaspartic acids differentiated by, 17:15
 CID and, in linear ion traps, protonated peptides, 20:1672
 decoupling, peptide analysis by CRCID and, 20:567
 doubly sodiated glycerophosphocholine lipids, 18:1783
 fragmentation by MS based on, protein ubiquitination analyzed by CID or, 20:1652
 iTRAQ with, quantification of proteins and peptides, 19:1255
 methylated polypeptides, 20:1349
 noncovalent complexes, electrostatic interaction site identified by ECD/ETD, 20:176
 of peptide anions, 16:880
 post-acquisition spectral processing of peptides, 20:1435
 protein conformations probed using, in top-down HDX MS, 20:1514
 sequencing of peptides containing oxidized cysteine and methionine, 18:1499
 sulfopeptide fragmentation in electron ECD and, 18:1617
- Electron-transfer dissociation-Orbitrap mass spectrometry, nano-flow LS, methylated arginine quantified with, 20:507
- Electron-transfer reactions, chemical, gas-phase molecular dications formed from neutral organics in solution via, by using electrospray ionization mass spectrometry, 5:689
- Electronic archiving of data, improved, networking mass spectrometer data systems for, 4:742
- Electronic excitation under fast-atom bombardment conditions by using a liquid matrix, M^+ ion formation and, 6:114
- Electronic sputtering, produced by fission fragments, condensed CO and CO_2 , 17:1120
- Electronic states
 long-lived excited
 of dichloroethylene cation isomers, charge exchange ionization of, 15:1266
 of vinylchloride, vinylbromide, vinyliodide, and acrylonitrile cations, 14:110
 metastable, of Fe^+ , collisional relaxation of, 6:543
- Electrons, superelastic collisions of, with excited MN^+ , 1:182

- Electrophoresis
capillary, -electrospray mass spectra of herbicides paraquat and diquat, 7:981
capillary zone-continuous flow fast atom bombardment mass spectrometry
an integral probe for, 3:198
identification of deoxynucleoside-polyaromatic hydrocarbon adducts by, 3:757
capillary zone/electrospray ionization mass spectrometry, application of, to the analysis of macrolide antibiotics, 3:563
capillary zone/electrospray ionization parameters for the mass spectrometry and tandem mass spectrometry analysis of peptides, optimization of, 3:289
- Electrophoresis capillary, *See* Capillary electrophoresis
- Electrophoresis Fourier transform ion cyclotron resonance mass spectrometry, capillary, with sustained off-resonance irradiation for the characterization of protein and peptide mixtures, 5:894
- Electrosonic spray ionization
comparison of nanoESI, ESI and, for dissociation constant determination in protein-ligand interactions, 19:332
G. biloba flavonoids and terpenoids in plasma, efficiency of determination by, 19:445
macromolecular ionization by nano-ESI, MCS, 19:1098
Venturi-assisted, comparison of internal energy deposition with V-assisted AMUSE, 19:1320
- Electrosonic spray ionization mass spectrometry
deprotonation reactions on globular and denatured proteins at atmospheric pressure, 19:455
eicosanoid characterization by ESI-FTICR, 19:569
mimetic derivatives of primaquine, 19:1476
screening of enzyme inhibitors immobilized on silica microspheres, 19:865
- Electrosonic spray ionization-quadrupole time-of-flight tandem mass spectrometry, analysis of bimetallic oxovanadium complexes, 19:1247
- Electrospray. *See also* Liquid chromatography-electrospray tandem mass spectrometry.
as chemical reactor, of anionic α -adducts in gas phase, 15:927
clustering mechanisms in, 12:1077
controlled-current electrolytic process inherent to, observation of gas-phase molecular dications formed from neutral organics in solution via, 7:157
gramicidin S ions generated by, in studies of ion trapping in FAIMS/FAIMS coupled to TOFMS, 12:1320
high-energy collision-induced dissociation of multiply charged polypeptides produced by, 4:652
of high molecular weight polymers, molecular entanglements in, 9:299
micro, zeptomole/attomole per microliter sensitivity for peptides, 5:867
nano, combined with a quadrupole ion trap for the analysis of peptides and protein digests, 7:150
from optical fibers coiled with platinum wire, 11:464
in perrhenate system characterization, 12:1137
- Electrospray and atmospheric pressure chemical ionization
MS/MS behavior of eight anabolic steroid glucuronides, 11:722
- Electrospray and tandem mass spectrometry, identification of components in waste streams by, 4:604
- Electrospray emitter
analyte electrolysis in, 11:951
current measurements within ES capillary, 18:735
differential, potential build-up and gradient mapping within, 15:1737
porous flow-through, analyte oxidation in electrospray ion source using, 15:1755
- Electrospray instrumentation, Fourier-transform, for tandem high-resolution mass spectrometry of large molecules, 4:557
- Electrospray interface, novel, design and performance of, 3:18
- Electrospray ion source
analyte oxidation in, using porous flow-through emitter, 15:1755
high resolution, for magnetic sector mass spectrometer with heated capillary inlet, design and performance of, 6:1256
for magnetic sector mass spectrometers, 2:205
nebulizer-assisted, improvement by atmospheric pressure ion lens, 13:906
- Electrospray ionization (ESI). *See also* Ion trap
acetonitrile, pyridine, and methanol generated by, gas phase reactions of, 8:1124
angiotensin II ($M + 2H$)²⁺ ions formed by, location of charge in, 11:40
API source with, 9:1196
arginine-containing peptide ions formed by, modeling the maximum charge state of, 7:972
automated orthogonal control system for, 15:1201
automated spectral deconvolution of, 9:225
cations and anions in, ion/ion reactions of, 13:614
of ceramides, 13:558
lithiated, 13:680
characterization of the opioid peptide-containing protein and of bovine alpha-lactalbumin by, 5:377
charge separation in AMUSE ion source for MS, 20:1684
charge state and proton transfer reactivity of peptide and protein ions formed by, 6:1086
cholesteryl neoglycolipids analyzed using CID MS/MS and, 18:294
of copper-glycine solutions, 7:25
as current density profiler, 16:312
declustering and fragmentation of protein ions from, 8:1053
DESI, remote MS sampling using air ejector, 18:1844
differentiation of isomeric photomodified oligodeoxynucleotides by fragmentation of ions produced by, 10:329
disintegration of solvent droplets during, 20:496
DNA from, charge detection mass determination of, 9:305
doubly protonated angiotensin II formed by, charge-separation reactions in tandem mass spectrometry of, experimental considerations and structural implications, 7:30
droplet surface partitioning coefficient in, 12:1002
electrochemical, initiated by a corona discharge, 20:1988
endgroup determination of synthetic polymers by, 11:536
explosives, ambient vapors detected by SESI, 20:287
fluorescence and fragmentation of ions in ultrahigh vacuum, 20:1731
focus on, 11:931
gas-phase ions formed by, hydration of, 10:958
interface region of, CID of bradykinin ions in, 12:772
internal, low magnetic field, -Fourier transform ion cyclotron resonance mass spectrometer, a high performance, 7:915
ion pull needed to escape droplet, 8:1147
ionization mechanism, analyte concentrations, and ion cool time effects on peptide ion internal energies produced by, 16:743
ionization suppression in, 11:942
large multiply charged polypeptides and proteins produced by, collisional activation and collision-activated dissociation of, 1:53
leveling response factors in, 16:772
and low energy tandem mass spectrometry of polyhydroxy unsaturated fatty acids, 7:140
on a magnetic sector mass spectrometer, improved detection limits for, by using an array detector, 5:194

- maximum charge state and proton transfer reactivity of peptide and protein ions formed by, 6:1086
- mechanism features affecting protein complexes, 15:1424
- metal nitrate negative ions produced by, MS/MS of, 14:671
- multiple charging of polypropylene glycol by binary mixtures of cations in, 16:152
- multistage LIT-MS with, long-chain unsaturated fatty acids, double-bond position assigned by, 19:1678
- nano, time scale of internal energy relaxation, by AP-MALDI and, in QITMS, 20:1801
- a new approach for the study of gas-phase ion-ion reactions using, 3:695
- a new fused silica micro-electrospray tip, 20:751
- of nitrocatechol glucuronides, 10:537
- oligosaccharides analysis using nanoESI-DMS-MS, 18:502
- peptide aggregate formation during, 12:1020
- peptide amide hydrogen migration during, 12:410
- peptide metal(II)-ion complexes formed by, use of scans at a constant ratio of B/E for studying, 6:608
- of pharmaceutical compounds, metal complexation vs, 9:463
- pneumatic splitter, nanoliter flow analysis of protein digests using, 8:703
- pneumatically assisted, microbore high-performance liquid chromatography-tandem mass spectrometry utilizing, quantitative determination of a nonpeptide antithrombotic in dog plasma by, 7:1238
- of poly(ethylene glycol), 12:565
- poly(methylacrylic acid) oligomers produced by, MS/MS, 20:25
- precursor ion scan, protein-ligand complex detection by, MS/MS, 20:227
- of proteins, 8:483
- proton affinity values of MALDI matrices under, estimation with kinetic method, 15:431
- pulsed dual, for ion/ion reactions, 16:1750
- quadrupole ion trap MS with, sulfatide studies using, 15:536
- reduction of copper(II) complexes in, 9:1217
- scans at a constant ratio of B/E for studying decompositions of peptide metal(II)-ion complexes formed by, 6:608
- shrinking droplets and effect on chemical equilibria, 18:385
- of some novel alkylamine thiohydantoin amino acid derivatives, 6:1211
- source for TCID with guided ion beam mass spectrometer, 18:1124
- special issues on (editorial), 4:523, 4:615
- in study of melittin bound to phospholipid micelles, 12:1247
- supercharging from aqueous solutions during, 20:1933
- in tetracycline characterization, 13:543
- of transition metal oligosaccharides, 8:32
- unsaturated glycerophospholipids characterized by multiple stage LIT-MS with, 19:1700
- Electrospray ionization/capillary zone electrophoresis parameters for the mass spectrometry and tandem mass spectrometry analysis of peptides, optimization of, 3:289
- Electrospray ionization-collision induced dissociation, isoquinoline-3-carboxamide dissociation pathway analyzed by, 20:2034
- Electrospray ionization-collision-induced dissociation-tandem mass spectrometry (ESI-CID-MS-MS)
- in characterization of artemisinin-heme noncovalent complexes, 15:1181
- of glycerophosphoethanolamine plasmalogen phospholipids, 15:1499
- tetronasin fragmentation studies using, 15:325
- Electrospray ionization (ESI) atmospheric pressure ion mobility (APIM) time-of-flight (TOF) mass spectrometry, in separation of sodiated isobaric saccharides, 16:660
- Electrospray ionization (ESI) Fourier transform (FT) mass spectrometry, of silsequioxane low molecular weight components, 16:524
- Electrospray ionization (ESI) Fourier transform ion cyclotron resonance (FTICR) mass spectrometry
- in chemical cross-linking of protein complexes, 16:1921
- negative ion, glycosphingolipid mixtures screening and sequencing by, 16:571
- Electrospray ionization (ESI) ion trap mass spectrometry
- in crown ether complex self-assembly detection, 16:1162
- of diazoxon fragmentation and reactions, 16:515
- Electrospray ionization (ESI) ion trap tandem mass (MS/MS) spectrometry, chiral resolution of arginine by, 16:825
- Electrospray ionization (ESI) low energy CID-MS/MS
- electrospray ionization Fourier transform ion cyclotron resonance, high magnetic field, trapping and detection of ions generated in, 3:615
- electrospray ionization ion cyclotron resonance, high-resolution accurate mass measurements of biomolecules using, 4:566
- electrospray time-of-flight, vacuum chamber of, ion-molecule collisions in, on mass spectra of proteins, 7:342
- external ion source Fourier transform ion cyclotron resonance, matrix-assisted laser desorption ionization on, high resolution end group determination of low molecular weight polymers by, 7:449
- four sector
- in in-line surface-induced dissociation device, design and performance of, 6:257
- interface for, with dual purpose collision cell, high transmission at low to intermediate energies, 6:175
- fourier transform
- low magnetic field internal electrospray ionization, a high performance, 7:915
- external source, oligosaccharide cations produced in, "slow" metastable decomposition of, 4:210
- methyl guanine isomer distinction by hydrogen/deuterium exchange using, 4:296
- small, characterization of long-chain carboxylic esters with $\text{CH}_3\text{OBOCH}_3^+$ in, 7:1138
- surface-induced dissociation in, iron pentacarbonyl ions undergoing, energy deposition in, 3:445
- fourier transform ion cyclotron resonance, concentric tube vacuum chamber for high magnetic field, 4:168
- glow discharge, high-resolution, operated with an argon/hydrogen mixture, relative ion yields measured with, 4:278
- high-resolution, the development of, a reminiscence, 2:447
- ion trap
- laser desorption in, sensitivity for, a method for increasing, 4:706
- multiparticle simulation of ion motion in, resonant and direct current pulse excitation, 6:57
- quadrupole
- coupling a particle beam directly interface to, 4:38
- high resolution on, 2:198
- positions, velocities, and kinetic energies of resonantly elicited ions in, by laser photodissociation, 4:792
- surface-induced dissociation of molecular ions in, 2:487
- magnetic sector
- electrospray ionization on, improved detection limits for, by using an array detector, 5:194
- with heated capillary inlet, high resolution electrospray ion source for, design and performance of, 6:1256
- membrane inlet, sample modulation at, a mathematical study of, potential application in analysis of mixtures, 7:93
- multichannel plasma-source, novel, 7:458
- pentaquadrupole, MS3 capabilities of, a case study in the use

- of, ion-molecule reactions and collision-activated dissociation of $C_4H_4^+$ isomers, 3:518
- of polyglycols, lithium and metal ions in, 12:832
- quadrupole
 - double, effects of target gas in collision-induced dissociation using, 5:845
 - high-mass-to-charge ratio, proton transfer reaction studies of multiply charged proteins in, 5:1064
 - triple, ion mobility in, conformer selection of protein ions by, 5:127
- quadrupole ion trap
 - internal pulsed valve sample introduction on, 6:976
 - isomerization of linear $C_3H_3^+$ in its reaction with acetylene, and collisional stabilization of the $[C_3H_5^+]$ * collision complex in, 3:631
- tandem
 - four-sector, interface region of, enhancement of ion transmission at low collision energies via, 5:460
 - for the study of molecular dissociations, 3:487
- tandem quadrupole
 - acceleration-deceleration, novel, for neutralization-reionization studies, 3:493
 - surface-induced dissociation in, a comparison of three designs, 3:27
- time-of-flight
 - linear, large molecules in, high-resolution mass spectrometry of, 4:2
 - linear orthogonal-acceleration, detector pulse width and jitter in, resolution limitations from, 5:772
 - magnetic-orthogonal acceleration, tandem double focusing, matrix-assisted laser desorption ionization employing, peptide sequence determination by, 7:1
 - tandem, high efficiency photo-induced dissociation of precursor ions in, 4:751
- VG analytical ZAB-2FQ, cluster ions on, dissociation studies of, collision-induced high and low energy, a compact high pressure ion source for, 3:33
- Electrospray ionization (ESI) mass spectrometry
 - of affinity constants and response factors of gramicidin noncovalent dimer, 16:1031
 - calmodulin-melittin complex interface probe by, 16:225
 - charge reduction, of intact proteins, 16:1876
 - of cyclo(Pro-Gly)₃ complexes, 16:1536
 - of derivatized oligosaccharides, 16:683
 - hydrogen/deuterium exchange, *t*-Boc group elucidation using, 16:553
 - Mycobacterium* lipoarabinomannans characterized by, 16:1109
 - negative
 - organic molecule responsiveness to, acidity and polarity in, 16:446
 - unsaturated acidic phospholipids structural characterization by, 16:46
 - of noncovalent myosin VI complexes, 16:1367
 - of *Ophiopogon japonicus* homoisoflavonoids, 16:234
 - protein-DNA binding affinity evaluated by, 16:116
 - quadrupole ion trap MS with, of cardiolipin, 16:491
 - of 20S proteasome, 16:998
 - sialic acid synthase characterization by, 16:324
 - of triacylglycerols, 16:1498
- Electrospray ionization (ESI) tandem mass spectrometry (MS/MS)
 - in ion trap MS confirmation of amoxicillin degradation products, 16:1670
 - of *meso*-tetrakis(heptafluoropropyl)prophyrin, 16:1915
 - of oligoribonucleotide gas phase dissociation, 16:1262
 - for peptide fragmentation, 16:505
 - of phosphatidylserine, 16:1510
 - of synthetic neoglycolipids, 16:565
 - triiodothyronine detection and quantification by, 16:1781
- Electrospray ionization-field asymmetric waveform ion mobility spectrometry–mass spectrometry, of leucine enkephalin, 10:492
- Electrospray ionization-Fourier transform ion cyclotron mass spectrometry (ESI-FTICR-MS)
 - of 3+ oxidation state iron -sulfur metalloprotein, 12:817
 - ion fragmentation in 2-D quadrupole ion trap for external ion accumulation in, 12:1312
 - using ion preselection and external accumulation, 12:38
- Electrospray ionization-Fourier transform ion cyclotron resonance
 - of cyclodextrin complex dissociation, 8:244
 - external ion accumulation for, 8:970
 - of lanthanide(III) porphyrin-phthalocyanine heteroleptic sandwich complexes, 8:161
 - Li^+ and H^+ affinity of sulfone and sufoxide, 8:262
- Electrospray ionization Fourier transform ion cyclotron resonance (ESI-FTICR), of large polymerase chain reaction products, 11:333
- Electrospray ionization Fourier transform ion cyclotron resonance mass spectrometer (ESI-FTICR-MS), high magnetic field, trapping and detection of ions generated in, 3:615
- Electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry (ESI-FTICR-MS)
 - of alkali metal-mediated resorcarene capsule complexes, 13:851
 - amino acids, gas-phase enantioselectivity and zwitterionic formation, 17:442
 - of ammonium ion-mediated resorcarene capsules, 14:143
 - of aromatic vs aliphatic phosphopeptide dissociation, 15:121
 - ArPd(PPh₃)₂I, gas phase fragmentation reactions, 17:1582
 - automated petroleum analysis by, 20:263
 - bimolecular DNA quadruplexes and transitions to higher order molecular structures, 18:1467
 - bivalent cations and formation of purine-rich triple-helix DNA, 20:1281
 - creatine kinase complexed with ADP or ATP, deconvolution for specific vs. nonspecific interactions, 17:1239
 - elemental compositions by isotope pattern evaluation, 17:1692
 - employing data-dependent external ion selection and accumulation, 13:144
 - of GlcNAc-6-O-sulfotransferase, 15:1400
 - HPLC, MALDI-TOF/TOF MS vs. comparison using HIV envelope glycoproteins, 19:1209
 - human urine biomarkers, modified nucleosides and ribosylated metabolites, ESI-IT-MS and, 19:1500
 - localization of noncovalent binding site between oleuropein and amyloid- β -peptide, 19:1078
 - M. tuberculosis* APS reductase noncovalent complexes analyzed using, 18:167
 - mono- and diphosphonate bridged cavitands, alcohol complexation properties, 17:213
 - positive mode nano, MS/MS and, SORI-CID structural analysis of sulfated oligosaccharides, 17:96
 - of sample preparations for PCR product mass measurements, 13:338
 - in sequence confirmation of modified oligonucleotides, 14:825
 - speciation of nitrogen-containing aromatics by APPI or, 18:1265
 - at 11.5 tesla: instrument design and initial results, 9:692
 - uranyl nitrate ions, ESI generated, detected by, 17:230
 - ion kinetic energy modulation for improved ion trapping in, 6:76
 - of recombinant human macrophage colony-stimulating factor b and derivatives, 11:237

- Electrospray ionization high field asymmetric waveform ion mobility spectrometry mass spectrometry (ESI-FAIMSMS) for tryptic peptide analysis, 13:1282
of ubiquitin elongated conformers, 12:894
- Electrospray ionization ion cyclotron resonance mass spectrometer
analysis of noncovalent chemokine ligand complexes with polyanionic molecules, 17:524
high-resolution accurate mass measurements of biomolecules using, 4:566
- Electrospray ionization-ion funnel-*rf* hexapole source, in-source fragmentation technique for production of thermalized ions, 20:2312
- Electrospray ionization-ion mobility spectrometry, high resolution, drift gas selectivity in, 13:300
- Electrospray ionization ion mobility spectrometry mass spectrometry, monitoring copopulated states during protein folding, 18:2180
- Electrospray ionization ion trap mass spectrometry (ESI-ITMS) of deoxyguanosine linkage to deuterated 2-hydroxyestradiol, 364
in discrimination of arginine dipeptide isomers, 15:840
fragmentation of photomodified oligodeoxynucleotides adducted with metal ions in, 12:1174
human urine biomarkers, modified nucleosides and ribosylated metabolites, ESI-FTICR-MS and, 19:1500
of isomeric *O*-alkyl *S*-(*s*-dialkylamino)ethyl methylphosphonothiolate fragmentation and reactions, 12:902
1-*O*-alk-1'-enyl-2-acyl and 1-*O*-alkyl-2-acyl glycerophospholipid differentiation by, 18:2065
in separation and identification of oxygen-centered radical DMPO adducts, 14:862
signal and noise evaluation in, 20:1859
of tetrahedral supramolecular Ti₄L₄ cluster, 9:1099
- Electrospray ionization-liquid chromatography, techniques of, application of to FTICR-MS, 9:333
- Electrospray ionization mass spectra, peptide and protein ions in, adducts (molecular mass=98 u) attached to, origin and removal of, 1:382
- Electrospray ionization mass spectrometry. *See also* Microelectrospray ionization mass spectrometry
2-(4,6-dimethoxy-pyrimidin-2-ylsulfanyl)-*N*-phenylbenzamide, gas-phase Smiles rearrangement reactions, 18:1813
of 1-anilino-8-naphthalene-sulfonate binding to proteins, 12:428
activation energy estimation using fragmentation yield:CID of iron(II)phenanthroline, 17:962
acyl carrier protein conformation by, 18:1525
adduct between 4-aza-3-ene-1,6-diyne and DNA, 17:1342
alkali cation affinities of polyoxyethylene dodecylethers, 18:1914
of amide anions in solution, 11:1061
ammonium-counter ion clusters studied by, 18:285
amyloid- β -peptide interaction with oleuropein, 17:568
in analysis of hexose monophosphates by ion-exchange chromatography and, 17:104
for analysis of low-molecular weight anticancer drugs and their analogues, 4:588
analysis of peptides and protein mixtures by, 9:1081
analysis of serum protein precipitated with antiserum, 6:854
antimony(III)-*D*, *L*-tartrates, enantioselective binding in solution and gas phase, 20:2100
applications of, to neutral organic molecules including fullerenes, 4:596
artifacts in spectra of saturated diacyl glycerophosphocholines, 17:384
- ATP binding to growth factors detected by MALDI-TOF MS and, 19:91
 β -cyclodextrin complexes with polyamide acids, ESI-MS, 17:9
biases in ion transmission through capillary inlet, 20:2265
in biomolecular characterization, 11:1
of biotin binding to streptavidin, 6:912
of bromo- and chloroacetic acids, 11:809
cadmium thiolate conjugate characterization by ESI and triple quadrupole MS MS, 17:1442
of calcium binding to EF-hand motifs, 12:1296
of calcium-induced calmodulin association with melittin, 8:809
calibration standards for, tris (perfluoroalkylethyl)silyl alkyl amines as, 12:1050
cardiolipin characterization as sodiated ions, positive ion ESI with *q*-ion trap MS, 17:1146
CEPH family 1362 STR data base verification of, 13:97
characterization of lanthionine polysulfides from peptide disulfides, 20:783
characterization of large, heterogeneous proteins by, 7:1219
characterization of self-assembled [Ga₄L₆] host-guest complexes, 17:292
charge competition and linear dynamic detection range in, 15:1416
charge discrimination with, applying to protein analyses, 6:1098
charge reduction, 19:629
charge state distribution and macromolecular structure, 19:1239
charge state distribution in, 8:155
charge state distribution of proteins in, 20:1617
in chiral recognition of antimony potassium tartrate association with bis(*L*-alaninate)ethylenediamine cobalt(III), 11:804
chirality in formation and stability of charged amino acid clusters, 17:275
CID with, in qualitative assessment of ionic terpolymer sample monomer ratios, 13:921
cluster division model, 17:151
cobalt(III)-hexammine and, RNA hairpins probed using, 17:1376
comparison of nanoESI, ESSI and, for dissociation constant determination in protein-ligand interactions, 19:332
of complex polyol mixtures, 12:55
of complexes, quantum calculations and, 12:304
comprehensive library of, 10:287, 10:565
in confirmation of Ca²⁺-induced tetramers of MRP8 and MRP14, 11:780
of copper adducts of protonated ubiquitin, 6:521
coupling liquid chromatography to, 13:265
in cyclodextrin complex binding constant evaluation, 14:1123
of cytochrome *c* structure, 15:1612
decomposition of chloride adduct ions in, 11:933
deconvolution by an entropy-based algorithm, 3:207
detection of ribonuclease-nucleotide complexes, 19:1199
determination of binding selectivities by, 9:1049
dichloro(ethylenediamine)palladium(II) complex formation in aqueous solutions, 18:769
diffusion measurements of solution phase noncovalent interactions by, 14:430
of diketone/metal ion complexes, 10:402
dimethyldioctadecylammonium ions as contamination source in, 19:666
divalent cation binding by calmodulin analyzed by, 20:1159
of DNA-bezoxazole or DNA-benzimidazole complex, 15:1593
DNA-endonuclease VIII repair complex characterization by, 11:505
dominulin A and B in cuticle and venom of social paper wasp

- analyzed by MALDI-TOF, MALDI-TOF/TOF and ESI-ion trap, 17:376
- dueling, 13:614
- effect of protein stabilization on CSD, 18:1605
- effect of tanshinone IIA on warfarin-HSA interactions, 19:1568
- electrochemically-induced reactions of hexafluorophosphate anions with water, ES-MS, 17:939
- of encapsulation of amino acids by cyclodextrins, 6:866
- energy-dependent, gas-phase reactivity of ruthenium carbonyl clusters, 20:658
- ESMS fragmentation study, *N,N'*-disubstituted imidazolium ionic liquids, 17:85
- in evaluating lariat ether amide selectivities, 14:1215
- in evaluating noncovalent chorismate mutase inhibitor binding, 14:70
- in evaluating photoinduced changes, 14:1110
- evaluation of flavonoid-DNA complexes, 19:914
- evaluation of metal-mediated DNA binding of benzoxazole ligands by, 19:209
- exact-mass data from, FT-ICR MS automation and, 10:1166
- extractive, sampling explosives on skin, using neutral desorption and, 20:719
- in flavonoid glycoside isomer differentiation, 14:1437
- flow injection, in bacterial identification, 13:118
- fragmentation of doubly-protonated tripeptides, formation of b_2^+ ions, 20:2135
- and fragmentation of *N*-linked carbohydrates derivatized at the reducing terminus, 11:900
- fundamentals, instrumentation, and applications of, 8:1191
- gas phase hydrogen/deuterium exchange in, as a practical tool for structure elucidation, 5:434
- gas-phase molecular dications formed from neutral organics in solution via chemical electron-transfer reactions by using, observation of, 5:689
- gas-phase stability of G-quadruplex DNA using MDS and, 18:1760
- glucose concentration in tears by, 18:332
- glutamine N-terminal, dehydration *versus* deamination in, 18:27
- H-D exchange combined with, in calcitonin folding analysis, 15:1556
- H/D exchange of hydrophobic peptides by, 13:1376
- heme-globin complex detection in, electrostatic interactions and, 12:1092
- of hemocyanins, 14:419
- hemoglobin folding and assembly monitored by ESI-MS, 18:8
- heterotetrameric sarcosine oxidase characterization by, 9:510
- high mass-to-charge ratio ions from, observations and implications of, 4:536
- high resolution, of large molecules, automated reduction and interpretation of, 11:320
- HPLC, arsenic-glutathione complex analysis by, 19:1559
- hydrogen/deuterium exchange, β_2 -M release from MHC-I analyzed by, 20:278
- hydrogen/deuterium exchange for probing the interaction of calmodulin with calcium, 10:711
- hydrolysis of PtenCl₂ investigated by DFT and, 20:1015
- in identification of cyclodextrin bearing by azobenzene group, 11:16
- improved signal intensity of oligonucleotides in, charge-state reduction with, 7:697
- in indirect assessment of hydrophobic ligand binding, 14:506
- in indirect detection of protein-metal binding, 15:1658
- interaction of angiotensin peptides and zinc metal ions by, 5:959
- interactions among polythymine oligonucleotides, ethidium bromide and mercury ions, 20:1834
- intrasource separation and selective ionization of glycerophospholipids, 17:264
- investigation of dimeric G-quadruplex inhibitors of HIV-1 integrase by, 19:550
- ion distribution within droplets, 17:318
- ionization and transmission efficiency in, 18:1582
- iron II and III gaseous complexes with BINOLate ligands, 19:121
- ITMS coupled to, phosphopeptide fragmentation by, 13:274
- LC, TPST-1 and TPST-2 analysis by, 19:1459
- of leech hemoglobin subunit structure, 15:22
- of Li⁺ adducts of unsaturated fatty acids, 10:600
- of local anesthetic-cyclodextrin non-covalent complexes, 13:928
- low-energy, geometrical isomers of α -linolenic acid methyl ester separated using, 20:1272
- low mass poly(methyl methacrylate) polymers and, 6:906
- low temperature aqueous, of noncovalent complexes, 9:580
- of lysoglycerophospholine lipid subclasses, 11:283
- M. bovis* BCG PIMS characterization by MALDI-TOF MS and, 18:466, 18:479
- and matrix-assisted ionization laser desorption mass spectrometry, characterization of mouse switch variant antibodies by, 7:707
- metal-complex formation in continuous-flow ligand-exchange reactors, 18:707
- metallo- β lactamase ternary inhibitor complexes, 17:1000
- Michael-Michael-retro Michael addition catalyzed by 9-amino-9-deoxyepiquinine, 18:2074
- micro, centrin-2 metal-binding properties determined by UV spectroscopy and, 17:1158
- microfabricated nanospray tips for, nib-like design for, 15:407
- microfabricated PDMS multichannel emitter for, 12:463
- microfluidic chip for millisecond time-scale kinetic studies, 20:124
- of microcystins, heptapeptide hepatotoxins, modulation of charge states and $[M + H]^+$ to $[M + Na]^+$ ratio, 10:1138
- MS/MS and, investigation of enantioselective multivalency by, 19:1634
- with multiple stage QIT MS, *E. coli* cardiolipin characterization by, 17:420
- multiple substrate, determination of enzyme-substrate specificity constants using, 15:233
- nano, automated, CphA, monitoring zinc affinity by, 17:180
- nanoscale liquid chromatography with, of proteolytic digests, exponential dilution gradient system for, 12:1205
- nanoscale separations combined with, sulfonamide determination, 3:139
- negative, diuretic fragmentation in, 14:658
- negative, IgG antibody disulfide linkages characterized by reversed phase HPLC with, 17:1590
- negative, MS/MS of singly and doubly charged ions in, 13:671
- nemorosone degradation profile by, 20:1688
- NeuAc oligomers associated with tetanus toxin C fragment, 17:967
- noise reduction by supplemental IR activation, 20:2206
- noncovalent complexes, weak interactions analyzed by, 20:303
- noncovalent complexes formed between duplex DNA and minor groove-binding molecules by, characterization of, 6:1154
- noncovalent DNA complexes with polybasic compounds, studied by nanoES-MS, 18:346
- nonspecific ligand binding identification using reporter molecule method, 20:1242
- of nucleobases and nucleosides, processes that affect, 7:1106

- observation of noncovalent complexes to the avidin tetramer by, 5:201
- observations of phenyl-fused porphorinoids using, 17:1306
- of oligonucleotides, 8:90
- of oligonucleotides, specific and nonspecific, dimer formation in, 6:159
- oligosaccharides labeled with multifunctioning tag, analysis by MALDI-TOF and, 17:194
- organotrifluoroborate mass determination using, 18:404
- ortho effect in *N*-acylanilines, 19:1114
- oxidized glutathione and contryphan, fragmentation of peptide disulfides from negative ion MS, 19:358
- PCR-LC, genomic DNA analysis by, 17:124
- of peptide-peptide interactions, 13:166
- of peptides and proteins, alkali-metal adduct ion formation in, 11:967
- of a per(onio)-substituted benzene, retention of a coulombic charge upon collisionally activated decomposition, 9:15
- perylene diimide and benzannulated perylene diimide ligand binding to DNA, 17:593
- of pH-dependent dissociation and denaturation processes of a heterodimeric protein, 11:54
- of phenylthiohydantoin amino acid derivatives, 8:1165
- of phosphopeptides isolated by on-line immobilized metal ion affinity chromatography, 4:662
- phosphorylated peptide residues in peptide-peptide formation, 19:1535
- of photo-oxidized ruthenium(II) complexes, 8:713
- pneumatically assisted, development and application of, 9:498
- polyamide binding to DNA evaluated by, 17:1742
- of poly(ethylene glycol) esters, 13:888
- positive and negative, Boc-protected peptides, 18:651
- positive and negative ion, of drug-DNA complexes, 15:1382
- positive ion mode, dicationic reagents in anion detection, 19:261
- positive mode, *meso*-tris-*N*-methylpyridinium-4-yl porphyrins reduction in, 18:762
- probing protein structure using site-specific 18-crown-6 attachment, 17:1209
- probing the helical content of growth hormone-releasing factor analogs using, 4:646
- processes that affect, nucleobases and nucleosides, 7:1106
- of protein, 13:107
- protein complex sizing by IMS and, 18:1206
- protein identification by, 9:194, 13:215
- protein ions produced by, charge states of, solvent effects on, 11:976
- protein noncovalent complexes, 20:593
- protein unfolding monitored by, 9:1248
- of proteins, mass error in, water cluster calibration and, 8:1158
- of proteins and protein mixtures, ion-impact energy measurement in, 12:826
- of proteins in methanol, 12:317
- of proton affinities, 10:483
- protonation in, 11:961
- Q-FT-ICR and, MMA, intact proteins and product ions, 18:1
- for quantitation of human serum albumin, 8:1247
- rapid squencing of peptides with a picomole-scale method, 17:995
- reproducibility using thermometer ion-based strategy, 20:398
- rhenium compound transformation during, 8:191
- role of basicity in determining selectivity in, 19:719
- of salt concentration effect on analyte response, 10:625
- of self-assembled 1,1-ferrocenedicarboxylic acid, 10:557
- of semiconductor nanoclusters, analysis of positive and negative ion mode, 11:338
- sheathless CE interface with, 8:561
- signal intensities in, ground loop current effects on, 15:1748
- silver(I) complexes produced in, 8:781
- sodium stabilization of iturin A2 during, 12:505
- sodium trifluoroacetate as tune/calibration compound for, 9:977
- of solution phase H/D exchange of non-covalent complexes, 12:795
- solvent-cation binding in, 10:254
- of Src SH₂-phosphopeptide complexes, 8:234
- study of anion- π interactions, 20:257
- study of noncovalent enzyme-inhibitor complexes and metal binding stoichiometry of matrolysin by, 6:1105
- a study of the thermal denaturation of ribonuclease S by, 5:614
- structural characterization of triacylglycerols by, 10:587
- structural elucidation by, 9:1267
- synchronized dual-polarity, 20:2254
- thermal denaturation of cytochrome *c* analyzed by, 20:819
- time-resolved, of pH effect on myoglobin unfolding, 11:312
- transition metal complexation by 3-azidopropionitrile, 18:453
- of tris(trimethoxyphenyl)phosphonium acetyl charged derivatives of peptides, 11:107
- tunable metal-ligand complexation for flavonoid diglycosides, 18:422
- of unsubstituted and disubstituted β -cyclodextrins, 14:124
- use of battery-powered two-electrode emitter cell, 17:953
- vinylpyridinium substituted porphyrins characterized by, 18:218
- zinc deposition during peptide-zinc complex analysis, 20:2199
- Electrospray ionization mass spectrometry/capillary zone electrophoresis, application of, to the analysis of macrolide antibiotics, 3:563
- Electrospray ionization mass spectrometry/gradient reverse-phase high-performance liquid chromatography, analysis with trifluoroacetic and other strong acid modifiers by postcolumn addition of propionic acid and isopropanol, signal enhancement of, 6:1221
- Electrospray ionization-mass spectrometry matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry-liquid chromatography, rapid analysis of antibiotic-containing mixtures from fermentation broths by using, 7:1227
- Electrospray ionization matrix-assisted laser desorption/ionization mass spectrometry, CID in, free-radical-induced site-specific peptide cleavage in gas phase, 18:807
- Electrospray ionization-MS/MS
- HPLC with, in on-line strong cation exchange, 14:719
- of protein modification by 4-hydroxy-2-nonenal and other short chain aldehydes, 14:215
- in trypsin-catalyzed ¹⁶O-to-¹⁸O exchange, for comparative proteomics, 14:704
- Electrospray ionization pseudo-MS, biopolymer sequencing by, 12:846
- Electrospray ionization-quadrupole-ion mobility time-of-flight mass spectrometry, trastuzumab heterogeneity profile, 20:2021
- Electrospray ionization quadrupole ion trap mass spectrometry, *A. thaliana* phosphatidylglycerol analysis by MS/MS, 18:783
- Electrospray ionization-quadrupole ion trap MS/MS, in differentiation of C8-substituted alkylaniline guanine adducts, 14:88
- Electrospray ionization-quadrupole MS/MS, on phosphatidylcholines, fragmentation processes in, 14:352
- Electrospray ionization quadrupole time-of-flight (ESI-Q-TOF) mass spectrometry, in quantifying methionine oxidation in calmodulin, 16:1470

- Electrospray ionization quadrupole time-of-flight MS fucosylation study, of low abundance species in complex mixtures, 12:964
- Electrospray ionization quadrupole time-of-flight tandem mass spectrometry, oversulfation of bovine aorta chondroitin sulfate and GAG analyzed using nano-ESI-Q-TOF-MS/MS and FTICR MS, 179
- Electrospray ionization-quadrupole time-of-flight tandem mass spectrometry (ESI-quadrupole TOF-MS/MS), negative mode, in sequencing multiantennary N-glycans, 13:1138
- Electrospray ionization source, small-bore, construction and characterization of, 4:964
- Electrospray ionization source H/D exchange, LC-MS and LC-MS/MS for characterization of metabolites, 13:345
- Electrospray ionization tandem mass spectrometry (ESI-MS/MS) of acidic heteroxylans uronosyl substitution patterns, 15:43
- analysis of peptides derived by enzymatic digestion of oxidized globin subunits, an improved method to determine amino acid substitution in the hemoglobin "core," 7:1040
- of cysteinyl eicosanoids, leukotriene, C4 and FOG7, 12:763
- of cytosine nucleosides self-assembling into dimers, 15:268, 15:436
- effect of cisplatin on gas phase dissociation of oligonucleotides, 20:792
- of flavonoid aglycones, 12:707
- fragmentation pathways of anabolic steroids, 17:477
- of glycerophosphocholine lipid lithiated adducts, 9:516
- HPLC-, phosphorylation of IRS-1 quantified by, 17:562
- of hydroperoxy bishemiacetal product derived from cholesterol ozonolysis, 15:194
- an identification of new metabolite of rapamycin and SDZ RAD, 11:516
- in identification of ozone oxidation products of amino acids and small peptides, 11:526
- of kynurenine fragmentation, 12:786
- LC-, identification of mammalian cell lines using MALDI-TOF and, 17:490
- Leishmania major* inositol phosphorylceramides, characterization by, 18:1591
- medicinal herb saponin structural analysis with, 15:133
- of metal-adducted oligodeoxynucleotide ions, 12:550
- of methylarginine-containing peptide fragmentation pathways, 15:142
- multistage, nLC, *C. quinoa* triterpene saponin mixture, 17:795
- of mutagenic nitropyrenes, 15:1853
- negative ion, alginate-derived oligosaccharide sequence analysis by, 17:621
- of oligonucleotide fragmentation, 12:193
- for oxidized deoxynucleoside LC, 12:80
- oxysterol analysis by, 17:341
- of phosphatidylinositol and its phosphates, 11:986
- of polyamine toxins of spiders, neighboring group participation in, alpha,omega-diaminoalkane compounds, 6:634
- poly(ethylene oxide)/polystyrene block copolymer, microstructural study by, 19:1163
- positive, odd-electron prodiginine fragments, 19:1857
- positive and negative mode, identification of phosphorylation sites in phosphopeptides by, 7:243
- positive ion, of cleavage reactions of complex ions, 15:689
- product ion isotope ratios from hemoglobin Camden, 18:1493
- pyrimidinal antiviral agents, CID mechanisms of ion formation, 18:1477
- resolvin D1, protectin D1, and related derived products of DHA, 18:128
- S- and N- nitrosopeptide analysis using, 17:1725
- of *Salmonella typhimurium* acylphosphatidylglycerols, 15:1
- screening threading bis-intercalator binding to DNA by, 18:311
- sialyl anomers distinction on FAB-MS/MS and, 20:394
- silver binding by polyamidoamine dendrimers, 17:676
- structural determination of picomole amounts of phospholipids via, 6:1202
- of sulfated disaccharides in chondroitin sulfate, 11:916
- triple quadrupole, in molecular characterization of thiols, 15:288
- without chromatography, of NSAIDs, 9:255
- Electrospray ionization time-of-flight mass spectrometry collisional damping interface for, 9:69
- CSD in, IgG1κ mAb characterization using, 20:1030
- heterotetrameric sarcosine oxidase characterization by, 9:510
- HPLC, improving mass accuracy of, antibodies, 17:867
- multianode detection in, 9:1325
- of protein kinase G, cGMP, and AMPNP, 15:1392
- sub-ppm mass accuracy on, use with drug discovery recovery libraries, 18:1612
- Electrospray ionization trap mass spectrometry fragmentation of analogs of saccharides from *V. cholerae* 0:1, 17:749
- low energy, isoleucine and leucine discrimination using, 18:57
- Electrospray ionization-collision-induced dissociation, in serum drug detection, 10:1028
- Electrospray ionization-Fourier transform mass spectrometry, of external ion accumulation of poly(ethylene)glycol, 10:1298
- Electrospray ionization-high field asymmetric waveform ion mobility spectrometry-mass spectrometry, in separation of leucine and isoleucine anions, 10:1279
- Electrospray ions mass and charge assignment for, by cation adduction, 4:828
- solvation studies of, 9:1241
- Electrospray liquid chromatography/mass spectrometry, selective detection of phosphopeptides in complex mixtures by, 4:710
- Electrospray liquid chromatography-tandem mass spectrometry, quantitation of SR 27417 in human plasma using, a study of ion suppression, 7:1099
- Electrospray low energy CID and MALDI PSD fragmentations, of protonated sulfinamide cross-linked peptides, 13:709
- Electrospray mass analysis, of chemical oxidation of 1,2-diarylcyclopropanes by Cu(II) salt, 9:966
- Electrospray mass spectra capillary electrophoresis of herbicides paraquat and diquat, 7:981
- of proteins, are they related to their aqueous solution chemistry?, 3:216
- Electrospray mass spectrometric analysis, of 5-hydroperoxy and 5-hydroxyeicosatetraenoic acids generated by lipid peroxidation of red blood cell ghost phospholipids, 9:527
- Electrospray mass spectrometry (EMS) for the analysis of opioid peptides and for the quantification of endogenous methionine enkephalin and beta-endorphin, 2:149
- application of, in probing protein-protein and protein-ligand noncovalent interactions, 4:624
- of borane salts, electrospray needle as an electrochemical cell, 6:860
- cyclodextrin inclusion complexes detected by, false positives and, 6:437
- droplets in break-up of, physical/chemical separations in, 12:343
- charge gradients in, 12:207
- evidence from, stepwise refolding of acid-denatured myoglobin, 4:638

- of gas phase ternary β -cyclodextrin inclusion complexes, 14:449
- glycoprotein profiling by, 15:750
- of 4-hydroxy-2-alkylquinolines from *Pseudomonas aeruginosa*, 15:862
- of ligands with retinoid X receptor α ligand binding domain, 16:1631
- liquid phase composition and salt cluster formation in, 12:1077
- of methanol and water solutions suppression of electric discharge with SF₆ gas, 2:497
- negative ion, of nucleotides, ionization from water solution with SF₆ discharge suppression, 4:289
- negative ion adduct stability in, multiple hydrogen bonding and, 13:1360
- negative ion formation in, 4:578
- negative mode, oligosaccharide analysis in, 16:60
- noncovalent adduction in, mass and charge state assignment for proteins and peptide mixtures via, 6:1175
- nonmetallized tip for, 10:72
- nonolite, long-lived metallized tips for, 7:1270
- plasma peptide mapping with, 10:45
- of protein electroeluted from sodium dodecylsulfate polyacrylamide gel electrophoresis gels, 10:453
- protein fluorescence measurements within, 12:716
- reversed-phase microbore liquid chromatography, complementary mobile phases, and sequential exoglycosidase digestion, characterization of protein N-glycosylation by, 5:350
- sequential, for structural elucidation of macrolide aglycones, 13:862
- solution flow rates in, analyte ion signals and, 14:492
- specificity of cyclodextrin complexes detected by, 13:946
- stable association complex, for cyanuric acid determination, 12:1085
- stereoselectivity and ion response in, correlation of, 6:207
- a study of some aqueous solutions of metal salts, 3:281
- Electrospray mass spectrometry and matrix-assisted laser desorption ionization time-of-flight mass spectrometry, application of, for molecular weight assignment of peptides in complex mixtures, 4:670
- Electrospray-MS/MS, nano-LC combined with, to detect estrogen DNA-adducts in breast tumor and healthy tissue, 14:482
- Electrospray-MS/MS investigations of morphinans, 14:1262
- Electrospray needle voltage, in CE-MS, 10:261
- Electrospray quadrupole time-of-flight mass spectrometry
- noise filtering techniques for, 766
 - rapid switchability with MALDI, for protein identification, 11:493
 - to study noncovalent associations of glutathione S-transferase with ligands, 11:606
- Electrospray-single photon ionization mass spectrometry, analysis of organic compounds by ES/VUV-SPI MS, 20:430
- Electrospray source
- a heated, for mass spectrometry of analytes from aqueous solutions, 5:791
 - heated capillary, charge-dependent/independent ion energy distribution from, selective generation of, 5:221
- Electrospray spectra
- native, charge state assignment of, protein complexes, 20:435
 - negative ion, production from carbohydrates
 - enzyme-catalyzed reactions, microwave-assisted, in solvent systems, 16:581
 - high mannose N-linked glycans in, 16:631
 - hybrid and complex N-linked glycans in, 16:645
 - nitrate and other anionic adducts in, 16:622
- Electrospray tandem mass spectrometric structural characterization, negative ion, of leukotriene B₄ (LTB₄) and LTB₄-derived metabolites, 7:129
- Electrospray tandem mass spectrometry
- of ceramides, lithiated, 13:680
 - in characterization of human Hb Ta-Li β 83 (EF7)Gly \rightarrow Cys variant by, 13:187
 - glutathione conjugates, EPR and, 20:2013
 - in locating abasic sites in oligodeoxynucleotides, 13:1418
 - of mixed-sequence RNA/DNA oligonucleotides, 13:936
 - structure analysis of, phenylglycidyl ether adducts of 29-deoxycytidine and 29-deoxyadenosine, 7:682
- Electrospray time-of-flight mass spectrometer, vacuum chamber of, ion-molecule collisions in, on mass spectra of proteins, 7:342
- Electrospray time-of-flight mass spectrometry (ESMS-TOF), of Ti₆O₄(O₂C₄H₅)₈(OCH₂CH₃)₈, 15:260
- Electrospray voltages, selection for gradient elution LC-MS, 20:682
- Electrosprayed ions, gas-phase, implications for the conformation of, defect formation on surfaces bombarded by energetic multiply charged proteins, 7:329
- Electrosprayed nucleotide ions, non-covalent adducts of, infrared photodissociation of, 7:209
- Electrosprayed proteins, gas-phase structure of, investigation of using ion-molecule reactions, 5:207
- Electrospraying, of water in air at atmospheric pressure, 15:253
- Electrosprays, oxidizing, desorption electrospray ionization enhanced signal using, 17:1091
- Electrostatic energy analyzer-time-of-flight mass spectrometer, for measurement of laser desorbed ion kinetic energies, design and calibration of, 6:619
- Electrostatic interactions
- and detection of heme-globin complex in ESI-MS, 12:1092
 - noncovalent complexes, site identified by ECD/ETD, 20:176
 - in stability of lithium atom complexes with formaldehyde and formalimine, 12:1229
- Electrostatic ion deflection, new method for, 1:28
- Electrostatic ion guide
- for external ion source Fourier transform ion cyclotron resonance mass spectrometry, ion trajectories in, 6:936
 - stacked-ring, 7:101
- Electrostatic ion traps, multiply reflecting, stability for, 10:241
- Element composition, isotope incorporation for, 11:835
- Elemental analysis, inductively coupled plasma-time-of-flight mass spectrometer for elemental analysis, 6:400, 411
- Elemental composition, by isotope pattern evaluation, ESI-FTICR MS, 17:1692
- Elemental distribution, in fluorinated amorphous carbon thin films, FIB-SIMS of, 16:126
- Elemental ion source, the pulsed glow discharge as, 1:138
- Elimination of methane from protonated acetaldehyde, the ab initio transition state, 5:1102
- Elimination of the concentration dependence in MIAMS of methyl palmitate using MAB, 12:754
- Elimination of water from the carboxyl group of GlyGlyH⁺, 1192
- 1,2-Eliminations, cations in the gas phase, 19:1491
- Elongated conformers of charge states +11 to +15 of bovine ubiquitin studied using ESI-FAIMS-MS, 12:894
- Elucidating the structures of isomeric silylenium ions (SiC₃H⁺9, SiC₄H⁺11, SiC₅H⁺13) by using specific ion/molecule reactions, 2:278
- Elucidation of the CID pathways of water and alcohol coordinated complexes containing the uranyl cation, 14:1205
- Elucidation of the initial step of oligonucleotide fragmentation in MALDI using modified nucleic acids, 12:744

- Elution time peptide tags, in proteome analyses, with capillary LC-TOFMS, 14:980
- Emergin pollutants, and communicating the science of environmental chemistry and MS, pharmaceuticals in the environment, 12:1067
- Emitter, electrospray, potential build-up and gradient mapping within, 15:1737
- Emitter cells, battery-powered two-electrode emitter cell, SES-MS, 17:953
- Emitter tips, conducting microfabricated nib-like nanoelectrospray design, 17:75
- Emulate ions, H/D exchange reactions of, with CH₃OD and C₂H₅OD, reactive collisions in quadrupole cells, 3:853
- Empirical approach to estimation of critical energies by using a quadrupole ion trap, 7:1116
- EMS. *See* Electrospray mass spectrometry
- Enantiomers, separation by MS, β -amino acids, 20:1235
- Enantioselective molecular recognition, investigation of, by ESI MS and MS/MS, 19:1634
- End group determination, high resolution, of low molecular weight polymers by matrix-assisted laser desorption ionization of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, 7:449
- End-stage renal disease, glycosylated globins, analysis by MALDI MS, 18:1018
- End user, putting mass spectrometry in hands of, 6:394
- Endgroup-assisted siloxane bond cleavage in gas phase, 14:1039
- Endgroup determination of synthetic polymers, by ESI-FTICR, 11:536
- Endothermic ion molecule reactions, 2:189
- Energetic multiply charged proteins, defect formation on surfaces bombarded by, implications for the conformation of gas-phase electrosprayed ions, 7:329
- Energy
channeling into dissociation, a photoionization study of the ion-neutral complexes [CH₃CH₃+ CH₂CH₃] and [CH₃CH₂CH⁺CH₃CH₃] in the gas phase, 7:73
critical, estimation of, an empirical approach to, by using a quadrupole ion trap, 7:1116
geometry and time, ion formation from charged droplets, 4:524
internal, distributions deposited in doubly and singly charged tungsten hexacarbonyl ions generated by charge stripping, electron impact, and charge exchange, 1:16
of large MALDI ions, 8:1094
- Energy analyzer, a high transmission, for ion spectrometry, 2:506
- Energy compensation, improved, for time-of-flight mass spectrometry, 5:779
- Energy-dependent electrospray ionization mass spectrometry, gas-phase reactivity of ruthenium carbonyl clusters, 20:658
- Energy deposition
during electron-induced dissociation, 4:145
in iron pentacarbonyl ions undergoing surface-induced dissociation in a Fourier transform mass spectrometer, 3:445
- Energy loss, to investigate ubiquitin cross-sections, 11:738
- Energy partitioning, differential overlap, modified neglect of, and ab initio and modified neglect of differential overlap bond orders, effect of protonation site on bond strengths in simple peptides, 5:704
- Energy-resolved mass spectrometry, fucosyl galactoside isomers analyzed by, 18:1873
- Energy spectra, translational, angle-resolved, of survivor ions, high-energy collisional activation studied via, 4:835
- Energy transfer, in inelastic molecular collisions, 14:1348
- Enhanced detection of phosphopeptides in MALDI-MS using ammonium salts, 10:35
- Enhancement of detection of amino acids or alcohols by electron capture negative ion chemical ionization mass spectrometry, pentafluorobenzyl chloroformate derivatization for, 6:525
- Enhancement of ion intensity in time-of-flight secondary-ionization mass spectrometry, 7:467
- Enhancement of ion transmission at low collision energies via modifications to the interface region of a four-sector tandem mass spectrometer, 5:460
- Enhancement of mass spectrometric detection of LTC₄, LTC₄, and PTE₄ by derivatization, 5:292
- Enhancement of the effective resolution of mass spectra of high-mass biomolecules by maximum entropy-based deconvolution to eliminate the isotopic natural abundance distribution, 8:659
- Enkephalin
leukine, dissociation of, 8:771
methionine, endogenous, electrospray mass spectrometry for, 2:149
reaction with NDA for LC/LIF/MS analysis, 18:1706
- Enkephalin derivatives, molecular radical cations of, formation via CID of ES-generated copper(II) complex ions, 12:1114
- Enlarged data base of electron-ionization mass spectra, 2:432
- Enlarged sampling orifice and offset ion lens, inductively coupled plasma mass spectrometry with
ion trajectories and detector performance, 4:16
polyatomic ion interferences and matrix effects, 4:28
- Enndo tricyclo[5.2.1.0^{2,6}]decene, double hydrogen migration in, 13:1186
- Enol ion, of methyl propionate heat of formation, 10:200
- Enolate anion, gas-phase base-catalyzed Claisen-Schmidt reactions of, with various para-substituted benzaldehydes, 7:82
- Enolic radical cations, 1,3-H shifts of, orbital symmetry restraint circumvention by, 15:972
- Enols, ionized reaction with tert-butanol, 12:938
- Enterobactin, siderophores, Fe(III) complex with, IRMPD, 18:632
- Enterococcus faecium*, vancomycin-susceptible, structural variation in peptidoglycan of, LC-MS, 19:1467
- Entropic effects, in alkali chloride cluster ion dissociation, 13:1388
- Entropy, apparent, meaning of, 13:435
- Entropy analysis, kinetic method with
to determine proton affinities of bromo- and iodoacetonitrile, 11:601
as statistically meaningful approach, 11:371
- Entropy-based algorithm, deconvolution by, electrospray ionization mass spectrometry, 3:207
- Environment, pharmaceuticals in, 12:1067
- Environmental and forensic applications of field-portable GC/MS, 12:683
- Environmental applications, of analytical MS, 13:898
- Environmental atmospheric pressure, and ionization pathways of polyfluorinated chemicals, 14:1177
- Environmental contaminants, mutagenic/carcinogenic potential of, estimation of, by ion-molecule reactions and tandem mass spectrometry, 1:110
- Environmental samples, trace monobromopolychlorodibenzo-p-dioxins in, application of gas chromatography high-resolution mass spectrometry to, 3:248
- Enzymatic digestion
fast-atom bombardment mass spectrometry and, recombinant human macrophage colony stimulating factor, disulfide linkage and disulfide bond scrambling in, 6:638
labeled oligonucleotide characterization using, 9:660

- of oxidized globin subunits, peptides derived by, electrospray ionization-tandem mass spectrometry analysis of, an improved method to determine amino acid substitution in the hemoglobin "core," 7:1040
- to study IgG nonenzymic glycation, 11:153
- Enzyme-inhibitor complexes, noncovalent, and metal binding stoichiometry of matrilysin by electrospray ionization mass spectrometry, 6:1105
- Enzyme inhibitors
- immobilized on silica microspheres, screening by ESI-MS, 19:865
 - protein structure and, 12:97
- Enzyme-substrate specificity constants, determination using multiple substrate ESI-MS assay, 15:233
- Enzymes
- monitoring reactions and screening inhibitors, MALDI-TOF-MS with oxidized carbon nanotube matrix, 17:1616
 - rapid inhibitor screening, label-free approach to, MALDI-TOF MS, 17:815
- Epidermal growth factor receptor, ligand-induced site-specific phosphorylation of, 14:1022
- Epitope mapping of monoclonal antibodies by mass spectrometry, identification of protein antigens in complex biological systems, 9:208
- Epitope peptides, identification of, by direct MALDI-MS/MS, 10:76
- Epoxides, Meerwein reaction of phosphonium ions with, in gas phase, 15:398
- EPR. *See* Electron paramagnetic resonance spectroscopy
- Equilibrium ion density distribution and trapping force, combined ion traps vs, 9:473
- Equilibrium space charge distribution in a quadrupole ion trap, 5:64
- Ergodic dissociation, peptide cation radicals, spin-trap capped peptides, electron capture in, 18:432
- Ergosterol, pathway inhibition in fungi, analysis by ASAP, 18:1274
- ERK2, deuterium exchange behavior of, pepsin mapping of, 10:685
- Errata, 16:797, 16:798, 16:799, 16:800, 16:801, 16:802
- Atmospheric pressure chemical ionization of alkanes, alkenes, and cycloalkanes, 5:1026
 - Complexes of Ion (II) with Cysteine-Containing Peptides in the Gas Phase, 8:199
 - Improving Fast Atom Bombardment Mass Spectra, The Influence of Some Controllable Parameters on Spectral Quality, 1:349
 - LC/MS and LC/MS Determination of Protein Tryptic Digests, 1:497
 - Low-Energy Tandem Mass Spectrometry of the Molecular Ion Derived from Fatty Acid Methyl Esters, A Novel Method for Analysis of Branched-Chain Fatty Acids, 4:834
 - New derivatization strategy for analysis of phosphopeptides by precursor ion scanning in positive ion mode, 14:83
 - Role of the Site of Protonation in the Low-Energy Decomposition of Gas Phase Peptide Ions, 7:758
 - A Study of the Gas-Phase Reaction Between Protonated Acetaldehyde and Methanol, 2:259
 - Transition-Metal Mediated Heteroatom Removal by Reactions of FeL^+ [$\text{L}^+ \text{O}$, C_4H_6 , $\text{c-C}_5\text{H}_6$, $\text{c-C}_5\text{H}_5$, H_6H_6 , C_3H_4 ($=\text{CH}_2$)] with Furan, Thiophene, and Pyrrole in the Gas Phase, 7:1177
- Errors, systematic, originating from Coulomb mutual interaction and image charge in Fourier transform ion cyclotron resonance precise mass difference measurements, analysis of estimation of, 4:855
- Erythromycin analogs, characterization by CAD and IRMPD, 13:630
- ES-MS. *See* Electrospray mass spectrometry; Electrospray-MS
- Escherichia coli*
- ribosome topography analyzed by proteolysis and MALDI-MS, 18:1304
 - Shigella* species and, protein expression monitored by LC/QTOF-MS, 19:1626
- ESI. *See* Electrospray ionization
- ESI-FAIMS-MS. *See* Electrospray ionization-high field asymmetric waveform ion mobility spectrometry-mass spectrometry.
- ESI-IMS-MS. *See* Electrospray ionization ion mobility spectrometry mass spectrometry
- ESI-LC. *See* Electrospray ionization-liquid chromatography; Liquid chromatography-(electrospray ionization) mass spectrometry
- ESI-MS. *See* Electrospray ionization mass spectrometry
- ESI-MS/MS. *See* Electrospray ionization tandem mass spectrometry
- ESMS. *See* Electrospray mass spectrometry
- ESR. *See* Electron spin resonance
- Essential oil components, GC-quadrupole MS identification of, 16:1902
- Esterification, and peptide ion detection by MALDI-MS, 16:1269
- Esters
- deprotonated phenol fatty acid, ion-dipole complex formation, evidenced by using gas-phase labeling combined with tandem mass spectrometry, 4:343
 - long-chain, comparison with, electron ionization induced fragmentation of macrocyclic amines $\text{C}_n\text{H}_{2n-1}\text{NH}_2$, evidence for the rearrangement of aminocarbene radical cations, 1:238
- Esters and acids, gas-phase methylation and dissociation of, effects of functional group interactions on, 7:565
- Estimating probabilities of correct identification from results of mass spectral library searches, 5:316
- Estimation of effective ion temperatures in a quadrupole ion trap, 9:845
- Estimation of the electron affinities of H_2O
- cirranulene and coronene by using the kinetic method, 7:619
 - corannulene, and coronene by using, 7:619
- 17 β -Estradiols, 11 α -OH and 11 β -OH, stereochemical effects on gas phase acidity, 20:2318
- Estrogen DNA-adducts, in breast tumor and healthy tissue, detection by combined nano-LC and nano-ES MS/MS, 14:482
- Estrogen receptor, zinc fingers of, cysteine reactivity in, 16:2017
- Estrogen receptor α , serine phosphorylation site in NTD, from MCF7 cells, 19:729
- Estrogens
- determination using SALDI-MS, 19:1343
 - ultrafiltration MS/MS of, 16:271
- Estrone and estradiol, glutathione conjugates of, structural isomers of, 9:612
- ETD. *See* Electron transfer dissociation
- Ethanol
- hydrogen atom abstraction from, phenyl radical selectivity in, 15:913
 - protonated, and its neutral counterparts, 2:459
- Ethanol-treated rat hepatocytes, novel 3-hydroxylated leukotriene B_4 metabolites from, 3:762
- Ethene elimination from $\text{CH}_3\text{CH}_2\text{NH}=\text{CH}_2^+$, reaction pathways at the boundary between stepwise and concerted processes, 9:138
- Ethene loss kinetics, of methyl 2-methyl butanoate ions studied

- by threshold photoelectron-photoion coincidence, enol ion of methyl propionate heat of formation, 10:200
- Etheno-DNA adducts, urinary, quantification by column switching LC/APCI-MS/MS, 17:605
- Ether adducts, phenylglycidyl, of 29-deoxycytidine and 29-deoxyadenosine, stability in solution and structure analysis by electrospray tandem mass spectrometry, 7:682
- Ether ions, dimethyl
 gas-phase reactions of amino alcohols with site-selective, 4:230
 selective ion-molecule reactions of, with lactams, 4:353
- Ethers
 crown, -metal ion complexes, direct desorption of, in liquid secondary ionization mass spectrometry, 5:638
 crown, orders of relative alkali metal ion affinities of, by the kinetic method, 3:543
 cyclic organic, ClNP^+ NCI reactivity toward, 10:379
 dimethyl, ion/molecule reaction with, selective detection of the tolyl cation among other $[\text{C}_7\text{H}_7]^+$ isomers by, 2:270
 ethylene glycol, monomethyl and dimethyl, functional group-selective ion-molecule reactions of, 4:97
 ionized aliphatic, ion-neutral-mediated alkane eliminations from, size effect in, ethylene glycol, monomethyl and dimethyl, functional group-selective ion-molecule reactions of, 2:261
 ionized phytyl methyl, $\text{C}_3\text{H90}^+$ formation from, the structure and mechanisms of, 7:205
 lariat crown, lysine probing by, in peptides, 13:493
 phenyl allenylmethyl, electron-impact-induced 3,3-sigmatropic rearrangement and cyclization in, 6:195
 phenyl n-propyl and methyl substituted analogs, chemical ionization of, propene loss initiated by competing proton transfer to the oxygen atom and the aromatic ring, 7:639
 trimethylsilyl, of cholesterol oxidation products, mass spectra of, trimethylsilyl group migration in, product ion characterization by linked-scan tandem mass spectrometry, 4:327
- Ethidium bromide, interactions among polythymine oligonucleotides, mercury ions and, ESI MS, 20:1834
- Ethyl structures, classical and nonclassical, tandem mass spectrometric investigation of $(\text{C}_2\text{H}_5)\text{X}^+$ ions ($\text{X}=\text{Cl}, \text{Br}, \text{I}$), 5:814
- Ethyl sulfate, forensic confirmatory analysis of, by LC-ESI-MSMS, 15:1644
- Ethyl thioglycolate, molecular ions of, metastable decomposition of, 8:801
- Ethylene
 catalyzed by Ti^+ in the gas phase, polymerization of, 3:464
 diacetylene radical cation reactions with, 15:114
 dynamic alignment investigated using linearly polarized femtosecond laser pulses, 17:1717
 selectivity of, methylene substitution reactions in a quadrupole ion trap, 3:39
- Ethylene glycol and its monomethyl and dimethyl ethers, functional group-selective ion-molecule reactions of, 4:97
- Ethylene oxide
 ion reactions with propenal and, selected ion flow tubes, 18:445
 selectivity of, methylene substitution reactions in a quadrupole ion trap, 3:39
- (Ethylendiamine)bis(2,2'-bipyridine)ruthenium(II) complexes, photo-oxidized, 8:713
- Ethylglucuronide, in urine, LC-ESI-MS-MS of, 15:188
- Ethylsilanes, gaseous, thermochemistry of, 9:597
 $[\text{Et}_2\text{S}^+\text{SEt}_2]^+$, structure and fragmentation of, 8:605
- Evaluating electron ionization mass spectral library search results, 7:313
- Evaluation of a dual ESI/atmospheric pressure chemical ionization source at low flow rates (50 $\mu\text{L}/\text{min}$) for the analysis of both highly and weakly polar compounds, 9:1196
- Evaluation of alkali and alkaline earth metal cation selectivities of lariat ether amides by ESI-MS, 14:1215
- Evaluation of alkali metal binding selectivities of caged azacrown ether ligands by MESI-MS, 11:884
- Evaluation of an improved general unknown screening procedure using LC-ES-MS by comparison with GC and HPLC-DAD, 14:14
- Evaluation of atmospheric pressure ionization techniques for the analysis of N-methyl carbamate pesticides by liquid chromatography mass spectrometry, 3:378
- Evaluation of automatically generated substructure identification rules from tandem mass spectra, 3:169
- Evaluation of carrier gases for use in FAIMS, 11:1125
- Evaluation of field desorption mass spectrometry for the analysis of polyethylene, 7:1070
- Evaluation of MALDI-MS for studying sec-butyllithium and nbutyllithium initiated ring-opening polymerization of hexamethylcyclotrisiloxane, 14:95
- Evaluation of matrix-assisted laser desorption ionization mass spectrometry for polymer characterization, 7:11
- Evaluation of matrix-assisted laser desorption ionization-time-of-flight mass measurement accuracy by using delayed extraction, 7:995
- Evaluation of metal complexation as an alternative to protonation for ESI of pharmaceutical compounds, 9:463
- Evaluation of noncovalent interactions between peptides and polyether compounds via energy-variable CAD, 14:1148
- Evaluation of photoinduced changes in stability constants for metal-ion complexes of crowned pirobenzopyran derivatives by ESI-MS, 14:1110
- Evaluation of role of multiple hydrogen bonding in offering stability to negative ion adducts in electrospray MS, 13:1360
- Evaluation of sample preparation techniques for mass measurements of PCR products using ESI-FTICR-MS, 13:338
- Evaluation of small mass spectrometer systems for permanent gas analysis, 13:1005
- Evaluation of steric and substituent effects in phenols by competitive reactions of dimethyl ether ions in a quadrupole ion trap, 6:627
- Evaluation of the true effect of experimental parameters on the reduction/oxidation processes observed in fast-atom bombardment/liquid secondary ion mass spectrometry, 4:482
- Everninomicin
 quadrupole ion trap multiple-stage MS analysis of, 13:1313
 structure of, by FAB-MS, 8:1134
- Evidence for a mechanism that involves secondary electron capture in the liquid secondary ionization mass spectrometry beam-induced dehalogenation of organic compounds, 7:1109
- Evidence for linkage position determination in cobalt-coordinated pentasaccharides by using ion trap MS, 9:1125
- Evidence for long-range glycosyl transfer reactions in gas phase, 13:325
- Evidence for the direct desorption of crown ether-metal ion complexes in liquid secondary ionization mass spectrometry, 5:638
- Evidence for the polymerization of ethylene catalyzed by Ti^+ in the gas phase, 3:464
- Evidence for thermal decomposition contributions to the mass spectra of chlorinated phenoxyacid herbicides obtained by particle beam liquid chromatography/mass spectrometry, 3:823
- Evidence of isomerization during ion isolation in the quadrupole ion trap, 3:680

- EX1, characterization of kinetics in by peak width analysis, H/D exchange MS, 17:1498
- Exact-mass measurements
from ESI-MS
FT-ICR MS automation and, 10:1166
on-line acquisition, and E-mailing of, 10:1212
of product ions for the structural elucidation of drug metabolites with tandem quadrupole orthogonal-acceleration TOFMS, 10:1305
using 7 tesla FT-ICR MS in a good laboratory practices-regulated environment, 10:1291
- ExB sector-field mass spectrometer, 12:619
- Exchange behavior, bimolecular hydrogen-deuterium, of protonated alkyl dipeptides in the gas phase, 6:466
- Exchange reactions, hydrogen/deuterium, complexation of protonated peptides with saccharides in gas phase decreases rates of, 6:1247
- Excitation, nonresonance, ion trap tandem mass spectrometry and selected ejection chemical ionization, 7:668
- Excitation frequencies
higher order quadrupole, in linear ion trap, 11:1016
of ions confined in a quadrupole field with quadrupole excitation, 11:10
- Excitation frequency, phase angle and, effect on dynamic CID in quadrupole ion trap, 18:749
- Excitation modes for Fourier transform-ion cyclotron resonance mass spectrometry, 4:433
- Excitation profiles, real-time, by SED FTICR-MS, 11:1009
- Excited ions, motion of, in Fourier transform ion cyclotron resonance cells, effect of frequency sweep direction on, 3:847
- Excited MN⁺, superelastic collisions of electrons with, 1:192
- Excited states of the tungsten hexacarbonyl dication, double-ionization energies to, 2:108
- Exhaustive enumeration, de novo peptide sequencing using, 17:1041
- β 1,4 Exoglucanase, *C. fimi*, xylobiose inhibitor binding to, MS/MS, 18:64
- Exoglycosidase digestion
neutral O-linked oligosaccharide structure and, 12:877
sequential, reversed-phase microbore liquid chromatography/electrospray mass spectrometry, complementary mobile phases, characterization of protein N-glycosylation by, 5:350
- Exonuclease digestion, photomodified oligodeoxynucleotide determination by, 12:1127
- Experimental and theoretical investigation of the main fragmentation pathways of protonated H-Gly-Gly-Sar-OH and H-Gly-Sar-Sar-OH, 14:1454
- Experimental and theoretical studies of diatomic gold halides, 13:485
- Experimental and theoretical studies of the basicity and proton affinity of SiF₄ and the structure of SiF₄H⁺, 10:848
- Experimental and theoretical study of gas phase interaction between ionized nitrile sulfides and pyridine, 14:241
- Experimental and theoretical study of the gas phase decomposition of monoprotonated peptide nucleic acids, 11:615
- Experimental design, of generic LC-MS, 13:2
- Experimental evaluation of a hyperbolic ion trap for Fourier transform ion cyclotron resonance mass spectrometry, 3:188
- Experimental parameters
pictograms for, 8:756
on the reduction/oxidation processes observed in fast-atom bombardment/liquid secondary ion mass spectrometry, evaluation of, 4:482
- Exploiting complementary nature of LC-MALDI-MS/MS and LC-ESI-MS/MS for increased proteome coverage, 14:971
- Exploring limits of bacterial identification, by ICM-MS, 12:49
- Explosives
ambient vapors detected by SESI, 20:287
electron capture ionization of, with a microflow rate particle beam interface, 7:753
low molecular weight, use of β -cyclodextrin in determining MWs of, MALDI-TOF MS, 17:189
nitro-aromatic and related compounds, femtosecond laser photoionization TOF MS, 20:419
RDX, PETN, and TNT, negative-ion formation in, by using the reversal electron attachment detection technique, 3:243
sampling on human skin using neutral desorption and EESI-MS, 20:719
- Exponential dilution gradient system for nanoscale liquid chromatography in combination with MALDI or nano-ESI-MS for proteolytic digests, 12:1205
- Expression profile analysis, Peptide Shifter enhancing separation reproducibility using, 18:1638
- Extended Cook's kinetic method, cysteine-polyalanine peptide gas-phase acidities determined by, 18:188
- Extended theoretical considerations for mass resolution in the resonance ejection mode of quadrupole ion trap mass spectrometry, 5:676
- External accumulation of ions for enhanced ESI-FTICR-MS, 8:970
- External ion accumulation of low molecular weight poly(ethylene glycol) by ESI-FTMS, 10:1298
- External ion injection in glow discharge-Fourier transform ion cyclotron resonance mass spectrometry, ion guide for, 4:387
- External ion source Fourier transform ion cyclotron resonance mass spectrometer, matrix-assisted laser desorption ionization on, high resolution end group determination of low molecular weight polymers by, 7:449
- External ion source Fourier transform ion cyclotron resonance mass spectrometry, electrostatic ion guide for, ion trajectories in, 6:936
- External ionization Fourier transform mass spectrometry, a low voltage ion transport system for, 5:398
- External secondary ion source for Fourier transform mass spectrometry, 1:208
- External source Fourier transform mass spectrometer, oligosaccharide cations produced in, "slow" metastable decomposition of, 4:210
- Extraction, delayed, evaluation of matrix-assisted laser desorption ionization-time-of-flight mass measurement accuracy by, 7:995
- Extraction and characterization of adenovirus proteins from sodium dodecylsulfate polyacrylamide gel electrophoresis by MALDI-MS, 11:356
- Extractive electrospray ionization mass spectrometry, sampling explosives on skin, using neutral desorption and, 20:719
- F
- F₂-isoprostanes, four regioisomers of, formed by free radical oxidation of arachidonic acid, mass spectrometric analysis of, 7:490
- FAB. *See* Fast atom bombardment
- FAB-MS. *See* Fast-atom bombardment-mass spectrometry
- FAB MS/MS. *See* Fast-atom bombardment tandem mass spectrometry
- FACSS Meeting, 22nd annual, Mass Spectrometric Highlights of, 7:507
- Factors governing solubilization of phosphopeptides retained on ferric NTA IMAC beads and their analysis by MALDI-TOF-MS, 13:1042

- Factors influencing determination of analyte ion surface partitioning coefficients in electrosprayed droplets, 12:1002
- Factors influencing the analytical performance of an ASGDI source as revealed via ionization dynamics modeling, 1315
- Factors that influence the observed fast fragmentation of peptides in matrix-assisted laser desorption, 7:225
- FAIMS. *See* Field asymmetric wave form ion mobility spectrometry; Field asymmetric waveform ion mobility spectrometry
- FAIMS. *See also* Field asymmetric waveform ion mobility spectrometry; High field asymmetric waveform ion mobility spectrometry
- FAIMS/FAIMS, coupled to TOFMS, ion trapping in, 12:1320
- False positive identification, risk of, using multiple ion monitoring in qualitative MS, 17:823
- False positives and the detection of cyclodextrin inclusion complexes by electrospray mass spectrometry, 6:437
- Familial adenomatous polyposis, diagnostic PCR and ESI-MS in, 8:737
- Fast, very fast, and ultra-fast gas chromatography-mass spectrometry of thermally labile steroids, carbamates, and drugs in supersonic molecular beams, 7:737
- Fast analysis of drugs in a single hair, 9:1311
- Fast atom bombardment
- of alkali-metal-carboxylate radical, observation of, 6:1252
 - coaxial, the application of normal phase liquid chromatography/mass spectrometry by using, 3:575
 - collision activated dissociation-B/E linked-scan mass spectrometry of aminoethyl-triphenylphosphonium derivatives of amino acids, charge-remote fragmentation during, 3:769
 - collision activated dissociation-mass spectrometry, characterization of diacylglycerolphosphocholine molecular species by, a general method not sensitive to the nature of fatty acyl groups, 3:71
 - collision activation of a series of homoconjugated octadecadienoic acids with, 1:183
 - collision-induced dissociation of prostaglandins and thromboxanes, some examples of charge remote fragmentation, 1:325
 - combined with mass spectrometry for characterization of polycyclic aromatic hydrocarbons, 1:405
 - combined with tandem mass spectrometry, determination of structure and properties of modified chlorophylls using, 1:72
 - conditions, electronic excitation under, by using a liquid matrix, M⁺ ion formation and, 6:114
 - continuous flow, capillary zone electrophoresis, identification of deoxynucleoside-polyaromatic hydrocarbon adducts by, 3:757
 - continuous flow, with packed microcolumns, a comparison of precolumn *versus* coaxial matrix delivery, 1:312
 - of digoxin and related cardiac glycosides, ions formed by, high resolution mass spectrometry and tandem mass spectrometry analysis of, mechanistic considerations of the protonation and fragmentation of highly functionalized molecules in fast atom bombardment, 1:455
 - of glycerol, background ions produced by, origins and structures of, 5:72
 - highly functionalized molecules in, protonation and fragmentation of, mechanistic considerations of, high resolution mass spectrometry and tandem mass spectrometry analysis of the ions formed by fast atom bombardment of digoxin and related cardiac glycosides, 1:455
 - and tandem mass spectrometry, of macrolide antibiotics, 5:151
 - by tandem mass spectrometry/liquid secondary ion mass spectrometry, beam-induced reactions occurring in, characterization of, 4:312
- Fast atom bombardment-desorbed (M + H)⁺, (M + Cat)⁺, and (M–H)[–] ions of aromatic beta-hydroxyoximes, charge-remote fragmentation and the two-step elimination of alkanols from, 4:819
- Fast-atom bombardment gun, saddle field, flux from, determination of, 5:100
- Fast atom bombardment-induced condensation of glycerol with ammonium surfactants
- regioselectivity of the adduct formation, 3:318
 - time dependence of mass spectra and tandem mass spectra, 5:92
- Fast atom bombardment mass spectra
- improving, the influence of some controllable parameters on spectral quality, 1:149
 - methods for the identification of carcinogen-nucleoside adducts, development of, 3:360
- Fast-atom bombardment mass spectra of azo dyes, effect of tautomerization on, 5:407
- Fast atom bombardment mass spectrometry
- of brassinosteroid analogs, 8:270
 - continuous-flow
 - an integral probe for, 3:198
 - of oligonucleotides, 2:157 - detection limits for, improved, a study of time-dependent desorption using a model pulsed bombardment ionization method, 3:637
 - differentiation of isomeric C8- and N²-deoxyguanosine adducts of 2-acetylaminofluorene by, 5:58
 - of everminomicin structure, 8:1134
 - high resolution, positive and negative ion, reference compound for, ultramark 1621 as, 3:842
 - injection of reagent ions into the selvedge region in, 4:259
 - ionization mechanism in, 8:337
 - of meso-phenyl-substituted tetrabenzoporphyrin adduct ions, 8:62
 - metal chelation of dinucleotide analogs in the gas phase by, 7:42
 - m-Nitrobenzyl alcohol electrochemistry in, 3:113
 - of mycobacterial glycopeptidolipid antigens, structural characterization by charge remote fragmentation, 3:656
 - negative ion
 - analysis of long-chain fatty acyl coenzyme A thioesters by, 5:416
 - of polyanionic analytes, enhancing sensitivity in, a class of cationic matrix additives for, N-alkylnicotinium halides, 5:935 - probing high order structure of proteins by, 5:19
 - redox processes by ruthenium(II) polypyridine complexes induced by, 4:306
 - salen complex analysis by, 8:996
 - sensitivity of, effect of varying counterions on, 1:233
 - a study of the solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes by, synthesis and characterization of molecular complexes, 4:255
 - a study of the solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes by 1, generation of molecular signals, 4:249
 - tandem, negative ion, to determine sulfate and linkage position in glycosaminoglycan-derived disaccharides, 3:797
 - time dependence of, 8:337
- Fast-atom bombardment/mass spectrometry, multisample probe for, 4:504
- Fast-atom bombardment mass spectrometry and low energy col-

- lision-induced tandem mass spectrometry, of tauroconjugated bile acid anions, 6:588
- Fast-atom bombardment mass spectrometry of enzymatic digests, recombinant human macrophage colony stimulating factor, disulfide linkage and disulfide bond scrambling in, characterization of, 6:638
- Fast-atom bombardment of the cyclic acetals, evidence indicating the predominant involvement of condensed-phase processes in ionization, 4:493
- Fast atom bombardment tandem mass spectrometry
analysis of phosphatidylcholine isolated from heart mitochondrial fractions by, evidence of incorporation of monohydroxylated fatty acyl moieties, 7:50
of [¹³C]arachidonic acid labeled phospholipid molecular species, 5:144
determination of the anomeric configuration of glycosyl esters of nucleoside pyrophosphates and polyisoprenyl phosphates by, 7:541
employing ion-molecule reactions for the differentiation of phospholipid classes, 2:470
isomer differentiation in 7/12-dimethyl[a]anthracene-pyridine adducts by, 2:256
of lipophilic phosphotriesters, 10L975
sialyl anomers distinction on ESI-MS/MS and, 20:394
- Fast direct current pulse, as pump and a laser probe, determination of ion frequencies in a quadrupole ion trap by, 5:29
- Fast excitation CID in quadrupole ion trap MS, 14:785
- Fatty acid esters, deprotonated phenol, ion-dipole complex formation from, evidence by using gas-phase labeling combined with tandem mass spectrometry, 4:343
- Fatty acid methyl esters
molecular ion derived from, low-energy tandem mass spectrometry of, a novel method for analysis of branched-chain fatty acids, 4:223
O⁻ and OH⁻ chemical ionization of some, 5:553
- Fatty acid picolinyl esters, GC-MS determination of, 14:58
- Fatty acids
aminoethyl-triphenylphosphonium derivatives of, FAB-CAD-B/E linked-scan mass spectrometry of, charge-remote fragmentation during, 3:769
branched-chain, analysis of, a novel method for, low-energy tandem mass spectrometry of the molecular ion derived from fatty acid methyl esters, 4:223
CID spectra of, fragment ions in, charge-associated mechanism in, 16:280
lithiated, CRF on TOF-TOF MS, 18:1967
long-chain unsaturated, double-bond position assigned by multistage LIT-MS with ESI, 19:1678
methyl esters of, isotopomer abundances of, fragmentation effects on, 10:430
monounsaturated, double bond migration during EI MS, 20:1997
modification in ghrelin, ECD identification and localization of, 13:1443
polyhydroxy unsaturated, electrospray ionization and low energy tandem mass spectrometry of, 7:140
polyunsaturated, methyl esters of, CI-MS/MS of, 10:1253
unsaturated, lithium adducts of, ESI-MS of using CAD, 10:600
- Fatty acids cationized with alkaline earth metal ions, charge remote fragmentation of, 1:320
- Fatty acyl coenzyme A thioesters, long-chain, analysis of, by negative ion fast-atom bombardment mass spectrometry and tandem mass spectrometry, 5:416
- Fatty acyl groups, a general method not sensitive to, characterization of diacylglycerolphosphocholine molecular species by FAB-CAD-MS/MS, 3:71
- Fatty acyl moieties, monohydroxylated, incorporation of, evidence of, analysis by fast-atom bombardment tandem mass spectrometry of phosphatidylcholine isolated from heart mitochondrial fractions, 7:50
- Fe⁺, cysteine-containing peptide interaction with, gas phase studies of, 9:1285
- Fe, Al, Co, Cu, ScO, M(acetone)₂⁺, collision-induced and infrared multiphoton dissociation studies on, in the gas phase, 4:135
- Fe(II) effect, tributyl phosphate on mineral surfaces, static secondary ionization mass spectrometry analysis of, 6:165
- Fe(III) and Mn(III) tetraarylporphyrin complexes, solution and gas phase chemistry of, by fast-atom bombardment mass spectrometry
generation of molecular signals, 4:249
synthesis and characterization of molecular complexes, 4:255
- Fe(III)-heme cation, gaseous, binding to biological molecules, 16:589
- Femtomole, levels of, peptide, detection of, by tandem magnetic sector/reflection time-of-flight mass spectrometry and matrix-assisted laser desorption ionization, 2:91
- Femtosecond and anosecond laser pulse ionization efficiency, 6:1059
- Femtosecond laser pulse ionization efficiency, for multiphoton ionization of molecules, 6:1059
- Femtosecond laser pulses, linearly polarized, dynamic alignment of C₂H₄ investigated using, 17:1717
- Fermentation broths, rapid analysis of antibiotic-containing mixtures from, by using liquid chromatography-electrospray ionization-mass spectrometry matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry, 7:1227
- Ferric chelates, in agricultural matrices, as fertilizers, LC-ESI-MS, 18:37
- Ferric NTA IMAC beads, solubilization of phosphopeptides retained on, MALDI-TOF-MS of, 13:1042
- Ferrichrome, surprising stability of a cyclic peptide-FeIII complex revealed by MS, 8:1070
- Ferrocene, protonation of, in the gas phase, 5:274
- Ferrocene cation, gas-phase, photodissociation of, 2:97
- Ferrocenedicarboxylic acid, ESI-MS of, 10:557
- Ferrocenoyl piperazide, using LC-EC-S, 15:1727
- Fertilizers, determination in agricultural matrices of synthetic ferric chelates as, LC-ESI-MS, 18:37
- Ferulic acids, deprotonated monomeric and dimeric, gas phase basicities of, 10:1111
- Fetuin, characterization using hybrid linear ion trap/FT-IRC MS, 17:168
- FF-MAS, quantification by LC-APPI-MS-MS in vitro, 15:1216
- Fibrinopeptide B
fragment intermediate lifetimes measured by DR ECD, 17:1605
negative ion MALDI-TOFMS post-source decay calibration of using, 9:540
- Field asymmetric waveform ion mobility spectrometry (FAIMS) analyzers for
design and operation of, 16:2
domed, compensation voltage peak shapes using, 16:948
characterization of temperature-controlled system, 18:1653
for gas flow profile and asymmetric waveforms, 16:1447
improving sensitivity with slit interfaces, 20:1768
of leucine enkephalin, 10:492
modeling resolution and sensitivity of, 15:1487
optimal waveforms for, 19:1286
planar, resolving power and sensitivity for, 18:1672
in separation of leucine and isoleucine anions, 10:1279
- Field desorption mass spectrometry, for the analysis of polyethylene, evaluation of, 7:1070

- Field-modulated selective ion storage in quadrupole ion trap, 6:928
- Field-portable
high speed GC/TOFMS, 12:648
and miniature mass spectrometers, from 12th Sanibel Conference, 12:617
- Field strength, of tryptic peptides, in separation by IM MS, 16:158
- Filter diagonalization method, use in space charge induced frequency modulations, FT-ICR-MS, 17:836
- Filter disks, nylon 66, cyclic polyamide oligomer contamination from, LC/MS, 17:652
- Fingerprint patterns, from laser-induced azido photochemistry of spin-labeled photoaffinity ATP analogs in MALDI-MS, 10:1337
- Fingerprinting
mammalian cell line identification using MALDI-TOF and LC-ESI-MS/MS, 17:490
of peptides
mass assignment from, 14:943
origins of, 14:931
- Fingerprinting proteins coupled with polymers by mass spectrometry, investigation of polyethylene glycol-conjugated superoxide dismutase, 6:478
- Finnigan TSQ-700 electrospray inlet system, iron carboxylate oxygen-centered-triangles detected during electrospray use of organic acid modifiers and, 6:1226
- First observation of 3+ oxidation state for a [4Fe-4S] metalloprotein, ESI-FTICR-MS of, 12:817
- First-stage pressure to reduce interference effects in inductively coupled plasma-mass spectrometry, selection of, 3:128
- Fitness for purpose, of MS methods, 14:528
- Five-membered ring H shift in the isomerization of $\text{CH}_3 + \text{HCH}_2\text{CH}_2$ to $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + \text{H}$ by ab initio theory, preference for ion-neutral complex-mediated pathway over, 6:1037
- FKBP-FK506 and FKBP-rapamycin complexes, noncovalent, detection of by capillary electrophoresis-mass spectrometry and capillary electrophoresis-tandem mass spectrometry, 6:85
- Flame, $\text{C}_2\text{H}_4/\text{N}_2\text{O}/\text{Ar}$, triple quadrupole mass spectrometry as applied to flame diagnostics, 4:152
- Flame diagnostics, triple quadrupole mass spectrometry as applied to, study of the $\text{C}_2\text{H}_4/\text{N}_2\text{O}/\text{Ar}$ flame, 4:152
- Flame-generated C_{10} to C_{160} polycyclic aromatic hydrocarbons, characterization of, by atmospheric-pressure chemical ionization mass spectrometry with liquid introduction via heated nebulizer interface, 7:276
- Flavonoid, transition metal complexes of, threshold dissociation and molecular modeling of, 16:139
- Flavonoid aglycones, LC-ESI-MS/MS of, 12:707
- Flavonoid diglycosides, tunable metal-ligand complexation for, ESI-MS, 18:422
- Flavonoid glycoside isomers, differentiation with metal complexation and ESI-MS, 14:1437
- Flavonoid glycosides, characterization of, by CAD of metal complexes, 12:537
- Flavonoid O-diglycosides, glucose loss from, 11:136
- Flavonoids
determining glycosylation site of, by metal complexation and MS-MS, 15:1287
evaluation of flavonoid-DNA complexes ESI-MS, 19:914
gas phase acidities of, 15:848
licorice, quantification in rat plasma, 18:778
plasma determination of terpenoids and, from *G. biloba*, efficiency of ESI, 19:445
- UGT1A1 regioselectivity towards, metal complexation and MS/MS, 19:246
- Flight time, of large MALDI ions, 8:1094
- Floating sampling interface, electrically, inductively coupled plasma mass spectrometry with, 4:733
- Flow analysis, nanoliter, of protein digests, 8:703
- Flow injection electrospray ionization mass spectrometry, of crude cell extracts for high throughput bacterial identification, 13:118
- Flow injection with chemical reaction interface-isotope ratio mass spectrometry, alternative to off-line combustion for detecting low levels of enriched ^{13}C in mass balance studies, 10:153
- Flow rate, in nano-ESI-MS, analyte ion signals and, 14:492
- Flowing afterglow technique for ion-molecule reaction studies, early development of, a personal history of, 3:479
- Fluid injection, direct supercritical, collision-induced dissociation mass spectrometry of nonionic surfactants following, 3:150
- Fluidics, induction-based, real-time dispense event and volume verification, MALDI MS, 20:1064
- Fluorescence
analysis using fluorescent dyes, MALDI MS, 19:1054
trapped gas-phase ions, effect of buffer gases on yield, 17:855
of trapped ions, measurement with zero background detection, 13:696
- Fluorescence resonance energy transfer (FRET), in gas phase ions, 16:1481
- Fluorinated benzene molecules, double-ionization energies of, 4:513
- Fluorinated chloroformates, comparison of, 16:803
- Fluorinated phenols, APCI-MS of, 10:1157
- Fluorinated self-assembled monolayer surfaces, chemical modification of by low energy reactive ion bombardment, 6:187
- Fluorobenzene and 1,4-Difluorobenzene, on the acidity of, gas-phase reactions between O^- and $\text{H}_6\text{H}_8\text{F}$, 4:73
- Fluorocarbons, SAMs of, methyl cation neutralization, 13:1151
- Fluorophenyl anions, gas-phase H/D exchange of, 10:840
- Fluorophenyl n-propyl ethers, chemical ionization of, 9:121
- Fluorotrimethylsilane, anionic nucleophile affinities of, 16:697
- 5-Fluorouracil, derivatization of, for LC/MS determination, 9:971
- Fluorovinyl anion, characterization of, reactions of strong bases with vinyl fluoride formation and, 3:91
- Flux from a saddle field fast-atom bombardment gun, determination of, 5:100
- Focused ion beam (FIB) secondary ion mass spectrometry (SIMS), of elemental distribution in fluorinated amorphous carbon thin films, 16:126
- FOG7. *See* 5-Oco-7-glutathionyl-8,11,14-eicosatrienoic acid.
- Folates, microbial, isotope ratio-based profiling, LC-MS/MS, 18:898
- Folding, peptide ions, kinetics measured by IR activated DR ECD, 19:780
- Folding analysis, of calcitonins, by ESI-MS with H-D exchange, 15:1556
- Food products, multiresidue screening in food and feed matrices, Oribtrap MS, 20:1464
- Forensics, field-portable GC/MS in, 12:683
- Formaldehyde and formalimine, lithium atom complexes with, stability of, 12:1229
- Formation and characterization of iron-oligonucleotide complex with MALDI FT-ICR MS, 10:941
- Formation and characterization of the sulfur-containing distonic radical anion, $\text{CH}_2\text{-S-CH-CH}_2$, in the gas phase, 6:71
- Formation and decomposition of chloride adduct ions $[\text{M} + \text{Cl}]^-$, in negative ion ESI-MS, 11:933

- Formation and decomposition of water clusters as observed in a triple quadrupole mass spectrometer, 9:814
- Formation and emission of tetraalkylammonium salt molecular ions sputtered from a gelatin matrix, 4:930
- Formation and fate of ion pairs during MALDI analysis, anion adduct generation as indicative tool to determine ionization processes, 13:1218
- Formation and fragmentation of gas-phase ion-molecule complexes of transition-metal ions with organic molecules containing two functional groups, 3:451
- Formation of alkali-metal adduct ions in ESI-MS of peptides and proteins, 11:967
- Formation of lithiated adducts of glycerophosphocholine lipids facilitates their identification by ESI-MS/MS, 9:516
- Formation of M^+ ions and electronic excitation under fast-atom bombardment conditions by using a liquid matrix, 6:114
- Formation of molecular radical cations of enkephalin derivatives via CID of ES-generated copper(II) complex ions of amines and peptides, 12:1114
- Formation of peptide aggregates during ESI, 12:1020
- Formation of the 3-pentanone ion from ionized propyl propanoate through ion-neutral complexes, 3:183
- Formic acid, protonated, decomposition of, 15:982
- Formulation of matrix solutions for use in MALDI of biomolecules, 9:166
- 5-Formyluracil, oligodeoxynucleotide fragmentation with, 13:1190
- Forthcoming events, 1:350
- Four decades of structure determination by MS, from alkaloids to heparin, 13:1254
- Four-sector mass spectrometer
in in-line surface-induced dissociation device, design and performance of, 6:257
interface for, with dual purpose collision cell, high transmission at low to intermediate energies, 6:175
- Four-sector tandem mass spectrometer
interface region of, enhancement of ion transmission at low collision energies via, 5:460
photon-induced dissociation with, 1:107
- Fourier-transform electrospray instrumentation for tandem high-resolution mass spectrometry of large molecules, 4:557
- Fourier transform (FT) mass spectrometry, top-down, in kinetic characterization of multiple protein forms, 16:1052
- Fourier transform ion cyclotron resonance (FTICR). *See also* Electrospray ionization-FTICR; Liquid chromatography-FTICR.
in analysis of metal-induced conformational mutations, 12:1272
collision-induced ion ejection in, 16:422
detection of multiphoton ionization spectroscopy, 7:1018
hydration of protonated 18-crown-6 by, 10:862
internal glow discharge mass spectrometry, 7:923
capacitive-coupling gated-trapping MALDI and, 8:195
of lanthanide(III) porphyrin-phthalocyanine heteroleptic sandwich complexes, 8:161
space-charge effects on, 8:319
sustained off-resonance in, 8:86
trapped negative ion cooling in, 8:793
motion of excited ions in, effect of frequency sweep direction on, 3:847
of multiply charged aggregates of small singly charged peptides formed by electrospray ionization, 10:347
in simulation of organic environment on Saturn's moon (Titan), 16:850
single crystal characterization by, 10:352
in study of O- reactions with methyl benzoate, 12:694
study of the gas-phase acidities of germane and methylgermane, bond dissociation energy of germane, 4:54
two-dimensional Coulomb-induced frequency modulation in, a mechanism for line broadening at high mass and for large ion populations, 4:909
- Fourier transform ion cyclotron resonance (FTICR) tandem mass spectrometry
H-D exchange integration with, in monitoring intermolecular migration during collisional activation, 15:639
identifying class I HLA-H*0201 peptide from vaccinia virus, 16:1812
- Fourier transform ion cyclotron resonance mass spectrometer-low magnetic field internal electrospray, a high performance, 7:915
- Fourier transform ion cyclotron resonance mass spectrometric automation of, 10:1166
-in-cell matrix-assisted laser desorption, 7:1026
capillary electrophoresis, with sustained off-resonance irradiation for the characterization of protein and peptide mixtures, 5:894
charged phenyl radical and biradical synthesis by laser photolysis in, 9:1135
in detection of small Ca^{2+} -induced conformational changes in the regulatory domain of human cardiac troponin C, 10:703
ESI with, instrument design and initial results of, 9:692
exact-mass measurement using, 10:1291
high-frequency, 4:177
in a high homogeneity 25 tesla resistive magnet, 10:265
high performance, via a single trap electrode, 6:812
high pressure chemistry in, split-pair magnet in, 10:918
hyperbolic ion trap for, experimental evaluation of, 3:188
improved isotopic abundances in, 9:799
laser desorption, ionization, collision-induced dissociation in, improved efficiency of, selective parent ion axialization for, 5:900
laser desorption in, 10:1105
matrix-shimmed cubic penning trap for, 10:759
probing trapped ion energies via ion-molecule reaction kinetics, 3:727
quadrupolar excitation in, simplified application of, 6:448
quantitation of ion abundances in, 9:1204
sensitivity of, 9:333
in structural validation of saccharomycins, 10:1285
for trapping and gas phase hydrogen/deuterium exchange of peptide ions, 9:1012
- Fourier transform ion cyclotron resonance mass spectrometric detection, a glow discharge ion source with, 3:122
- Fourier transform ion cyclotron resonance mass spectrometry (FTICR-MS), 12:243. *See also* Electrospray ionization-Fourier transform ion cyclotron mass spectrometry; Electrospray ionization-Fourier transform ion cyclotron resonance mass spectrometry
of the 3,5-didehydrophenyl cation, 12:258
analysis of complex peptide mixtures using differential MS and, 18:226
APPI proton transfer investigated by, 18:1682
artificial neural networks for ppb mass accuracy in, 20:1303
 β -amino acid diastereochemical differentiation, 18:1038
biomolecule analysis using IR-MALDESI coupled to, 20:667
central trapping ring in, 12:30
in composition-based sequencing of peptides, mass determination by, 15:703
concentric tube vacuum chamber for high magnetic field, 4:168
cryogenic, first signal on, 18:2090

- determination of unique AA substitutions, peptide mass mapping, 17:508
- disulfur monoxide-water charged complexes, isotope exchanges, CI-CAD, TQ, 18:1664
- DREAMS in, increased proteome coverage with, in peptide abundance measurements, 13:954
- dynamic ion cooling technique for, 12:1169
- ECD efficiency in, 16:1060
- ECD implementation progress in, 19:762
- eicosanoid characterization by ESI-FTICR, 19:569
- electrically-compensated trap designed to 8th order for, 19:1281
- electrodynamic ion funnel for, 11:19
- electron capture dissociation by, 12:245
- endgroup determination of synthetic polymers by, 11:536
- excitation modes for, 4:433
- external ion source, matrix-assisted laser desorption ionization on, high resolution end group determination of low molecular weight polymers by, 7:449
- fast rpLC reduces back exchange in, 20:520
- filter diagonalization method in space charge induced frequency modulations, 17:836
- in H/D exchange of positively charged mononucleotides, 12:268
- high magnetic field electrospray ionization, trapping and detection of ions generated in, 3:615
- high-resolution, photosystem II structure and dynamics by, 19:1004
- higher accuracy of with multiply charged ions, 11:416
- identification of analytes with two basic functional groups, 20:1251
- improved ion trapping and detection capabilities, 20:755
- improved, of large biomolecules, 4:190
- incorporation of flared inlet capillary tube, 17:772
- of in-source H/D exchange and ion molecule reactions, 13:308
- in-trap cleanup in, in competitive binding to oligopeptide-binding protein OppA, 11:1023
- ion charge states of peptides and proteins, 17:903
- isotope-labeled cross-linkers and, analysis of protein/peptide complex, 17:1100
- kinetic energy orifice as probe of metastable dissociation in, 11:1118
- LIAD with CID in, peptide sequencing using, 18:525
- low-noise, wide band preamplifier for, 18:2233
- of macrophage colony-stimulating factor b and derivatives, 11:237
- MALDSI coupled to, detection of multiply-charged proteins and peptides by, 17:1712
- mass-based classification of peptides, 19:1808
- mass measurement errors in, 13:99
- nanoLC
 - high-throughput comparative proteomics using, 18:1332
 - parallel processing of large data sets from, 18:152
- nanoLC-FTICR MS, ruthenium arene complex reactions with glutathione, 19:544
- in *Neisseria gonorrhoeae* lipooligosaccharide characterization, 13:571
- observation of heterogeneous gene products by, 19:103
- oversulfation of bovine aorta chondroitin sulfate and GAG analyzed using nano-ESI-Q-TOF-MS/MS and, 18:179
- peak interference in, 13:387
- of poly(ethylene glycol), 12:565
- peptide fragment ion structures, analyzed by SORI-HDX in, 19:1771
- polyphosphoester characterization by, 20:2238
- protein complexes, effects of single amino acid substitution on dissociation in gas phase, 18:688
- in protein identification from phenylisothiocyanate-derivatized electrosprayed tryptic peptides, 12:288
- protein identification via SID in, 17:700
- of proton-bound heterodimers and homodimers of amides and amines, in gas phase, 13:1273
- quantification of isotopic pairs, proteins, 19:973
- in resolution, identification, and screening of non-covalent complexes of Hck Src homology 2 domain receptor and ligands from a 324-member peptide combinatorial library, 13:1162
- SED with, real-time excitation profiles by, 11:1009
- signal filtering method for quantification and noise discrimination, 20:1087
- small, characterization of long-chain carboxylic esters with $\text{CH}_3\text{OBOCH}_3^+$ in, 7:1138
- SORI-CAD and, direct dynamic combinatorial library screening, 17:1074
- SORI-CID, hawthorn leaf α -glucosidase inhibitors characterization using LC-DAD-MS and, 20:1496
- spontaneous loss of coherence catastrophe, 20:247
- of stereoselective dissociations of hexosamine monosaccharides, 12:517
- in study of melittin bound to phospholipid micelles, 12:1247
- sub part-per-million mass accuracy, 17:1681
- trapped-ion cell with improved harmonicity, 19:586
- ultrahigh resolution, in He@C60 detection and quantitation, 13:1349
- in validation of exact mass measurements, 16:1100
- zwitterion formation in gas-phase H/D-exchange with CH_3OD , 18:512
- Fourier transform ion cyclotron resonance mass spectrometry-glow discharge, external ion injection in, ion guide for, an improved, 4:387
- Fourier transform ion cyclotron resonance mass spectrometry of aromatic hydrocarbons, ammonia laser desorption/chemical ionization with ammonium bromide, 2:299
- Fourier transform ion cyclotron resonance mass spectroscopy, remeasurement at high resolving power, 6:564
- Fourier transform ion cyclotron resonance-MS
 - of ammonium ion-mediated resorcinarene capsules, 14:143
 - infrared multiphoton dissociation, in determining activation energy for noncovalent gas phase peptide unimolecular dissociation, 14:1282
 - with nanoLC/microelectrospray ionization and MALDI: analytical performance in peptide mass fingerprint analysis, 553
 - in probing phosphorylated carbohydrate isomeric differences, 323
 - in quantification of phosphorylated positional isomers, 14:916
- Fourier transform ion cyclotron resonance precise mass difference measurements, coulomb mutual interaction and image charge in, systematic errors originating from, analysis and estimation of, 4:855
- Fourier transform-ion cyclotron resonance spectra, relative peak heights in, space charge effects on, 3:734
- Fourier-transform/laser-induced thermal desorption mass spectrometry, for surface analysis, deflection of laser-produced ions in, 5:583
- Fourier transform laser microprobe mass spectrometry, for molecular identification of inorganic compounds, 9:482
- Fourier transform mass spectrometer
 - external source, oligosaccharide cations produced in, "slow" metastable decomposition of, 4:210
 - methyl guanine isomer distinction by hydrogen/deuterium exchange using, 4:296
 - surface-induced dissociation in, iron pentacarbonyl ions undergoing, energy deposition in, 3:445

- Fourier transform mass spectrometry (FTMS). *See also* Matrix-assisted laser desorption/ionization Fourier transform mass spectrometry (MALDI-FTMS)
- for analysis of complex organic mixtures, 1:431
 - analysis of hydrocarbon dendrimers by, 5:731
 - collisionally-activated dissociation and 193-nm photodissociation of peptide ions in, efficiency of, 1:288
 - electric ion resonance cell design for, 10:1000
 - external ionization, a low voltage ion transport system for, 5:398
 - an external secondary ion source for, 1:208
 - gated trapping strategy in, 12:296
 - high mass detection in, ion cloud manipulation using the radiofrequency-only mode as an improvement for, 4:186
 - high pressure, ganglioside analysis by, 13:402
 - high pressure trapping in, a radiofrequency-only-mode event, 3:590
 - of hydrocarbon polymers, 8:225
 - ion excitation for, linear response theory of, 2:483
 - ionization, ion cyclotron, reagent ions in, potassium halide adducts as, 3:301
 - of ionized ethylated acetone and deuterated analogs, 8:327
 - isomer differentiation in, 193 nm laser photoionization and photodissociation for, 1:361
 - for large targets with diverse surfaces, MALDI ion source for, 15:128
 - laser desorption
 - analysis of copolymers by, 1:66
 - analysis of hydrocarbon polymers by, silver nitrate chemical ionization for, 4:453
 - matrix-assisted structural characterization of normal and modified oligonucleotides by, 2:402
 - of LHRH-metal ion complexes, 10:19
 - of long-range glycosyl transfer reactions, 13:325
 - low-voltage on-resonance ion selection for, 9:931
 - peak position determination in, 15:457
 - for peptides, UV matrix-assisted laser desorption, investigation for, 2:22
 - poor high mass performance in, a mechanism for, 3:460
 - pressure-limited SORI-CAD in, 10:15
 - surface-induced dissociation of peptide ions in, 1:413
 - tandem, with Californium-252 plasma desorption, 1:427
 - verification of $(M + Fe-5H)^{2+}$ peptide ion composition by, 11:83
- Fourier-transform mass spectrometry trapped ion cells, elongated, trapping motion in, direct observation of, 1:351
- Fourier transform-MS
- in analysis of whole cell bacteria, 14:1306
 - in comparison of laser desorption and MALDI for ruthenium and osmium trisbipyridine complexes, 14:393
- Fragment ion type
- in the collision-induced dissociation spectra of peptides, formation of α_2-16 ions, 4:424
 - new, in the collision-induced dissociation spectra of peptides, identification of, formation of α_2-16 ions, 4:424
- Fragment ions, $[M-35]^+$, novel, in the mass spectra of 2-alkyl-4,6-dinitrophenols, characterization of, sequential ortho effects, 2:69
- Fragmentation
- of 1- and 3-methoxypropene ions, another part of the $C_4H_8O^+$ potential energy surface, 3:99
 - of 4-azasteroids, 12:385
 - of ACE inhibitors, MS in, 15:1300
 - of alkenamides, by collisional activation in triple quadrupole mass spectrometer, 15:1462
 - of amidinated peptide ions, 15:158
 - analysis of bradykinin by ^{252}Cf -plasma desorption mass spectrometry, 2:379
 - analyte, Influence of matrix on, atmospheric pressure MALDI, 17:1005
 - of b2 ions in peptide mass spectra, 11:427
 - of β -substituted *meso*-tetraphenylporphyrins induced by high energy collisional activation, 13:1427
 - of cationized phosphotyrosine containing peptides by atmospheric pressure MALDI-ion trap spectrometry, 14:581
 - characteristics of glutathione conjugates activated by high-energy collisions, 3:815
 - of charge-derivatized peptides, 12:399
 - charge remote
 - of fatty acids cationized with alkaline earth metal ions, 1:320
 - some examples of, fast atom bombardment and collision-induced dissociation of prostaglandins and thromboxanes, 1:325
 - structural characterization by, fast atom bombardment mass spectrometry of mycobacterial glycopeptidolipid antigens, 3:656
 - "charge-remote," implications of the charge site in, 3:776
 - charge-remote
 - during FAB-CAD-B/E linked-scan mass spectrometry of aminoethyl-triphenyl-phosphonium derivatives of fatty acids, 3:769
 - in hybrid (BEqQ) spectrometer to determine isotopic purity in selectively polydeuterated surfactants, 1:85
 - and the two-step elimination of alkanols from fast atom bombardment-desorbed $(M + H)^+$, $(M + Cat)^+$, and $(M-H)^-$ ions of aromatic beta-hydroxyoximes, 4:819
 - chemical, mass spectrometric, electron capture negative, and calculated internal energies for polychlorinated biphenyls, correlation between, 5:44
 - of conjugate bases of esters derived from multifunctional alcohols including triacylglycerols, 6:498
 - of cyclopropyltetrahydropyridine, 8:724
 - competing processes of, in MS-MS of heparin-like glycosaminoglycans, 15:1534
 - CSR for ionization mode and, 12:571
 - of deprotonated ions of oligodeoxynucleotides carrying 5-formyluracil or 2-aminoimidazolone, 13:1190
 - of deprotonated N-benzolypeptides, 15:446
 - of deprotonated peptides, 12:1
 - phenylalanine and, 13:1242
 - of derivatized glutamine, 16:1192
 - of derivatized N-linked glycans, 12:926
 - of derivatized N-linked carbohydrates, 11:900
 - of diacyl glycerophosphatidic acids
 - charge-driven, 11:797, 11:892
 - charge-remote, 11:892
 - dissociation, APCI low energy collision-induced, of protonated ortho silicates, McLafferty or ion-neutral complex rearrangements, 5:544
 - efficiency in linear ion trap, 16:1342
 - of electrospray-produced oligodeoxynucleotide ions adducted to metal ions, 12:550
 - gas-phase, $ArPd(PPh_3)_2I$ complexes, ESI-FTICR-MS, 17:1582
 - of glycosidic bonds, in gas phase, direction by non-covalent complexes, 15:715
 - high-energy, of protonated peptides, using a BEqQ hybrid mass spectrometer, 4:470
 - of highly functionalized molecules in fast atom bombardment, mechanistic considerations of, high resolution mass spectrometry and tandem mass spectrometry analysis of the ions formed by fast atom bombardment of digoxin and related cardiac glycosides, 1:455

- induced by ion-atom reactions, 8:587
- initial, of oligodeoxynucleotides in quadrupole ion trap, 16:1853
- of ions, charge-associated mechanism of, in fatty acid CID spectra, 16:280
- of kyurenine, ESI-MS/MS of, 12:786
- of $[M + Cu]^+$ peptide ions containing an *N*-terminal arginine, 11:626
- of $[M + Ni]^+$ adducts of aliphatic amino acids, 8:749
- of macrocyclic amines $C_nH_{2n+1}NH_2$ electron ionization induced, evidence for the rearrangement of aminocarbene radical cations and a comparison with long-chain esters, 1:238
- mass spectral, novel, of heptafluorobutyryl derivatives of acyl analogues of platelet-activating factor, 2:476
- of metastable molecular ion of methyl lactate, formation of oxygen-protonated methanol $[CH_3OH_2]^+$ involving double hydrogen atom transfers, 6:202
- of molecules sliding along surfaces in the speed range above thermal and below Bohr velocity, 10:471
- of multiply protonated peptides, charge derivatization and, 13:47
- of negative ions from carbohydrates
 high mannose *N*-linked glycans in, 16:631
 hybrid and complex *N*-linked glycans in, 16:645
 nitrate and other anionic adducts in, 16:622
- of noncovalent sugar-sugar complexes by infrared atmospheric pressure MALDI, 14:1158
- of a novel marine peptide, plicatamide, unusual gas phase intramolecular rearrangement involved in, 12:470
- of oligodeoxynucleotides
 effects of replacing phosphates with methylphosphonates and thymines with other bases in T-rich sequences, 12:580
 studied by H/D exchange and ESI-MS/MS, 12:193
- of oligonucleotide/metal complexes, CAD of, 16:28
- of oligonucleotides, using modified nucleic acids, 12:744
- oligopeptide, viewed as constant neutral loss, 1:336
- of organosulfur compounds upon electron impact, metastable decomposition of molecular ions of methyl thioglycolate and ethyl thioglycolate, 8:801
- of oxofatty acids via high-energy collisional activation, 9:620
- peptide, obtained from a reflector matrix-assisted laser desorption-ionization time-of-flight and a tandem four sector mass spectrometer, comparison of, 6:822
- of peptide ions, vacuum ultraviolet light-induced pathways in, 16:1384
- of peptides
 fast, observed, in matrix-assisted laser desorption, factors that influence, 7:225
 MALDI of, 8:501
- in phosphatidylglycerol studies using triple quadrupole MS/MS, 12:1036
- of phosphopeptides, manipulation of, gas phase boron derivatization in, 16:1905
- of phosphopeptides by atmospheric pressure MALDI and ESI-ion trap MS, 13:274
- of photomodified oligodeoxynucleotides adducted with metal ions in an ESI-ITMS, 12:1174
- postsources decay, of carbohydrates from ovalbumin, 11:572
- proferrioxamine, collision-activated, evidence for a succinimide-mediated mechanism, 6:836
- of propofol and structurally related phenols, 16:814
- of protonated amides through intermediate ion-neutral complexes, neighboring group participation, 9:454
- of protonated glycine, computational study of, 11:687
- of protonated ions of cysteine-containing peptides, 15:697
- of protonated oligoalanines, 15:1810
- of protonated *O,O*-diethyl *O*-aryl phosphorothionates in tandem mass spectral analysis, 7:189
- of protonated *O,O*-dimethyl *O*-aryl phosphorothionates in tandem mass spectral analysis, 6:488
- of protonated thioether conjugates of acrolein using low collision energies, 8:727
- remote, of protonated aromatic carbonyl compounds via internal reactions in intermediary ion-neutral complexes, 3:417
- retro-diels-alder, stereoselectivity of, 12:1011
- selective gas-phase reactions, mechanisms for, methionine, 17:1631
- of some cyclohexene-fused 2-*N*-phenylimin operhydro-3,1-oxazines and related thiazines, effects of *N*-substitution on, 2:125
- studies on mechanisms of, picolinyl esters, 17:1620
- of substituted oxonium ions: role of ion-neutral complexes, 10:386
- of tetronasin, ESI-CID-MS-MS studies of, 15:325
- of vitamin B12 in aerosol MALDI, 8:140
- Fragmentation and reactions, of two isomeric *O*-alkyl *S*-(2-dialkylamino) ethyl methylphosphonothiolates studied by ESIITMS, 12:902
- Fragmentation mechanism of the adducts, ion-molecule reaction of alkanenitriles and transition-metal ions in the gas phase, 2:120
- Fragmentation pathway
 of oligonucleotides in MALDI-MS by charge tagging and H/D exchange, 13:1407
 methylarginine-containing peptides, ESI-MS-MS of, 15:142
- Fragmentation studies, metastable and collision-induced, of all $C_4H_{12}Si^+$ isomers, systematic study of structure-reactivity relations, 5:537
- Fragmentation yield:CID, of iron(II)phenanthroline, activation energy estimation using, ESI-MS, 17:962
- Francisella tularensis novicida*, lipid A, environmentally-regulated remodeling, MS/MS, 18:1080
- Frank-Condon factors, study of keV $CO_2^+ \bullet / He$ collisions by collision-induced photon emissions, 19:1551
- Fraying and electron autodetachment dynamics of trapped gas phase oligonucleotides, 14:1330
- Free-electron laser, tunable mid-infrared, infrared matrix-assisted laser desorption and ionization by using, 7:1187
- Free energy determination, protein unfolding, accuracy of MALDI-MS and SUPREX, 17:1535
- Free fatty acid transport, in adipocytes, multi imaging MS of, 15:1572
- Free radical and acylium ion reactivity, in single molecule, 11:697
- Free radical oxidation of arachidonic acid, four regioisomers of F2-isoprostanes formed by, mass spectrometric analysis of, 7:490
- Frequency modulation
 space charge induced, filter diagonalization method in, FT-ICR-MS, 17:836
 two-dimensional Coulomb-induced, in Fourier transform ion cyclotron resonance, a mechanism for line broadening at high mass and for large ion populations, 4:909
- Frequency shifts due to the interference of resolved peaks in magnitude mode FTICR-MS, 13:387
- Frequency sweep direction, effect of, on motion of excited ions in Fourier transform ion cyclotron resonance cells, 3:847
- FT-ICR. *See* Fourier transform ion cyclotron resonance mass spectrometry
- FT-ICR MS. *See* Fourier transform ion cyclotron resonance mass spectrometry
- FTICR. *See* Fourier transform ion cyclotron resonance

- FTICR mass measurements, without internal standards using multiply charged ions, 11:416
- FTICR-MS. *See* Fourier transform ion cyclotron resonance mass spectrometry; Fourier transform ion cyclotron resonance-MS
- FTMS. *See* Fourier transform mass spectrometry; Fourier transform-MS
- Fucosyl galactoside, isomers analyzed by ERMS, 18:1873
- Fullerene trications, derivatized, proton transfer reactions of, 9:114
- Fullerenes
 - derivatized, alkali cation attachment to, MALDI of, 13:1448
 - H60, gas-phase alkylation of, by ion-molecule reaction under chemical ionization, 4:894
 - mass spectrometry and, 3:268
 - matrix-assisted laser desorption/ionization tandem reflectron time-of-flight mass spectrometry of, 7:590
 - methylated, MS of, using in-beam electron impact technique, 16:1311
 - neutral organic molecules including, applications of electro-spray ionization mass spectrometry to, 4:596
- Fulvene structure, isomerization to 1,2-hydrogen shifts in, benzene radical cation, *Ab initio* calculations on, 6:513
- Functional group interactions, effects of
 - on the bimolecular and dissociation reactions of diols, 3:535
 - on the gas-phase methylation and dissociation of acids and esters, 7:565
- Functional group-selective ion-molecule reactions of ethylene glycol and its monomethyl and dimethyl ethers, 4:97
- Fundamentals of applying electrospray ionization mass spectrometry to low mass poly(methyl methacrylate) polymers, 6:906
- Fundamentals of the application of matrix-assisted laser desorption-ionization mass spectrometry to low mass poly(methylmethacrylate) polymers, 7:287
- Fungi, ergosterol pathway inhibition in, analysis by ASAP, 18:1274
- Furan, transition-metal mediated heteroatom removal by reactions of FeL^+ [L^+O , C_4H_6 , $\text{c-C}_5\text{H}_6$, H_6H_6 , $\text{C}_5\text{H}_4(=\text{CH}_2)$] with, in the gas phase, 7:938
- G
- G-quadruplexes, in deoxyguanosine clusters, ion mobility MS of, 16:989
- [Ga_4L_6] host-guest complexes, self-assembled, ESI MS characterization of, 17:292
- Galectin-1, protein interface footprinting quantified by hydroxyl radical oxidation and MDS, 19:1686
- Gallium ions, energetic, silver surface bombardment by, 16:1677
- Ganglia, invertebrate, spatial profiling, 14:752
- Ganglioside analysis, by high pressure MALDI-FTMS, 13:402
- Gangliosides
 - analysis by TLC-MALDI-o TOF-MS, 16:1552
 - from human granulocytes, 12:964
 - GM1, IRMPD, ECD, and EDD of, 16:752
 - underivatized and permethylated, matrix-assisted laser desorption ionization time-of-flight mass spectrometry of, 3:785
- Ganoderma lucidum*, triterpenoids, analysis using LC-ESI/MS, 18:927
- Garcinia* spp., screening of PPAPs using PID and UPLC-ESI-Q-TOF, 20:1846
- Gas
 - as carriers, in FAIMS, 11:1125
 - pulsed, introduction into quadrupole ion traps, 1:308
 - reagent, isotopically labeled, chemical ionization mass spectrometry using, 5:596
 - viscous, flow through capillaries, ion transport by, 5:873
- Gas analysis, small MS systems for, 13:1005
- Gas chromatograph-toroidal ion trap mass spectrometer, hand portable for hazardous compound detection, 19:1425
- Gas chromatographic-mass spectrometric method for the assessment of oxidative damage to double-stranded DNA by quantification of thymine glycol residues, 4:336
- Gas chromatography
 - capillary/ion trap mass spectrometry quantitative methods, combined, using labeled or unlabeled internal standards, 1:265
 - combined with mass spectrometry, structural determination of long-chain aldehydes by, pyridine-containing Schiff base derivatives for, 2:245
 - high-temperature/mass spectrometry with a polarizable stationary phase, negative ion chemical ionization, and mass-reversed chromatography, high-resolution triglycerol mixture analysis using, 7:350
 - isotope-ratio-monitoring/mass spectrometry, nitrogen isotope analyses by, 5:387
 - on-line isotopic dilution cryofocusing/mass spectrometry, on-line, automated high-speed analysis of selected organic compounds in urban air by, 5:1001
 - purge and trap/mass spectrometry, measuring volatile organic compounds in human blood by, interferences in, removing of, importance of enhanced mass resolution in, 3:831
 - suspended trapping/Fourier transform mass spectrometry for analysis of complex organic mixtures, 1:431
 - toxicological screening using, 14:14
- Gas chromatography/chemical reaction interface mass spectrometry, selective detection of sulfur-containing compounds by, 2:250
- Gas chromatography Fourier transform mass spectrometry (GCFTMS), preconstructor coupled to, 15:1191
- Gas chromatography (GC) quadrupole mass spectrometry AP LC/MS ion source combined with, advantages of, 16:1730
- identifying essential oil components by, 16:1902
- Gas chromatography inductively coupled plasma mass spectrometry, HF-LPME with, PBDE analysis in soil, dust and human serum, 18:1740
- Gas chromatography/ion mobility spectrometry (GC/IMS), in U.S. space program, 12:656
- Gas chromatography/ion trap mass spectrometry, alternating RF/DC isolations for quantitation with coeluting internal standards in, 3:85
- Gas chromatography ionization mass spectrometry, relative sensitivities in, correlations of, with molecular parameters, 1:397
- Gas chromatography mass spectrometry
 - in acute poisoning, scope and limitations of, in general unknown screening, 6:995
 - in analysis of hydrocarbons, Cu^+ chemical ionization for, 9:731, 9:981
 - early, 4:367
 - fast, very fast, and ultra-fast, of thermally labile steroids, carbamates, and drugs in supersonic molecular beams, 7:737
 - high-resolution, application to the determination of trace monobromopolychlorodibenzo-p-dioxins in environmental samples, 3:248
 - in identifying minor compounds in complex mixtures, COMSPARI and, 15:580
 - of neuromelanin pyrolysis products, 15:920
 - in phenyl- and cyclopentylimino derivatization for double bond location in unsaturated $\text{C}_{37}\text{-C}_{40}$ alkenones, 15:1161
 - isotope dilution, for platinum determination in urine, 2:85
 - and laser desorption, in supersonic molecular beams, 9:628
 - location of alkenone double bonds, 17:710

- β -methylamino-L-alanine ethylchloroformate derivative characterization by, 18:817
- nitramine anion fragmentation, 18:835
- overlapping Isotopologue separation and quantitation, 17:1028
- practical guide to, 8:294
- pyrolytic, analysis of melanin from epidermal melanocytes, 20:464
- quantitative signatures of steroids using, 20:1626
- quadrupole, fast scanning of, 8:539
- selective ion monitoring mode in, direct chemical analysis of UV laser ablation products of organic polymers by using, 5:106
- solid-phase microextraction in-line coupled to, pesticides in aqueous samples determined by, 6:1119
- spectrum extraction and compound identification data from, 10:770
- of wasp cuticular hydrocarbons, 9:320
- Gas chromatography/mass spectrometry, early, 4:367
- Gas chromatography/mass spectrometry and particle beam liquid chromatography/mass spectrometry, poly(amic acid) and polyimide characterization using, 5:299
- Gas chromatography/mass spectrometry components, overlapping, spectral deconvolution for, 3:558
- Gas chromatography/mass spectrometry (GC/MS)
- in clinical chemistry, 12:127
- current practice of, 12:1348
- field-portable, environmental and forensic applications of, 12:683
- of platelet activating factor, 11:176
- of polycyclic aromatic hydrocarbons in ancient terrestrial rocks, 12:989
- Gas chromatography-MS, of fatty acid picolinyl esters by direct Curie-point pyrolysis of whole bacterial cells, 14:58
- Gas chromatography-MS/MS, of diazinon photocatalytic degradation products, 14:803
- Gas chromatography/time-of-flight mass spectrometry (GC/TOFMS), field-portable high speed spectrometer for, 12:648
- Gas flow profile, FAIMS for, 16:1447
- Gas phase
- α -cyanoethylindole isomeric differentiation in, 16:388, 16:801 (erratum)
- 3,4-dihydro-2,4-dioxo-2H-1,3,5-oxadiazinium ion synthesis in, 16:1602
- acidity, of glycine and alanine, theoretical calculations of, 10:318
- acidity in, stereochemical effects of 11 α -OH and 11 β -OH 17 β -estradiols, 20:2318
- agostic effects in, SiCl₃⁺ and SiCl⁺ affinities for pyridines determined by using the kinetic method with multiple stage mass spectrometry, 7:198
- alkanenitriles and transition-metal ions in, ion-molecule reaction of, a study on fragmentation mechanism of the adducts, 2:120
- anionic α -adducts in, using electrospray ion source as chemical reactor, 15:927
- aromatic amino acid complexation with Na⁺ and K⁺, cation- π effects in, 11:1037
- basicity, of deprotonated monomeric and dimeric ferulic and sinapic acids, 10:1111
- bimolecular hydrogen-deuterium exchange behavior of protonated alkyl dipeptides in, 6:466
- boron derivatization in, in phosphopeptide fragmentation pattern manipulation, 16:1905
- bradykinin fragment 1-5 ions in, IM-MS of, 16:893
- bradykinin ions in, H/D exchange of, in linear quadrupole ion trap, 14:85
- cations in, 1,2-eliminations, 19:1491
- chiral recognition in, mass spectrometric studies of diastereomeric cobalt complexes, 5:452
- C₅H₅RF⁺ (R=F, Cl, Br, I, O, OH, OCH₃, H₆H₅) ionic, structure of, and generation of their neutral counterparts by neutralization-reionization mass spectrometry, 6:1143
- chromium oxyanions in, ion-molecule reactions of, 14:1067
- cluster ions in, thermochemical stabilities and structures of, 16:1760
- collision-induced and infrared multiphoton dissociation studies on M(acetone)₂⁺ (M=Al, Fe, Co, Cu, ScO) in, 4:135
- complexation of protonated peptides with saccharides in, decreases rates of hydrogen/deuterium exchange reactions, 6:1247
- complexes of iron(II) with cysteine-containing peptides in, 7:977
- conformations of biological molecules, the hydrogen/deuterium exchange mechanism, 10:9
- conformations of deprotonated trinucleotides in, zwitterion formation, 14:161
- Cr(CO)₅⁻, relaxation of, 10:950
- cyclodextrin complexes in, zwitterion formation in, 16:166
- cyclopropylphenyl benzamide cyclization in, 14:956
- cytochrome c ions in, in Paul trap, 16:1493
- dehydration of peptide [M + H]⁺ ions in, 4:477
- dinucleotide analogs in, metal chelation of, by fast-atom bombardment mass spectrometry, 7:42
- dissociation of multiply charged peptide clusters in, 14:1373
- dissociation of oligonucleotides in, effect of cisplatin on, ESI-MS/MS, 20:792
- disulfide bond scrambling during MALDI MS, 20:1603
- of doubly charged alkaline earth and transition metal complexes of acetonitrile, pyridine, and methanol generated by ESI, 8:1124
- ecotin thermal dissociation in, 13:1432
- endgroup-assisted siloxane bond cleavage in, 14:1039
- flavonoid acidities in, 15:848
- fragmentation of long-lived radical cations, cysteine, 20:985
- fragmentation of protonated oligomers in, 20:60
- glycosidic bond fragmentation in, direction by non-covalent complexes, 15:715
- gramicidin S and linear analog conformational preferences in, IM-MS of, 15:870
- of growth hormone secretagogue, rearrangement after MS/MS, 13:371
- guanosine H₂O adduct formation in, CID study of, 16:1291
- H/D exchange and collision cross-sections, hemoglobin, 20:484
- H/D exchange kinetics in, DI versus D₂O, 11:167
- H/D exchange of peptides and peptide dimers in, acidic residues and sodium ion adduction in, 14:1477
- H/D exchange reactions of fluorophenyl anions, 10:840
- halid S_N2 reactions with halodimethylamine in, 15:673
- heme-globin complex dissociation in, 8:519
- hydrogen/deuterium exchange of protonated amino acids in, 16:1846
- insertion reaction of metal carbonyl anions with methyl formate in, as revealed by ¹³C- and D-labeling, 5:525
- ion chemistry, 10:797
- ion chemistry of methylbenz(is)oxazole isomers in, 15:1005
- ion/ion interactions, peptide AA composition and role in charge inversion, 20:180
- ion-ion reactions in, proton transfer reactions of protonated pyridine with multiply charged oligonucleotide anions, 6:529
- ion-molecule reactions in, 8:610

- ionic and neutral $[\text{HS-P-OH}]^{+/\bullet}$ and $\text{S}=\text{P}(\text{OH}_2+/\bullet)$ generation and characterization in, 16:1353
- ionic and neutral $\text{P}(\text{OH})_2+/-$ in, MS/MS and computational chemistry of, 13:250
- ionized, 1,5-hexadiene structure in, 12:840
- ions formed by ESI, hydration of, 10:958
- ions in, FRET in, 16:1481
- isomeric arylglycoside reactivity toward amines in, 15:244
- isomeric dihydroxybenzoic acid basicities in and gas phase acidities of their radical cations, 11:544
- long-range glycosyl transfer reactions in, 13:325
- loss of nucleobases from deprotonated oligonucleotide anions in, Arrhenius activation parameters for, 15:55
- Meerwein reaction of phosphonium ions with epoxides and thioepoxides in, 15:398
- Meisenheimer complexes bonded in, 15:998
- metal and proton complexes of di-*o*-benzoyl-tartaric acid dibutyl ester and L-tryptophan in, chiral recognition characteristics of, 14:571
- metal oxide surface fabrication by reactive landing of ions, 20:915
- multiply charged negative clusters of adenosine 5'-monophosphate in, 16:1840
- N-heterocycle synthesis in, in arylnitrenium ion generation, 15:1679
- Ni(II) complex reactions with acetonitrile, 15:1128
- noncovalent interactions between vancomycin-group antibiotics and bacterial cell-wall precursor peptides probed by hydrogen/deuterium exchange, 9:1255
- noncovalent peptide in, unimolecular dissociation of, FTICR of, 14:1282
- oligonucleotide duplexes in, stability determination of, 15:1354
- oxidation and reduction of some 1-(2-benzothiazolyl)-3,5-diphenyl formazans, complex formation with transition metals under laser desorption ionization, 10:339
- of oxidized peptides containing cysteine, fragmentation of, 12:228
- paralytic shellfish poison behavior in, 15:462
- peptide acidity from adduct dissociation in, 15:616
- phospholipid headgroup activities in, 16:926
- of polyether ionophore antibiotics, stabilities by CID, 16:779
- polymerization of ethylene catalyzed by Ti^+ in, evidence for, 3:464
- proton-bound heterodimers and homodimers of amides and amines in, FTICR-MS of, 13:1273
- protonated serine and derivative fragmentation reactions in, 11:1047
- protonated tryptophan reactions in, 15:65
- protonation of ferrocene in, 5:274
- reactive and dissociative behavior of transition metal oxide cluster ions in, characterization of, 2:212
- reactivity and dissociation of $[\text{M} + 4\text{H}]^{4+}$ ions from insulin chain B, cysteic acid group effect on, 10:928
- reactivity, protonation and deprotonation and, fluorinated 1,2,4-triazines, ESI MS, 19:686
- retro-*Michael* fragmentation reactions in, of 2-hydroxybenzyl-*N*-pyrimidinylamine derivatives, 16:1552
- Ru^+ , activation of C_2H_6 , C_3H_8 , $\text{HC}(\text{CH}_3)_3$, and $c\text{-C}_3\text{H}_6$ by, 10:821
- salt complexes in, CAD of, stereoselectivity in, 15:1509
- Shiga toxin B subunit stability in, nanoESI-FTICR MS of, 16:1957
- sodiated peptide formation in, 15:607
- sodium controlled reactivity of protonated deoxyoligonucleotides in, MALDI, 20:689
- solvation of propanediol ions by water molecules in, 15:1123
- stability of double-stranded oligodeoxynucleotides and their noncovalent complexes with DNA-binding drugs as revealed by collisional activation in anion trap in, 11:450
- on the stabilization of carbanions by adjacent phenyl, cyano, methoxy-carbonyl, and nitro groups in, 5:159
- stereoelectronic effects in, of pyridine affinity of trifluoride cation, 8:68
- substituted cyclopentadienylmanganese cations in, metastable ion study of, 4:216
- ter-body intermediates in, 12:938
- ternary β -cyclodextrin inclusion complexes in, ES-MS of, 14:449
- thiol and thioester nitrosation in, 15:344
- transformation of deprotonated polypeptides, transition metal complex cations in, 20:1718
- transition-metal mediated heteroatom removal by reactions of FeL^+ [L^+O , C_4H_6 , $c\text{-C}_5\text{H}_6$, H_6H_6 , $\text{C}_5\text{H}_4(=\text{CH}_2)$] with furan, thiophene and pyrrole in, 7:938
- trapped oligonucleotides in, fraying and electron autodetachment dynamics of, 14:1330
- ubiquitin conformers in, multidimensional separations of, 16:1427
- uracil and uracil analog acidity in, 13:985
- ureas and thioureas in, cyclization of substituted *N* in, 16:1739
- Gas phase acidities, of monosaccharides, ranking of, 12:1193
- Gas-phase acidities, apparent, of multiply protonated peptide ions, ubiquitin, insulin B, and renin substrate, 7:1211
- Gas-phase acidities of diols, 4:723
- Gas-phase acidities of germane and methylgermane, Fourier transform-ion cyclotron resonance study of, bond dissociation energy of germane, 4:54
- Gas phase acidity, of methylthioacetic acid, determining with Cooks' kinetic method, 16:535
- Gas-phase aldose-ketose isomerization, mechanism of, a study using tandem mass spectrometry and theoretical calculations, 4:943
- Gas-phase anionic complexes of alkali metal ions and peptides, structure and collision activated decompositions, 5:137
- Gas-phase base-catalyzed Claisen-Schmidt reactions of the acetone enolate anion with various para-substituted benzaldehydes, 7:82
- Gas-phase basicities and ammonium ion affinities of polyethers, orders of, comparison of, by kinetic method and ligand exchange technique, 4:718
- Gas-phase basicities of histidine and lysine and their selected di- and tripeptides, 7:1203
- Gas-phase basicities of serine and dipeptides of serine and glycine, 5:718
- Gas phase basicities of polyamines, 9:1043
- Gas phase bimolecular chemistry of isomeric $\text{C}_3\text{H}_6\text{Br}^+$ cations, 6:11
- Gas-phase bond strengths of CO_2 and N_2O , comparative study of, with the halide ions, 4:58
- Gas-phase cationization and protonation of neutrals generated by matrix-assisted laser desorption, 4:393
- Gas phase cations, proton transfer from, to reference bases, 12:359
- Gas-phase C_7F_7^+ ions, structures of, 4:813
- Gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes, by fast-atom bombardment mass spectrometry generation of molecular signals, 4:249
- synthesis and characterization of molecular complexes, 4:255
- Gas phase cleavage, of phenylisothiocyanate-derivatized electrosprayed tryptic peptides in an FTICR trapped-ion cell: mass-based protein identification without LC separation, 12:288
- Gas phase clusters, between halide ions and olefin, 12:144
- Gas phase CNF bond cleavage, in perfluorohexane using W^- ,

- Si-, P-, Br-, and I-containing ions: comparisons with reactions at fluorocarbon surfaces, 9:1158
- Gas-phase complexation of polyethers with halide ions, 4:242
- Gas phase conformations of cationized poly(styrene) oligomers, 13:499
- Gas phase cyclodextrin-amino acid complexes, chiral recognition in, 12:278
- Gas phase dissociation, of oligoribonucleotides, ESI-MS/MS of, 16:1262
- Gas-phase dissociation, MS/MS, crosslinked Lys-Lys peptide influence of cross-linker identity on, 17:395
- Gas-phase dissociations of protonated nitriles, ion neutral complexes in, $n\text{-C}_3\text{H}_7\text{CNH}^+$ and $i\text{-C}_3\text{H}_7\text{CNH}^+$, 2:178
- Gas phase dissociative electron transfer, perturbation molecular orbital theory of, 8:510
- Gas phase electrophoretic mobility molecular analyser, nanoES, PDMA and, nanobioparticle analysis using, 19:1062
- Gas-phase electrophoretic mobility molecular analysis, protein complex sizing by ESI and IMS, 18:1206
- Gas-phase electrosprayed ions, implications for the conformation of, defect formation on surfaces bombarded by energetic multiply charged proteins, 7:329
- Gas-phase enantioselectivity, zwitterionic formation and, amino acids, 17:442
- Gas-phase ferrocene cation, photodissociation spectroscopy of, 2:97
- Gas phase fragmentation of the Ag^+ -phenylalanine complex, cation- π interactions and radical cation formation, 13:408
- Gas-phase fragmentation reactions
mechanisms for, peptide ions containing S-alkyl cysteine sulfonide, 18:1690
selective. mechanisms for, methionine, 17:1631
- Gas phase H/D exchange
of model peptide: FTICR and computational analyses and metal-induced conformational mutations, 12:1272
of positively charged mononucleotides by use of FTICR-MS, 12:268
- Gas phase H/D exchange kinetics in, of polyamine complexes, 11:711
- Gas-phase halo alkylation of H_6O -fullerene by ion-molecule reaction under chemical ionization, 4:894
- Gas phase hydrates
of alkylammonium ions, observations of magic numbers in, 8:1175
in gramicidin S, 8:565
- Gas-phase hydrogen/deuterium exchange as a molecular probe for the interaction of methanol and protonated peptides, 5:526
- Gas phase hydrogen-deuterium exchange in electrospray ionization mass spectrometry as a practical tool for structure elucidation, 5:434
- Gas phase intramolecular rearrangement, in plicatamide fragmentation, 12:470
- Gas phase ion/ion chemistry, study via mass spectrometry, 19:173
- Gas-phase ion-ion reactions
new approach for the study of, using electrospray ionization, 3:695
study of, using electrospray ionization, a new approach for, 3:695
- Gas-phase ion-molecule chemistry of borate and boronate esters, 5:169
- Gas-phase ion-molecule complexes of transition-metal ions, formation and fragmentation of, with organic molecules containing two functional groups, 3:451
- Gas-phase ion molecule reactions, selective metal ion source for, resonant laser ablation as, 7:664
- Gas-phase ion-molecule reactions of the CH_3O^+ cation, 3:515
- Gas phase ion-molecule reactions of transition metal complexes, effect of different coordination spheres on complex reactivity, 13:813
- Gas-phase ionic species, generation and characterization of, multiple stage pentaquadrupole mass spectrometry for, case of the PyC_2H_5^+ isomers, 7:1126
- Gas phase ions
structures, proline-containing b_2 ion, 17:20
trapped, effect of buffer gas on fluorescence yield, 17:855
- Gas-phase labeling combined with tandem mass spectrometry, ion-dipole complex formation from deprotonated phenol fatty acid esters evidenced by, 4:343
- Gas-phase lithium-peptide adducts, investigations of, tandem mass spectrometry and semiempirical studies, 1:473
- Gas-phase methylation and dissociation of acids and esters, effects of functional group interactions on, 7:565
- Gas-phase molecular dications
formed from neutral organics in solution via chemical electron-transfer reactions by using electrospray ionization mass spectrometry, 5:689
formed from neutral organics in solution via the controlled-current electrolytic process inherent to electrospray, 7:157
- Gas phase peptide-cyclodextrin complexes, noncovalent binding interactions in, 8:244
- Gas-phase peptide ions, low energy decompositions of, role of the site of protonation in, 7:522
- Gas-phase polyether/ammonium ion complexes, hydrogen-bonding interactions in, 5:260
- Gas-phase proton transfer reactions, involving multiply charged cytochrome c ions and water under thermal conditions, 3:624
- Gas-phase protonated and methylated lactones, dissociation behavior of, characterization of, 3:47
- Gas-phase reaction, between protonated acetaldehyde and methanol, 1:481
- Gas phase reactions
of charged phenyl radicals with neutral biomolecules evaporated by laser-induced acoustic desorption, 13:192
of chloramine with anionic nucleophiles, nucleophilic substitution at neutral nitrogen, 12:139
of hydrated alkaline earth metal ions with benzene, 12:250
of O_2II with alkyl and aryl esters of benzenedicarboxylic acids, 5:990
of Sc^+ , Y^+ , and Lu^+ with alcohols, effects of the class and chain length of alcohols on the nature of primary products, 7:1157
of trimethyl borate with phosphates and their non-covalent complexes, 13:1088
- Gas-phase reactions between O^- and $\text{H}_6\text{H}_5\text{F}$, on the acidity of fluorobenzene and 1,4-difluorobenzene, 4:73
- Gas-phase reactions of amino alcohols with dimethyl ether ions, site-selective, an investigation of, 4:230
- Gas phase reactions of the ion $\text{C}_5\text{H}_5\text{Fe}^+$, 9:314
- Gas phase reactivity
and molecular modeling studies on triply protonated dodecapeptides that contain four basic residues, 9:716
of rare earth cations with phenol: competitive activation of CNO, ONH, and CNH bonds, 9:1035
- Gas phase RNA and DNA ions, conformational dependence of gas phase H/D exchange of AMP and dAMP, 12:780
- Gas phase $\text{RSC}_6\text{H}_5\text{Cr}^+\text{L}$ ($\text{R}=\text{C}_6\text{H}_5$; L^+ arene) and $\text{C}_6\text{H}_5\text{SSH}_6\text{H}_5\text{Cr}^+$ ions, a tandem mass spectrometry study of, 5:928
- Gas phase separations of protease digests, 9:1213
- Gas phase SN_2 reactivity of dicoordinated borinium cations using pentaquadrupole mass spectrometry, 12:948

- Gas-phase stability of cluster ions SF_m^+ (SF_6)_n with $m=0.25$ and $n=1-23$, 6:1137
- Gas-phase structure, of electrosprayed proteins using ion-molecule reactions, investigation of, 5:207
- Gas phase studies, of cytochrome *c* ions, temperature-dependent H/D exchange in, 13:506
- Gas phase studies of the interactions of Fe^{2+} with cysteine-containing peptides, 9:1285
- Gas-phase studies of valinomycin-alkali metal cation complexes, attachment rates and cation affinities, 5:632
- Gas-phased oligosaccharide nonreducing end, sequencing and structural analysis by RP-HPLC-MS, 20:1821
- Gas reactions, in APCI-MS sources, 15:311
- Gas stage ions, applications of multiple stage mass spectrometry in a pentaquadrupole, polar [4+2+] Diels-Alder cycloaddition to, 6:1
- Gaseous carbanions, hydrogen/chlorine exchange reactions of, 16:2045
- Gaseous Fe(III)-heme cation, binding to biological molecules, 16:589
- Gaseous myoglobin ions stored at greater than 300 K, 5:324
- Gaseous protein ion structure, hydroiodic acid attachment kinetics of, 10:552
- Gases
3D concentration mapping using portable MS, 19:1411
heavy, effects of, on the tandem mass spectra of peptide ions in the quadrupole ion trap, 7:1194
- Gas(es), heavy, using static pressures of, in quadrupole ion trap, 1099
- Gate valve design, minimum thickness, for MS, 16:441
- Gated electrostatic mass spectrometer, 19:1384
- Gated trapping, capacitive coupling with, in MALDI-FTICR, 8:195
- Gated trapping strategy with a two-time constant and a delay for catching in-field generated ions that range over three decades in mass-to-charge and two decades in velocity in FTMS, 12:296
- GC. *See* Gas chromatography
- GC/IMS. *See* Chromatography/ion mobility spectrometry.
- GC-MS. *See* Gas chromatography-mass spectrometry
- GC/MS. *See* Gas chromatography/mass spectrometry
- GC-TMS. *See* Gas chromatograph-toroidal ion trap mass spectrometer
- Gel filtration, automated in-line, of proteins for native state MS, 19:239
- Gel microfluidic chips, capillary, interfacing with infrared LDI MS, 17:469
- Gel permeation chromatography-MALDI-TOFMS, of poly(dimethylsiloxanes), 14:195
- Gel permeation chromatography-matrix-assisted laser desorption ionization, α , ω -bis(4-hydroxybutyl)poly(dimethylsiloxane) analysis using, 13:170
- Gelatin matrix, tetraalkylammonium salt molecular ions sputtered from, formation and emission of, 4:930
- Gelation films, doped, as a model matrix for molecular secondary ion mass spectrometry studies of biological soft tissue, 4:419
- GEMMA. *See* Gas phase electrophoretic mobility molecular analyzer; Gas-phase electrophoretic mobility molecular analysis
- Gene products, heterogeneous, observation by FT-ICR MS, 19:103
- General method for precalculation of parameters for sustained off resonance irradiation/collision-induced dissociation, 13:318
- General unknown screening by gas chromatography-mass spectrometry in acute poisoning, scope and limitations of, 6:995
- Generation and characterization of ionic and neutral $P(OH)_2$ +/- in gas phase by MS/MS and computational chemistry, 13:250
- Generation of average mass values and end group information of polymers by means of a combination of matrix-assisted laser desorption/ionization-mass spectrometry and liquid secondary ion-tandem mass spectrometry, 8:76
- Generation of electrospray from a solution predeposited on optical fibers coiled with a platinum wire, 11:464
- Generation of substructure identification rules using feature-combinations from tandem mass spectra, 3:159
- Generation of the m/z 69 ion $C_3HO_2^+$ by electron ionization of some aryl-substituted β -diketones, 9:29
- Genetic algorithms, feature extraction, alignment of LC-MS and LC-MS/MS data sets, 18:1835
- Genetic diagnosis by polymerase chain reaction and electrospray ionization mass spectrometry, detection of five base deletion from blood DNA of familial adenomatous polyposis patient, 8:737
- Genotyping
ApoE gene SNPs and indels by MS, 17:3
ESI-MS in, online data base for, 13:87
semi-quantitative, of single nucleotide polymorphisms, LC-MS for, 15:1897
- Geometry, energy, and time, roles of, ion formation from charged droplets, 4:524
- Germane, bond dissociation energy, Fourier transform-ion cyclotron resonance study of gas-phase acidities of germane and methylgermane, 4:54
- Ghrelin, fatty acid modification in, identification and localization by ECD, 13:1443
- Giant organic molecules, synthetic polymers and, improvements in analysis by solvent-free MALDI-MS, 17:661
- GIB-MS/MS. *See* Guided ion beam tandem mass spectrometry
- Ginkgo biloba*, flavonoids and terpenoids in plasma, efficiency of ESI determination in, 19:445
- GLaD. *See* Guanidino-labeling derivitization
- GlcNAcylation sites of peptides and α -crystallin, Q-TOFMS in identification of, 12:1106
- Gln-Gly cleavage, correlation between CID and biological degradation in, 12:337
- Globin subunits, oxidized, enzymatic digestion of, peptides derived by, electrospray ionization-tandem mass spectrometry analysis of, an improved method to determine amino acid substitution in the hemoglobin "core," 7:1040
- Globins, glycosylated, analysis by MALDI MS, nephropathic patients, 18:1018
- Glow discharge
dc, effect on the performance of a particle beam liquid chromatography/mass spectrometry interface, investigation of, 5:908
modeling of, 8:1021
pulsed, as an elemental ion source, 1:138
radiofrequency, effects of target gas in collision-induced dissociation using, 5:845
- Glow discharge-Fourier transform ion cyclotron resonance mass spectrometry, external ion injection in, ion guide for, 4:387
- Glow discharge in atmospheric gases, structures of NO_3^+ formed via, 1:217
- Glow discharge ion source for mass spectrometric analysis of organic compounds, characterization of, 5:305
- Glow discharge ion source with Fourier transform ion cyclotron resonance mass spectrometric detection, 3:122
- Glow discharge mass spectrometer, high-resolution, operated with an argon/hydrogen mixture, relative ion yields measured with, 4:278

- Glow discharge mass spectrometry
analysis of solution residues by, 4:47
relative ion yields for, theoretical calculations of, 3:79
- Glow discharge plasma, characterization of, as a function of sampling orifice potential, 1:448
- Glucagon, gas-phase conformation probed by CID, 17:786
- Glucagon-like peptide-1 derivative, *N*-acylated, ECD of, 16:548
- Glucokinase, from *Bacillus stearothermophilus*, ESI-FTMS-IRMPD of, 9:1222
- Glucopyranosides, FAB MS/MS of, 10:975
- Glucosamines, location of ester groups on, 11:301
- Glucose
concentration in tears, by ESI MS, 18:332
loss of, from flavonoid *O*-diglycosides, 11:136
- Glucose disaccharide-glucosyl complexes-diautomeric cobalt, tandem mass spectrometry of, differentiation of stereochemistry of glycosidic bond configuration, 7:953
- α -Glucosidase inhibitors, hawthorn leaf, characterization using LC-DAD-MS and SORI-CID FTICR MS, 20:1496
- Glucosyl-glucose disaccharide complexes-diautomeric cobalt, tandem mass spectrometry of, differentiation of stereochemistry of glycosidic bond configuration, 7:953
- Glucuronides, anabolic steroid, ESI and APCI MS/MS of, 11:722
- Glutamic acid, gas-phase acidity, cluster structure and kinetic method measurements, 19:1888
- Glutamine
derivatized, fragmentation of, 16:1192
N-terminal
dehydration *versus* deamination in ESI MS, 18:27
studies on a prion protein peptide, tandem mass spectrometry of peptides with, 1:258
- Glutamine lipids, ornithine lipids and, from *R. sphaeroides* cell membranes, MS/MS, 20:198
- Glutathione
conjugates, EPR and ES-MS/MS study, 20:2013
oxidized contryphan and, fragmentation of peptide disulfides from negative ion mass spectrometry, 19:358
ruthenium arene complex reactions with, nanoLC-FTICR MS, 19:544
- Glutathione conjugate of 2-furamide, mass spectrometric screening and characterization of, tandem, hybrid, multiple scan modes in, 2:55
- Glutathione conjugates, activated by high-energy collisions, fragmentation characteristics of, 3:815
- Glutathione conjugates of estrone and estradiol, structural isomers of, 9:612
- Glutathione S-transferase, noncovalent associations with ligands, ESI-quadrupole-TOFMS of, 11:606
- Glycans
HCV E2 envelope glycoprotein, MS characterization, 19:428
high mannose *N*-linked, fragmentation from carbohydrates, 16:645
isomeric *N*-Glycan, identification by CID MS/MS with 157nm laser-induced photofragmentation, 19:1027
N-linked, assigning structural isomers from ovalbumin, IMS-MS, 19:1711
N-linked, ionization and CID of, 12:926
N-linked derivatized, fragmentation of, 11:900
sialylation effect on negative ion dissociation, 17:844
sialylated, oligosaccharides and, analysis using DHB/An and DHB/DMA matrices, MALDI-TOF MS, 19:1138
sulfated, analysis by sequential double-permethylation, MALDI MS, 20:1660
- N*-Glycans, generation of structure databases, 20:1739
- Glycation sites, bovine insulin, characterization, MS/MS, 20:1319
- Glycerine, dehydration of, 9:945
- Glycerol
condensation of, with ammonium surfactants
fast atom bombardment-induced, regioselectivity of the adduct formation, 3:318
fast-atom bombardment-induced, time dependence of mass spectra and tandem mass spectra, 5:92
fast-atom bombardment of, background ions produced by, origins and structures of, 5:72
- Glycerol-amine solutions, solution chemistry and secondary ion emission from, 2:33
- Glycerol phosphatidylcholine phospholipids, characterization of, constant neutral loss scanning for, 9:1189
- Glycerophosphocholine lipids
doubly sodiated, ETD of, 18:1783
lithiated adducts of, ESI-MS/MS of, 9:516
- Glycerophosphocholines, arachidonate containing, molecular species analysis of, by tandem mass spectrometry, 2:45
- Glycerophosphoethanolamine plasmalogen phospholipids, ES-IMS/MS of, 15:1499
- Glycerophospholipids
1-*O*-alk-1'-enyl-2-acyl and 1-*O*-alkyl-2-acyl, differentiation by ESI-IT-MS, 18:2065
electrospray intrasource separation and selective ionization, 17:264
in situ characterization of sulfatides and, brain, using MALDI-MS/MS, 18:17
unsaturated, characterization by multiple stage LIT-MS with ESI, 19:1700
- Glycine
dipeptides of, gas-phase basicities of, 5:718
gas-phase acidity of, theoretical calculations of, 10:318
protonated, fragmentation pathways of, 11:687
- Glycine-copper solutions, electrospray ionization of, 7:25
- Glycoalkaloids, and their aglycons, analysis of, comparison of high- and low-energy collision-induced dissociation tandem mass spectrometry in, 7:173
- Glycoconjugate mixtures, cross-ring cleavage in, 15:1863
- Glycoconjugates, acidic, MALDI-TOF of, 9:1293
- Glycomics, platform for analysis of permethylated oligosaccharides, 18:1799
- Glycopeptide mixtures, heterogeneous, precursor ion scanning of, 13:1065
- Glycopeptides
gp120, from HIV, LC-MS of, 15:1545
high resolution scanning of, 14:777
by liquid chromatography/mass spectrometry with collision-induced dissociation, 3:804
O-linked, structural elucidation of, by high energy collision-induced dissociation, 7:319
- Glycopeptidolipid antigens, mycobacterial, fast atom bombardment mass spectrometry of, structural characterization by charge remote fragmentation, 3:656
- Glycoproteins
analysis of, Q-trap mass spectrometer in, 15:413
carbohydrates and, distinguishing phosphorylation and sulfation, ion-pairing and MS, 17:1282
characterization using hybrid linear ion trap/FT-IRC MS, 17:168
composition of, 11:564
electrospray MS profiling of, 15:750
HIV envelope, comparison of HPLC/ESI-FTICR MS *vs.* MALDI-TOF/TOF MS, 19:1209
monosaccharide analysis by LC-MS/MS, 20:1224
postsurface decay fragmentation of, 11:572
quantify glycosylation changes by label-free quantitation, 20:1048
sulfated glycosylation sites in, detection of, 16:340

- Glycosaminoglycan-derived disaccharides, sulfate and linkage position in, negative fast-atom bombardment tandem mass spectrometry to determine, 3:797
- Glycosaminoglycan tetrasaccharides, EDD analysis of, 19:1449
- Glycosaminoglycans
- bovine aorta, analysis of oversulfation using nano-ESI-Q-TOF-MS/MS and FTIRCR MS, 18:179
 - effect of charged state and sodium cationization on EDD and IRMRD, 19:790
 - heparin-like, MS/MS of, competing fragmentation processes in, 15:1534
 - tetrasaccharides, EDD of, 18:234
- Glycosides
- cardiac, fast atom bombardment of, ions formed by, high resolution mass spectrometry and tandem mass spectrometry analysis of, mechanistic consideration of the protonation and fragmentation of highly functionalized molecules in fast atom bombardment, 1:455
 - flavonoid, characterization of, by CAD of metal complexes, 12:537
- Glycosidic bond configuration, stereochemistry of, differentiation of, tandem mass spectrometry of diastereomeric cobalt-glycosyl-glycose disaccharide complexes, 7:953
- Glycosidic bond fragmentation, in gas phase, direction by non-covalent complexes, 15:715
- Glycosidic bonds
- anomericity of, by quadrupole ion trap, 10:269
 - CID of, alkyl derivatives in, 14:63
- Glycosphingolipid mixtures, complex, screening and sequencing by negative ion ESI-FTICR MS, 16:571
- Glycosphingolipids
- analysis of, quadrupole TOF mass spectrometer fitted with MALDI ion source for, 12:1220
 - electron ionization-tandem mass spectrometry of, identification of compound-specific sequence ions in the collision-induced dissociation spectra of the immonium ions of two isomeric hexaglycosylceramides, 3:353
 - as lithiated adducts, ESI-MS and CAD of, 12:61
 - NBD-tagged, UV-MALDI MS analysis, 19:923
 - neutral, structure determination of, by collisional activation of $[M + Li]^+$ ions, 3:260
 - permethylated, high-energy collision-induced dissociation of ceramide ions from, 5:558
- Glycosyl esters of nucleoside pyrophosphates, anomeric configuration of, by fast-atom bombardment tandem mass spectrometry, 6:516
- Glycosyl esters of nucleoside pyrophosphates and polyisoprenyl phosphates, anomeric configuration of, by fast-atom bombardment tandem mass spectrometry, 7:541
- Glycosyl transfer reactions, long-range, in gas phase, 13:325
- Glycosylation
- glycoproteins quantify changes by label-free quantitation, 20:1048
 - trastuzumab, heterogeneity profile, ESI-Q-IM-TOF MS, 20:2021
- N-Glycosylation, recombinant IgGs, quantification by LC or LC/MS, 19:1648
- Glycosylation site, of flavonoids, determining by metal complexation and MS/MS, 15:1287
- α -Glycyl cation, radical, and anion ($H_2NCH^+ / ^- / -COOH$), generation and characterization in the gas phase, 15:1241
- GlyGlyH⁺ carboxyl group, eliminating water from, 14:1192
- Glyquest, generation of N-glycan structure databases, 20:1739
- Gold
- CHCA-modified Au nanoparticles for peptide analysis, LDI-MS, 20:1530
 - nanoparticles as assisted matrices for biomolecules, MALDI-TOF-MS, 20:875
 - silver-coated nanoparticles as a matrix in SALDI MS analysis of aminoglycosides, 20:1925
- Gold ions, comparison of gold and cobalt ion cluster projectiles, molecular depth profiling, SIMS, 18:406
- Gold-silver clusters, bimetallic, MALDI of, 15:1455
- GONE. *See* Gas-phased oligosaccharide nonreducing end gp120 glycopeptides, from HIV, LC-MS of, 15:1545
- GPC. *See* Gel permeation chromatography entries.
- GPC-MALDI. *See* Gel permeation chromatography-matrix-assisted laser desorption ionization.
- Gradient chromatofocusing-mass spectrometry, protein analysis by, 19:1132
- Gradient reverse-phase high-performance liquid chromatography-electrospray ionization mass spectrometry, analysis with trifluoroacetic and other strong acid modifiers by postcolumn addition of propionic acid and isopropanol, signal enhancement of, 6:1221
- Gramicidin
- dimers show EX1 and EX2 H/D exchange mechanisms, 20:1813
 - noncovalent dimer of, affinity constants and response factors of, ESI-MS and mathematical modeling of, 16:1031
- Gramicidin S, gas phase ions in, hydration of, 8:565
- Gramicidin S and linear analogs, conformational preferences in gas phase, IM-MS of, 15:870
- Gramicidin S ions, electrospray-generated, in studies of ion trapping in FAIMS/FAIMS coupled to TOFMS, 12:1320
- Granulocytes
- calcyclin isolated from, nonspecific cleavage of, 12:1180
 - gangliosides from, 12:964
- Graphitized carbon black, oxidized nanoparticles, signal enhancement by, SALDI-TOF-MS, 20:1207
- Green tea, catechins, isomeric differentiation using gas-phase H/D exchange reactions, 18:1749
- Gridless ion acceleration systems for time-of-flight mass spectrometry, 4:782
- Groove-binding molecules, minor, and duplex DNA, noncovalent complexes formed between, by electrospray ionization-mass spectrometry, 6:1154
- Gross, Michael L., 16:1727
- Ground and excited states of the tungsten hexacarbonyl dication, double-ionization energies to, 2:108
- Ground loop current, signal intensities and, in ESI-MS, 15:1748
- Ground state Cu^+ and Fe^+ , the reactions of, with the 20 common amino acids, 7:722
- Group 15 tetrachloride anions, thermochemistry of, 13:469
- Group-selective ion-molecule reactions, functional, of ethylene glycol and its monomethyl and dimethyl ethers, 4:97
- Growth factors, ATP binding to, detected by MALDI-TOF and ESI-MS, 19:91
- Growth hormone-releasing factor analogs, helical content of, probing, using electrospray ionization mass spectrometry, 4:646
- Growth hormone secretagogue, MS/MS of, 13:371
- Guanidinium, complexes between ssDNA and derivatives of, MALDI characterization of, 17:283
- Guanidino-labeling derivitization, Tau-related peptide quantification using LC-IT-TOF MS with, 18:201
- Guanine, C8-substituted alkylaniline adducts of, differentiation by ESI and quadrupole ion trap MS/MS, 14:1488
- Guanosine
- in cluster with sinapinic acid, electron transfer vs proton transfer in, 12:176
 - H_2O adducts of, CID study of, 16:1291
 - in-source collision induced dissociation, 17:1050
- Guided ion beam mass spectrometer, ESI source for TCID with, 18:1124

- Guided ion beam MS/MS, to determine thermochemistry, 13:419
- Guided ion beam tandem mass spectrometry, mechanism of decomposition and, protonated asparagine, 20:852
- Gun, fast-atom bombardment saddle field, flux from, 5:100
- H
- H. *See* Hydrogen, H_3O^+ and $\text{H}_3\text{O}^+ (\text{H}_2\text{O})^3$ ion isotopic composition, in exchange reactions with water vapor, SIFT-MS analysis of, 11:866
- H^+ , sulfone and sulfoxide affinity of, ESI-FTICR of, 8:262
- H_6O , cirranulene and coronene, electron affinities of, estimation of, by using the kinetic method, 7:619
- $\text{H}_6\text{-C}_{23}$ linear alkenes, carbon-carbon double bonds in, from carbon addition reactions in a quadrupole ion trap equipped with a pulsed sample-inlet system, 2:105
- H/D exchange, gas phase
- DI *versus* D_2O , 11:167
- of polyamine complexes, 11:711
- H/D exchange. *See* Hydrogen/deuterium exchange
- H/D exchange MS. *See* Hydrogen/deuterium exchange mass spectrometry
- H-Gly-Gly-Sar-OH and H-Gly-Sar-Sar-OH, protonated, fragmentation pathways of, 14:1454
- H-POPs. *See* Halogenated persistent organic pollutants
- H-Resin, hydrogen silsesquioxane characterization by MS, 17:1437
- H side chains, fragmentation of deprotonated peptides containing, 12:1
- H-transfer and C-C bond formation between partners, and channeling of energy into dissociation, a photoionization study of the ion-neutral complexes $[\text{CH}_3\text{CH}_3 + \text{CH}_2\text{CH}_3]$ and $[\text{CH}_3\text{CH}_2\text{CH}^+\text{CH}_3\text{CH}_3]$ in the gas phase, 7:73
- H-transfers
- ion/ion, in linear ion traps, 18:882
- onium reactions and, characterization in closed *versus* open shell ions, 18:270
- ring-opening, in ionized cyclopentanone, Ab initio characterization of, similarity to ion-neutral complex-mediated alkane eliminations, 7:1251
- Hadamard transform (HT) time-of-flight (TOF) mass spectrometry
- peak height precision in, 16:1117
- two-channel, duty cycle and modulation efficiency of, 16:1888
- Hadamard transform TOFMS, sequence length, sequence frequency, and data acquisition rate effects on, 12:1302
- Hadamard transform-TOFMS, modulation defects and, 14:278
- Haemophilus ducreyi*, lipooligosaccharides from, poly lactosamine repeats in, 13:724
- Hair, drugs in, fast analysis of, 9:1311
- Halide ion-olefin clusters, 12:144
- Halide ions
- comparative study of the gas-phase bond strengths of CO_2 and N_2O with, 4:58
- gas-phase complexation of polyethers with, 4:242
- Halides
- diatomic gold, experimental and theoretical studies of, 13:485
- metallic, reactivity and binding energy with benzene, 13:477
- N-alkylnicotinium, class of cationic matrix additives for enhancing the sensitivity in negative ion fast-atom bombardment mass spectrometry of polyanionic analytes, 5:935
- $\text{S}_{\text{N}}2$ reactions with halodimethylamine, in gas phase, 15:673
- Halo alkylation, gas-phase, of C_{60} -fullerene by ion-molecule reaction under chemical ionization, 4:894
- Halodimethylamine, halide $\text{S}_{\text{N}}2$ reactions with, in gas phase, 15:673
- Halogenated persistent organic pollutants, analyzed by pLVI-GC/ECNI-qMS, 18:1375
- Halonitrobenzenes, nucleophilic substitution reactions of dinitrobenzene and, with carbanions, gas phase, 18:1351
- Hanuš, Vladimír, obituary, 20:2352
- Harsh environments
- aquatic, membrane inlet MS for online and in situ analysis of methane, 19:1395
- 3D gas concentration mapping using portable MS, 19:1411
- development of a palm portable MS, 19:1442
- gated electrostatic mass spectrometer, 19:1384
- hand portable GC-TMS for hazardous compound detection, 19:1425
- implementation of DASH and DESI MS on fieldable MS, 19:1419
- lake water testing of chemistry with portable and AUV-based MS, 19:1403
- mass spectrometry focus, editorial, 19:1375
- miniaturized mass analyzers, novel ion traps using planar resistive electrodes, 19:1435
- HASTE. *See* High amplitude short time excitation
- Hawthorn leaf, α -glucosidase inhibitors, characterization using LC-DAD-MS and SORI-CID FTICR MS, 20:1496
- Hck Src homology 2 domain receptor, identification and screening of, FTICR-MS in, 13:1162
- HCNH^+ association adducts with C_2H_2 and C_2H_4 , 12:557
- HD desktop, platform for analysis of H/D exchange data, 20:601
- HDX. *See* Hydrogen/deuterium exchange
- HDX-MS. *See* Hydrogen/deuterium exchange-electrospray mass spectrometry
- Heart, troponin C in, small Ca^{2+} -induced conformational changes in, 10:703
- Heart mitochondrial fractions, phosphatidylcholine isolated from, analysis by fast-atom bombardment tandem mass spectrometry, evidence of incorporation of monohydroxylated fatty acyl moieties, 7:50
- Heart tissue, MudPIT of, 16:1207
- Heated capillary electrospray source, charge-dependent/independent ion energy distribution from, selective generation of, 5:221
- Heated capillary inlet, magnetic sector mass spectrometer with, high resolution electrospray ion source for, design and performance of, 6:1256
- Heated electrospray source, for mass spectrometry of analytes from aqueous solutions, 5:791
- Heated nebulizer interface, liquid introduction with, atmospheric-pressure chemical ionization mass spectrometry, characterization of flame-generated C_{10} to C_{160} polycyclic aromatic hydrocarbons by, 7:276
- Heavy-atom substituents, effects of, on matrices used for matrix-assisted laser desorption-ionization mass spectrometry, 5:800
- Heavy gases, effects of, on the tandem mass spectra of peptide ions in the quadrupole ion trap, 7:1194
- He@C60, detection and quantitation by ultrahigh resolution FTICR-MS, 13:1349
- Helical content of growth hormone-releasing factor analogs, probing, using electrospray ionization mass spectrometry, 4:646
- Helical unfolding in peptides, the mass spectrometry of, 5:425
- Helium
- effect on fluorescence yield of trapped gas-phase ions, 17:855
- quantitative analysis of, by quadrupole ion trap MS, 13:1120
- Helium and air, effects of, on the performance of the ion trap detector when used for real-time monitoring of some atmospheric pollutants, 4:774

- Helium and argon as collision gases in the high energy collision-induced decomposition of MH^+ ions of peptides, comparison of, 3:502
- Helium-argon inductively coupled plasma mass spectrometry, matrix-induced interferences in, investigation of, 2:355
- Helix-turn-helix motifs, folding and unfolding in gas phase, 18:1239
- Heme-artemisinin noncovalent complexes, characterization by ESI-MS and CID-MS, 15:1181
- Heme binding in myoglobin, hemoglobin, and cytochrome c, by ion spray mass spectrometry, 4:631
- Heme-globin complex
detection in ESI-MS, electrostatic interactions and, 12:1092
dissociation of, by BIRD, 8:519
- Heme proteins, naked, metal atom oxidation state in, 11:120
- Hemispherical energy analyzer, high, for ion spectrometry, 2:506
- Hemocyanins, ESI-MS of, 14:419
- Hemoglobin
Camden variant, product ion isotope ratios from, ESI-MS/MS, 18:1493
folding and assembly monitored by ESI-MS, 18:8
gas phase H/D exchange and collision cross-sections, 20:484
Hb askew variant analysis by ECD, 20:1707
human, use of combined mass spectrometry methods for, the double mutant hemoglobin Villeparisis $\beta 77(EF1)$ His \rightarrow Tyr, $\beta 80(EF_4)$ Asn \rightarrow Ser, 7:163
from leech, subunit structure of, ESI-MS of, 15:22
of *Lumbricus terrestris*, MS of, 8:352
myoglobin and cytochrome c, heme binding in, by ion spray mass spectrometry, 4:631
of single intact cell, CE/ESI-TOFMS of, 10:184
subunit analysis by ESI-MS, 18:1279
structure probed by traveling-wave ion mobility MS, 20:625
- Hemoglobin α chain, ion trap CID of cations from, 17:923
- Hemoglobin "core," amino acid substitution in, an improved method to determine, electrospray ionization-tandem mass spectrometry analysis of peptides derived by enzymatic digestion of oxidized globin subunits, 7:1040
- Hemoglobin Ta-Li $\beta 83(EF7)$ Gly \rightarrow Cys variant, ES MS/MS of, 13:187
- Hemoglobin Villeparisis, double mutant, $\beta 77(EF1)$ His \rightarrow Tyr, $\beta 80(EF_4)$ Asn Ser, use of combined mass spectrometry methods for the characterization of a new variant of human hemoglobin, 7:163
- Hemolymph, crustacean, neuropeptide measurement by MALDI MS, 20:708
- Heparin
3-O-sulfated disaccharide isomers of, analysis of MS/MS, 20:652
structure determination of, by MS, 13:1254
- Heparin disaccharides, dissociation mechanisms of, ion trap MS/MS of, 15:1274
- Heparin-like glycosaminoglycans, MS/MS of, competing fragmentation processes in, 15:1534
- Heparin octasaccharide, effects of sulfate position on binding to CCL2, MS/MS, 17:1114
- Hepatitis C virus, E2 envelope glycoprotein glycosylation, MS characterization, 19:428
- Hepatocytes, rat, ethanol-treated, novel 3-hydroxylated leukotriene B_4 metabolites from, 3:762
- Hepatotoxins, cyclic heptapeptide, ESI-MS of, 10:1138
- Heptafluorobutyl derivatives of acyl analogues of platelet-activating factor, mass spectral fragmentation of, novel, 2:476
- Herbicides
chlorinated phenoxyacid, mass spectra of, obtained by particle beam liquid chromatography/mass spectrometry, evidence for thermal decomposition contributions to, 3:823
rearrangement reactions of, in MALDI-FTMS, 15:1820
- Herbicides paraquat and diquat, capillary electrophoresis-mass spectra of, 7:981
- Herbs, medicinal, saponins from, ESI-MS/MS structural analysis of, 15:133
- Heteroatom removal, transition-metal, by reactions of $FeL^+[L^+O, C_4H_6, c-C_5H_6, H_6H_6, C_5H_4(=CH_2)]$ with furan, thiophene and pyrrole in the gas phase, 7:938
- Heterocyclic isomers
characterization and differentiation of, mass spectrometry and molecular orbital calculations on pyrrolo[1,2-a][1,4]-benzodiazepin-4-one, -6-one, and -4,6-dione, 7:653
characterization and differentiation of tandem mass spectrometry and molecular orbital calculations on 3-methylisoxazolo- and 2-methylisoxazopyridines, 6:962
- Heterodimers, cationized, of amino acids, decompositions of, in relation to charge location in peptide ions, 4:461
- Heterodimers and homodimers, of amides and amines, proton-bound in gas phase, FTICR-MS of, 13:1273
- Heterogeneity, thiaminase I (42 kDa), sequence refinement, and active site location from high-resolution tandem mass spectrometry, 6:981
- Heteroxylans, acidic, uronosyl substitution patterns in, ESI-MS/MS of, 15:43
- Hewlett-Packard mass-selective detector, implementation of the chemical reaction interface mass spectrometry technique on, 5:765
- Hewlett-Packard particle-beam liquid chromatography-mass spectrometry interface by statistical experimental design, optimization of, 6:507
- Hexacarbonyl dication, tungsten, ground and excited states of, double-ionization energies to, 2:108
- Hexacarbonyl ions, tungsten, internal energy distributions, after neutralization-reionization, 5:1093
- 1,5-Hexadiene, ionized, structure in gas phase, 12:840
- 1,5-Hexadiyne radical cation, isomerization of, 8:599
- Hexafluorophosphate anions, electrochemically-induced reactions with water, ES-MS, 17:939
- Hexglycosylceramides, isomeric, two, immonium ions of, collision-induced dissociation spectra of, identification of compound-specific sequence ions in, electron ionization-tandem mass spectrometry of glycosphingolipids, 5:353
- Hexamethylcyclotrisiloxane, polymerization of, 14:95
- Hexapole accumulation, dual ESI source with, for mass measurement, 11:835
- Hexapole fields, added, linear quadrupoles with, 17:1063
- 1,2,4,5-Hexatetraene, isomerization of, 8:594
- Hexosamine monosaccharides, cobalt-coordinated, stereoselective dissociations of, 12:517
- Hexose monophosphates, analysis by ion-exchange chromatography and ESI-MS, 17:104
- HF-LPME. *See* Hollow fiber-liquid phase microextraction
- High- and low-energy collisionally activated decompositions of octaethylporphyrin and its metal complexes, 9:797
- High accuracy measurement of isotope ratios of molybdenum in some terrestrial molybdenites, 3:10
- High accuracy molecular weight determination and variation characterization of proteins up to 80 ku by ionspray mass spectrometry, 2:387
- High amplitude short time excitation, CID, detection of low mass product ions, quadrupole ion trap MS, 17:81
- High and low energy collision-induced dissociation studies of cluster ions on VG analytical ZAB-2FQ, compact high pressure ion source for, 3:33

- High efficiency photo-induced dissociation of precursor ions in a tandem time-of-flight mass spectrometer, 4:751
- High-efficiency surface-induced dissociation of rhenium oxide surface, 4:869
- High energy collision experiments, electron loss from multiply protonated lysozyme ions in, 12:889
- High energy collision-induced dissociation of ceramide ions from permethylated glycosphingolipids, 5:558
- of multiply charged polypeptides produced by electrospray, 4:652
- structural elucidation of O-linked glycopeptides by, 7:319
- tandem mass spectra of polycyclic aromatic hydrocarbon diolepoxide adducted peptides, characteristics of, 5:649
- tandem mass spectrometry, charged derivatives for, comparison of, 6:428
- High-energy collision-induced dissociation in tandem mass spectrometry, low-mass ions produced from peptides by, 4:882
- High-energy collision-induced dissociation mass spectrometry, of synthetic mannose-6-phosphate oligosaccharides, 7:182
- High-energy collision-induced dissociation spectra of peptides, uninterpreted, search of, with sequence databases, 7:1089
- High-energy collision-induced dissociation spectra using artificial neural networks, peptide sequence determination from, 6:947
- High-energy collisional activation, delayed dissociation spectra of survivor ions from, 4:845
- High energy collisional activation of multiply protonated ions, complete sequence confirmation of large peptides by, 6:234
- High-energy collisional activation studied via angle-resolved translational energy spectra of survivor ions, 4:835
- High-energy fragmentation of protonated peptides, determination of, using a BEqQ hybrid mass spectrometer, 4:470
- High field asymmetric waveform ion mobility spectrometry (FAIMS)
- carrier gases for, 11:1125
- cylindrical geometry prototypes for, of differing electrode diameters, 16:349
- for differentiating disaccharide anomers, linkage, and position isomers, 14:265
- of ubiquitin cross-sections, 11:738
- of ubiquitin elongated conformers, 12:894
- High field asymmetric waveform ion mobility spectrometry-mass spectrometry: investigation of leucine enkephalin ions produced by ESI, 10:492
- High-field FTICR-MS for simultaneous trapping and gas phase hydrogen/deuterium exchange of peptide ions, 9:1012
- High-frequency Fourier transform ion cyclotron resonance mass spectrometry, 4:177
- High magnetic field, concentric tube vacuum chamber for, high-pressure ionization in a Fourier transform ion cyclotron resonance mass spectrometer, 4:168
- High mass, line broadening at, and for large ion populations, a mechanism for, two-dimensional Coulomb-induced frequency modulation in Fourier transform ion cyclotron resonance, 4:909
- High mass detection in Fourier transform mass spectrometry, ion cloud manipulation using the radiofrequency-only mode as an improvement for, 4:186
- High mass performance, poor, in Fourier transform mass spectrometry, a mechanism for, 3:460
- High mass ranges, calibration of, 10:273
- High mass-to-charge-ratio ions from electrospray ionization mass spectrometry, observations and implications of, 4:536
- High mass-to-charge ratio quadrupole mass spectrometer, multiply charged proteins in, proton transfer reaction studies of, 5:1064
- High mobility group 1 proteins, study on acetylation, LC-MS/MS, 18:1569
- High molecular weight peptides and proteins in a two-stage linear time-of-flight mass spectrometer, photodissociation of, 6:578
- High-molecular-weight polycyclic aromatic hydrocarbons in soils by particle beam high-performance liquid chromatography-mass spectrometry, measurement of, 6:597
- High-order structure and dissociation of gaseous peptide aggregates that are hidden in mass spectra, 9:743
- High-order structure of proteins, probing by fast-atom bombardment mass spectrometry, 5:19
- High performance Fourier transform ion cyclotron resonance mass spectrometry via a single trap electrode, 6:812
- High performance liquid chromatography
- pharmaceuticals in surface waters, matrix effects in HPLC-MS/MS and UPLC-MS/PM analyses, 19:713
- wheat gluten proteins, HPLC/MALDI-TOF-MS, 19:1542
- High-performance liquid chromatography, gradient reverse-phase, electrospray ionization mass spectrometry analysis with trifluoroacetic and other strong acid modifiers by postcolumn addition of propionic acid and isopropanol, signal enhancement of, 6:1221
- High-performance liquid chromatography atmospheric pressure chemical ionization mass spectrometry, quantitative analysis by, the determination of renin inhibitor CP-80,794 in human serum, 2:164
- High performance liquid chromatography/atmospheric pressure microwave-induced plasma ionization mass spectrometry (HPLC/AP-MIPI-MS), of underivatized amino acids, 12:117
- High performance liquid chromatography-DAD-tandem mass spectrometry, hydrolytic and photodegradation of tribenuron methyl, 19:1221
- High performance liquid chromatography-diode array detection, toxicological screening using, 14:14
- High performance liquid chromatography-ESI-MS/MS, on-line strong cation exchange, 14:719
- High performance liquid chromatography-ESR
- multidimensional nano, for protein complex analysis, 14:1003
- in separation and identification of oxygen-centered radical DMPO adducts, 14:862
- High performance liquid chromatography (HPLC)
- in confirmation/quantitation of cocaine and benzoylecgonine in urine, 11:257
- identification of new metabolite of rapamycin and SDZ RAD by, 11:516
- protein identification by, 12:215
- reversed-phase, ion trap MS with, as monoclonal antibody trap, 16:307, 16:797 (erratum)
- High performance liquid chromatography (HPLC) diode array detection (DAD), of *Ophiopogon japonicus* homoisoflavonoids, 16:234
- High performance liquid chromatography (HPLC) electrospray ionization (ESI) mass spectrometry, hydrogen/deuterium exchange, *t*-Boc group elucidation using, 16:553
- High performance liquid chromatography-mass spectrometry, SSI vs APCI as interface of, in drug analysis, 13:204
- High-performance liquid chromatography-mass spectrometry of porphyrins by using an atmospheric pressure interface, 7:965
- High-performance liquid chromatography mobile phase components, effect of, on sensitivity in negative atmospheric

- pressure chemical ionization liquid chromatography-mass spectrometry, 7:1059
- High-performance liquid chromatography particle beam mass spectrometry, investigation of enhanced ion abundances from a carrier process in, 1:92
- High-performance liquid chromatography-particle beam-mass spectrometry, determination of polycyclic aromatic hydrocarbons by, 4:898
- High performance liquid chromatography tandem mass spectrometry, in quantitation of zidovudine, lamivudine, and stavudine metabolites in peripheral mononuclear blood cells of HIV-infected patients, 11:1134
- High performance liquid chromatography-tandem mass spectrometry
 nitrogen-containing intracellular metabolites in *S. enterica*, 17:37
 of plasma atorvastatin and active metabolites, 10:55
 thiolated deoxyoligonucleotide characterization and sequence variation, 17:1396
- High performance low magnetic field internal electrospray ionization-Fourier transform ion cyclotron resonance mass spectrometer, 7:915
- High pressure chemistry in a FT-ICRNMS using a split-pair magnet, 10:918
- High pressure drift reaction cell to study thermal energy reactions of mass-selected ions, a hybrid double-focusing mass spectrometer, 1:197
- High pressure ion source, a compact, for high and low energy collision-induced dissociation studies of cluster ions on a VG analytical ZAB-2FQ, 3:33
- High-pressure ionization in a Fourier transform ion cyclotron resonance mass spectrometer, concentric tube vacuum chamber for high magnetic field, 4:168
- High pressure liquid chromatography (HPLC), of histone modifications in leukemia, 15:77
- High pressure MALDI-FTMS for minimization of ganglioside fragmentation, 13:402
- High-pressure quadrupole collision cell, characterization of, for low-energy collision-induced dissociation, 5:1042
- High pressure segmented quadrupole collision cell (HP-SQCC), as ion molecule reactor, 16:957
- High pressure trapping in Fourier transform mass spectrometry, a radiofrequency-only-mode event, 3:590
- High-resolution accurate mass measurements of biomolecules using a new electrospray ionization ion cyclotron resonance mass spectrometer, 4:566
- High resolution and tandem Fourier-transform mass spectrometry with Californium-252 plasma desorption, 1:427
- High resolution characterization of DNA fragment ions produced by US MALDI using linear and reflecting TOFMS, 10:423
- High resolution electrospray ion source for magnetic sector mass spectrometer with heated capillary inlet, design and performance of, 6:1256
- High resolution end group determination of low molecular weight polymers by matrix-assisted laser desorption ionization of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, 7:449
- High-resolution ESI-FTMS with infrared multiphoton dissociation of glucokinase from *Bacillus stearothermophilus*, 12:22
- High-resolution glow discharge mass spectrometer operated with an argon/hydrogen mixture, relative ion yields measured with, 4:278
- High-resolution imaging mass spectrometry, automated image alignment of biological samples, 19:823
- High-resolution ion isolation with the ion cyclotron capacitively coupled open cell, 6:533
- High resolution ion mobility time-of-flight mass spectrometry, of synthetic failures in combinatorial libraries, 11:352
- High-resolution mass spectrometer, the development of, a reminiscence, 2:447
- High resolution mass spectrometry, for synthetic chemistry, 10:546
- High-resolution mass spectrometry, tandem, of large molecules, Fourier-transform electrospray instrumentation for, 4:557
- High resolution mass spectrometry and tandem mass spectrometry, analysis of the ions formed by fast atom bombardment of digoxin and related glycosides, mechanistic considerations of the protonation and fragmentation of highly functionalized molecules in fast atom bombardment, 1:455
- High-resolution mass spectrometry of large molecules in a linear time-of-flight mass spectrometer, 4:2
- High resolution on a quadrupole ion trap mass spectrometer, 2:198
- High resolution power, in Fourier transform ion cyclotron resonance mass spectroscopy, remeasurement at, 6:564
- High-resolution tandem mass spectrometry, thiaminase I (42 kDa) heterogeneity and sequence refinement, active site location and, 6:981
- High-resolution triacylglycerol mixture analysis using high-temperature gas chromatography/mass spectrometry with a polarizable stationary phase, negative ion chemical ionization, and mass-resolved chromatography, 7:350
- High sensitivity analysis of phenylthiohydantoin amino acid derivatives by ESI-MS, 8:1165
- High-speed analysis, automated, of selected organic compounds in urban air by on-line isotopic dilution cryofocusing gas chromatography/mass spectrometry, 5:1001
- High-speed quantitative mass spectrometry, with MALDI ionization, design considerations, 17:1129
- High-temperature gas chromatography/mass spectrometry with a polarizable stationary phase, negative ion chemical ionization, and mass-resolved chromatography, high-resolution triglycerol mixture analysis using, 7:350
- High throughput approach to quantitative proteomic analysis, expression-dependent protein identification by MS, 12:1238
- High-throughput sample preparation and analysis using 96-well membrane solid-phase extraction and LC-MS/MS for the determination of steroids in human urine, 10:1322
- High throughput screening, for binding to RNA, 15:884
- High transmission hemispherical energy analyzer, for ion spectrometry, 2:506
- High voltage RF oscillator for driving multipole ion guides, 13:1370
- Highly functionalized molecules in fast atom bombardment, protonation and fragmentation of, mechanistic considerations of, high resolution mass spectrometry and tandem mass spectrometry analysis of the ions formed by fast atom bombardment of digoxin and related cardiac glycosides, 1:455
- Histidine
 effect of oxidation on peptide ion dissociation patterns, MS/MS, 18:553
 ternary copper (II) complexes, macrocyclic ligand effects on electron and proton transfers, 20:972
- Histidine adducts, in cytochrome c modification by 4-hydroxy-2-nonenal, 15:1136
- Histidine and lysine, gas-phase basicities of, and their selected di- and tripeptides, 7:1203

- Histidine residues, position of, effect on CID of triply protonated tryptic peptides, 12:1262
- Histone, modifications in leukemia, HPLC and MS of, 15:77
- Histone H4 acetylated isoforms, from lymphosarcoma, peptide mass mapping of, 16:1641
- Hites, Ronald A., award for outstanding research publication, editorial, 19:1577, 19:1578
- HIV
- binding sites of TAR and TAR mutants for peptide ligands, LILBID, 19:1605
 - envelope glycoproteins, comparison of HPLC/ESI-FTICR MS vs. MALDI-TOF/TOF MS, 19:1209
 - p24, epitope mapping using chemical modification and MS, 17:1560
 - packaging signal, mapping noncovalent ligand binding to stemloop domains, MS-MS, 17:1401
- HIV-1 integrase, investigation of dimeric G-quadruplex inhibitors by ESI MS, 19:550
- HIV-infected patients, zidovudine, lamivudine, and stavudine metabolites in peripheral mononuclear blood cells, HPLC-MS/MS of, 11:1134
- H₂O adducts, of guanosine, CID study of, 16:1291
- Hollow fiber-liquid phase microextraction, with GC-ICP-MS, PBDE analysis in soil, dust and human serum, 18:1740
- Homoconjugated octadecadienoic acids, collisional activation of a series of, with fast atom bombardment and tandem mass spectrometry, 1:183
- Homoisoflavonoids, HPLC-DAD-ESI MS analysis of, 16:234
- Homolytic cleavages in pyridinium ions, an excited state process, 790
- Homomyoglobin, ion-ion reactions of, 8:637
- Honeybee venom, detection in envenomed tissues, MALDI MSI, 20:112
- Hop extract, bitter acids from, LC-APCI-MS of, 15:180
- HOPG, surface-induced dissociation and chemical reactions on stainless steel, carbon and diamonds, 20:927
- Hormones, veterinary drug residue analysis and, UPLC-Orbitrap MS, 20:451
- Host-guest complexation, binding selectivity determination in, ESI-MS and quadrupole ion trap in, 9:1049
- Hot-wire chemical vapor deposition, LIEI application in, SiH₄-NH₃ gas mixture, 18:1950
- How do dimethyl oxalate ions CH₃O-C(=O)-C(=O)-OCH₃+ break in half? Loss of CH₃• + CO₂ versus CH₃O-C=O, 10:869
- How to measure the kinetic energy distribution of the precursor ion main beam in mass-analytical ion kinetic energy spectrometry experiments, 6:1232
- HPLC. *See* High performance liquid chromatography
- HPLC-MS/MS. *See* High performance liquid chromatography-tandem mass spectrometry
- [HS-P-OH]^{+/*}, ionic and neutral, generation and characterization in gas phase, 16:1353
- HT2a agonist and metabolite, in plasma and brain, LC-MS/MS of, 8:371
- Human adipose tissue, highly chlorinated dimethanofluorenes in, 1:99
- Human hemoglobin, new variant of, characterization of, use of combined mass spectrometry methods for, the double mutant hemoglobin Villeparisis β77(EF1) His → Tyr, β80 (EF₄) Asn → Ser, 7:163
- Human immunodeficiency virus-gp120 epitopes, MALDI-MS of, 9:157
- Human immunodeficiency virus (HIV), gp120 glycopeptides from, LC-MS of, 15:1545
- Human leukocyte antigen A*0201 class I, from vaccinia virus, identification by mass FTICR MS/MS, 16:1812
- Human plasma
- chemically oxidized dexamethasone in, detection of, electron capture negative ionization in, comparison of tandem and conventional mass spectrometry using, 1:341
 - SR 27417 in, quantitation of, using electrospray liquid chromatography-tandem mass spectrometry, a study of ion suppression, 7:1099
- Human serum albumin, warfarin interactions with, effect of tan-shinone IIA on, ESI MS, 19:1568
- HWCVD. *See* Hot-wire chemical vapor deposition
- HX-Express, data processing of hydrogen exchange mass spectra using, 17:1700
- HX MS. *See* Hydrogen exchange mass spectrometry
- Hyaluronic acid, distinguished from N-acetylheparosan by MS/MS, 19:82
- Hybrid (BEqQ) mass spectrometer, charge-remote fragmentation in, to determine isotopic purity in selectively polydeuterated surfactants, 1:85
- Hybrid double-focusing mass spectrometer, high pressure drift reaction cell to study thermal energy reactions to mass-selected ions, 1:197
- Hybrid linear ion trap/Fourier transform-ion cyclotron resonance mass spectrometry, glycoprotein characterization using, 17:168
- Hybrid tandem mass spectrometric screening and characterization of the glutathione conjugate of 2-furamide, multiple scan modes in, 2:55
- Hydrated ion nanocalorimetry, nonergodicity in ECD investigated by, 18:1217
- Hydration
- intrinsic, of monopositive uranyl hydroxide, nitrate, and acetate cations, 15:777
 - mechanisms of, deproteinated amino acids, 20:632
 - microhydration of protonated nucleic acid bases, 20:1900
- Hydration energies, from gas-phase equilibria measurements, deprotonated AAs, 19:1091
- Hydration of gas phase gramicidin S (M + 2H)²⁺ ions formed by electrospray: transition from solution to gas phase structure, 8:565
- Hydration of gas-phase ions formed by ESI, 10:958
- Hydrazine, complexes with ketones, mobility resolution and mass analysis of ions from, 18:940
- Hydrocarbon dendrimers, analysis of, by laser desorption time-of-flight and Fourier transform mass spectrometry, 5:731
- Hydrocarbon diolepoxide adducted peptides, polycyclic aromatic, characteristics of high energy collision-induced dissociation tandem mass spectra of, 5:649
- Hydrocarbon emission species, automobile, analysis, nitric oxide-assisted atmospheric pressure corona discharge ionization for, 5:1107
- Hydrocarbon mass spectrometry, in situ silver cationization for, 16:437
- Hydrocarbon plaques, MALDI-FTICR and reflection TOF MS of, 16:2009
- Hydrocarbon polymers
- analysis of, by laser desorption Fourier transform mass spectrometry, silver nitrate chemical ionization for, 4:453
 - MALDI-FMS of, 8:225
- Hydrocarbon radical cations, polycyclic aromatic, reactions with model biological nucleophiles, 7:628
- Hydrocarbons
- aromatic, Fourier transform ion cyclotron resonance mass spectrometry of, ammonia laser desorption/chemical ionization with ammonium bromide, 2:299
 - CH₃OCH₂⁺ reactions with, 13:1015
 - GC-MS analysis of Cu⁺ chemical ionization for, 9:731, 9:981
 - non-polar isomeric, APCI studies of, 13:1078

- polynuclear aromatic, nitro-substituted, in airborne particulate matter and soil, sub-parts-per-billion determination by electron capture-tandem mass spectrometry, 7:1255
- pure, in Titan's atmosphere, ion-molecule reactions in, 14:900
- SAMs of, methyl cation neutralization at, 13:1151
- saturated, molecular mass determination of, 14:51
- Hydrocarbons polycyclic aromatic. *See* Polycyclic aromatic hydrocarbon
- C_{10} to C_{160} , flame-generated, characterization of, by atmospheric-pressure chemical ionization mass spectrometry with liquid introduction via heated nebulizer interface, 7:276
- characterization of, fast atom bombardment combined with mass spectrometry for, 1:405
- determination of, by high-performance liquid chromatography-particle beam-mass spectrometry, 4:898
- Hydrofluorocarbon-245fa, EPA recommendation of, ion chemistry in air plasma and reactions with atmospheric ions, 16:1081
- Hydrofluoroether-7100, EPA recommendation of, ion chemistry in air plasma and reactions with atmospheric ions, 16:1081
- Hydrogen
- measurement of, CRIMS in, 11:400
 - migration upon electron transfer, arginine-containing peptide ions, 19:1726
 - peptide amide, migration of, during ESI, 12:410
 - quantitative analysis of, by quadrupole ion trap MS, 13:1120
 - scrambling, anionic peptides, collisional activation by MALDI TOF/TOF MS, 19:1719
 - synchronized concerted 1,4- H_2 elimination from $CH_3CH=N^+HCH_3$, 19:27
- Hydrogen abstraction
- diuretic fragmentation and, in negative ESI-MS, 14:658
 - long-lived ions from ECD show radical migration via, 17:576
 - in N—C bond cleavage of peptide backbone, 12:1044
- Hydrogen/argon mixture, high-resolution glow discharge mass spectrometer operated with, relative ion yields measured with, 4:278
- Hydrogen atom
- abstraction from ethanol, phenyl radical selectivity in, 15:913
 - host-guest transfer induced by ECD, 20:639
- Hydrogen bonding
- multiple, in negative ion adduct stability in electrospray MS, 13:1360
 - role in peptide cation survival after electron capture, 17:1675
- Hydrogen bonding in transient bifunctional hypervalent radicals by neutralization-reionization mass spectrometry, 6:1004
- Hydrogen-bonding interactions in gas-phase polyether/ammonium ion complexes, 5:260
- Hydrogen bonds in gas phase clusters between halide ions and olefins, 12:144
- Hydrogen/chlorine exchange, of gaseous carbanions, 16:2045
- Hydrogen cyanide, in situ generation for MS, 17:51
- Hydrogen/deuterium exchange, 10:9
- accuracy of MALDI-MS and SUPREX, protein unfolding free energy determination, 17:1535
 - amide, modeling data from, in protein-ligand interactions, 15:388
 - arginine containing di- and tri- peptides, 20:2049
 - β_2 -M release from MHC-I, analyzed by HDX-ESI-MS, 20:278
 - in bacterial cell wall precursor peptide interaction with vancomycin antibiotics, 9:1255
 - between [m-1] ions of chlorinated benzenes and D_2O or ND_3 , mechanistic study of, 5:282
 - electrospray ionization source, LC-MS and LC-MS/MS of metabolites, 13:345
 - ESI-MS with, in calcitonin folding analysis, 15:1556
 - EX1 and EX2 exchange mechanisms, for gramicidin dimers, 20:1813
 - EX1 characterization of kinetics in by peak width analysis, MS, 17:1498
 - of fixed charge derivative aspartic acid peptides, 16:1067
 - in fragmentation pathway studies of oligonucleotides in MALDI-MS, 13:1407
 - gas phase
 - 5'- and 3'-mononucleotides, quadrupole ion trap, 18:724
 - collision cross-sections and, hemoglobin, 20:484
 - dinucleotide negative ions, 18:1827
 - in electrospray ionization mass spectrometry as a practical tool for structure elucidation, 5:434
 - green tea catechin isomeric differentiation using, 18:1749
 - as a molecular probe for the interaction of methanol and protonated peptides, 5:526
 - sodiated amino acids, 17:29
 - of gas phase bradykinin ions in a linear quadrupole ion trap, 85
 - of gas phase peptides and peptide dimers, acidic residues and sodium ion adduction in, 14:1477
 - HD desktop platform for analysis, 20:601
 - HDX MS, protein conformations probed using ETD in, 20:1514
 - of hydrophobic peptides in model membranes by ESI-MS, 13:1376
 - in-source, MALDI-FTICR-MS of, 13:308
 - in indirect assessment of hydrophobic ligand binding, 14:506
 - local protein hydrogen exchange data from HDX CAD MS using deconvolution, 17:1543
 - mass spectra data processing using *HX-Express*, 17:1700
 - mass spectrometry and, historical perspective, 17:1481
 - methyl guanine isomer distinction by, using a Fourier transform mass spectrometer, 4:296
 - MS/MS with, in sulfadiazine metabolite structural characterization, 16:1687
 - in multidimensional separations of ubiquitin conformers in gas phase, 16:1427
 - of peptide ions, simultaneous trapping with, FTICR-MS for, 9:1012
 - of proteins
 - of calmodulin interaction with calcium, 10:711
 - continuous *versus* pulsed labeling/MS of, 10:675
 - of fluorophenyl anions, 10:840
 - gas-phase, of fluorophenyl anions, 10:840
 - of peptide conformational changes, 10:719
 - in solution, MS of, 10:671, 10:672
 - of protonated amino acids, in gas phase, 16:1846
 - RXR ligand interactions, MS and, 17:1510
 - SIMSTEX application to oligomerization of insulin analogs and mutants, 17:1526
 - solution phase, local stability of *R. capsulatus* cytochrome c_2 probed by MS and, 17:1518
 - solvent accessibility of protein surfaces, MALDI-TOF MS, 17:1490
 - to study DNA decay under UV MALDI, 9:866
 - temperature-dependent, in gas phase studies of cytochrome c ions, 13:506
 - in tetracycline characterization, 13:543
 - on yeast ATPase supramolecular protein complex analyzed at high sensitivity by MALDI-MS, 14:471
- Hydrogen deuterium exchange, FTICR MS/MS integration with, in monitoring intermolecular migration during collisional activation, 15:639
- Hydrogen-deuterium exchange behavior, bimolecular, of protonated alkyl dipeptides in the gas phase, 6:466

- Hydrogen/deuterium exchange-electrospray mass spectrometry, triaxial probe for proteolysis coupled with, 18:208
- Hydrogen/deuterium exchange high performance liquid chromatography (HPLC) electrospray ionization (ESI) mass spectrometry, *t*-Boc group elucidation using, 16:553
- Hydrogen/deuterium exchange infrared spectrometry, zwitterion formation in gas-phase H/D-exchange with CH₃OD, 18:512
- Hydrogen/deuterium exchange mass spectrometry
 efficiency of MS-based screening assay for detection of ligand binding, 19:1303
 oligonucleotide gas-phase conformers, use of ion mobility techniques and, 19:938
 of protein-ligand binding, 16:869
 SAX for studying protein/DNA interactions by, 19:887
- Hydrogen/deuterium exchange reactions
 complexation of protonated peptides with saccharides in gas phase decreases rates of, 6:1247
 of emolate ions with CH₃OD and C₂H₅OD, reactive collisions in quadrupole cells, 3:853
- Hydrogen/deuterium (H/D) exchange
 data from, automatic analysis of, 12:1153
 gas phase
 of AMP and dAMP, conformational dependence of, 12:780
 of model peptide, 12:1272
 at non-labile sites, a new reaction facet with broad implications for structural and dynamic determinations, 12:1163
 of positively charged mononucleotides, FTICR-MS in, 12:268
 of positively charged mononucleotides by use of FTICR-MS, 12:268
 solution phase, of non-covalent complexes, ESI-MS of, 12:795
 in study of melittin bound to phospholipid micelles, 12:1247
 in study of oligonucleotide fragmentation, 12:193
- Hydrogen exchange, to locate regional changes in oxidized and reduced cytochrome *c*, 8:1039
- Hydrogen exchange mass spectrometry, improved by UPLC, 17:163
- Hydrogen exchange mass spectrometry study, of diphosphorylated ITAM binding to spleen tyrosine kinase, conformational dynamics in, 16:1039
- Hydrogen migration, double, in endo tricyclo [5.2.1.0_{2,6}] decene, 13:1186
- Hydrogen peroxide, laser flash photolysis of, for microsecond photochemical oxidation of proteins, 16:2057
- Hydrogen protein exchange rates, changes in, that result from point mutations, mass spectrometric measurement of, 7:515
- Hydrogen rearrangements, in ECD, peptides, 18:113
- Hydrogen silsesquioxanes, characterization by MS, 17:1437
- Hydroiodic acid attachment kinetics, as a chemical probe of gaseous protein ion structure: bovine pancreatic trypsin inhibitor, 10:552
- Hydrolysis, PtenCl₂, investigated by electrospray MS and DFT, 20:1015
- Hydroperoxides, lipid, and long-chain conjugated keto acids, analysis of, by negative ion electrospray mass spectrometry, 6:1190
- 5-Hydroperoxy and 5-hydroxyeicosatetraenoic acids, ESI-MS of, 9:527
- Hydroperoxy bishemiacetal product, derived from cholesterol ozonolysis, ESI-MS/MS of, 15:194
- Hydrophilic compounds, ultratrace determination of, 10:1328
- Hydrophobic ligand binding, indirect assessment of, ESI-MS and H/D exchange in, 14:506
- Hydrophobic peptides, H/D exchange of, by ESI-MS, 13:1376
- Hydrophobic surface chemistry and morphology, effect on digestion of surface-bound proteins, 14:203
- Hydrophobic surfaces, conductive, liquid matrix deposition on, UV-MALDI MS, 18:693
- Hydrosphere, chemical analysis of, underwater MS for, 12:677
- 4-Hydroxy-2-alkylquinolines, from *Pseudomonas aeruginosa*, ESILC-MS of, 15:862
- 4-Hydroxy-2-nonenal
 cytochrome *c* modification by, 15:1136
 protein modification by, ESI-MS/MS of, 215
- Hydroxy-carboxylic acid, TMS derivatives of oxo-carboxylic acid and, fragmentation of, EIMS, 19:66
- Hydroxy fatty acids as pentafluorobenzyl ester, trimethylsilyl ether derivatives by electron ionization gas chromatography/mass spectrometry, analysis of, 6:40
- Hydroxy radicals, oxidation of, protein interface footprinting quantified by MDS and, 19:1686
- 2-Hydroxybenzyl-*N*-pyrimidinylamine derivatives, gas phase retro-*Michael* fragmentation reactions of, 16:1552
- 3-Hydroxycoumarin, new matrix for MALDI-TOF MS, 17:1665
- Hydroxyl radical adducts, MS of, 12:1214
- Hydroxyl radical footprinting, proteins, aliphatic peptidyl hydroperoxides as source of secondary oxidation, 20:1123
- Hydroxyl radical probe, of calmodulin-melittin complex interface, 16:225
- 2-Hydroxyestradiol, deuterated, deoxyguanosine linkage to, LC/ESI-ion trap MS of, 14:364
- Hydroxymethylpropanal, analytical and atmospheric chemistry of, ITMS in, 13:530
- Hydroxyoximes, beta, aromatic (M + H)⁺, (M + Cat)⁺, and (M-H)⁻ ions of, charge-remote fragmentation and the two-step elimination of alkanols from, 4:819
- 2-Hydroxyxylchalcone, protonated 2-methoxychalcone and, Nazarov cyclization, 20:805
- 2-Hydroxyoxolan-2-yl radical, modeling by neutralization-reionization MS, 15:1055
- 3-Hydroxyoxolan-3-yl radical, modeling by neutralization-reionization MS, 15:1068
- Hyperbolic ion trap, for Fourier transform ion cyclotron resonance mass spectrometry, experimental formation of, 3:188
- Hyperthermal ion beams, at self-assembled monolayer surfaces, X3F⁺, IF⁺, and other unusual ions generated by, 4:938
- Hypervalent pyrrolidinium radicals, by neutralization-reionization mass spectrometry: metastability and radical leaving group abilities, 9:242
- I
- IBF. *See* Induction-based fluidics
- ICAT MS. *See* Isotope coded affinity technology mass spectrometry
- ICM-MS. *See* Intact cell-mass spectrometry (ICM-MS)
- ICP. *See* Inductively coupled plasma entries
- ICP-DRC-MS. *See* Inductively coupled plasma mass spectrometry
- ICPMS. *See* Inductively coupled plasma mass spectrometry
- ICR-MS. *See* Fourier transform ion cyclotron resonance mass spectrometry
- ICR MS. *See* Ion cyclotron resonance mass spectrometer
- ICR-MS/MS. *See* Ion cyclotron resonance tandem mass spectrometry
- ICRMS/MS, new approach for effecting surface-induced dissociation in, a modeling study, 11:1107
- Ideal velocity focusing in a reflection TOFMS, 10:992
- Identification, correct, estimating probabilities of, from results of mass spectral library searches, 5:316
- Identification and localization of fatty acid modification in ghrelin by ECD, 13:1443
- Identification of a new fragment ion type in the collision-in-

- duced dissociation spectra of peptides, formatin of alpha-2-16 ions, 4:424
- Identification of a new metabolite of macrolide immunosuppressant, like rapamycin and SDZ RAD, using HPLC and electrospray MS/MS, 11:516
- Identification of an in vitro IRS-1 phosphorylation site by negative-ion μ LC/ES-API-CID-MS hybrid scan technique, 14:401
- Identification of components in waste streams by electrospray and tandem mass spectrometry, 4:604
- Identification of deoxynucleoside-polyaromatic hydrocarbon adducts by capillary zone electrophoresis-continuous flow-fast atom bombardment mass spectrometry, 3:757
- Identification of diethylene glycol monobutyl ether as a source of contamination in an ion trap mass spectrometer, 14:1247
- Identification of face-to-face inclusion complex formation of cyclodextrin bearing an azobenzene group by ESI-MS, 14:1116
- Identification of GlcNAcylation sites of peptides and α -crystallin using Q-TOFMS, 12:1106
- Identification of in vitro metabolites of Indinavir by intelligent automated LC-MS/MS (INTAMS) utilizing triple quadrupole MS/MS, 10:175
- Identification of noncovalent dimeric complexes of the recombinant murine S100 protein CP10 by ESI-MS and chemical cross-linking, 9:533
- Identification of oxidation products and free radicals of tryptophan by MS, 14:406
- Identification of phosphorylation sites in phosphopeptides by positive and negative mode electrospray ionization-tandem mass spectrometry, 7:243
- Identification of photocatalytic degradation products of diazinon in TiO₂ aqueous suspensions using GC-MS/MS and LC-MS with quadrupole TOFMS, 14:803
- Identification of single stranded regions of DNA by enzymatic digestions with MALDI analysis, 10:521
- Identification of tetracycline antibiotics by ESI in a quadrupole ion trap, 9:1089
- Identification of vertebrate type steroid hormones in the shrimp *Penaeus japonicus* by tandem mass spectrometry and sequential product ion scanning, 8:365
- Identifying sites of protein modification by using matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry and on-line membrane preconcentration-capillary electrophoresis-tandem mass spectrometry, 8
- IDMS. *See* Isotope dilution mass spectrometry
- IF⁺, XeF⁺, and other unusual ions generated by reactions of hyperthermal ion beams at self-assembled monolayer surfaces, 4:938
- Illicit drugs, LC-MS/MS bioanalysis of, matrix effect in, 14:1290
- ILMs. *See* Ionic liquid matrices
- ILPR. *See* Insulin-linked polymorphic region
- IM-MALDI. *See* Ion mobility matrix-assisted laser desorption ionization.
- Image alignment, automated, of biological samples, high-resolution imaging MS, 19:823
- Image analysis, polymer characterization with, 13:1195
- Image charge and Coulomb mutual interaction, in Fourier transform ion cyclotron resonance precise mass difference measurements, systematic errors originating from, analysis and estimation of, 4:855
- Imaging
 large insulating samples, by using secondary ion tandem mass spectrometry, charge compensation for, 5:37
 organic ion, beyond the limit of static secondary ion mass spectrometry, 6:1047
 secondary ion, of developing brain, 15:1116
 a wide-angled secondary ion probe for, 2:362
- Imaging mass spectrometry, spatial, spectral and conformational resolution, 20:1006
- Imaging/profiling mass spectrometry, preparation of single cells for, 19:1230
- Imidazolium, N,N'-disubstituted, ESMS fragmentation study of ionic liquids, 17:85
- Imine/methylol adduction, peptides with N-terminal dimethylation and, MS/MS, 17:859
- Immobilized metal affinity chelates, comparison of TiO₂ surfaces and, in protein phosphorylation studies, 18:1932
- Immonium ions in the gas stage, applications of multiple stage mass spectrometry in a pentaquadrupole, polar [4+2⁺] Diels-Alder cycloaddition to, 6:1
- Immonium ions of two isomeric hexaglycosylceramides, collision-induced dissociation spectra of, identification of, electron ionization-tandem mass spectrometry of glycosphingolipids, 3:353
- Immunoassays, based on single-particle mode detection by ICP MS, 20:1096
- Immunoglobulin G
 nonenzymic glycation of, MALDI-MS, enzymatic digestion, and molecular modeling in study of, 11:153
 recombinant, quantification of N-glycosylation by LC or LC/MS, 19:1648
- Immunoglobulin γ , antibodies disulfide linkages characterized by reversed phase HPLC with negative ESI MS, 17:1590
- Immunoglobulin G1 monoclonal antibodies
 IgG1 κ mAb, characterization using CSD in ESI-TOF-MS, 20:1030
 photo-induced methionine oxidation, MS, 20:525
- Immunoreceptor tyrosine-based activation motif (ITAM) diphosphorylated, spleen tyrosine kinase conformational dynamics on binding to, 16:1039
- Immunosuppressant, macrolide, new metabolite of, identification by HPLC and electrospray MS/MS, 11:516
- Immunosuppressant metabolism, ESI-MS of, 9:1267
- Implementation of the chemical reaction interface mass spectrometry technique on a Hewlett-Packard mass-selective detector, 5:765
- Implication of the charge site in "charge-remote" fragmentations, 3:776
- Importance of charge-separation reactions in tandem mass spectrometry of doubly protonated angiotensin II formed by electrospray ionization, experimental considerations and structural implications, 7:30
- Importance of enhanced mass resolution in removing interferences when measuring volatile organic compounds in human blood by using purge-and-trap gas chromatography/mass spectrometry, 3:831
- Importance of ions and ion-molecule reactions to plasma-surface interface reactions, 13:518
- Importance of matrix:analyte ratio for buffer tolerance using 2,5-dihydrobenzoic acid as a matrix in a MALDI-FT-MS and MALDI-TOF, 9:805
- Importance of preferential solvation of CN ligands in electron and proton transfers observed for *cis*[Ru(CN)₂(bpy)₂] under ion bombardment, 13:1209
- Improved detection limits for electrospray ionization on a magnetic sector mass spectrometer by using an array detector, 5:194
- Improved detection limits for fast atom bombardment mass spectrometry, a study of time-dependent desorption using a model pulsed bombardment ionization method, 3:637

- Improved energy compensation for time-of-flight mass spectrometry, 5:779
- Improved Fourier-transform ion-cyclotron-resonance mass spectrometry of large biomolecules, 4:190
- Improved ion guide for external ion injection in glow discharge-Fourier transform ion cyclotron resonance mass spectrometry, 4:387
- Improved isotopic abundance measurements for high resolution FTICR-MS via time-domain data extraction, 9:799
- Improved mass accuracy in matrix-assisted laser desorption/ionization time-of-flight mass spectrometry of peptides, 5:955
- Impulse-driven droplet deposition system, interface combining LC with MALDI MS and MS/MS, 17:325
- IMS. *See* Ion mobility mass spectrometry; Ion mobility spectrometry mass spectrometry
- In-beam electron impact technique, in MS of methylated fullerenes, 16:1311
- In-cell matrix-assisted laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry, 7:1026
- In-gel digestion, protein identification by, 12:215
- In³⁺ ion, binding to transferrin, indirect detection with ESI-MS, 15:1658
- In-line surface-induced dissociation device in a four-sector mass spectrometer, design and performance of, 6:257
- In-source decay, top-down determination of PEGylation sites using MALDI ISD MS, 20:326
- In-source decay characteristics of peptides of MALDI-TOFMS, 12:420
- In-source decay of hyperbranched polyesteramides in MALDI-OFMS, 11:218
- In-source H/D exchange and ion molecule reactions using MALDI-FTICR-MS with pulsed collision and reaction gases, 13:308
- In vitro rat liver, and microsomal metabolism of bromotrichloromethane and carbon, tandem mass spectrometry study of C-phenyl-N-tert-butyl nitron spin adducts, 6:847
- Inclusion complexes, cyclodextrin, detection of by electrospray mass spectrometry, false positives and, 6:437
- Increased proteome coverage for quantitative peptide abundance measurements based upon high performance separations and DREAMS FTICR-MS, 13:954
- Increasing the dynamic range of a transient recorder by using two analog-to-digital converters, 7:107
- Incremented alkyl derivatives enhance collision-induced glycosidic bond cleavage in MS of disaccharides, 14:63
- Indanediols, dimethyl ether ion interaction with, 8:610
- Indels, *ApoE* SNPs and, genotyping by MS, 17:3
- Indinavir, metabolites of, INTAMS of, 10:175
- Indirect assessment of small hydrophobic ligand binding to a model protein using a combination of ESI-MS and H/D exchange/ESI-MS, 14:506
- Indole, substituted, CAD of, by quadrupole TOFMS, 12:1145
- Induced signal, in a cubic trap, effect of ion initial axial position on, simulated ion trajectory and induced signal in ion cyclotron resonance ion traps, 5:807
- Induced signal and simulated ion trajectory in ion cyclotron resonance ion traps, effect of ion initial axial position on ion coherence, induced signal, and radial or z ejection in a cubic trap, 5:807
- Induced signal in ion cyclotron resonance ion traps, 5:238
- Induction-based fluidics
real-time dispense event and volume verification, MALDI MS, 20:1064
signal intensity and sensitivity improvement using nanoliter spotting by, MALDI MS, 19:1086
- Inductively coupled plasma mass spectrometer, with axial field in quadrupole reaction cell, 13:1176
- Inductively coupled plasma mass spectrometry
ASV with, NCBFE and NCMFE for, 17:945
determination of chlorine isotope ratios by, alleviation of polyatomic ion interferences for, 1:284
determination of trace, rare earth elements using MA-PTFE fibers as sorbent, 17:1258
dynamic reaction cell for, interference reduction in, 10:1083
laser ablation, analyte responses, 17:641
single-particle mode detection by, immunoassays based on, 20:1096
- Inductively coupled plasma mass spectrometry (ICP-MS), laser ablation for, 15:769
- Inductively coupled plasma-mass spectrometry (ICP-MS), 12:1226
- Inductively coupled plasma mass spectrometry with an electrically floating sampling interface, 4:733
- Inductively coupled plasma mass spectrometry with an enlarged sampling orifice and offset ion lens
ion trajectories and detector performance, 4:16
polyatomic ion interferences and matrix effects, 4:28
- Inductively coupled plasma MS
carbon isotope ratio determination with, 14:295
with quadrupole mass filter operated in the third stability region, 8:1230
- Inductively coupled plasma sector-field mass spectrometers, simultaneous or scanning data acquisition for, 14:227
- Inductively coupled plasma-time-of-flight mass spectrometer, 6:400, 6:411, 6:920
- Inductively coupled plasma-time-of-flight mass spectrometer for elemental analysis, optimization and characteristics, 5:1008
- Industrial effluent, alkyl and chloroalkyl phosphates from, multispectral identification of, 2:419
- Industrial proteomics
improved coverage of, nano LC/MS/MS for, 13:760
MALDI-quadrupole TOF-MS of peptides in, 13:772
MALDI-TOF-TOF in peptide sequencing, 13:784
novel precursor ion discovery method on a hybrid quadrupole orthogonal acceleration TOF-MS for studying protein phosphorylation, 13:792
in vivo labeling: glimpse of dynamic proteome and additional constraints for protein identification, 13:804
- Influence of co-matrix proton affinity on oligonucleotide ion stability in MALDI-TOFMS, 9:668
- Influence of cysteine to cysteic acid oxidation on the collision-activated decomposition of protonated peptides, evidence for intraionic interactions, 3:337
- Influence of electrostatic interactions on the detection of hemoglobin complexes in ESI-MS, 12:1092
- Influence of peptide side chains on the metal ion binding site in metal ion-cationized peptides, participation of aromatic rings in metal chelation, 6:1079
- Influence of the length and position of the collision cell on the resolving power attainable from collisionally activated processes in sector instruments, 6:212
- Informatics, focus on, 13:1
- Information management
building a database of bifunctional probes for MS3D investigation of nucleic acids, 17:1570
generation of N-glycan structure databases, 20:1739
- Information on the location of carbon-carbon double bonds in H₆-C₂₃ linear alkenes from carbon addition reactions in a quadrupole ion trap equipped with a pulsed sample-inlet system, 2:305

- Infra red multiphoton dissociation, glycosaminoglycans, effect of charged state and sodium cationization on EDD, 19:790
- Infrared. *See also* Blackbody infrared radiative dissociation
- Infrared and collision-induced multiphoton dissociation studies on $M(\text{acetone})_2^+$ ($M=\text{Al, Fe, Co, Cu, ScO}$) in the gas phase, 4:135
- Infrared atmospheric pressure MALDI, in study of noncovalent sugar-sugar complexes, 14:1158
- Infrared dissociation, blackbody, of leucine enkephalin, 8:771
- Infrared laser desorption/ionization Fourier transform ion cyclotron resonance mass spectrometry, reagent ions in, potassium halide adducts as, 3:301
- Infrared laser post-ionization mass spectrometry, of large biomolecules, 15:934
- Infrared MALDI-MS
of aqueous solutions, 13:354
laser pulse duration role in, Infrared MALDI-MS
mean initial ion velocities in, 13:209
of protein, 9:1, 9:879
- Infrared matrix-assisted laser desorption and ionization by using a tunable mid-infrared free-electron laser, 7:1187
- Infrared matrix-assisted laser desorption electrospray ionization, coupled to FT-ICR MS, biomolecule analysis using, 20:667
- Infrared multiphoton dissociation. *See also* Electron capture dissociation and infrared multiphoton dissociation
B. anthracis petrobactin, 18:842
in external ion reservoir, 14:1413
Fe(III) complex with enterobactin siderophores, 18:632
neutral and sialylated oligosaccharides, electron detachment dissociation, 18:2162
oligosaccharide characterization with boronic acid derivatization using QIT and, 18:2094
in quadrupole ion trap, erythromycin analog characterization by, 13:630
- Infrared multiphoton dissociation-FTICR-MS, indetermining activation energy for noncovalent gas phase peptide unimolecular dissociation, 14:1282
- Infrared multiphoton dissociation (IRMPD)
of aromatic vs aliphatic phosphopeptide dissociation, 15:121
of ganglioside GM1, 16:752
of phosphopeptides, 15:1581
- Infrared multiphoton dissociation of *Bacillus stearothermophilus* glucokinase, 9:1222
- Infrared multiphoton dissociation tandem mass spectrometry, in structural validation of saccharomycins, 10:1285
- Infrared multiple photon dissociation
anionic group II metal nitrate cluster ions, 20:772
improved, using N-terminal phosphonite derivatization, peptides, 20:377
improving efficiency of, QIT, 20:1127
IRMPD and CID of peptides in ion traps, 20:349
oxazolone structure of b_2 fragment ions from tri-alanine, 20:334
studying amide position protons, peptide ions, 20:1841
- Infrared multiple photon dissociation in the quadrupole ion trap via a multipass optical arrangement, 5:886
- Infrared photodissociation of non-covalent adducts of electrosprayed nucleotide ions, 7:209
- Infrared SALDI-MS, of protein and peptide frozen aqueous solutions, 9:912
- Inhibitors, rapid screening, label-free approach to, MALDI-TOF MS, 17:815
- Initial implementation of an electrodynamic ion funnel with FTICR-MS, 11:19
- Initial velocity of ions generated by matrix-assisted laser desorption/ionization and its effect on the calibration of delayed extraction time-of-flight mass spectra, 8:209
- Initiation factors, eukaryotic, protein conformation using T-WAVE ion mobility MS, 20:1699
- Injection of reagent ions into the seldedge region in fast-atom bombardment mass spectrometry, 4:259
- Innovative method for measuring hydrogen and deuterium, chemical reaction interface MS with nitrogen reactive gas, 11:400
- Inorganic compounds, FT LMMS of, 9:482
- Inositol phosphorylceramides, *Leishmania major*, characterization by ESI-MS/MS, 18:1591
- Insertion reactions of metal carbonyl anions with methyl formate in the gas phase as revealed by ^{13}C - and D-labeling, 5:525
- Insights into analyte electrolysis in an electrospray emitter from chronopotentiometry experiments and mass transport calculations, 11:951
- Instrumentation
for ESI-MS, 8:1191
gas phase ion/ion chemistry study via mass spectrometry, 19:173
hybrid mass spectrometers for MS/MS, 19:161
for mass spectrometry, billionfold data increase from, 8:1
- Insulating samples, large, imaging, by using secondary ion tandem mass spectrometry, charge compensation for, 5:37
- Insulin
bovine, characterization of glycation sites, MS/MS, 20:1319
ILPR binding sites in IGF-2 and, MALDI MS/MS, 20:1974
mapping disulfide bonds with Route 66 method, 20:157
SIMSTEX application to oligomerization of insulin analogs and mutants, 17:1526
- Insulin B, apparent gas-phase acidity of, 7:1211
- Insulin chain B, gas-phase reactivity and dissociation of $[M + 4\text{H}]^4$ ions from, cysteic acid group effect on, 10:928
- Insulin-like growth factors, ILPR binding sites in insulin and, MALDI MS/MS, 20:1974
- Insulin-linked polymorphic region, binding sites in insulin and IGF-2, MALDI MS/MS, 20:1974
- Insulin receptor substrate-1, phosphorylation of, quantified by HPLC-ESI-MS/MS, 17:562
- Insulin receptor substrate-1 phosphorylation site, identification negative-ion $\mu\text{LC}/\text{ES-API-CID-MS}$ hybrid scan technique, 14:401
- Intact cell-mass spectrometry (ICM-MS), for bacterial identification, limits of, 12:49
- INTAMS. *See* Intelligent automated liquid chromatography-tandem mass spectrometry
- Integral probe for capillary zone electrophoresis/continuous-flow fast atom bombardment mass spectrometry, 3:198
- Integrated method for spectrum extraction and compound identification from GC/MS data, 10:770
- Intelligent automated liquid chromatography-tandem mass spectrometry, indinavir metabolite identification by, 10:175
- Intensity facing, noncovalent interactions, exploring by MALDI-TOF-MS, 18:359
- Inter- and intramolecular migration of peptide amide hydrogens during ESI, 12:410
- Interaction of angiotensin peptides and zinc metal ions by electrospray ionization mass spectrometry, 5:959
- Interaction of first row transition metals with $\alpha 1\text{N}3, \alpha 1\text{N}6$ mannotriose and conserved trimannosyl core oligosaccharides, comparative electrospray ionization study of doubly and singly charged complexes, 8:32
- Intercomparison study on accurate mass measurement of small molecules in mass spectrometry, 14:1086
- Interface
for a four-sector mass spectrometer with dual purpose colli-

- sion cell, high transmission at low to intermediate energies, 6:175
- mass spectrometry, atmospheric pressure chemical ionization liquid chromatography, optimization of, 7:69
- particle beam liquid chromatography/mass spectrometry, performance of, effect of a dc glow discharge on, investigation of, 5:908
- thermospray, with carbamate pesticides, comparative study of different, influence of the ion source geometry, 6:656
- Interface region of a four-sector tandem mass spectrometer, enhancement of ion transmission at low collision energies at, 5:460
- Interference effects, in inductively coupled plasma-mass spectrometry, solvent load and first-stage pressure to reduce, selection of, 3:128
- Interference reduction, in ICP-DRC-MS, 10:1083
- Interferences
- matrix-induced, in mixed-gas helium-argon inductively coupled plasma mass spectrometry, investigation of, 2:355
 - removal of, when measuring volatile organic compounds in human blood by using purge-and-trap gas chromatography/mass spectrometry, importance of enhanced mass resolution in, 3:831
- Interferences from spectra of unknowns, software-based mass spectral enhancement to remove, 7:598
- Interleukin-3, mouse, baculovirus expressed, application to. Liquid secondary ion mass spectrometry analysis of permethylated, n-hexylamine derivatized oligosaccharides, 3:345
- Intermediary ion-neutral complexes, internal reactions in, remote fragmentations of protonated aromatic carbonyl compounds via, 3:417
- Internal electrospray ionization, low magnetic field, Fourier transform ion cyclotron resonance mass spectrometer, a high performance, 7:915
- Internal energies, calculated, for polychlorinated biphenyls, correlation between electron capture negative chemical ionization mass spectrometric fragmentation, 5:44
- Internal energy deposition in chemical ionization/tandem mass spectrometry, 14:102
- Internal energy distributions deposited in doubly and singly charged tungsten hexacarbonyl ions generated by charge stripping, electron impact, and charge exchange, 1:16
- Internal energy distributions of tungsten hexacarbonyl ions after neutralization-reionization, 5:1093
- Internal energy relaxation, time scale of, AP-MALDI and nano-ESI ions in QITMS, 20:1801
- Internal glow discharge-Fourier transform ion cyclotron resonance mass spectrometry, 7:923
- Internal glucose residue loss in protonated O-diglycosyl flavonoids upon low-energy collision-induced dissociation, 11:136
- Internal monosaccharide, losses in the collisionally activated dissociation mass spectra of anthracycline aminodisaccharides, 9:710
- Internal pulsed valve sample introduction on a quadrupole ion trap mass spectrometer, 6:976
- Internal standard correction of results obtained by atmospheric pressure chemical ionization mass spectrometry, 2:427
- Internal standards, labeled or unlabeled, combined capillary gas chromatography/ion trap mass spectrometry quantitative methods using, 1:265
- Interpretation, automatic, algorithms for, high-resolution mass spectra, 17:459
- Intraionic interactions, evidence for, influence of cysteine to cysteic acid oxidation on the collision-activated decomposition of protonated peptides, 3:337
- Intramolecular cyclization and electron impact ionization mass spectrometry, in 2-substituted pyrimidin-4(3H)-ones, 5:113
- Intramolecular electron transfer, in generation of tripeptide radical cations, 16:763
- Intrinsic basicity of oligomeric peptides that contain glycine, alanine, and valine—the effects of the alkyl side chain on proton transfer reactions, 6:91
- Intrinsic Ca²⁺ affinities of peptides, application of the kinetic method to analogs of calcium-binding site III of rabbit skeletal troponin C, 11:770
- Intrinsic hydration, of monovalent uranyl hydroxide, nitrate, and acetate cations, 15:777
- Invertebrate ganglia, spatial profiling, 14:752
- Investigation of dipeptides containing polar and nonpolar side groups by Curie-point pyrolysis tandem mass spectrometry, 3:750
- Investigation of bovine ubiquitin conformers separated by high field asymmetric waveform ion mobility spectrometry, cross section measurements of using energy-loss experiments with a triple quadrupole mass spectrometer, 11:738
- Investigation of calcium-induced, noncovalent association of calmodulin with melittin by ESI-MS, 8:809
- Investigation of cell culture media infected with viruses by pyrolysis MS, implications for bioaerosol detection, 10:502
- Investigation of charge derivatization influence of the fragmentation of multiply protonated peptides, 13:47
- Investigation of drift gas selectivity in high resolution ion mobility spectrometry with MS detection, 13:300
- Investigation of formation process of MCs⁺-molecular ions during sputtering, 11:650
- Investigation of low-voltage on-resonance ion selection for FT-MS, 9:931
- Investigation of matrix-induced interferences in mixed-gas helium-argon inductively coupled plasma mass spectrometry, 2:355
- Investigation of metal attachment to polystyrenes in MALDI, 9:724
- Investigation of metastable decay of DNA under UV MALDI conditions with post-source-decay analysis and hydrogen/deuterium exchange, 9:866
- Investigation of regio- and stereoselectivity of deoxyguanosine linkage to deuterated 2-hydroxyestradiol using LC/ESI-ion trap MS, 14:364
- Investigation of site-selective gas-phase reactions of amino alcohols with dimethyl ether ions, 4:230
- Investigation of the effect of a dc glow discharge on the performance of a particle beam liquid chromatography/mass spectrometry interface, 5:908
- Investigation of the energies of peptide ion dissociation by laser desorption chemical ionization Fourier transform mass spectrometry, 6:1069
- Investigation of the formation of MS⁺-molecular ions in sputtering processes, 10:246
- Investigation of the gas-phase structure of electrosprayed proteins using ion-molecule reactions, 5:207
- Investigation of the initial reactions of the calcote mechanism for soot formation, 7:559
- Investigation of the mechanism of intracuster proton transfer from sinapinic acid to biomolecular analytes, 9:1060
- Investigation of tris(trimethoxyphenyl)phosphonium acetyl charged derivatives of peptides by ESI-MS and MS/MS, 11:107
- Investigation of UV matrix-assisted laser desorption Fourier transform mass spectrometry for peptides, 2:22
- Investigations of matrix-assisted laser desorption/ionization sample preparation by TOF secondary ion MS, 10:104

- Investigative proteomics, identification of an unknown plant virus from infected plants using MS, 14:736
- Involatile compounds, detection with laser desorption electron attachment TOFMS, 12:1339
- Iodide, poly(ethylene glycol) cation reactions with, 9:957
- Iodination, deciphering the code for, peptides, 20:965
- Iodoacetonitrile, proton affinities of, using kinetic method with full entropy analysis, 11:601
- Ion
- complex, cleavage reactions of, positive ion ESI-MS/MS of, 15:689
 - generated in a high magnetic field electrospray ionization Fourier transform ion cyclotron mass spectrometer, trapping and detection of, 3:615
 - isolation, in the quadrupole ion trap, evidence of isomerization during, 3:680
 - metastable, 15:1261
 - protonated, of cysteine-containing peptides, fragmentation of, 15:697
- Ion abundance quantitation, in FTICR-MS, 9:1204
- Ion abundances, enhanced, from a carrier process, investigation of, in high-performance liquid chromatography particle beam mass spectrometry, 1:82
- Ion acceleration systems, gridless, for time-of-flight mass spectrometry, 4:782
- Ion accumulation, for LC-ESI-FTICR, using central trapping electrode, 11:592
- Ion activation
- alternative approaches to infrared multiphoton dissociation in an external ion reservoir, 14:1413
 - in chemistry and biochemistry, mechanisms, dynamics, and applications of, 14:1329
 - collisions of slow polyatomic ions with surfaces, scattering method and results, 14:1360
 - direct dynamics study of N-protonated diglycine surface-induced dissociation, influence of collision energy, 14:1402
 - fraying and electron autodetachment dynamics of trapped gas phase oligonucleotides, 14:1330
 - gas phase dissociation pathways of multiply charged peptide clusters, 14:1373
 - model for energy transfer in inelastic molecular collisions applicable at steady state or non-steady state and for an arbitrary distribution of collision energies, 14:1348
 - peptide sequencing using a patchwork approach and surface-induced dissociation in sector-TOF and dual quadrupole MS, 14:1387
 - surface-induced dissociation of peptide ions, 14:1340
- Ion adduct, negative, stability in electrospray MS, multiple hydrogen bonding and, 13:1360
- Ion association by time-of-flight mass spectrometry, a study of V-P-O catalysts, 1:129
- Ion beam images, direct generation of with a two-dimensional charge injection device, 6:140
- Ion beam MS/MS, guided, to determine thermochemistry, 13:419
- Ion beams
- afterglow-guided, in MS/MS, 8:688
 - hyperthermal, reactions of, at self-assembled monolayer surfaces, X3F⁺, IF⁺, and other unusual ions generated by, 4:938
 - improving quality exiting quadrupole ion guide, 17:335
- Ion bombardment, low, energy reactive, chemical modification of fluorinated self-assembled monolayer surfaces by, 6:187
- Ion calorimetry, using MS to measure melting points, 18:74
- Ion cells, elongated, Fourier transform mass spectrometry, trapped trapping motion in, direct observation of, 1:351
- Ion charge states, peptide and protein, FTICR, 17:903
- Ion chemistry
- of deprotonated phenylthiocarbonyl-phenylalanine, 8:1078
 - of methylbenz(is)oxazole isomers in gas phase, 15:1005
- Ion cloud manipulation, using the radiofrequency-only mode as an improvement for high mass detection in Fourier transform mass spectrometry, 4:186
- Ion cloud signals, individual, time-resolved measurements of, to investigate space-charge effects in plasma mass spectrometry, 7:362
- Ion coherence, in a cubic trap, effect of ion initial axial position on, simulated ion trajectory and induced signal in ion cyclotron resonance ion traps, 5:807
- Ion cool times, effect on peptide ion internal energies produced by ESI and AP-MALDI, 16:743
- Ion cooling, for FTICR-MS, 12:1169
- Ion coupling, strategies for microengineered quadrupole mass filters, 20:146
- Ion cyclotron capacitively coupled open cell, high-resolution ion isolation with, 6:533
- Ion cyclotron resonance. *See* Fourier transform ion cyclotron resonance.
- electrospray ionization mass spectrometer, high-resolution accurate mass measurements of biomolecules using, 4:566
 - Fourier transform
 - detection of multiphoton ionization spectroscopy, 7:1018
 - internal glow discharge mass spectrometry, 7:923
 - two-dimensional Coulomb-induced frequency modulation in, a mechanism for line broadening at high mass and for large ion populations, 4:909
- Ion cyclotron resonance ion traps, simulated ion trajectory and induced signal in, 5:238
- effect of ion initial axial position on ion coherence, induced signal, and radial or z ejection in a cubic trap, 5:807
- Ion cyclotron resonance mass spectrometer
- external ion source Fourier transform, matrix-assisted laser desorption ionization on, high resolution end group determination of low molecular weight polymers by, 7:449
 - Fourier transform
 - concentric tube vacuum chamber for high magnetic field, 4:168
 - low magnetic field internal electrospray, a high performance, 7:915
 - small, characterization of long-chain carboxylic esters with CH₃OBOCH₃⁺ in, 7:1138
 - realistic simulation of, 10:136
- Ion cyclotron resonance mass spectrometry
- capillary electrophoresis Fourier transform, with sustained off-resonance irradiation for the characterization of protein and peptide mixtures, 5:894
 - Fourier transform
 - excitation modes for, 4:433
 - high-frequency, 4:177
 - ionization, reagent ions in, potassium halide adducts as, 3:301
 - remeasurement at high resolving power, 6:564
 - glow discharge-Fourier transform, external ion injection in, ion guide for, an improved, 4:387
 - ionization Fourier transform, in-cell matrix-assisted laser desorption, 7:1026
 - probing trapped ion energies via ion-molecule reaction kinetics, 3:727
 - quadrupolar excitation in, simplified application of, 6:448
 - study of ion-molecule reactions relevant to Titan's atmosphere, 17:544
- Ion-cyclotron-resonance mass spectrometry, Fourier-transform, improved, of large biomolecules, 4:190

- Ion cyclotron resonance precise mass difference measurements, Fourier transform, Coulomb mutual interaction and image charge in, systematic errors originating from, analysis and estimation of, 4:855
- Ion cyclotron resonance spectra, Fourier transform, relative peak heights in, space charge effects on, 3:734
- Ion cyclotron resonance study-Fourier transform, of the gas-phase acidities of germane and methylgermane, bond dissociation energy of germane, 4:54
- Ion cyclotron resonance tandem mass spectrometry (ICR-MS/MS), surface-induced dissociation in, 11:1107
- Ion deceleration lenses, understanding, what are the simplicity/performance trade offs?, 2:322
- Ion decompensation *versus* molecular size probed by vacuum ultraviolet photoionization, 2:103
- Ion deflection system, a pulsed, for background reduction in ²⁵²Cf-plasma desorption mass spectrometry, 3:706
- Ion-dipole complex formation, from deprotonated phenol fatty acid esters evidenced by using gas-phase labeling combined with tandem mass spectrometry, 4:343
- Ion dissociation and resonance ejection, competition between, during resonant excitation in a quadrupole ion trap, 5:1031
- Ion ejection
collision induced, in FTICR cell, 16:422
from linear quadrupole ion trap, 14:1130
- Ion emission, secondary, and solution chemistry, from amine-glycerol solutions, 2:33
- Ion energies, trapped, probing, via ion-molecule reaction kinetics Fourier transform ion cyclotron resonance mass spectrometry, 3:727
quadrupole ion trap mass spectrometry, 3:716
- Ion energy, in quadrupole MS, 15:48
- Ion energy distribution, charge-dependent/independent, from a heated capillary electrospray source, selective generation of, 5:221
- Ion exchange chromatography, ESI-MS and, in analysis of hexose monophosphates, 17:104
- Ion excitation
in linear quadrupole ion trap, 16:835
linear response theory of, for Fourier transform mass spectrometry, 2:483
- Ion extraction, improved, from linear octopole ion trap, SIMION analysis and experimental demonstration of, 13:1304
- Ion flow tube studies, CH₃OCH₂⁺ reactions with hydrocarbons in, 13:1015
- Ion flow tubes, selected, ion reactions with ethylene oxide and propenal, 18:445
- Ion formation from charged droplets, roles of geometry, energy, and time, 4:524
- Ion formation of N-methyl carbamate pesticides in thermospray mass spectrometry, the effects of additives to the liquid chromatographic eluent and of the vaporizer temperature, 5:913
- Ion formation processes in the afterpeak time regime of pulsed glow discharge plasmas, 4:727
- Ion fragmentation
in 2-D quadrupole ion trap for external ion accumulation in ESI FTICR-MS, 12:1312
charge-associated mechanism of, in fatty acid CID spectra, 16:280
- Ion frequencies in a quadrupole ion trap, determination of, by using a fast direct current pulse as pump and a laser probe, 5:29
- Ion funnel, electrodynamic
evaluation of low *m/z* transmission, 17:586
for FTICR-MS, 11:19
- Ion funnel interfaces, high-pressure electrodynamic, MS sensitivity improved using, 17:1299
- Ion funnels, for masses, 16:1708
- Ion guide
improved, for external ion injection in glow discharge-Fourier transform ion cyclotron resonance mass spectrometry, 4:387
multipole, driving by high voltage RF oscillator, 13:1370
stacked-ring electrostatic, 7:101
- Ion images
acquisition and quantification of, with a camera-based detections system and classical quantification algorithms, 1:37
secondary, of rodent brain, 11:191
- Ion-impact energy measurement, in ESI-MS of proteins and protein mixtures, 12:826
- Ion initial axial position, effect of, on ion coherence, induced signal, and radial or z ejection in a cubic trap, 5:807
- Ion intensity, enhancement of, in time-of-flight secondary-ionization mass spectrometry, 7:467
- Ion interferences, polyatomic, and matrix effects, inductively coupled plasma mass spectrometry with an enlarged sampling orifice and offset ion lens, 4:28
- Ion/ion interactions
data acquisition from, linear ion-trap MS, 19:1821
gas phase
peptide AA composition and role in charge inversion, 20:180
study via mass spectrometry, 19:173
IT-IM TOF mass spectrometry analysis of, 20:1549
- Ion-ion proton transfer reactions, of bio-ions involving noncovalent interactions: homomyoglobin, 8:637
- Ion/ion proton transfer reactions, mutual storage mode, in hybrid linear ion trap, 16:71
- Ion-ion reactions
gas-phase, study of, using electrospray ionization, a new approach for, 3:695
in gas phase, proton transfer reactions of protonated pyridine with multiply charged oligonucleotide anions, 6:529
- Ion/ion reactions
of ESI cations and anions, 13:614
for oligopeptide mixture analysis: application to mixtures comprised of 0.5–100-kDa components, 9:585
pulsed dual ESI for, 16:1750
- Ion-ion recombination, role of, for alkali chloride cluster formation in liquid secondary ion mass spectrometry, 4:410
- Ion isolation, high-resolution, with the ion cyclotron capacitively coupled open cell, 6:533
- Ion isolation/rejection, selective, over a broad mass range in the quadrupole ion trap, 2:11
- Ion kinetic energies, laser desorbed, electrostatic energy analyzer-time-of-flight mass spectrometer for measurement of laser desorbed ion kinetic energies, design and calibration of, 6:619
- Ion kinetic energy spectra, mass-analyzed, main beam in, low energy side of, origin of the tailing signal on, 3:644
- Ion kinetic energy spectrometry experiments, mass-analytical, kinetic energy distribution of precursor ion main beam in, measurement of, 6:1232
- Ion kinetic modulation, for improved ion trapping in electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry, 6:76
- Ion lens, offset, inductively coupled plasma mass spectrometry with an enlarged sampling orifice and
ion trajectories and detector performance, 4:16
polyatomic ion interferences and matrix effects, 4:28
- Ion mirror
simple cylindrical, 11:759
simple geometry gridless, 11:764

- Ion mobility
 dynamic method for measurement of, 20:2070
 high-field, resolution equations for, 15:1320
 of Zn-ligand-hexose diastereomers, 13:284
- Ion mobility cells, combined with collision cell and TOF/MS, 17:691
- Ion mobility collision-induced desorption mass spectrometry, peptide identification using, 20:907
- Ion mobility (IM) mass spectrometry
 applied to cyclic peptide analysis, 15:870
 in brain tissue phospholipid analysis, 16:158
 of charge-solvated vs salt bridge structures of protonated trimers, 16:1009
 collision-induced protein ion unfolding analyzed by, 20:1851
 field strength of tryptic peptides by, 16:158
 of G-quadruplexes in deoxyguanosine clusters, 16:989
 of gas phase bradykinin fragment 1–5 ions, 16:893
 in metallothionein investigations, 16:1787
 molecular framework for collision of peptide ions, MALDI TOF MS, 20:1593
 organoruthenium anticancer complexes, 20:1119
 of 20S proteasome, 16:998
 T-WAVE, protein conformation using, 20:1699
 traveling-wave, hemoglobin structure probed by, 20:625
 use of shift reagents in, 20:1
- Ion mobility in a triple quadrupole mass spectrometer, conformer selection of protein ions by, 5:127
- Ion mobility matrix-assisted laser desorption ionization, of non-covalent complexes, 13:167
- Ion mobility spectrometry
 ambient pressure, carbohydrate isomer resolution by ESI-APIMS-TOF-MS, 18:1163
 APCI of fluorinated phenols in, 10:1157
 in APCI studies of non-polar isomeric hydrocarbons, 13:1078
 atomic transition metals as ion mobility source for guided ion beam experiments, 18:1196
 high resolution, drift gas selectivity in, 13:300
 human plasma proteome mapping using SCX-LC-IMS-MS, 18:1249
 incorporation with LC-CID-TOFMS techniques, 14:1424
 orthogonal extraction, 17:500
 protein complex sizing by ESI and, 18:1206
 ubiquitin elongated states, resolution and structural transitions, 18:322
- Ion mobility spectrometry mass spectrometry
 assigning N-linked glycan structural isomers from ovalbumin, 19:1711
 characterization of proton-bound acetate dimers, 19:1361
 CID fragmentation of multiple ions, 19:411
- Ion mobility spectrometry-mass spectrometry, using electrodynamic ion funnels and elevated drift gas pressures, 18:1176
- Ion mobility tandem mass spectrometry, variance of a- and b-type peptide fragment ions, 19:609
- Ion mobility time-of-flight mass spectrometry
 chemical warfare agent degradation products detected by, 17:241
 high resolution, of synthetic failures in combinatorial libraries, 11:352
 phospholipid analysis by, 19:1660
- Ion-molecule chemistry, gas-phase, of borate and boronate esters, 5:169
- Ion-molecule collisions in the vacuum chamber of an electrospray time-of-flight mass spectrometer, effect of, on mass spectra of proteins, 7:342
- Ion-molecule equilibria, how and why, 3:1
- Ion-molecule reaction
 of alkanenitriles and transition-metal ions in the gas phase, a study on fragmentation mechanism of the adducts, 2:120
 early development of the flowing afterglow technique for, 3:479
- Ion/molecule reaction, with dimethyl ether, selective detection of the tolyl cation among other $[C_7H]^+$ isomers by, 2:270
- Ion-molecule reaction kinetics, probing trapped ion energies via Fourier transform ion cyclotron resonance mass spectrometry, 3:727
- Ion-molecule reaction studies, flowing afterglow technique for, early development of, a personal history of, 3:479
- Ion-molecule reaction under chemical ionization, gas-phase halo alkylation of H_6O -fullerene by, 4:894
- Ion molecule reactions
 for the differentiation of phospholipid classes, fast atom bombardment tandem mass spectrometry employing, 2:470
 endothermic, 2:189
 estimation of mutagenic/carcinogenic potential of environmental contaminants by, 1:110
 functional group-selective, of ethylene glycol and its monomethyl and dimethyl ethers, 4:97
 gas-phase, of the CH_3O^+ cation, 3:515
 of gas phase chromium oxyanions, 14:1067
 investigation of the gas-phase structure of electrosprayed proteins using, 5:207
 in methylamine and dimethylamine and trimethylamine systems for, 2:220
 in probing phosphorylated carbohydrate isomeric differences, 14:323
 as a quantification method for phosphorylated positional isomers, FTICR approach to, 14:916
 selective, of lactams with dimethyl ether ions, 4:352
 in-source, MALDI-FTICR-MS of, 13:308
- Ion/molecule reactions
 isomeric recognition by, case of ionized phenol-cyclohexadienone, 19:126
 specific, elucidating the structures of isomeric silylenium ions (SiC_3H^+9 , SiC_4H^+ , SiC_5H^+13) by, 2:278
- Ion-molecule reactions and collision-activated dissociation of $C_4H_4^+$ isomers, a case study in the use of the MS3 capabilities of a pentaquadrupole mass spectrometer, 3:518
- Ion-molecule reactions in mixtures of trimethylaluminum and methylamines, 4:111
- Ion-molecule reactions in the gas phase, reactivity of the cis and trans-indanedions with dimethyl ether ions, 8:610
- Ion-molecule reactions in Titan's atmosphere, 12:557
- Ion-molecule reactions of oxygenated chemical ionization reagents with vincamine, 7:250
- Ion molecule reactor, HP-SQCC as, 16:957
- Ion monitoring, multiple, risk of false positive identification using, qualitative MS, 17:823
- Ion monitoring mode, selective, in gas chromatography/mass spectrometry, direct, 5:106
- Ion motion, in quadrupole mass spectrometer, analytical description of, 14:818
- Ion nanocalorimetry, effects of electron kinetic energy and ion-electron inelastic collisions. ECD, 19:772
- Ion-neutral complex-mediated alkane eliminations from ionized aliphatic ethers, size effects in, 2:261
- Ion-neutral complex-mediated pathway over a five-membered ring H shift in the isomerization of $CH_3^- + HCH_2CH_2$ by ab initio theory, preference for, 6:1037
- Ion-neutral complexes
 the decompositions of N-(substituted benzalamino)phthalimide radical cations embody, 5:837
 formation of the 3-pentanone ion from ionized propyl propanoate through, 3:183

- intermediary, internal reactions in, remote fragmentations of protonated aromatic carbonyl compounds via, 3:417
- isomeric, generated from ionized 2-methylpropanol and n-butanol, the effect of the polarity of the neutral partner on complex-mediated reactions, 3:409
- Ion-neutral complexes $[\text{CH}_3\text{CH}^+\text{CH}_3\text{?CH}_2\text{CH}_3]$ and $[\text{CH}_3\text{CH}_2\text{CH}^+\text{CH}_3\bullet\text{CH}_3]$ in the gas phase, a photoionization study of, formation, H-transfer and C-C bond formation between partners, and channeling of energy into dissociation, 7:73
- Ion-neutral complexes in gas-phase dissociations of protonated nitriles, $n\text{-C}_3\text{H}_7\text{CNH}^+$ and $i\text{-C}_3\text{H}_7\text{CNH}^+$, 2:178
- Ion-neutral or McLafferty complex arrangements, APCI low energy collision-induced dissociation fragmentation of protonated ortho silicates, 5:544
- Ion optical elements, for TOF-MS, conductive carbon-filled polymeric electrodes as, 13:1170
- Ion optics, in minimum thickness gate valve design, for MS, 16:441
- Ion pairing, isobaric monosaccharides distinguished by, 16:1827
- Ion pairing and observed charge states, of diquatery ammonium salts in electrospray ionization mass spectrometry, effects of solvent and counterion on, 7:1050
- Ion pairs, anion adduct generation and, during MALDI, 13:1218
- Ion populations
- large, line broadening at, a mechanism for, two-dimensional Coulomb-induced frequency modulation in Fourier transform ion cyclotron resonance, 4:909
 - and monoisotopic masses for large biomolecules from resolved isotopic distributions, determination of, 6:229
 - probe, secondary, a wide-angle, for organic ion imaging, 2:362
- Ion preselection, for ESI-FTICR-MS, 12:38
- Ion production, irradiation effects in, 2,5-dihydroxybenzoic acid monocrystals in, 14:893
- Ion reservoir, external, infrared multiphoton dissociation in, 14:1413
- Ion resonance cell, for FTMS, 10:1000
- Ion response, and stereoselectivity in electrospray mass spectrometry, correlation of, and electrospray ionization-mass spectrometry as tool to predict chemical behavior, 6:207
- Ion scanning in positive mode, for phosphopeptide analysis, derivatization strategy for, 13:996
- Ion selection, low-voltage, on-resonance, for FTMS, 9:931
- Ion selection and accumulation, data-dependent external, in ESI-FTICR-MS, 13:144
- Ion size, the effect of, on rate of dissociation, RRKM calculations on model large polypeptide ions, 4:11
- Ion soft-landing
- chiral enrichment of serine via, 15:1360
 - into liquids, 15:1874
- Ion source
- an electrospray, for magnetic sector mass spectrometers, 2:205
 - a compact high pressure, for high and low energy collision-induced dissociation studies of cluster ions on a VG analytical ZAB-2FQ, 3:33
 - dielectric barrier discharge, for ambient mass spectrometry, 18:1859
 - elemental, the pulsed glow discharge as, 1:138
 - emittance influence on the transmission of a quadrupole operated in the second stability region, 9:1074
 - glow discharge, for mass spectrometric analysis of organic compounds, characterization of, 5:305
 - ion-neutral reactions in, SFCMS, 18:2173
 - isolated droplets as, 11:393
 - for medium pressure laser ionization, 15:1885
 - pulsed triple, source for sequential ion/ion reactions, LIT MS/MS, 18:369
 - space-charge dominated mass spectrometry, modeling and sensitivity, 2:1
- Ion source geometry, comparative study of different thermospray interfaces with carbamate pesticides and, 6:656
- Ion source problem, why are electron capture negative ion mass spectra not reproducible?, 4:270
- Ion source residence time, effective, estimating, a simple method for, 4:787
- Ion spectrometry, a high transmission hemispherical energy analyzer for, 2:506
- Ion spray mass spectrometry, studies on heme binding in myoglobin, hemoglobin, and cytochrome c by, 4:631
- Ion-spray mass spectrometry
- high accuracy molecular weight determination and variation characterization of proteins up to 80 ku by, 2:387
 - a model investigation using, observation of noncovalent complexes between margatoxin and the KV1.3 peptide ligands, 7:1075
- Ion spray-tandem mass spectrometry, of supramolecular coordination complexes, 5:748
- Ion storage, selective field-modulated, in quadrupole ion trap, 6:928
- Ion storage in a quadrupole ion trap, nonlinear resonance effects during, 4:917
- Ion structural analogs, formation by solution phase chemistry, 16:607
- Ion structures, nonclassical, and 1,3-hexadien-5-yne radicalcation to the benzene structure, isomerization of, Ab initio calculations of, 7:731
- Ion study, metastable, of substituted cyclopentadienylmanganese cations in the gas phase, 4:216
- Ion suppression, study of, quantitation of SR 27417 in human plasma using electrospray liquid chromatography, 7:1099
- Ion tandem mass spectrometry, secondary, imaging large insulating samples by, charge compensation for, 5:37
- Ion temperature, effective, in quadrupole ion trap, 9:845
- Ion thermochemistry, focus on, 13:417
- Ion tomography, laser-induced fluorescence for, in Penning trap, 9:925
- Ion trajectories and detector performance, inductively coupled plasma mass spectrometry with an enlarged sampling orifice and offset ion lens, 4:16
- Ion trajectory
- in an electrostatic ion guide for external ion source Fourier transform ion cyclotron resonance mass spectrometry, 6:936
 - simulated and induced signal in ion cyclotron resonance ion traps, 5:238
 - effect of ion initial axial position on ion coherence, induced signal, and radial or z ejection in a cubic trap, 5:807
 - single, in Paul-type ion traps, computer simulation of, 4:687
- Ion trajectory simulation program, use for electrode configurations with arbitrary geometries, 17:1216
- Ion transmission at low collision energies, enhancement of, via modifications to the interface region of a four-sector tandem mass spectrometer, 5:460
- Ion transport by viscous gas flow through capillaries, 5:873
- Ion transport system, a low voltage, for external ionization Fourier transform mass spectrometry, 5:398
- Ion trap. *See also* Laser desorption-ion trap mass spectrometry; Penning trap
- quadrupole
 - accurate mass measurements in, 11:876
 - of alkali-cationed polysaccharides, 8:987
 - combined, equilibrium ion density distribution and trapping force vs, 9:473

- competitive reactions of dimethyl ether ions in, steric and substituent effects in phenols evaluated by, 6:627
- equilibrium space charge in, 5:64
- equipped with a pulsed sample-inlet system, carbon reactions in, information on the location of carbon-carbon double bonds in C₆-C₂₃ linear alkenes from, 2:305
- in FAIMS/FAIMS coupled to TOFMS, 12:1320
- field-modulated selective ion storage in, 6:928
- FT-ICR-MS, improved ion trapping and detection capabilities, 20:755
- gap-tripole, computer simulation of, 19:1367
- high mannose glycans characterized by ion trap MS, bovine ribonuclease B, 20:539
- hybrid linear, mutual storage mode ion/ion reactions in, 16:71
- hyperbolic, for Fourier transform ion cyclotron resonance mass spectrometry, experimental evaluation of, 3:188
- infrared multiple photon dissociation in, via a multipass optical arrangement, 5:886
- ion collision cross section on, 8:697
- ion cyclotron resonance, simulated ion trajectory and induced signal in, 5:238
- effect of ion initial axial position on ion coherence, induced signal, and radial or z ejection in a cubic trap, 5:807
- ion frequencies in, determination of, by using a fast direct current pulse as pump and a laser probe, 5:29
- ion isolation in, evidence of isomerization during, 3:680
- of ion loss, 8:1141
- ion storage in, nonlinear resonance effects during, 4:917
- IRMPD and CID of peptides in, 20:349
- linear
- for ESI-MS of triacylglycerols, 16:1498
 - fragmentation efficiency in, 16:1342
 - higher order quadrupole excitation frequencies in, 11:1016
 - ion quadrupole excitation resonance shifts in, 13:577
 - off-resonance excitation, 20:443
 - resonant excitation in, 14:622
- linear octopole, ion extraction from, SIMION analysis and experimental demonstration of, 13:1304
- linear quadrupole, fast scanning of, 8:539
- with linear TOF-MS, development and characterization of, 8:1085
- linked scans of
- artifact peaks in, 8:554
 - parent ion resolution in, 8:545
- mass analysis at high pressure, 20:2144
- matrix-shimmed, for excitation, detection, quadrupolar axialization, and trapping, 10:759
- measuring fluorescence of molecular ions trapped in, with zero background detection, 13:696
- methylene substitution reactions in, selectivity of ethylene, ethylene oxide, and dimethyl ether reactive ions, 3:39
- miniaturized mass analyzers, novel ion traps using planar resistive electrodes, 19:1435
- MS/MS of, polycyclic aromatic hydrocarbon analysis by, 8:183
- nano electrospray combined with, for the analysis of peptides and protein digests, 7:150
- Paul-type, single-ion trajectories in, computer simulation of, 4:687
- peptide ions in, tandem mass spectra of, effects of heavy gases on, 7:1194
- peptides in, MS/MS spectra of, 9:341
- performance, the effects of air and helium on, when used for real-time monitoring of some atmospheric pollutants, 4:774
- polyether-transition metal ion complex in, 8:620
- preparative scale, 8:943
- pulsed introduction of buffer gas in, 8:532
- quadrupole ion trap, 8:569
- an empirical approach to estimation of critical energies by using, 7:1116
- quadrupole, pulsed gas introduction into, 1:308
- radio frequency, collisional activation in, 15:1616
- conformational differences of peptide ions in, 12:1331
 - measurement of collision-induced dissociation of tantalum oxide ions in, 11:1072
- with reflectron TOF mass detector, for protein digest analysis, 8:1237
- resonance excitation in, competition between resonance ejection and ion dissociation during, 5:1031
- selective ion isolation/rejection over a broad mass range in, 2:11
- single crystal characterization by, 10:352
- in study of stereoselective dissociations of hexosamine monosaccharides, 12:517
 - trapped negative ion cooling in, 8:793
- Ion trap collision-induced dissociation, multiply deprotonated RNA, 19:1832
- Ion trap collisional activation, relative dissociation energy measurements using, 5:250
- Ion trap collisional activation of deprotonated deoxymononucleoside and deoxydinucleoside monophosphates, 6:102
- Ion trap detector, an automatic reaction control chemical ionization technique in, for quantitative plasma profiling of arecoline in treated Alzheimer patients, 2:168
- Ion trap/ion mobility/time-of-flight mass spectrometer, nano-flow LC with, to analyze protein mixtures, 15:1341
- Ion trap-ion mobility time-of-flight mass spectrometry, analysis of ion/ion interactions, 20:1549
- Ion trap mass spectrometer
- diethylene glycol monobutyl ether contamination of, 14:1247
 - laser desorption in, sensitivity for, a method for increasing, 4:706
 - multiparticle simulation of ion motion in, resonant and direct current pulse excitation, 6:57
- quadrupole
- coupling a particle beam interface directly to, 4:38
 - high resolution on, 2:198
 - positions, velocities, and kinetic energies of resonantly elicited ions in, determination of, by laser photodissociation, 4:792
 - surface-induced dissociation of molecular ions in, 2:487
 - self-ionization of arylglycosides in, 14:851
- Ion trap mass spectrometric analysis, Tenax collection with jet-separator enrichment and, air analysis using, 7:1172
- Ion trap mass spectrometry. *See also* Quadrupole ion trap mass spectrometry
- in advances in analytical and atmospheric chemistry of 2-hydroxy-2-methylpropanal, a proposed photooxidation product of 2-methyl-3-buten-2-ol, 13:530
 - in confirmation of amoxicillin degradation products, ESI-TOF mass determination in, 16:1670
 - cyclic peptide sequencing in, multistep MS/MS for, 10:732
 - practical aspects of, 10:80
 - deoxynucleoside adducts with heterocyclic aromatic amines, analysis by, 18:2107
 - determination by, chemometrics in, 9:938
 - to determine linkage position in cobalt-coordinated pentasaccharides, 9:1125
- DNA, fragmentation of isomeric intrastrand cross-link lesions, 20:611
- ESI coupled to, phosphopeptide fragmentation by, 13:274
- of isomeric O-alkyl S-(2-dialkylamino) ethyl methylphosphotriolate fragmentation and reactions, 12:902
- laser desorption, of self-assembled monolayers, 4:968

- linkage and branched determination, N-linked oligosaccharides, 18:248
- miniature toroidal, 17:916
- methyl orange photolytic degradation in water, 19:997
- multiple stage quadrupole, *E. coli* cardiolipin characterization by ESI with, 17:420
- multiplexed multilinear MS, co-occurrence of boundary and resonance ejection, 17:56
- negative, in determining linkages of linear and branched oligosaccharides, 13:1322
- neutral lactooligosaccharide fragmentation in neutral ion mode UV-MALDI-TOF and UV-MALDI ion trap/TOF MS, 17:67
- orthogonal injection, TOF-MS, 20:1342
- personal perspective on, 13:589
- PFTBA ions/analyte molecule reaction, tobacco pyridine alkaloid determination by, 18:1774
- phosphopeptide fragmentation in, 9:1175
- plasma source, enhanced abundance sensitivity by resonant ejection of atomic ions, 7:1161
- quadrupole
- negative chemical ionization in, effects of applied voltages and reaction times, 5:757
 - plasticizer contamination from vacuum system O-rings in, 13:901
 - probing trapped ion energies via-molecule reaction kinetics, 3:716
 - in quantitative analysis of H₂, He, O₂, and Ar, 13:1120
 - resonance ejection mode of, mass resolution in, extended theoretical considerations for, 5:676
- reversed-phase HPLC with, as monoclonal antibody trap, 16:307, 16:797 (erratum)
- secondary ion microprobe, 13:1099
- selective chemical ionization of nitrogen and sulfur heterocycles in petroleum fractions by, 4:322
- in simulation of organic environment on Saturn's moon (Titan), 16:850
- simulations in micrometer-sized cylindrical ion trap, 17:430
- in situ SIMS analysis and reactions of surfaces prepared by SL of ions, 20:949
- tandem, quantitative analysis of crude plasma extracts by, 3:372
- of Ti₄L₄ cluster, 9:1099
- ZoomQuant software with, ¹⁸O labeling on, quantification using, 16:916
- Ion trap mass spectrometry/gas chromatography/alternating RF/DC isolations for quantitation with coeluting internal standards in, 3:85
- Ion trap mass spectrometry (IT-MS), instructional analysis of chromophore-labeled disaccharides, 11:292
- Ion trap mass spectrometry quantitative methods/capillary gas chromatography, combined, using labeled or unlabeled internal standards, 1:265
- Ion trap-MS
- cationized phosphotyrosine fragmentation by, 14:581
 - in determining polycyclic aromatic hydrocarbons in solid matrices, 14:562
 - in trypsin-catalyzed ¹⁶O-to-¹⁸O exchange, for comparative proteomics, 14:704
- Ion trap quadrupole. *See* Quadrupole ion trap.
- Ion trap reflection time-of-flight mass spectrometry, capillary liquid chromatography MS/MS on, for peptide sequencing and protein identification, 11:127
- Ion trap-secondary ion mass spectrometer, characterization of mercury nitrate surface bis(alkylamine)mercury cations by, 9:35
- Ion trap secondary ionization mass spectrometry (IT-SIMS), of VX on concrete, 11:69
- Ion trap source time-of-flight mass spectrometry, a segmented ring, cylindrical, 7:1009
- Ion trap tandem mass spectrometry
- applied to small multiply charged oligonucleotides with a modified base, 5:740
 - data-dependent, in proteomics, 15:150
 - of dexamethasone transformation products on light-activated TiO₂ surface, 12:1286
 - of heparin disaccharide dissociation mechanisms, 15:1274
- Ion trap tandem mass spectrometry and selected ejection chemical ionization, application of nonresonance excitation to, 7:668
- Ion trapping, improved, on kinetic energy modulation for, electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry and, 6:76
- Ion trapping mass spectrometers, a pulsed-leak valve for use with, 7:118
- Ion type, new fragment, in the collision-induced dissociation spectra of peptides, identification of, formation of alpha-2-16 ions, 4:424
- Ion velocities, mean initial, in infrared MALDI-MS, 13:209
- Ion yields, relative
- for glow discharge mass spectrometry, theoretical calculations of, 3:79
 - measured with a high-resolution glow discharge mass spectrometer operated with an argon/hydrogen mixture, 4:278
- Ionic diffusion, in SIFT-MS, 12:863
- Ionic liquid matrices, peptides, proteins and carbohydrate analysis by MALDI MS, 20:1790
- Ionic-liquid matrices, for quantitative analysis by MALDI-TOFMS, 15:1833
- Ionic liquid matrix
- metastable decay of peptides and oligonucleotides and stabilization of phospholipids induced by, 16:2000
 - for phospholipid analysis by MALDI-TOF MS, 16:679
- Ionic liquids
- LIFDI-MS of, 18:2254
 - molecular ions in gas phase, 19:1347
 - N,N'-disubstituted imidazolium, ESMS fragmentation study, 17:85
- Ionic matrices, ME-SALDI MS, 20:2229
- Ionic stable species, ionic, in peptide spectra, 6:1165
- ionic terpolymer samples, monomer ratios in, qualitative assessment by ESI-MS with CID, 13:921
- Ionization
- analyte, effect of solvent and surface effects on, DIOS MS, 19:8
 - atmospheric pressure chemical, reduction in, 13:109
 - chemical
 - atmospheric pressure, of alkanes, alkenes, and cycloalkanes, 5:177; Erratum, 5:1026
 - the early days of, a reminiscence, 1:277
 - iron-molecule reaction under, gas-phase halo alkylation of H₆O-fullerene by, 4:894
 - O⁻ and OH⁻, of some fatty acid methyl esters and triacylglycerols, 5:553
 - of peptides, Fe⁺, 4:106
 - of phenyl n-propyl ether and methyl substituted analogs, propene loss initiated by competing proton transfer to the oxygen atom and the aromatic ring, 7:639
 - silver nitrate, for analysis of hydrocarbon polymers by laser desorption Fourier transform mass spectrometry, 4:453 - competitive, of tetraphenylporphyrin in a laser-generated metal ion plasma, 4:159

- condensed-phase processes in, evidence indicating, fast-atom bombardment of the cyclic acetals, 4:493
- detection of femtomole and sub-femtomole levels of peptides by, 2:91
- matrix design for, sensitive determination of PAH-DNA adducts, 5:1021
- electron, induced fragmentation of macrocyclic amines $C_nH_{2n-1}NH_2$, evidence for the rearrangement of amino-carbene radical cations and a comparison with long-chain esters, 1:238
- electron capture of explosives with a microflow rate particle beam interface, 7:753
- electrospray
- of copper-glycine solutions, 7:25
 - and low energy tandem mass spectrometry of polyhydroxy unsaturated fatty acids, 7:140
 - tandem mass spectrometry analysis of peptides derived by enzymatic digestion of oxidized globin subunits, an improved method to determine amino acid substitution in the hemoglobin "core," 7:1040
- electrospray/capillary zone electrophoresis parameters for the mass spectrometry and tandem mass spectrometry analysis of peptides, optimization of, 3:289
- enhanced efficiency, metal-metal interface, MALDI MS, 17:737
- Fourier transform ion cyclotron resonance/infrared laser desorption mass spectrometry, reagent ions in, potassium halide adducts as, 3:301
- high precision ambient sources, construction of, MALDESI MS, 19:1527
- high-pressure, in a Fourier transform ion cyclotron resonance mass spectrometer, concentric tube vacuum chamber for high magnetic field, 4:168
- laser desorption, matrix-assisted, of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, high resolution end group determination of low molecular weight polymers by, 7:449
- low magnetic field internal electrospray, Fourier transform ion cyclotron resonance mass spectrometer, a high performance, 7:915
- matrix-assisted laser desorption
- employing a tandem double focusing magnetic-orthogonal acceleration time-of-flight mass spectrometer, peptide sequence determination by, 7:1
 - time of flight mass measurement accuracy, evaluation of, by using delayed extraction, 7:995
- mechanisms of NI-APPI and NI-DART, 20:42
- mechanisms related to negative ion APPI, APCI and DART, 20:1518
- medium pressure laser, ion source for, 15:1885
- negative, electron capture, in the detection of chemically oxidized dexamethasone in human plasma, comparison of tandem and conventional mass spectrometry using, 1:341
- negative chemical
- with oxygen, dibenzothiophene subjected to, oxidation reactions of, 4:949
 - in quadrupole ion trap mass spectrometry, effects of applied voltages and reaction times, 5:757
- nitric oxide-assisted atmospheric pressure corona discharge, for the analysis of automobile hydrocarbon emission species, 5:1107
- primary, factors affecting, DA-APPI, 17:130
- selected ejection chemical, and ion trap mass spectrometry, application of nonresonance excitation to, 7:668
- soft, of organic compounds, PERCI for, 15:424
- suppression of, in ESI, 11:942
- Ionization and collision-induced fragmentation of N-linked and related carbohydrates using divalent cations, 12:926
- Ionization and fragmentation of neutral and acidic glycosphingolipids with a Q-TOF mass spectrometer fitted with a MALDI ion source, 12:1220
- Ionization and infrared matrix-assisted laser desorption by using a tunable mid-infrared free-electron laser, 7:1187
- Ionization cavity, thermal, for MS, 10:1008
- Ionization efficiency, laser pulse, femtosecond *vs.* nanosecond, for multiphoton ionization of molecules, 6:1059
- Ionization electrospray. *See* Electrospray ionization
- arginine-containing peptide ions formed by, modeling the maximum charge state of, 7:972
 - characterization of the opioid peptide-containing protein and of bovine alpha-lactalbumin by, 5:377
 - doubly protonated angiotensin II formed by, charge-separation reactions in tandem mass spectrometry of, the importance of, experimental considerations and structural implications, 7:30
 - large multiply charged polypeptides and proteins produced by, collisional activation and collision-activated dissociation of, 1:53
 - on a magnetic sector mass spectrometer, improved detection limits for, by using an array detector, 5:194
 - mass spectrometry
 - capillary zone electrophoresis, application of, to the analysis of macrolide antibiotics, 3:563
 - deconvolution by an entropy-based algorithm, 3:207
 - nanoscale separations combined with, sulfonamide determination, 3:139
 - of nucleobases and nucleosides, processes that affect, 7:1106 - a new approach for the study of gas-phase ion-ion reactions using, 3:695
 - pneumatically assisted, microbore high-performance liquid chromatography-tandem mass spectrometry utilizing, quantitative determination of a nonpeptide antithrombotic in dog plasma by, 7:1238
 - special issues on (editorials), 4:523, 4:615
- Ionization energies, double
- of fluorinated benzene molecules, 4:513
 - to ground and excited states of the tungsten hexacarbonyl dication, 2:108
- Ionization Fourier transform ion cyclotron resonance-laser desorption mass spectrometry, collision-induced dissociation in, improved efficiency of, selective parent ion axialization for, 5:900
- Ionization from water solution with SF₆ discharge suppression, negative ion electrospray mass spectrometry of nucleotides, 4:289
- Ionization-laser desorption mass spectrometry, matrix-assisted, matrices used for, effects of heavy-atom substituents on, 5:800
- Ionization mass spectrometry
- electrospray
 - gas-phase molecular dications formed from neutral organics in solution via chemical electron-transfer reactions by using, observation of, 5:689
 - improved signal intensity of oligonucleotides in, charge-state reduction with, interaction of angiotensin peptides and zinc metal ions by, 5:959, 7:697
 - of oligonucleotides, specific and nonspecific, dimer formation in, 6:159
 - study of noncovalent enzyme-inhibitor complexes and metal, binding stoichiometry of matrolysin by, 6:1105
 - a study of the thermal denaturation of ribonuclease S by, 5:614 - electrospray/gradient reverse-phase high-performance liquid chromatography analysis with trifluoroacetic and other

- strong acid modifiers by postcolumn addition of propionic acid and isopropanol, signal enhancement of, 6:1221
- laser desorption, aerosol matrix-assisted, 5:1
- liquid secondary, direct desorption of crown ether-metal ion complexes in, evidence for, 5:638
- matrix-assisted, for polymer characterization, evaluation of, 7:11
- matrix-assisted laser desorption, and electrospray, characterization of mouse switch variant antibodies by, 7:707
- static secondary, detection of cyclohexamine on soil surfaces exposed to laboratory air, 7:168
- static secondary analysis, of tributyl phosphate on mineral surfaces, effect of Fe(II), 6:165
- Ionization matrix-assisted. *See* Matrix-assisted laser desorption/ionization matrix-assisted laser desorption
- Ionization mechanism, effect on peptide ion internal energies produced by ESI and AP-MALDI, 16:743
- Ionization mechanism in fast-atom bombardment involving ion/molecule reactions, reagent ions in, 8:337
- Ionization method, bombardment, model pulsed, a study of time-dependent desorption using, improved detection limits for fast atom bombardment mass spectrometry, 3:637
- Ionization mode
 - effects on CSR fragmentation reactions, 12:571
 - in MS profiling of whole bacteria, 10:747
- Ionization pathways, of polyfluorinated chemicals, environmental atmospheric pressure and, 14:1177
- Ionization reagents, oxygenated chemical, with vincamine, ion-molecule reactions of, 7:250
- Ionization-related conformational differences of peptide ions in a quadrupole ion trap, 12:1331
- Ionization source, electrospray, small-bore, construction and characterization of, 4:964
- Ionization spectroscopy, multiphoton, Fourier transform ion cyclotron resonance detection of, 7:1018
- Ionization tandem mass spectrometry
 - desorption chemical, of polyprenyl and dolichyl phosphates, 7:958
 - electrospray, structural determination of picomole amounts of phospholipids via, 6:1202
- Ionization tandem reflection time-of-flight mass spectrometry/matrix-assisted laser desorption of fullerenes, 7:590
- Ionization techniques, atmospheric pressure, for the analysis of N-methyl carbamate pesticides by liquid chromatography mass spectrometry, an evaluation of, 3:378
- Ionization time-of-flight mass spectrometry, matrix-assisted laser desorption, and electrospray mass spectrometry, application of, for molecular weight assignment of peptides in complex mixtures, 4:670
- Ionization time-of-flight/matrix-assisted laser desorption mass spectrometry of peptides, improved mass accuracy in, 5:955
- Ionized 1,5-hexadiene, structure in gas phase, 12:840
- Ionized aliphatic ethers, ion-neutral-mediated alkane eliminations from, size effects in, 2:261
- Ionized methylene transfer, from the distonic radical cation +CH₂-O-CH₂ to dicyclic compounds, pentaquadrupole mass spectrometric study, 6:554
- Ionized nitrile sulfides, gas phase interaction with pyridine, 14:241
- Ionized phytyl methyl ether, C₃H₅O⁺ formation from, the structure and mechanisms of, 7:205
- Ionized propyl propanoate, formation of the 3-pentanone ion from, through ion-neutral complexes, 3:183
- Ion(s)
 - atom reactions with, fragmentation from, 8:587
 - containing S•bonds, MS/MS of, 8:605
 - excitation frequencies of, in quadrupole field, 11:10
 - external accumulation of, for ESI-FTICR, 8:970
 - gas phase gramicidin S, hydration of, 8:565
 - gaseous multiply charged, electron capture dissociation of, by FTICR-MS, 12:245
 - high energy, from chemical ionization source, 8:1262
 - initial velocity of, MALDI-generated, TOF-MS effects of, 8:209
 - ion reactions with, in homomyoglobin, 8:637
 - mass-selected, radial separation of, in FTICR, 8:86
 - mobility of, long-range potential and, 8:275
 - multiply charged, cleavage mechanisms in, 12:565
 - parent, resolution of, in linked scans, 8:545
 - polyiodide, thermochemistry of, 8:688
 - pull needed to escape from droplet, 8:1147
 - thermochemical stabilities and structures of, in gas phase, 16:1760
- Ions. *See* specific element or ion type
- Ions
 - 193-nm photodissociation of, from saturated and unsaturated aliphatic molecules, 7:114
 - axial, selective ejection from linear quadrupoles with added octopole fields, 19:270
 - background, produced by fast-atom bombardment of glycerol, origins and structures of, 5:72
 - CID fragmentation, IMS MS, 19:411
 - collisional energy and fragmentation, peptides, 20:469
 - controversial, (m/z 59, 5:60, 5:77, and 119) in the thermospray reagent plasma from ammonium acetate, on the origin of some, 5:186
 - copper phthalocyanine, heterolytic dissociation of negative ions into counter ions, 19:375
 - effect of cross-contribution to intensity of, quantitative GC MS, 19:598
 - formation mechanism in laser desorption ionization, 19:389
 - fragmentation and fluorescence of in ultrahigh vacuum, ESI, 20:1731
 - fragmentation in low-pressure linear ion trap, 18:1459
 - gas-phase protein conformations from different solution conformations, 19:1907
 - gas stage, applications of multiple stage mass spectrometry in a pentaquadrupole, polar [4+2+] Diels-Alder cycloaddition to, 6:1
 - incoherent production reactions, MALDI MS, 20:1078
 - multiply charged, resolved isotopic peaks from, charge states from, automated assignment of, 6:52
 - multiply protonated, high energy collisional activation of, complete sequence confirmation of large peptides by, 6:234
 - organic, relative collision cross sections of organic ions, 6:803
 - peptide and protein, proton transfer reactivity, formed by electrospray ionization, 6:1086
 - positive, mechanisms in APCI, 19:1927
 - protein, charge state distribution shifting of, observed in matrix-assisted laser desorption ionization mass spectrometry, 6:1183
 - quadrupole excitation in linear quadrupole traps, 19:510
 - suppression and enhancement of secondary formation, SIMS, 18:1559
 - tetrameric, of noncovalent streptavidin complexes formed by electrospray ionization, dissociation of, 6:459
 - thermalized, in-source fragmentation technique for production of, 20:2312
- IRMM-007 series, synthetic isotope mixture, mass discrimination using MC-ICPMS, 17:1412
- IRMPD. *See* Infrared multiphoton dissociation
- IRMS. *See* Isotope ratio mass spectrometry

Iron

- cysteine-containing peptide complexes with, 8:199
ground state and Fe⁺, the reactions of, with the 20 common amino acids, 7:722
- II and III, gaseous complexes with BINOLate ligands, 19:121
transition-metal mediated heteroatom removal by reactions of FeL⁺ [L⁺O, C₄H₆, c-C₅H₆, H₆H₆, C₅H₄(=CH₂)] with furan, thiophene and pyrrole in the gas phase, 7:938
- Iron carboxylate oxygen-centered-triangle complexes, detected during electrospray use of organic acid modifiers with a comment on the Finnigan TSQ-700 electrospray inlet system, 6:1226
- Iron chelation, MS of, 8:1070
- Iron-oligonucleotide complex, MALDI FT-ICR MS of, 10:941
- Iron oxide nanoparticles, analysis of peptides and proteins bound to, MALDI-MS, 18:910
- Iron pentacarbonyl ions undergoing surface-induced dissociation in a Fourier transform mass spectrometry, energy deposition in, 3:445
- Iron-sulfur metalloprotein, 3+ oxidation state for, 12:817
- Iron(II), complexes of, with cysteine-containing peptides in the gas phase, 7:977
- Iron(II)phenanthroline, activation energy estimation using fragmentation yield:CID, ESI-MS, 17:962
- Irradiation, sustained off-resonance, capillary electrophoresis Fourier transform ion cyclotron resonance mass spectrometry with, for the characterization of protein and peptide mixtures, 5:894
- Irradiation effects in MALDI, ablation, ion production, and surface modifications 2,5-dihydroxybenzoic acid monocrystals, 14:893
- IRS-1. *See* Insulin receptor substrate-1
- Islands of stability, mass analysis with linear quadrupoles and octopole fields, 18:826
- Isoaspartic acid, aspartic acid and, differentiated by electron transfer dissociation, 17:15
- Isoaspartyl residues, quantification in deaminated proteins by ECD, 18:48
- Isobaric interferences
in DE-MALDI-TOFMS of complex polymer mixtures, 12:911
deconvolution of, in mass spectra, 15:654
- Isobaric monosaccharides, distinguishing by ion pairing and MS/MS, 16:1827
- Isobaric tags for relative and absolute quantification, with ETD, quantification of proteins and peptides, 19:1255
- Isobutylene/ μ -methylstyrene copolymers, compositional analysis by MALDI-MS, 15:621
- Isoocyanic acid, neutral loss of, CID of peptides, 20:723
- Isoelectric focusing, capillary, MS with, on-line desalting in, 8:743
- Isoindoloquinazolines, stereoisomeric, mass spectra fragmentations in, 14:1049
- Isoleucine, leucine and, discrimination using low energy ESI-TRAP MS, 18:57
- Isomer differentiation by charge inversion tandem mass spectrometry, an investigation into the structure of the ionic products from an Sn(ANRORC) reaction, 3:467
- Isomer differentiation in 7,12-dimethyl[a]anthracene-pyridine adducts by fast atom bombardment tandem mass spectrometry, 2:256
- Isomer differentiation in Fourier-transform mass spectrometry, 193 nm laser photoionization and photodissociation for, 1:361
- Isomeric C8- and N²-deoxyguanosine adducts of 2-acetylaminofluorene, differentiation of, by fast-atom bombardment and tandem mass spectrometry, 5:58
- Isomeric cations, C₃H₆Br⁺, gas phase bimolecular chemistry of, 6:11
- Isomeric dibenzopyrene-adenine adduct, MS/MS determination of, 8:977
- Isomeric dipeptides, differentiation and quantitation of, 12:490
- Isomeric ion-neutral complexes generated from ionized 2-methylpropanol and n-butanol, the effect of the polarity of the neutral partner on complex-mediated reactions, 3:409
- Isomeric mixtures, structural analysis of, by nano-electrospray MS, 13:1341
- Isomeric O-alkyl S-(2-dialkylamino)ethyl methylphosphonothiolates, fragmentation and reactions of, ESI-ITMS of, 12:902
- Isomeric parent ions, reactions from, in the dissociation of dimethylpyrroles, 7:930
- Isomeric peptides, binary and ternary mixtures of, kinetic method in recognition and quantification of, 14:152
- Isomeric silylenium ions (SiC₃H⁺9, SiC₄H⁺1a, SiC₅H⁺13) structures of, elucidating of, by using specific ion/molecule reactions, 2:278
- Isomeric species
differentiation by two-stage PTR-TOF MS, 19:325
recognition by ion/molecule reactions, case of ionized phenol-cyclohexadienone, 19:126
- Isomeric tryptophans, modified, structural characterization and differentiation of, 13:1298
- Isomerization
of 1,5-hexadiyne radical cation, 8:599
aldose-ketose, gas-phase, mechanism of, a study using tandem mass spectrometry and theoretical calculations, 4:943
of alkene radical ions, 10:577
of CH₃+HCH₂CH₂ to CH₃CH₂CH₂OH⁺, five-membered ring H shift in by ab initio theory, preference for ion-neutral complex-mediated pathway over, 6:1037
to fulvene structure, 1,2-hydrogen shifts in, benzene radical cation, Ab initio calculations on, 6:513
of H₆H₆ radical cations, 8:594
during ion isolation in the quadrupole ion trap, evidence of, 3:680
of linear C₃H₃⁺ in its reaction with acetylene, and collisional stabilization of the [C₅H₅⁺] collision complex in a quadrupole ion trap mass spectrometer, 3:631
- Isomerization of CH₃O+=CHCH₃ to CH₂=O+CH₂CH₃, 130
- Isomerization of the 1,3-hexadien-5-yne radical cation to the benzene structure and to nonclassical ion structures, Ab initio calculations of, 7:731
- Isomerization proclivities and stabilities studied by dissociation and neutralization-reionization, the distonic ion CH₂CH=CHOH⁺, and related C₃H₆O⁺ radical cations, 7:573
- Isomers
absolute configuration assignment of, by MS, 16:431
geometrical, α -linolenic acid methyl ester, separated using low energy ESI/MS, 20:1272
phosphorylated positional, quantification by FTICR, 14:916
- Isopropanol, postcolumn addition of, signal enhancement for gradient reverse-phase high-performance liquid chromatography-electrospray ionization mass spectrometry analysis, trifluoroacetic and other strong acid modifiers in, 6:1221
- Isoprostanes, F₂, four regioisomers of, formed by free radical oxidation of arachidonic acid, mass spectrometric analysis of, 7:490
- Isoquinoline-3-carboxamides, dissociation pathway analyzed by ESI-CID, 20:2034
- Isoquinolines, gas phase reaction to carboxylic acids after ESI and CID, 19:151
- Iso(thio)cyanates, acylium ions interaction with, 16:1602

- Isotope, stable, analyzing labeling data in proteomics, 16:1181
- Isotope abundance
 deconvolution of, 8:659
 mass peak profile generation model of, 8:170
- Isotope abundance measurement, improved, for FTICR-MS, 9:799
- Isotope analysis, SIFT-MS for, 11:866
- Isotope beating effects in the analysis of polymer distributions by FTMS, 10:1074
- Isotope cluster calculation, by probability grouping, polypeptides, 20:295
- Isotope coded affinity technology mass spectrometry, protein disulfide isomerase redox sensitive thiols identified using, 18:260
- Isotope depletion of large biomolecules: implications for molecular mass measurements, 9:149
- Isotope dilution gas chromatography/mass spectrometry for platinum determination in urine, 2:85
- Isotope dilution mass spectrometry, protein cleavage coupled with, ALiPHAT evaluation of LOD correlation and, 20:2006
- Isotope dilution single detector inductively coupled plasma magnetic sector MS (ID-ICD-SMS), applied to isotopically enriched cadmium, 16:708
- Isotope-edited product ion assignment by α -N labeling of peptides with [2H₃(50)]₂,4-dinitrofluorobenzene, 10:448
- Isotope exchanges, disulfur monoxide-water charged complexes, FT-ICR, CI-CAD and TQ, 18:1664
- Isotope incorporation, for element composition, 11:835
- Isotope label quantitation, ZoomQuant for, 16:302
- Isotope labeling, in phosphopeptide quantification, 15:363
- Isotope pattern evaluation, elemental composition by, ESI-FTICR MS, 17:1692
- Isotope ratio mass spectrometry, detecting ¹³C by, 10:153
- Isotope-ratio-monitoring gas chromatography/mass spectrometry, nitrogen isotope analyses by, 5:387
- Isotope ratios
 of carbon dioxide in expired air from mice and men following the administration of ¹³C-methyl methionine, the quadrupole-based mass spectrometric evaluation of, 3:742
 of molybdenum, in terrestrial molybdenites, measurement of, high accuracy, 3:10
- Isotope ratios and abundance sensitivity obtained with an inductively coupled plasma-time-of-flight mass spectrometer, 6:920
- Isotope-resolved mass spectra, algorithm for charge state determination from PTFT, 19:46
- Isotopes
 carbon and oxygen ratios in whole foliage and cellulose, *Pinus* spp., 19:1330
 isotopic fine molecular structures, hierarchical algorithm for calculation, 19:1868
 labeling, a_n and a_n^* peptide fragments, MS/MS, density functional theory calculations and, 19:1788
 labeling, RABA, quantitative proteomics, 20:1366
 peptide, method for prediction of distributions, 19:703
 stable labeling, a- and b-type peptide fragment ions, MS/MS, 19:1781
- Isotopic dilution cryofocusing gas chromatography/mass spectrometry, on-line, automated high-speed analysis of selected organic compounds in urban air by, 5:1001
- Isotopic distribution calculations, in analysis of H/D exchange mass spectra of peptides and proteins, 12:1153
- Isotopic distributions, resolved, monoisotopic masses and ion populations for large biomolecules from, determination of, 6:229
- Isotopic peaks
 calculation of accurate masses of, 17:415
 individual, dissociation of, 14:310
 isotopic composition and accurate masses of, 15:12
 resolved, for multiply charged ions, charge states from, automated assignment of, 6:52
- Isotopic purity, in selectively polydeuterated surfactants, charge-remote fragmentation in a hybrid (BEqQ) spectrometer to determine, 1:85
- Isotopic standard, a particulate, of uranium and plutonium in an aluminosilicate matrix, 5:852
- Isotopic standard of plutonium in an aluminosilicate matrix, particulate, 2:81
- Isotopically labeled reagent gas, chemical ionization mass spectrometry using, 5:596
- Isotopologues, overlapping, GC/MS separation and quantitation, 17:1028
- Isotopomer abundances, fragmentation effects on, 10:430
- IT-MS. *See* Ion trap mass spectrometry.
- IT-SIMS. *See* Ion trap secondary ionization mass spectrometry.
- Iterative data analysis, is the key for exhaustive analysis of peptide mass fingerprints separated by two-dimensional electrophoresis, 14:943
- ITMS. *See* Ion trap mass spectrometry.
- iTRAQ. *See* Isobaric tags for relative and absolute quantification
- Iturin A₂, sodium stabilization of, during ESI-MS, 12:505
- J
- JASMS, twentieth birthday of, editorial, i (Jan)
- Jet disrupter electrode, automatic gain control in MS using, 16:244
- Jet-separator enrichment and ion trap mass spectrometric analysis, tenax collection with, air analysis using, 7:1172
- Jitter in a linear orthogonal-acceleration time-of-flight mass spectrometer, resolution limitations from, 5:772
- Journal of the American Society for Mass Spectrometry*, Critical Insights feature, 19:627
- K
- K⁺ ionization of desorbed species mass spectrometric technique, characterization of the primary thermal degradation processes of peptides using, 5:554
- Kassel statistical theory, 8:103
- Keap-1. *See* Kelch-like ECH-associated protein 18:1
- Kelch-like ECH-associated protein 1, alkylation sites by natural chemoprevention agents, 18:2226
- Kemptide, ADP-ribosylated, MS/MS, 20:477
- Ketalization of gaseous acylium ions, 12:150
- Ketalization of phosphonium ions by 1,4-dioxane, selective detection of chemical warfare agent stimulant DMMP in mixtures using ion/molecule reactions, 14:182
- Keto acids, long-chain conjugated, analysis of by negative ion electrospray mass spectrometry, 6:1190
- Keto-enol tautomerism, effect on ESI-MS of diketone/metal ion complexes, 10:402
- Ketones, ammonia and hydrazine complexes with, mobility resolution and mass analysis of ions from, 18:940
- Ketose-aldehyde isomerization, gas-phase, mechanism of, study using tandem mass spectrometry and theoretical calculations, 4:943
- K⁺IDS, K⁺ ionization of desorbed species mass spectrometric technique, characterization of the primary thermal degradation processes of peptides using, 5:554
- KIEs. *See* Kinetic isotope effects
- Kinetic and spectroscopic characterization of the isomers of the allyl bromide molecular ion, 2:371
- Kinetic constant determinations, of yeast hexokinase, using MS-based assays, 14:173
- Kinetic energies, positions, velocities and, of resonantly elicited

- ions in the quadrupole ion trap mass spectrometer, determination by laser photodissociation, 4:792
- Kinetic energy distribution of precursor ion main beam in massanalytical ion kinetic energy spectrometry experiments, measurement of, 6:1232
- Kinetic energy orifice, as probe of metastable dissociation, in FTICR-MS, 11:1118
- Kinetic isotope effects, deuterium
 microsolvated gas-phase E2 reactions, 19:1296
 in microsolvated gas-phase E2 reactions, 18:1046
- Kinetic method
 determination of orders of relative alkali metal ion affinities of crown ethers and acyclic analogs by, 3:543
 to determine mononucleotide gas phase proton affinities, 11:24
 in determining proton affinity of nucleic acid molecules, 15:1080
 in discrimination of arginine dipeptide isomers, 15:840
 electron affinity of 1,3,5,7-cyclooctatetraene by, 9:1141
 entropy measurements and, 11:371
 in estimating proton affinity values of MALDI matrices under ESI conditions, 15:431
 in examination of alkali chloride cluster ion dissociation, 13:1388
 microcanonical analysis of, 13:435
 in recognition and quantification of binary and ternary mixtures of isomeric peptides, 14:152
- Kinetic method and ligand exchange technique, comparison of the orders of gas-phase basicities and ammonium ion affinities of polyethers by, 4:718
- Kinetic method with multiple stage mass spectrometry, SiCl_3^+ and SiCl^+ affinities for pyridines determined by using, agostic effects in the gas-phase, 7:198
- Kinetic studies, millisecond time-scale, microfluidic chip for, ESI-MS, 20:124
- Kinetics and mechanism of the collision-activated dissociation of the acetone cation, 3:427
- Kinetics-based protocol for instrument-independent measurements of collision-activated dissociation in RF-only quadrupoles, precision and accuracy in tandem mass spectrometry measurements, 1:272
- Koala, rapid analysis of dissolved gases, lake water testing of chemistry with portable and AUV-based MS, 19:1403
- Krypton, collision cross sections of myoglobin and cytochrome c with, 8:681
- Kv1.3 peptide ligands and margatoxin, noncovalent complexes of, model investigation using ion-spray mass spectrometry, 7:1075
- Kynurenine fragmentation, ESI-MS/MS in, 12:786
- L
- l-(2-benzothiazolyl)-3,5-diphenyl formazans, gas-phase oxidation and reduction of, 10:339
- LA-ICP-MS. *See* Laser ablation inductively coupled plasma mass spectrometry
- LA-MS. *See* Laser ablation mass spectrometry
- Label-free quantitation, glycoproteins quantify glycosylation changes by, 20:1048
- Labeling in vivo, for protein identification, 13:804
- Laboratory air, soil surfaces exposed to, cyclohexamine on, static secondary ionization mass spectrometry detection of, 7:168
- Laboratory practices, good, mass spectrometry and, 7:211
- Lactalbumin, alpha-bovine, characterization of, by electrospray ionization and liquid secondary ion mass spectrometry, 5:377
- Lactams, selective ion-molecule reactions of, with dimethyl ether ions, 4:353
- Lactate dehydrogenase, classification by LC-MS and multivariate analysis, 14:236
- Lactacin 481, ECD-SORI-CID study of, 16:1595
- β -Lactoglobulin
 conformation of gas-phase ions, 19:1907
 glycosylated peptides, following gastrointestinal digestion, MS, 19:927
- Lactones, gas-phase protonated and methylated, dissociation behavior of, characterization of the, 3:47
- Lactooligosaccharides, neutral, fragmentation in negative ion mode, UV-MALDI-TOF and UV-MALDI ion trap/TOF MS, 17:67
- Lake Louise Workshop on Tandem Mass Spectrometry, The Eighth, 7:611
- Lake Ontario sediments, plutonium 240/239 ratios in, determination of, 2:240
- Lake water, testing of chemistry with portable and AUV-based MS, 19:1403
- Lamivudine, use of PEG as shift reagent, IMS, 20:1
- Lamivudine metabolites, in peripheral mononuclear blood cells of HIV-infected patients, HPLC-MS/MS of, 11:1134
- Lamps, VUV, mobile phases and, detection of PAH compounds, APPI-MS, 18:589
- Langmuir-Blodgett multilayers, model for molecular depth profiling by MS, 19:96
- Lanosterol, quantification by LC-APPI-MS/MS in vitro, 15:1216
- Lanthanide(III) porphyrin-phthalocyanine heteroleptic sandwich complex, ESI-FTICR studies of, 8:161
- Lanthionine polysulfides, characterization of lanthionine trisulfides and tetrasulfides from peptide disulfides, ESI MS, 20:783
- Large biomolecules, Fourier-transform ion-cyclotron resonance mass spectrometry of, improved, 4:190
- Large insulating samples, imaging, by using secondary ion tandem mass spectrometry, charge compensation for, 5:37
- Large ion populations, line broadening at, a mechanism for, two-dimensional Coulomb-induced frequency modulation in Fourier transform ion cyclotron resonance, 4:909
- Large molecules
 in a linear time-of-flight mass spectrometer, high-resolution mass spectrometry of, 4:2
 photoionization of, 6:883
 tandem high-resolution mass spectrometry of, Fourier-transform electrospray instrumentation for, 4:557
- Large peptides, complete sequence confirmation of, by high energy collisional activation of multiply protonated ions, 6:234
- Large volume injection gas chromatography, pulsed, H-POPs analyzed by pLVI-GC/ECNI-qMS, 18:1375
- Lariat crown ethers, lysine probing by, in peptides, 13:493
- Lariat ether amides, selectivities of, ESI-MS evaluation of, 14:1215
- Laser
 tunable mid-infrared free-electron, infrared matrix-assisted laser desorption and ionization by using, 7:1187
 UV, ablation products of organic polymers, direct chemical analysis of, by using selective ion monitoring mode in gas chromatography/mass spectrometry, 5:106
- Laser ablation
 for ICP-MS, 15:769
 resonant, as a selective metal ion source for gas-phase ion molecule reactions, 7:664
- Laser ablation inductively coupled plasma mass spectrometry, metal imaging on micro and nanoelectronic surfaces by, 20:883
- Laser ablation mass spectrometry
 activation processes and polyethylene formation on Phillips model catalyst analyzed by LDI-MS, s-SIMS and, 17:406
 cluster ions of inorganic transition metal compounds, 19:488

- Laser beam, features smaller than, MALDI-MS of, 16:1654
- Laser desorbed ion kinetic energies, electrostatic energy analyzer-time-of-flight mass spectrometer for measurement of laser desorbed ion kinetic energies, design and calibration of, 6:619
- Laser desorption
- chemical ionization Fourier transform mass spectrometry, energetics of peptide ion dissociation by, 6:1069
 - electron attachment TOFMS, new approach to detection of involatile compounds, 12:1339
 - fast gas chromatography-mass spectrometry in supersonic molecular beams, 9:628
 - for ruthenium and osmium trisbipyridine complexes, FTMS evaluation of, 14:393
 - and TOF-MS
 - direct *versus* matrix-assisted, laser microprobe mass spectrometry of quaternary phosphonium salts, 4:798
 - infrared/ionization Fourier transform ion cyclotron resonance mass spectrometry, reagent ions in, potassium halide adducts as, 3:301
 - infrared matrix-assisted, and ionization by using a tunable mid-infrared free-electron laser, 7:1187
 - ion trap mass spectrometry, of electrodeposited polythiophene film, 8:1203
 - matrix assisted. (*See* Matrix-assisted laser desorption/ionization)
 - matrix assisted-ionization mass spectrometry, application to low mass poly(methylmethacrylate) polymers, fundamentals of, 7:287
 - matrix-assisted/ionization tandem reflectron time-of-flight mass spectrometry of fullerenes, 7:590
 - neutrals generated by, gas-phase cationization and protonation of, 4:393
 - observed fast fragmentation of peptides in, factors that influence, 7:225
 - substrate effects of, 8:525
 - and TOF-MS
 - of bacterial lipooligosaccharides, 8:645
 - of nitrated polycyclic aromatic hydrocarbons, 8:630
 - in transmission geometry inside a FT-ICR mass spectrometer, 10:1105
 - Laser desorption-chemical ionization, in detection of pharmaceutical compounds in tissue, 10:1315
 - Laser desorption Fourier transform mass spectrometry
 - analysis of copolymers by, 1:66
 - matrix-assisted, structural characterization of normal and modified oligonucleotides by, 2:402
 - for peptides, UV matrix-assisted, investigation of, 2:22
 - Laser desorption in an ion trap mass spectrometer, sensitivity for, a method for increasing, 4:706
 - Laser desorption ionization
 - ammonia/chemical, with ammonium bromide, Fourier transform ion cyclotron resonance mass spectrometry of aromatic hydrocarbons, 2:299
 - ion formation mechanism in, 19:389
 - mass spectrometry, matrix-assisted, and electrospray ionization mass spectrometry, characterization of mouse switch variant antibodies by, 7:707
 - matrix-assisted
 - time of flight mass measurement accuracy, evaluation of, by using delayed extraction, 7:995
 - time-of-flight mass spectrometry, liquid chromatography-electrospray ionization, rapid analysis of antibiotic-containing mixtures from fermentation broths by using, 7:1227
 - of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, high resolution end group determination of low molecular weight polymers by, 7:449
 - detection of femtomole and sub-femtomole levels of peptides by, 2:91
 - employing a tandem double focusing magnetic-orthogonal acceleration time-of-flight mass spectrometer, peptide sequence determination by, 7:1
 - matrix design for, sensitive determination of PAH-DNA adducts, 5:1021
 - sub-micrometer resolved, SMALDI for, 13:735
 - Laser desorption/ionization
 - 10:l-(2-benzothiazolyl)-3,5-diphenyl formazan complex formation with transition metals under, 10:339
 - two-step, aerosol TOFMS with, for analysis of individual organic particles, 9:1068
 - Laser desorption ionization and MALDI-TOFMS for low molecular mass polyethylene analysis, 12:1186
 - Laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry
 - collision-induced dissociation in, improved efficiency of, selective parent ion axialization for, 5:900
 - in-cell matrix-assisted, 7:1026
 - Laser desorption/ionization (LDI) time-of-flight (TOF) mass spectrometry, probing biotic/abiotic boundary of plants with, 16:107
 - Laser desorption ionization mass spectrometric analysis of proteins immobilized on nylon-based membranes, direct matrix-assisted, 5:230
 - Laser desorption ionization mass spectrometry
 - aerosol matrix assisted, 5:1
 - CHCA-modified Au nanoparticles for peptide analysis, 20:1530
 - detection of gaseous $W_{12}O_{41}^{\bullet}$ and other tungstate anions by, 20:1782
 - matrix-assisted
 - for polymer characterization, evaluation of, 7:11
 - quantitative assessment of cysteine and cystine in peptides and proteins following organomercurial derivatization and analysis by, 5:359
 - surface-induced dissociation analyzed using SPALDI and, peptides, 20:957
 - Laser desorption-ionization mass spectrometry, matrix-assisted, matrices used for, effects of heavy-atom substituents on, 5:800
 - Laser desorption/ionization mass spectrometry
 - activation processes and polyethylene formation on Phillips model catalyst analyzed by LA-MS, s-SIMS and, 17:406
 - ballpoint dye photofading on paper analyzed by MALDI MS and, 17:297
 - infrared, capillary gel microfluidic chips interfacing with, 17:469
 - peptide fingerprinting on porous silica and alumina, 19:632
 - Laser desorption ionization time-of-flight mass spectrometry, matrix-assisted, of underivatized and permethylated gangliosides, 3:785
 - Laser desorption/ionization time-of-flight mass spectrometry, titanium oxide characterization of nanoclusters, MALDI-TOF-MS and, 18:517
 - Laser desorption/ionization time-of-flight mass spectrometry of peptides, matrix-assisted, improved mass accuracy in, 5:955
 - Laser desorption/laser photoionization ion trap-MS, in determining polycyclic aromatic hydrocarbons in solid matrices, 14:562
 - Laser desorption mass spectrometry
 - Fourier transform analysis of hydrocarbon polymers by, silver nitrate chemical ionization for, 4:453

- ion trap, of self-assembled monolayers, 4:968
- ionization
- matrix-assisted, with 2-(4-hydroxyphenylazo)benzoic acid matrix, 4:399
 - time-of-flight, matrix-assisted, and electrospray mass spectrometry, application of, for molecular weight assignment of peptides in complex mixtures, 4:670
 - matrix-assisted, of biotransformation products of dynorphin A *in vitro*, 5:10
 - negative-ion matrix-assisted, analysis of oligodeoxynucleotides by, 4:955
- Laser desorption postionization mass spectrometry, of adsorbed and covalently bound bisphenol A diglycidyl methacrylate, 18:1097
- Laser desorption time-of-flight mass spectrometry, analysis of hydrocarbon dendrimers by, 5:731
- Laser flash photolysis, of hydrogen peroxide, for microsecond photochemical oxidation of proteins, 16:2057
- Laser focus, volumetric intensity dependence within, 16:82
- Laser-generated metal ion plasma, competitive ionization of tetraphenylporphyrin in, 4:159
- Laser-induced acoustic desorption
- charged phenyl radicals evaporated by, gas-phase reactions of, 13:192
 - with CID in FT-ICR-MS, peptide sequencing using, 18:525
- Laser-induced azido photochemistry, fingerprint patterns from, 10:1337
- Laser induced electron impact ionization, application in HWCVD process, SiH₄-NH₃ gas mixture, 18:1950
- Laser-induced fluorescence for ion tomography in a Penning trap, 9:925
- Laser-induced liquid bead ion desorption, binding sites of TAR and TAR mutants for peptide ligands, 19:1605
- Laser-induced liquid bead ion desorption mass spectrometry, membrane protein analysis by, 18:1429
- Laser-induced thermal-desorption/Fourier-transform mass spectrometry for surface analysis, deflection of laser-produced ions in, 5:583
- Laser ionization, medium pressure, ion source for, 15:1885
- Laser ionization orthogonal time-of-flight mass spectrometry, multielemental analysis of biological samples by, 20:1355
- Laser microdissected cells, MALDI-MS analysis of, 13:1292
- Laser microprobe mass spectrometry, of inorganic compounds, 9:482
- Laser microprobe mass spectrometry of quaternary phosphonium salts, direct *versus* matrix-assisted laser desorption, 4:798
- Laser photodissociation
- determination of positions, velocities, and kinetic energies of resonantly elicited ions in the quadrupole ion trap mass spectrometer by, 4:792
 - positions, velocities, and kinetic energies of resonantly elicited ions in the quadrupole ion trap mass spectrometer, 4:792
- Laser photoionization and dissociation, 193 nm, for isomer differentiation in Fourier-transform mass spectrometry, 1:361
- Laser photoionization time-of-flight mass spectrometry, femtosecond, nitro-aromatic explosives and related compounds, 20:419
- Laser photolysis, charged phenyl radical and biradical synthesis by, in FTICR mass spectrometer, 9:1135
- Laser probe, determination of ion frequencies in a quadrupole ion trap by, 5:29
- Laser-produced ions, deflection of, in laser-induced thermal-desorption/Fourier-transform mass spectrometry for surface analysis, 5:583
- Laser pulse, duration of, in infrared MALDI-MS, 13:978
- Laser pulse ionization efficiency, femtosecond *vs.* nanosecond, for multiphoton ionization of molecules, 6:1059
- Laser spray, denaturation of lysozyme and myoglobin in, 16:860
- Laser spray mass spectrometry, binding affinities of protein-mutant DNA complexes, 17:611
- Laskin, Julia, 2008 Bieman medal recipient, editorial, i (June)
- LC. *See* Liquid chromatography entries; Liquid chromatography
- LC-ESI-IT MS. *See* Liquid chromatography-electrospray ionization-ion trap-mass spectrometry
- LC-ESI-MS, spin-trapped radicals determination by, 2:413
- LC-(ESI)MS. *See* Liquid chromatography-(electrospray ionization) mass spectrometry
- LC-MS. *See* Liquid chromatography mass spectrometry
- LC/MS. *See* Liquid chromatography/mass chromatography; Liquid chromatography/mass spectrometry
- LC/MS and LC/MS/MS determination of protein tryptic digests, 1:158
- LC-MS/MS. *See* Liquid chromatography-tandem mass spectrometry
- LC/MS/MS determination of protein tryptic digests, 1:158
- LC-RAM-MS. *See* Liquid chromatography radiochemical activity monitor mass spectrometry
- LC-TSP-MS, spin-trapped radicals determination by, 2:413
- LD/CI. *See* Laser desorption-chemical ionization
- LDI. *See* Laser desorption ionization; Laser desorption/ionization
- LDI MS. *See* Laser desorption ionization mass spectrometry
- LDI-TOF-MS. *See* Laser desorption/ionization time-of-flight mass spectrometry
- LDITMS. *See* Laser desorption ion trap mass spectrometry
- Lead, gas-phase interactions between lead(II) ions and thiouracils, 20:359
- Leaving group and gas phase neighboring group effects in side chain losses from protonated serine and its derivatives, 11:1047
- Leech, hemoglobin from, subunit structure of, ESI-MS of, 15:22
- Leishmania major*, inositol phosphorylceramides, characterization by ESI-MS/MS, 18:1591
- Lenses, ion deceleration, what are the simplicity/performance trade offs?, 2:322
- Letters, 11:182
- Letters to the editor, 2:349, 3:471, 4:363, 10:661
- on computer software to calculate weight in Daltons of ions, 15:626
- Leucine, isoleucine and, discrimination using low energy ESI-TRAP MS, 18:57
- Leucine and isoleucine anions, separation by ESI-FAIMS-MS, 10:1279
- Leucine enkephalin
- ESI-F AIMS-MS of, 10:492
 - fragmentation with laser fluence, MALDI-TOF-TOF-MS, 18:607
 - low-energy sequential decomposition pathways of, using multiple-resonance CID in quadrupolar ion guide, 15:1794
- Leucine enkephalin-alka(e)nal-glycerophosphatidylcholine adducts, identification by MS/MS, 17:657
- Leukemia, acute myeloid and chronic lymphocytic, histone modifications in, HPLC and MS of, 15:77
- Leukine enkephalin, dissociation of, 8:771
- Leukotriene A₄, collision-induced decomposition of, 13:1227
- Leukotriene B₄ (LTB₄) and LTB₄-derived metabolites, negative ion electrospray tandem mass spectrometric characterization of, 7:129
- Leukotriene B₄ metabolites, 3:3-hydroxylated, novel, from ethanol-treated rat hepatocytes, 3:762
- Leukotriene C₄, analysis of isomers using CID and 157nm photofragmentation, 19:14

- Leukotriene C₄, ESI and MS/MS of, 12:763
- Leukotriene E₄ and metabolites, in human urine, identification of, negative ion tandem mass spectrometry of, 2:314
- LHRH-metal ion complexes. *See* Luteinizing hormone-releasing hormone-metal ion complexes
- Li-Mg alloy liquid metal ion source, cluster ions ejected from, observation of Mg₂²⁺ and Mg₃²⁺, 2:76
- Library, mass spectral, 10:287, 10:565
- Library search, mass spectral
chemical substructure identification by, 6:644
estimating probabilities of correct information from, 5:316
- Library search algorithms, mass spectral, for compound identification, optimization and testing of, 5:859
- Library search results, electron ionization mass spectral, evaluating, 7:313
- Licorice flavonoids, quantification in rat plasma, 18:778
- LIEL. *See* Laser induced electron impact ionization
- LIFDI-MS. *See* Liquid injection field desorption/ionization mass spectrometry
- LIFT-TOF/TOF, of CHCA clusters, 15:336
- Ligand binding
noncovalent, to stemloop domains, HIV-1 packaging signal mapping, MS-MS, 17:1401
nonspecific, identification using reporter molecule method, ES-MS, 20:1242
by proteins, efficiency of MS-based screening assay for detection of, 19:1303
- Ligand exchange technique and kinetic method, comparison of the orders of gas-phase basicities and ammonium ion affinities of polyethers by, 4:718
- Ligand-induced site-specific phosphorylation of EGF receptor, 14:1022
- Ligand-protein and protein-protein noncovalent interactions, probing, application of electrospray mass spectrometry in, 4:624
- Light
proton transfer induced by, from gas phase cations to reference bases, 12:359
TiO₂ surface activated by, dexamethasone transformation products on, ion trap MS/MS of, 12:1286
- LILBID. *See* Laser-induced liquid bead ion desorption
- LILBID MS. *See* Laser-induced liquid bead ion desorption mass spectrometry
- Limit, of static secondary ion mass spectrometry, organic ion imaging beyond, 6:1047
- Limitations of a general unknown screening by gas chromatography-mass spectrometry, in acute poisoning, 6:995
- Limits of detection correlation, ALiPHAT evaluation of PC-IDMS and, 20:2006
- Line broadening at high mass and for large ion populations, mechanism for, two-dimensional Coulomb-induced frequency modulation in Fourier transform ion cyclotron resonance, 4:909
- Linear and cyclic polyglycols, [M + X]⁺ ions (X=H, Li, Na, K) from, dissociation characteristics of, 5:1081
- Linear dynamic detection range, in ESI-MS, 15:1416
- Linear ion trap-based tandem mass spectrometry, pulsed triple ionization source for sequential ion/ion reactions, 18:369
- Linear ion trap Fourier transform ion cyclotron resonance mass spectrometry, mechanistic studies on air amplifier, 18:1909
- Linear ion trap mass analyzer, novel, composed of four planar electrodes, 17:631
- Linear ion trap mass spectrometer, 8:962
- Linear ion-trap mass spectrometry
for data acquisition from ion/ion reactions, 19:1821
multistage, with ESI
long-chain unsaturated fatty acids, double-bond position assigned by, 19:1678
unsaturated glycerophospholipids characterized by, 19:1700
- Linear ion trap/orbitrap mass spectrometry, nefazodone liver microsome metabolism analysis by, 17:363
- Linear ion-trap tandem mass spectrometry, peptide sequence information from, 19:1813
- Linear ion traps
axially-resonant excitation, duty cycle enhancement, TOF-MS, 17:1669
CID and ETD in, protonated peptides, 20:1672
ion/ion proton transfer reactions in, 18:882
low-pressure, ion fragmentation in, 18:1459
mass selective axial ejection from, 20:1132
mass selective ejection by ARE, 17:685
MS/MS of ions in, 20:1714
off-resonance excitation, 20:443
study of dipolar resonance excitation enhancement by, 18:578
wavelength-tunable UVPD in, heparin-derived disaccharides, 20:1645
- Linear orthogonal-acceleration time-of-flight mass spectrometer, detector pulse width and jitter in, resolution limitations from, 5:772
- Linear quadrupole ion trap mass spectrometry, comparison with Orbitrap MS for peptide and protein identification, 20:1405
- Linear quadrupole mass filters, islands of stability, mass analysis with linear quadrupoles and octopole fields, 18:826
- Linear quadrupole mass spectrometer, fast scanning of, 8:539
- Linear quadrupoles, with added hexapole fields, 17:1063
- Linear response theory of ion excitation for Fourier transform mass spectrometry, 2:483
- Linear time-of-flight mass spectrometer
large molecules in, high resolution mass spectrometry of, 4:2
postsources pulse focusing conditions and mass scale, 6:889
two-stage, photodissociation of high molecular weight peptides and proteins in, 6:578
- Linearity and quadrupolarity of tetragonal and cylindrical Penning traps of arbitrary length-to-width ratio, 8:283
- Linkage position, assignment of, influence of the collision energy and of the collision gas for, collision-induced dissociation of alkali metal cationized and permethylated oligosaccharides, 4:197
- Linked-scan tandem mass spectrometry, product ion characterization by, trimethylsilyl group migration in the mass spectra of trimethylsilyl ethers of cholesterol oxidation products, 4:327
- Linked scans
artifact peaks in, 8:554
parent ion resolution in, 8:545
- α-Linolenic acid methyl ester, geometrical isomers separated using low energy ESI/MS, 20:1272
- Lipases
activity determination by LC MS, 18:1579
Candida rugosa, immobilized on PDMS membrane, use in MIMS, 18:973
- Lipid, silver complex formation with, CIS-MIS analysis of, 14:872
- Lipid A, *F. tularensis* subspecies *novicida*, environmentally-regulated remodeling, MS/MS, 18:1080
- Lipid chemistry, MALDI-FTMS of, strategy and data analysis techniques for, 15:1665
- Lipid hydroperoxides and long-chain conjugated keto acids, analysis of, by negative ion electrospray mass spectrometry, 6:1190
- Lipid peroxidation
of erythrocyte ghost phospholipids, 9:5-hydroperoxy and

- 5-hydroxyeicosatetraenoic acids generated by, ESI-MS, 9:527
- product of, peptides modified with, MS/MS of, 10:613
- Lipidomes, ornithine and glutamine lipids from *R. sphaeroides* cell membranes, MS/MS, 20:198
- Lipids
- LipidQA software for identifying and quantifying complex mixtures, 18:1848
- mapping of neuropeptides and, by 3D MSI, crustacean brain, 20:1068
- Lipoarabinomannans, from *Mycobacterium*, ESI-MS characterization of, 16:1109
- Lipooligosaccharides
- bacterial, LDI TOF-MS of, 8:645
- from *Haemophilus ducreyi*, poly lactosamine repeats in, 13:724
- from *Neisseria gonorrhoeae*, FTICR-MS in characterization of, 13:571
- Lipophilic phosphotriesters, FAB MS/MS of, 10:975
- Liquid chromatography
- coupling EIMS to, 13:265
- of flavonoid aglycones, 12:707
- impulse-driven droplet deposition system interface combining LC with MALDI MS and MS/MS, 17:325
- of oxidized deoxynucleosides, ESI-MS/MS for, 12:80
- of proteolytic digests, exponential dilution gradient system for, 12:1205
- radiochemical activity monitor mass spectrometry (LC-RAM-MS), of packed capillary, 13:82
- recombinant, quantification of *N*-glycosylation by LC/MS or, 19:1648
- reversed phase, reduces back exchange in FT-ICR MS, 20:520
- with tandem mass spectrometry
- chemical ionization, negative atmospheric pressure, -mass spectrometry, sensitivity in, effect of high-performance liquid chromatography mobile phase components on, 7:1059
- electrospray, -tandem mass spectrometry, quantification of SR 27417 in human plasma, study of ion suppression, 7:1099
- electrospray/mass spectrometry, selective detection of phosphopeptides in complex mixtures by, 4:710
- high performance, mobile phase components, sensitivity in negative atmospheric pressure chemical ionization liquid chromatography-mass spectrometry, 7:1059
- high-performance-particle beam-mass spectrometry, determination of polycyclic aromatic hydrocarbons by, 4:898
- of HT2a antagonist and metabolite in, 8:371
- with mass spectrometry, with collision-induced dissociation of conjugated metabolites of benzol[a]pyrene, 8:50
- microbore high-performance-tandem mass spectrometry utilizing pneumatically assisted electrospray ionization, quantitative determination of a nonpeptide antithrombotic in dog plasma by, 7:1238
- nanoscale packed capillary (75 m id), application of, to the analysis of macrolide antibiotics, 3:563
- normal phase/mass spectrometry, the application of, by using coaxial continuous flow fast atom bombardment, 3:575
- particle beam/mass spectrometry and gas chromatography/mass spectrometry, poly(amic acid) and polyimide characterization using, 5:299
- particle beam/mass spectrometry interface, performance of, effect of a dc glow discharge on, investigation of, 5:908
- reversed phase, of peptides, 8:495
- reversed phase-microbore/electrospray mass spectrometry, 5:350
- steroid determination by, 8:1010
- with tandem mass spectrometry of complex protein digest mixtures, 8:1059
- Liquid chromatography and electrospray mass spectrometric mapping of peptides from human plasma filtrate, 10:45
- Liquid chromatography atmospheric pressure chemical ionization mass spectrometry, quantitative analysis by, determination of the renin inhibitor CP-80,794 in human serum, 2:164
- Liquid chromatography-atmospheric pressure chemical ionization mass spectrometry (LC-APCI-MS)
- of bitter acids in crude hop extract, 15:180
- of mutagenic nitropyrenes, 15:1853
- Liquid chromatography-atmospheric pressure photoionization-tandem mass spectrometry (LC-APPI-MS/MS)
- lanosterol and FF-MAS quantification by, in vitro, 15:1216
- of mutagenic nitropyrenes, 15:1853
- Liquid chromatography collision-induced dissociation tandem mass spectrometry, protein identification by LC-ECD-MS/MS or, 18:891
- Liquid chromatography-electrochemical mass spectrometry (LCEC-MS), ferrocenoyl piperazide using, 15:1727
- Liquid chromatography electron capture dissociation tandem mass spectrometry, protein identification by LC-ESI-MS/MS or, 18:891
- Liquid chromatography-electrospray ionization-Fourier transform ion cyclotron resonance mass spectrometry (LC-ESI-FTICR-MS), with internal bore spectrometer, magnetic field focused ion accumulation for, 11:592
- Liquid chromatography-electrospray ionization-ion trap-mass spectrometry, multistage MALDI-IT/RTOF-MS and, moenomycin characterization by, 17:1081
- Liquid chromatography electrospray ionization mass spectrometry
- Ganoderma lucidum* triterpenoids analyzed using, 18:927
- rhubarbs, analysis of phenolic compounds using, 18:82
- Liquid chromatography-(electrospray ionization) mass spectrometry, of diquat and paraquat in water by, 9:830
- Liquid chromatography-electrospray ionization-mass spectrometry matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry, rapid analysis of antibiotic-containing mixtures from fermentation broths by using, 7:1227
- Liquid chromatography electrospray ionization tandem mass spectrometry
- nano, protein-RNA interactions detected by, 18:869
- negative, response enhancement, alkyl methylphosphonic acids, 18:1821
- Liquid chromatography-electrospray ionization-tandem mass spectrometry (LC-ESI-MS/MS)
- of ethyl sulfate as marker of alcohol consumption, 15:1644
- of ethylglucuronide in urine, 15:188
- miniaturized, DNA-phosphate adduct analysis with, 15:593
- Liquid chromatography-electrospray mass spectrometry (LC-ESMS), negative ion, neutral O-linked oligosaccharide alditol structural determination by, 15:659
- Liquid chromatography-electrospray-MS, toxicological screening by, 14:14
- Liquid chromatography-electrospray tandem mass spectrometry (LC-ES-MS/MS), miniaturized, of melphalan-modified 2'-oligodeoxynucleotides, 15:568
- Liquid chromatography-ESI-ion trap MS, of deoxyguanosine linkage to deuterated 2-hydroxyestradiol, 14:364
- Liquid chromatography-ESI-MS/MS
- in identification of two-dimensional gel proteins, 14:956
- LC-MALDI-MS/MS complementation of, in proteomics, 14:971
- of nonylphenol carboxylates in water and sludge, 14:516

- Liquid chromatography-Fourier transform ion cyclotron resonance (LC-FTICR), for proteomics, 15:212
- Liquid chromatography-IMS-CID-TOFMS techniques, incorporation of, 14:1424
- Liquid chromatography ion trap time-of-flight mass spectrometry, tau-related peptide quantification using GLaD with, 18:201
- Liquid chromatography/ion trap/time-of-flight mass spectrometry (LC/ion trap/TOFMS), for peptide sequencing and protein identification, 11:127
- Liquid chromatography laser-induced fluorescence mass spectrometry, enkephalin reaction with NDA for, 18:1706
- Liquid chromatography (LC), elution time of, in complex proteome analysis, 16:1239
- Liquid chromatography (LC) ion trap mass spectrometry, in calibration experiments of trilinolenoylglycerol, 16:388, 16:801 (erratum)
- Liquid chromatography (LC) mass spectrometry carbohydrate quantitative analysis by, 16:1805 of polypeptide fragments, 16:38 stable gradient nanoflow, on QqTOF instrument, 16:1545
- Liquid chromatography (LC) matrix-assisted laser desorption/ionization (MALDI) tandem mass spectrometry (MS/MS), microwave-assisted acid hydrolysis of protein with, 16:471
- Liquid chromatography (LC) tandem mass spectrometry (MS/MS) mercapturic acid detection by, 16:1976 for peptide fragmentation, 16:505 of polymer surface color bodies, 16:295 in residue determination of compounds, 16:1619
- Liquid chromatography-MALDI-MS/MS, LC-ESI-MS/MS complementation of, in proteomics, 14:971
- Liquid chromatography mass spectrometry (LC-MS) for the analysis of N-methyl carbamate pesticides, evaluation of atmospheric pressure ionization techniques for, 3:378 of electrospray ionization source H/D exchange metabolites, 13:345 generic, using experimental design, 13:2 genetic algorithm feature extraction, alignment of data sets from LC-MS/MS and, 18:1835 global view of, 10:78 gradient elution, electrospray voltage selection for, 20:682 high performance, of porphyrins by using an atmospheric pressure interface, 7:965 for high throughput microsomal stability screening of compound libraries, 13:155 of HIV gp120 glycopeptides, 15:1545 human plasma proteome mapping using SCX-LC-IMS-MS, 18:1249 of 4-hydroxy-2-alkylquinolines from *Pseudomonas aeruginosa*, 15:862 in identifying minor compounds in complex mixtures, COMSPARI and, 15:580 lipase activity determination by, 18:1579 mass spectrometric analysis of nitrogen-and phosphorus-containing pesticides by, 5:655 micro high-performance, analysis of coumarins by, with a particle beam interface, 6:132 postcolumn derivatization, chemical weapons-related compound detection using, 10:440 reagents for chemical noise reduction, 20:105 in semi-quantitative genotyping of single nucleotide polymorphisms, 15:1897
- Liquid chromatography/mass spectrometry atmospheric pressure ionization with, 9:187 capillary-scale particle-beam interface with, 9:3 cyclic polyamide oligomer contamination from nylon 66 filter disks, 17:652 derivatization of 5-fluorouracil for, 9:971 high dead volume pumps, delay time reduction by short-term low-ratio split flow, 18:245 rapid protein identification by, 9:194
- Liquid chromatography mass spectrometry for the analysis of N-methyl carbamate pesticides with collision-induced dissociation, the determination of glycopeptides by, 3:804 particle beam mass spectra of chlorinated phenoxyacid herbicides obtained by, evidence for thermal decomposition contributions to, 3:823 mass transport and calibration in, 3:662
- Liquid chromatography mass spectrometry interface, atmospheric pressure chemical ionization, optimization of, 7:69
- Liquid chromatography-mass spectrometry interface, Hewlett-Packard particle-beam by statistical experimental design, optimization of, 6:507
- Liquid chromatography/mass spectrometry (LC/MS), to determine molecular component weights, MoWeD computer program in, 12:599
- Liquid chromatography/mass spectrometry (LS/MS), with supersonic molecular beams, 11:587
- Liquid chromatography mass spectroscopy arylamine-modified oligonucleotide sequence verification by, 19:1147 MAPK-activated PK-2, limited proteolysis determined by, 19:841 *Ostreopsis ovata* palytoxin and ovatoxin-a analyzed by, 19:111 recombinant, quantification of N-glycosylation by LC or, 19:1648 structural variation of peptidoglycan from vancomycin-susceptible *E. faecium*, 19:1467
- Liquid chromatography-matrix-assisted laser desorption/ionization-time-of-flight/time-of-flight (LC-MALDI-TOF/TOF), in protein complex analysis, 15:803
- Liquid chromatography-MS of diazinon photocatalytic degradation products, 14:803 in lactate dehydrogenase classification, 14:236
- Liquid chromatography-MS/MS in bioanalysis of illicit drugs, matrix effect in, 14:1290 isotope dilution, in arylamine DNA adduct detection, 14:1057
- Liquid chromatography nanospray mass spectrometry, response normalized, 18:1891
- Liquid chromatography particle beam mass spectrometry, high-performance, investigation of enhanced ion abundances from a carrier process in, 1:92
- Liquid chromatography particle beam-mass spectrometry, with massive cluster impact, 7:293
- Liquid chromatography-photodiode array detection mass spectrometry, hawthorn leaf α -glucosidase inhibitors characterization using SORI-CID FTICR MS and, 20:1496
- Liquid chromatography-quadrupole time-of-flight mass spectrometry, protein expression by *E. coli* and *Shigella* species monitored by, 19:1626
- Liquid chromatography tandem mass spectrometry an analytic strategy of quaternary ammonium neuromuscular blocking agents, 17:1456 archaeal tetraether lipids analyzed by, 20:51 aristolochic acid DNA adducts, 18:642 in automated de novo DNA sequencing, 15:32 to detect steroids in human urine, 10:1322 genetic algorithm feature extraction, alignment of data sets from LC-MS and, 18:1835 glycoprotein monosaccharides, 20:1224

- intelligent automated, indinavir metabolite identification by, 10:175
- isotope ratio-based profiling of microbial folates, 18:898
- lanosterol and FF-MAS quantification by, *in vitro*, 15:1216
- licorice flavonoid quantification in rat plasma, 18:778
- nano, prion protein detection of attomoles by, 18:1070
- of N-terminal phospholipase A2 processing and amyloid precursor protein potential cleavage sites, 15:1780
- Liquid chromatography/tandem mass spectrometry analysis of peptide mixtures, data-controlled automation of, 7:532
- Liquid chromatography tandem mass spectrometry (LC-MS/MS)
- of electrospray ionization source H/D exchange metabolites, 13:345
 - variable flow nanoscale, for improved proteome coverage, 13:760
- Liquid chromatography-TOFMS, capillary, proteome analyses with, 14:980
- Liquid flow-injection isotope ratio mass spectrometry, direct analysis of carbon isotope variability in albumins by, 7:605
- Liquid injection field desorption/ionization mass spectrometry, of ionic liquids, 18:2254
- Liquid introduction via heated nebulizer interface, atmospheric-pressure chemical ionization mass spectrometry with, characterization of flame-generated C₁₀ to C₁₆₀ polycyclic aromatic hydrocarbons by, 7:276
- Liquid matrices, deposition on conductive hydrophobic surfaces, UV-MALDI MS, 18:693
- Liquid matrix, in electronic excitation under fast-atom bombardment conditions, M⁺ ion formation and, 6:114
- Liquid metal ion source, Li-Mg, cluster ions ejected from, observation of Mg₂²⁺ and Mg₃²⁺, 2:76
- Liquid phase, cyclopropylphenyl benzamide cyclization in, 12:956
- Liquid phase composition, in ESI, 12:1077
- Liquid reactions, in APCI-MS sources, 15:311
- Liquid secondary ion
- with mass spectrometry, nanoliter flow analysis of protein digests using, 8:703
 - with tandem mass spectrometry, of synthetic polymers, 8:76
- Liquid secondary ion mass spectrometry
- alkali chloride cluster formation in, ion-ion recombination for, role of, 4:410
 - analysis of permethylated, n-hexylamine derivatized oligosaccharides, application to baculovirus expressed mouse interleukin-3, 3:345
 - characterization of the opioid peptide-containing protein and of bovine alpha-lactalbumin by, 5:377
 - detection of thermally unstable intermediate in Wittig reaction by, 9:1168
- Liquid-secondary-ion mass spectrometry, beam-induced reaction between meta-nitrobenzyl alcohol and dipyrro cyanine dyes in, 5:1017
- Liquid secondary ion mass spectrometry/fast-atom bombardment, reduction/oxidation processes observed in, true effect of experimental parameters on, evaluation of, 4:482
- Liquid secondary ion mass spectrometry/fast-atom bombardment by tandem mass spectrometry, beam-induced reactions occurring in, characterization of, 4:312
- Liquid secondary ionization mass spectrometry, direct desorption of crown ether-metal ion complexes in, evidence for, 5:638
- Liquid secondary ionization mass spectrometry beam-induced dehalogenation of organic compounds, secondary electron capture in, evidence for a mechanism that involves, 7:1109
- Liquid surface, surface-induced dissociation from, 4:769
- Liquids, direct analysis by DSI-MS, 20:10
- Liquid(s), ion soft-landing into, 15:1874
- LIT MS/MS. *See* Linear ion trap-based tandem mass spectrometry
- Lithiated adducts, glycosphingolipids as, ESI-MS and CAD of, 12:61
- Lithiated valine, water binding energies to, 15:1014
- Lithium adducts of unsaturated fatty acids, ESI-MS of, using CAD, 10:600
- Lithium and transition metal ions enable low energy CID of polyglycols in ESI-MS/MS, 12:832
- Lithium atoms, complexes with formaldehyde and formaldehyde, stability of, 12:1229
- Lithium-attachment ions, poly(ethylene glycol), tandem mass spectrometry of, 5:1072
- Lithium-attachment ions from polyglycols, tandem mass spectrometry of, 3:225
- Lithium-cation and proton affinities of sulfoxides and sulfones: a Fourier transform ion cyclotron resonance study, 8:262
- Lithium-peptide adducts, gas-phase, investigation of, tandem mass spectrometry and semiempirical studies, 1:473
- Liver, nefazodone metabolism, analysis by linear ion trap/orbitrap MS, 17:363
- Living block copolymer, poly(ethylene oxide)/polystyrene, microstructural study by ESI MS/MS, 19:1163
- LMMS. *See* Laser microprobe mass spectrometry
- L2MS. *See* Two-step laser desorption/ionization
- Location of abasic sites in oligodeoxynucleotides by MS/MS and by chemical cleavage initiated by an unusual reaction of the ODN with MALDI matrix, 13:1418
- LOD. *See* Limits of detection
- Long-chain aldehydes, structural determination of, by gas chromatography combined with mass spectrometry, pyridine-containing Schiff base derivatives for, 2:245
- Long-chain carboxylic esters, characterization of, with CH₃OBOCH₃⁺ in a small Fourier-transform ion cyclotron resonance mass spectrometer, 7:1138
- Long-chain conjugated keto acids, negative ion electrospray mass spectrometry, 6:1190
- Long-chain esters, electron ionization induced fragmentation of macrocyclic amines C_nH_{2n}-1NH₂, evidence for the rearrangement of aminocarbene radical cations, 1:238
- Long-chain fatty acyl coenzyme A thioesters, by negative ion fast-atom bombardment mass spectrometry and tandem mass spectrometry, 5:416
- Long-lived metallized tips for nanoliter electrospray mass spectrometry, 7:1270
- Loo, Joe, Critical Insights feature, 19:627
- Loss of internal 1–6 substituted monosaccharide residues from underivatized and per-O-methylated trisaccharides, 8:43
- Loss of ions during forward and reverse scans in a quadrupole ion trap mass spectrometer and how to reduce them, 8:1141
- Loss of selenium from selenoproteins: conversion of selenocysteine to dehydroalanine *in vitro*, 14:593
- Low collision energies, enhancement of ion transmission at, via modifications to the interface region of a four-sector tandem mass spectrometer, 5:460
- Low-energy collision-induced dissociation, characterization of a high-pressure quadrupole collision cell for, 5:1042
- Low energy collision-induced dissociation fragmentation of protonated ortho silicates, APCI, McLafferty or ion-neutral complex rearrangements, 5:544
- Low energy collision-induced dissociation studies of cluster ions on a VG analytical ZAB-2FQ, a compact high pressure ion source for, 3:33

- Low energy collision-induced tandem mass spectrometry and fast-atom bombardment mass spectrometry of tauroconjugated bile acid anions, 6:588
- Low energy decompositions of gas-phase peptide ions, role of the site of protonation in, 7:522
- Low energy ion beams, real-time images of, monitor for displaying, 2:226
- Low energy reactive ion bombardment, chemical modification of fluorinated self-assembled monolayer surfaces by, 6:187
- Low-energy side of the main beam in mass-analyzed ion kinetic energy spectra, origin of the tailing signal on, 3:644
- Low energy tandem mass spectrometry and electrospray ionization of polyhydroxy unsaturated fatty acids, 7:140
- Low energy tandem mass spectrometry of deoxynucleoside adducts of polycyclic aromatic hydrocarbon dihydrodiol-epoxides, 6:248
- Low-energy tandem mass spectrometry of the molecular ion derived from fatty acid methyl esters, a novel method for analysis of branched-chain fatty acids, 4:223
- Low flow high-performance liquid chromatography solvent delivery system designed for tandem capillary liquid chromatography-mass spectrometry, 6:571
- Low magnetic field internal electrospray ionization-Fourier transform ion cyclotron resonance mass spectrometer, a high performance, 7:915
- Low-mass ions produced from peptides by high-energy collision-induced dissociation in tandem mass spectrometry, 4:882
- Low mass poly(methyl methacrylate) polymers, electrospray ionization mass spectrometry and, 6:906
- Low-molecular weight anticancer drugs and their analogues, analysis of, electrospray ionization mass spectrometry for, 4:588
- Low molecular weight polymers, high resolution end group determination of, by matrix-assisted laser desorption ionization of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, 7:449
- Low nanogram per liter determination of halogenated nonylphenols, nonylphenol carboxylates, and their non-halogenated precursors in water and sludge by LC-ES-MS/MS, 14:516
- Low temperature aqueous electrospray ionization mass spectrometry of noncovalent complexes, 9:580
- Low voltage ion transport system for external ionization Fourier transform mass spectrometry, 5:398
- Low-voltage on-resonance ion selection, for FT-MS, 9:931
- LSI-MS/MS. *See* Liquid secondary ion, with tandem mass spectrometry
- LTC₄, mass spectrometric detection of, by derivitization, 5:292
- LTD₄, mass spectrometric detection of, by derivitization, 5:292
- LTE₄, mass spectrometric detection of, by derivitization, 5:292
- LTQ MS. *See* Linear quadrupole ion trap mass spectrometry
- LTQ Orbitrap hybrid MS, dynamic range of mass accuracy in, 17:977
- LU⁺ Sc⁺, and Y⁺, gas-phase reactions with alcohols, effects of the class and chain length of alcohols on the nature of primary products, 7:1157
- Lumbricus terrestris hemoglobin, MS of, 8:352
- Lung tumor, in low molecular weight serum proteome quantitative analysis, 16:1221
- Luteinizing hormone-releasing hormone-metal ion complexes, MALDI-FTMS of, 10:19
- Lymphoblastoid cells, thymosin β -4 identification from, SELDITOF MS in, 14:760
- Lymphocytic leukemia, chronic, histone modifications in, HPLC and MS of, 15:77
- Lymphosarcoma, acetylated histone H4 isoforms from, peptide mass mapping of, 16:1641
- Lysine
chemical cross-linking with tyrosine, 15:1604
lariat crown ether probing of, in peptides, 13:493
- Lysine acetylation
bacteriotoxin E9:Im9 complex analysis using nano-LC-MALDI-MS/MS and, 17:983
resolution of Efb complex with complement C3d domain using MS and, 19:55
- Lysine adducts, in cytochrome c modification by 4-hydroxy-2-nonenal, 15:1136
- Lysine and histidine, gas-phase basicities of, and their selected di- and tripeptides, 7:1203
- Lysine-specific mass tagging, in peptide sequencing and protein quantification, in human proteome, 14:1
- Lysoglycerophosphocholine lipid subclasses, ESI-MS of, 11:283
- Lysozyme, denaturation of, in laser spray, 16:860
- Lysozyme ions, multiply protonated, electron loss from, 12:889
- M
- [M + Cu]⁺ peptide ions, containing an N-terminal arginine, fragmentation chemistry of, 11:626
- (M + Fe-5H)²⁺ peptide ion composition, verification of, by FTMS and MS/MS, 11:83
- [M + H]⁺ formation, mechanism of, in photoionization MS, 15:1521
- [M + 4H]₄⁺ ions from insulin chain B, gas-phase reactivity and dissociation from, cysteic acid group effect on, 10:928
- [M + Ni]⁺ adducts of aliphatic amino acids, fragmentation of, 8:749
- μ -ESI-MS. *See* Microelectrospray ionization mass spectrometry.
- M-2H anion generation, benzoylnitrene radical anion as new reagent for, 18:2014
- μ -Liquid chromatography-ES-API-CID-MS hybrid scan technique, to identify IRS-1 phosphorylation site, 14:401
- μ -L²MS. *See* Microprobe two-step laser desorption/laser ionization
- M₂⁺ molecular ions
in quantitation of major elements with secondary ion mass spectrometry, 9:638
in sputtering processes, formation of, 10:246
m/z transmission, low, electrodynamic ion funnel evaluation of, 17:586
- MAB. *See* Metastable atom bombardment
- M(acetone)₂⁺ (M=Al, Fe, Co, Cu, ScO) in the gas phase, collision induced and infrared multiphoton dissociation studies on, 4:135
- Macrolide aglycones, structural elucidation of, 13:862
- Macrolide antibiotics
analysis of, application of nanoscale packed capillary liquid chromatography (75 μ m id) and capillary zone electrophoresis/electrospray ionization mass spectrometry to, 3:563
fast-atom bombardment and tandem mass spectrometry of, 5:151
- Macrolide immunosuppressant, new metabolite of, identification by HPLC and electrospray MS/MS, 11:516
- Macromolecule mass spectrometry, citation mining of user documents in, 15:281
- Macromolecules
ionization by nano-ESI, MCS, MCS, 19:1098
structure, charge state distributions, ESI MS, 19:1239
- Macrophage colony-stimulating factor β and derivatives, ESI FTICR-MS of, 11:237
- Magnet
high homogeneity resistive, FT-ICR MS in, 10:265
split pair, high pressure chemistry in FT-ICR MS using, 10:918

- Magnetic beads, aminophenylboronic acid immobilization on, by MALDI, 16:1456
- Magnetic field, high, concentric tube vacuum chamber for, high-pressure ionization in a Fourier transform ion cyclotron resonance mass spectrometer, 4:168
- Magnetic field focused ion accumulation for an internal bore LC-ESI-FTICR using a central trapping electrode, 11:592
- Magnetic nanoparticles, zirconium phosphonate-modified, selective phosphopeptide capture by, 19:1176
- Magnetic-orthogonal acceleration time-of-flight mass spectrometer, tandem double focusing, matrix-assisted laser desorption ionization employing, peptide sequence determination by, 7:1
- Magnetic sector mass spectrometer, electrospray ionization on, improved detection limits for, by using an array detector, 5:194
- Magnetic sector mass spectrometer with heated capillary inlet, high resolution electrospray ion source for, design and performance of, 6:1256
- Magnetic sector mass spectrometers, an electrospray ion source for, 2:205
- Major histocompatibility complex 1 molecules, β_2 -M release from, analyzed by HDX-ESI-MS, 20:278
- Makarov, Alexander, award for distinguished contribution in mass spectrometry, editorial, i (August)
- M=Al, Fe, Co, Cu, ScO, M(acetone)₂⁺, collision-induced and infrared multiphoton dissociation studies on, in the gas phase, 4:135
- Malaria, solid artesunate antimalarials, quantization of active ingredients by DESI MS, 19:380
- MALDI. *See* Matrix-assisted laser desorption/ionization; Matrix-assisted laser desorption/ionization detection of intact double-stranded DNA by, 6:972
- MALDI-IT/RTOF-MS. *See* Matrix-assisted laser desorption/ionization-ion trap/reflectron time-of-flight mass spectrometry
- MALDI MS. *See* Matrix-assisted laser desorption ionization mass spectrometry
- MALDI-MS, preservation and detection of specific antibody-peptide complexes by, 11:746
- MALDI-MS. *See* Matrix-assisted laser desorption/ionization mass spectrometry; Matrix-assisted laser desorption/ionization mass spectrometry (MALDI-MS); Matrix-assisted laser desorption/ionization-MS
- MALDI-MS/MS. *See* Matrix-assisted laser desorption/ionization tandem mass spectrometry
- MALDI-PSD. *See* Matrix-assisted laser desorption/ionization post-source decay
- MALDI-PSD spectra. *See* Matrix-assisted laser desorption/ionization postsource decay spectra
- MALDI time-of-flight mass spectrometry, Braun's lipoprotein and determination of its attachment sites to peptidoglycan by, characterization of, 6:892
- MALDI-TOF. *See* Matrix-assisted laser desorption/ionization-time-of-flight
- MALDI-TOF MS. *See* Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry
- MALDI-TOF/TOF MS. *See* Matrix assisted laser desorption/ionization-time-of-flight/time-of-flight mass spectrometry
- MALDI-TOFMS, practical quantitative biomedical applications of, 13:1015
- Maleic acid, PTFE fibers grafted with, trace rare earth element determination using, ICP-MS, 17:1258
- Mammalian cell lines, identification using MALDI-TOF and LC-ESI-MS/MS, 17:490
- Manganese ions, preferential solvation in water/methanol clusters, 19:520
- Mannose-6-phosphate oligosaccharides, synthetic, high-energy collision-induced dissociation mass spectrometry of, 7:182
- Mannose glycans, high, characterized by ion trap MS, bovine ribonuclease B, 20:539
- Mannotriose, of transition metals, electrospray ionization study of, 8:32
- MARCKS. *See* Myristoylated alanine-rich C kinase protein
- Marcus statistical theory, 8:103
- Margatoxin and the Kv1.3 peptide ligands, noncovalent complexes of, model investigation using ion-spray mass spectrometry, 7:1075
- Marine peptide, fragmentation of, gas phase intramolecular rearrangement in, 12:470
- Mass accuracy
dynamic range in LTQ Orbitrap hybrid MS, 17:977
high measured, of peptide fragments, CID spectra, 19:891
improved
in matrix-assisted laser desorption/ionization time-of-flight mass spectrometry of peptides, 5:955
for MS/MS, 16:463
PPB, artificial neural networks for, in FT-ICR/MS, 20:1303
- Mass analysis
high pressure ion trap, 20:2144
with linear quadrupoles and octopole fields, islands of stability, 18:826
shifting unoccupied spectral space in mass spectrum of peptide fragment ions, 20:2124
- Mass-analytical ion kinetic energy spectrometry experiments, kinetic energy distribution of precursor ion main beam in, measurement of, 6:1232
- Mass-analyzed ion kinetic energy spectra, main beam of, low-energy side of, origin of the tailing signal on, 3:644
- Mass analyzers, miniaturized, novel ion traps using planar resistive electrodes, 19:1435
- Mass and charge assignment for electrospray ions by cation adduction, 4:828
- Mass and charge state assignment for proteins and peptide mixtures via noncovalent adduction in electrospray mass spectrometry, 6:1175
- Mass correlated acceleration in a reflectron MALDI-TOF mass spectrometer, an approach for enhanced resolution over a broad mass range, 13:135
- Mass-correlated pulsed extraction, theoretical analysis and implementation with a linear MALDI-TOFMS, 11:841
- Mass determination of megadalton-DNA electrospray ions using charge detection mass spectrometry, 9:305
- Mass difference measurements, precise, Fourier transform ion cyclotron resonance, Coulomb mutual interaction and image charge in, systematic errors originating from, analysis and estimation of, 4:855
- Mass discrimination, in SIFT-MS, 12:863
- Mass error, in ESI-MS of proteins, water cluster calibration and, 8:1158
- Mass isotopic distributions, exact calculation of, 18:1511
- Mass isotopomer abundance mass spectrometry (MIAMS), of methyl palmitate, eliminating concentration dependence using MAB, 12:754
- Mass measurement
accurate, in MS, 14:1086
exact, FTICR-MS validation of, 16:1100
by FTICR, without internal standards using multiply charged ions, 11:416
in ion trap, 11:876
by MALDI-TOFMS, of low molecular weight compounds, 11:458
of stable isotopes, 11:835

- for target compound verification by quadrupole TOF MS, 16:406
- time-of-flight, -matrix-assisted laser desorption ionization, accuracy, evaluation of, by using delayed extraction, 7:995
- by TOFMS, of bromo- and chloroacetic acid, 11:809
- unresolved interferences in, peak shape analyses in, 9:789
- Mass measurement accuracy, intact proteins and product ions, ESI and Q-FT-ICR, 18:1
- Mass measurement errors caused by δ local δ frequency perturbations in FTICR-MS, 13:99
- Mass measurements, high-resolution accurate, of biomolecules using a new electrospray ionization ion cyclotron resonance mass spectrometer, 4:566
- Mass peak profile generation model to facilitate determinations of elemental compositions of ions based on exact masses and isotopic abundances, 8:170
- Mass range, broad, in the quadrupole ion trap, selective ion isolation over, 2:11
- Mass resolution
- of 11,000 to 22,000 with a multiple pass quadrupole mass analyzer, 11:407
 - enhanced, in removing interferences when measuring volatile organic compounds in human blood by using purge-and-trap chromatography/mass spectrometry, 3:831
 - resonance ejection mode of quadrupole ion trap mass spectrometry, extended theoretical considerations for, 5:676
- Mass-resolved chromatography, negative ion chemical ionization, polarizable stationary phase, and mass-resolved chromatography, high-temperature gas chromatography/mass spectrometry with, high-resolution triglycerol mixture analysis using, 7:350
- Mass-selected ions, thermal energy reactions of, high pressure drift reaction cell to study, hybrid double-focusing mass spectrometer, 1:197
- Mass selective axial ion ejection from a linear quadrupole ion trap, 14:1130
- Mass selective detector, Hewlett-Packard, implementation of the chemical reaction interface mass spectrometry technique on, 5:765
- Mass selective ejection, linear ion trap, by ARE, 17:685
- Mass spectra
- of 2-alkyl-4,6-dinitrophenols, novel $[M-35]^+$ fragment ions in characterization of, sequential ortho effects, 2:69
 - of alkyl diselenide and selenosulfenate, 15:1325
 - of copolymers with compositional drifts or sequence constraints, 15:374
 - deconvolution of isobaric interferences in, 15:654
 - electron capture negative ion, an ion source problem, 4:270
 - electron-ionization, data base of, an enlarged, 2:432
 - electrospray
 - capillary electrophoresis of herbicides paraquat and diquat, 7:981
 - of proteins, are they related to their aqueous solution chemistry?, 3:216
 - fast-atom bombardment, of azo dyes, effect of tautomerization on, 5:407
 - gaseous peptide aggregates hidden in, 9:743
 - high resolution
 - algorithms for automatic interpretation, 17:459
 - new matching algorithm for, 15:1173
 - of intact proteins, cumulative distribution functions in, 13:40
 - normalization of, 15:385
 - pure component, reconstruction with weighted two-band target entropy minimization, 14:1295
 - reference, to identify unknowns, 10:1229
 - tandem
 - collision-induced dissociation, high energy, of polycyclic aromatic hydrocarbon diepoxide adducted peptides, characteristics of, 5:649
 - feature-combinations from, generation of substructure identification rules using, 3:159
 - of peptide ions in the quadrupole ion trap, effects of heavy gases on, 7:1194
 - substructure identification rules from, automatically generated, evaluation of, 3:169
 - tandem high energy collision-induced dissociation, peptide sequencing from, pattern-based algorithm for, 3:326
 - time dependence of, II, fast atom bombardment-induced condensation of glycerol with ammonium surfactants, 5:92
- Mass spectra/MS, electron ionization, calibration point for, measured with multi-quadrupole mass spectrometers, 5:576
- Mass spectra of proteins, effect of ion-molecule collisions in the vacuum chamber of an electrospray time-of-flight mass spectrometer on, 7:342
- Mass spectra of trimethylsilyl ethers of cholesterol oxidation products, trimethylsilyl group migration in, product ion characterization by linked-scan tandem mass spectrometry, 4:327
- Mass spectral analysis, tandem, fragmentation of protonated O,O-diethyl O-arylphosphorothionates in, 7:189
- Mass spectral characterization of tetracyclines by ESI, H/D exchange, and multiple stage MS, 13:543
- Mass spectral data, tandem, of peptides, an approach to correlate with amino acid sequences of a protein database, 5:976
- Mass spectral data bases, comparative evaluations of, 2:438:441
- Mass spectral electron ionization, library search results, evaluating, 7:313
- Mass spectral enhancement, software-based, to remove interferences from spectra of unknowns, 7:598
- Mass spectral fragmentation
 - novel, of heptafluorobutyl derivatives of acyl analogues of platelet-activating factor, 2:476
 - reactions of a therapeutic 4-azasteroid and related compounds, 12:385
- Mass spectral identification and positional mapping of aflatoxin B₁-guanine adducts in oligonucleotides, 9:676
- Mass spectral imaging, 3D, crustacean brain, neuropeptide and lipid mapping by, 20:1068
- Mass spectral library, 10:287, 10:565
- Mass spectral library search algorithms, optimization and testing of, for compound identification, 5:859
- Mass spectral library search algorithms for compound identification, optimization and testing of, 5:859
- Mass spectral library searches, estimating probabilities of correct information from, 5:316
- Mass spectral library searching, chemical substructure identification by, 6:644
- Mass spectral methods, fast atom bombardment, for the identification of carcinogen-nucleoside adducts, development of, 3:360
- Mass spectrograph
 - a new, 4:372
 - plasma-source, 8:307
- Mass spectrometer. *See also* Ion trap
 - an inductively coupled plasma-time-of-flight, for elemental analysis, part I, optimization and characteristics, 5:1008
 - BEQ hybrid, determination of high-energy fragmentation of protonated peptides using, 4:470
 - triple quadrupole, collisional activation in, alkenamide fragmentation y, 15:1462
- Mass spectrometer data systems, networking, for improved productivity and electronic archiving of data, 4:742

- Mass spectrometer systems, small, for gas analysis, 13:1005
- Mass spectrometers
- development of a palm portable, 19:1442
 - field-portable
 - environmental and forensic applications of, 12:683
 - high speed GC/TOFMS, 12:648
 - ICP sector-field, simultaneous or scanning data acquisition for, 14:227
 - ion trapping, a pulsed-leak valve for use with, 7:118
 - magnetic sector, an electrospray ion source for, 2:205
 - miniature
 - EI/Q/oa TOF, 12:641
 - ExB sector-field, 12:619
 - from 12th Sanibel conference, 12:617
 - trade-offs in, 12:633
 - multiquadrupole, calibration point for electron ionization MS/MS spectra measured with, 5:576
 - triple quadrupole, bipolymer sequencing in, 12:846
- Mass spectrometric 3-D analysis, building a database of bifunctional probes for investigation of nucleic acids, 17:1570
- Mass spectrometric analysis
- direct matrix-assisted laser desorption ionization, of proteins immobilized on nylon-based membranes, 5:230
 - of four regioisomers of F₂-isoprostanes formed by free radical oxidation of arachidonic acid, 7:490
 - ion trap, tenax collection with jet-separator enrichment and, air analysis using, 7:1172
 - of leukotriene A₄ and other chemically reactive metabolites of arachidonic acid, 13:1227
 - of nitrogen- and phosphorus-containing pesticides by liquid chromatography-mass spectrometry, 5:655
 - of organic compounds, glow discharge ion source for, 5:305
- Mass spectrometric and ab initio study of the pathways for dehydration of simple glycine and cysteine-containing peptide [M+H]⁺ ions, 9:945
- Mass spectrometric and quantum mechanical analysis of gas phase formation, structure, and decomposition of various b₂ ions and their specifically deuterated analogs, 9:1002
- Mass spectrometric and tandem mass spectrometric behavior of nitrocatechol glucuronides: comparison of APCI and ESI, 10:537
- Mass spectrometric characterization of a series of adenosylated peptides acting as bisubstrate analogs of protein kinases, 5:443
- Mass spectrometric detection, Fourier transform ion cyclotron resonance, a glow discharge ion source with, 3:122
- Mass spectrometric detection of LTC₄, LTD₄, and PTE₄, enhancement of, by derivitization, 5:292
- Mass spectrometric evaluation, the quadrupole-based, of isotope ratios of carbon dioxide in expired air from mice and men following the administration of ¹³C-methyl methionine, 3:742
- Mass spectrometric evidence for mechanisms of fragmentation of charge-derivatized peptides, 12:399
- Mass spectrometric fragmentation, electron capture negative chemical ionization, and calculated internal energies for polychlorinated biphenyls, correlation between, 5:44
- Mass spectrometric-gas chromatographic method for the assessment of oxidative damage to double-stranded DNA by quantification of thymine glycol residues, 4:336
- Mass spectrometric information, from known amino acid sequences, computer program (COMPOST) for, 2:174
- Mass spectrometric investigation, a tandem, of (C₂H₅)X⁺ ions (X=Cl, Br, I), A, the involvement of classical and nonclassical ethyl structures, 5:814
- Mass spectrometric measurement of changes in protein hydrogen exchange rates that result from point mutations, 7:515
- Mass spectrometric methods for distinguishing structural isomers of glutathione conjugates of estrone and estradiol, 9:612
- Mass spectrometric screening and characterization, tandem, hybrid, of the glutathione conjugate of 2-furamide, multiple scan modes in, 2:55
- Mass spectrometric strategy for characterization of lipooligosaccharides from *Neisseria gonorrhoeae* using FTICR, 13:571
- Mass spectrometric structural characterization, negative ion electrospray tandem, of leukotriene B₄ (LTB₄) and LTB₄-derived metabolites, 7:129
- Mass spectrometric studies of diastereomeric cobalt complexes, chiral recognition in the gas phase, 5:452
- Mass spectrometric technique, K⁺IDS, K⁺ ionization of desorbed species, characterization of the primary thermal degradation processes of peptide using, 5:554
- Mass spectrometrically measured molecular weight, determination of oligonucleotide composition from, 4:204
- Mass spectrometry. *See also* Liquid chromatography-mass spectrometry
- /electron ionization gas chromatography, analysis of hydroxy fatty acids as pentafluorobenzyl ester, trimethylsilyl ether derivatives by, 6:40
 - in absolute configuration assignment of isomers, 16:431
 - accurate base composition of double-strand DNA by, 7:1266
 - in ACE inhibitor fragmentation, 15:300
 - of advanced glycation end products/peptides, 15:496
 - analysis of an Egyptian mummy resin by, 3:582
 - analysis of peptides, optimization of capillary zone electrophoresis/electrospray ionization parameters for, 3:289
 - in analysis of the editosome and other multiprotein complexes in *Trypanosoma brucei*, 14:728
 - of analytes from aqueous solutions, heated electrospray source for, 5:791
 - analytical, principles of collisional activation in, 3:599
 - analytical chemistry and, 6:993
 - angle-resolved reionization-neutralization, 2:492
 - atmospheric pressure ionization
 - chemicaldelete, results obtained by, internal standard correction of, 2:427
 - detection mass bias in, 2:232
 - automated high resolution, for synthetic chemistry, 10:546
 - automatic gain control in, using electrode, 16:244
 - of binding of aldose reductase inhibitors, 10:635
 - of bovine serum albumin three-dimensional structure, 15:1237
 - capillary isoelectric focusing with, on-line desalting in, 8:743
 - of CD4/lc-copurifying proteins, 15:558
 - ²⁵²Cf plasma desorption
 - adsorption/desorption in, substrates for, small molecules as, 2:29
 - background reduction in, a pulsed ion deflection system for, 3:706
 - fragmentation analysis of bradykinin by, 2:379
 - in characterization of transferrins and their fragments derived by reduction of disulfide bonds, 14:635
 - chemical ionization
 - atmospheric-pressure, with liquid introduction via heated nebulizer interface, characterization of flame-generated C₁₀ to C₁₆₀ polycyclic aromatic hydrocarbons by, 7:276
 - utilizing an isotopically labeled reagent gas, 5:596
 - cluster chemical ionization and deuterium exchange, in supersonic molecular beams, 7:550
 - collision-induced dissociation
 - high-energy, of synthetic mannose-6-phosphate oligosaccharides, 7:182

- of nonionic surfactants following direct supercritical fluid injection, 3:150
- combined with gas chromatography, structural determination of long-chain aldehydes by, pyridine-containing Schiff base derivatives for, 2:245
- continuous-fast atom bombardment, of oligonucleotides, 2:157
- in depth profiling of cubic silver halide microcrystals, 10:1016
- in detection of affinity purified cross-linked peptides, 15:832
- in determination of association constants of adenylate kinase with two noncovalent inhibitors, 14:442
- of deuterium labeling regiospecificity, 15:1366
- in diagnosis of disease, 10:561
- for diastereomeric *N*-acetylhexosamine quantification, 11:1086
- in differentiation of diexo- and diendo-fused isomers of norbornane/ene-condensed thiouracil and thiazinopyrimidine derivatives, 12:1011
- of the effects of hydrophobic surface chemistry and morphology on the digestion of surface-bound proteins, 14:203
- electrochemistry combined with. (*See* Electrochemical mass spectrometry)
- electron capture, oxygen addition-induced dechlorination of PCBs and DDT metabolites in, regioselectivity of, 7:66
- electron capture negative ion chemical ionization, amino acid/alcohol detection via, pentafluorobenzyl chloro-formate derivatization for, enhancement of, 6:525
- electron capture-tandem, sub-parts-per-billion determination of nitro-substituted polynuclear aromatic hydrocarbons in airborne particulate matter and soil by, 7:1255
- electron impact, of alkanes in supersonic molecular beams, 6:120
- electron impact ionization, intramolecular cyclization in 2-substituted pyrimidin-4(3H)-ones, 5:113
- electron-induced, of 1,2-dihydropyridine derivatives, 8:1255
- electrospray
- for the analysis of opioid peptides and for the quantification of endogenous methionine enkephalin and b-endorphin, 2:149
 - application of, in probing protein-protein and protein-ligand noncovalent interactions, 4:624
 - borane salts, electrospray needle as an electrochemical cell, 6:860
 - evidence from, stepwise refolding of acid-denatured myoglobin, 4:638
 - identification of components in waste streams by, 4:604
 - and matrix-assisted laser desorption ionization time-of-flight, application of, for molecular weight assignment of peptides in complex mixtures, 4:670
 - of methanol and water solutions suppression of electric discharge with SF₆ gas, 2:497
 - nanoliter, long-lived metallized tips for, 3:281
 - negative ion
 - analyte charge state, signal intensity, and stability in, solvent effect on, implications for the mechanism of negative ion formation, 4:546
 - of nucleotides, ionization from water solution with SF₆ discharge suppression, 4:289
 - negative ion formation in, 4:578
 - reversed-phase microbore liquid chromatography, complementary mobile phases, and sequential exoglycosidase digestion, characterization of protein N-glycosylation by, 5:350
 - stereoselectivity and ion response in, correlation of, 6:207
 - study of some aqueous solutions of metal salts, 7:1270
- electrospray ionization
- for analysis of low-molecular weight anticancer drugs and their analogues, 4:588
 - applications of, to neutral organic molecules including fullerenes, 4:596
 - of biotin binding to streptavidin, 6:912
 - charge discrimination with, applying to protein analyses, 6:1098
 - of copper adducts of protonated ubiquitin, 6:521
 - deconvolution by an entropy-based algorithm, 3:207
 - gas phase hydrogen/deuterium exchange in, as a practical tool for structure elucidation, 5:434
 - high mass-to-charge ratio ions from, observations and implications of, 4:536
 - improved signal intensity of oligonucleotides in, charge-state reduction with, 7:697
 - interaction of angiotensin peptides and zinc metal ions by, 5:959
 - low mass poly(methyl methacrylate) polymers and, 6:906
 - nanoscale separations combined with, sulfonamide determination, 3:139
 - noncovalent complexes formed between duplex DNA and minor groove-binding molecules by, characterization of, 6:1154
 - observation of noncovalent complexes to the avidin tetramer by, 5:201
 - of oligonucleotides, specific and nonspecific, dimer formation in, 6:159
 - of phosphopeptides isolated by on-line immobilized metal ion affinity chromatography, 4:662
 - probing the helical content of growth hormone-releasing factor analogs using, 4:646
 - study of noncovalent enzyme-inhibitor complexes and metal, 6:1105
 - binding stoichiometry of matrolysin by, 6:1105
 - a study of the thermal denaturation of ribonuclease S by, 5:614
- electrospray ionization/capillary zone electrophoresis, application of, to the analysis of macrolide antibiotics, 3:563
- electrospray ionization/gradient reverse-phase high-performance liquid chromatography analysis with trifluoroacetic and other strong acid modifiers by postcolumn addition of propionic acid and isopropanol, signal enhancement of, 6:1221
- electrospray ionization quadrupole, molecular mass measurement of intact ribonucleic acids via, 6:27
- electrospray ionization tandem, of polyamine toxins of spiders, neighboring group participation in. Part 1, alpha, omega-diaminoalkane compound in, 6:634
- enhanced sample loading by, 8:15
- ESI. (*See* Electrospray ionization-mass spectrometry)
- fast atom bombardment
- continuous-flow, an integral probe for, 3:198
 - continuous flow, -capillary zone electrophoresis, identification of deoxynucleoside-polyaromatic adducts by, 3:757
 - detection limits for, improved, a study of time-dependent desorption using a model pulsed bombardment ionization method, 3:637
 - of mycobacterial glycopeptidolipid antigens, structural characterization by charge remote fragmentation, 3:656
- fast-atom bombardment (*See* Fast-atom bombardment-mass spectrometry)
- differentiation of isomeric C8- and N²-deoxyguanosine adducts of 2-acetylaminofluorene by, 5:58
 - high-resolution, positive and negative ion, reference compound for, ultramark 1621 as, 3:842
 - injection of reagent ions into the selvedge region in, 4:259
 - metal chelation of dinucleotide analogs in the gas phase by, 7:42

- negative ion, analysis of long-chain fatty acyl coenzyme A thioesters by, 5:416
- probing high order structure of proteins by, 5:19
- redox processes by ruthenium(II) polypyridine complexes induced by, 4:306
- study of the solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes, generation of molecular signals, 4:249
- fast atom bombardment-collision activated dissociation-B/E linked-scan, of amino ethyl-triphenylphosphonium derivatives of fatty acids, charge-remote fragmentation during, 3:769
- fast-atom bombardment/liquid secondary ion, reduction/oxidation processes observed in, true effect of experimental parameters on, evaluation of, 4:482
- fast-atom bombardment of enzymatic digests, recombinant human macrophage colony stimulating factor, disulfide linkage and disulfide bond scrambling in, characterization of, 6:638
- fast-atom bombardment tandem, determination of the anomeric configuration of glycosyl esters of nucleoside pyrophosphates by, 6:516
- field desorption, for the analysis of polyethylene, evaluation of, 7:1070
- field-portable miniature, 11:468
- fingerprinting proteins coupled with polymers by, investigation of polyethylene glycol-conjugated superoxide dismutase, 6:478
- focus on harsh environments, editorial, 19:1375
- Fourier transform
- analysis of hydrocarbon dendrimers by, 5:731
 - external ionization, a low voltage ion transport system for, 5:398
 - high mass detection in, ion cloud manipulation using the radiofrequency-only mode as an improvement for, 4:186
 - high pressure trapping in, a radio frequency-only-mode event, 3:590
 - ion excitation for, linear response theory of, 2:483
 - ionization, ion cyclotron resonance, reagent ions in, potassium halide ions as, 3:301
 - laser desorption
 - analysis of hydrocarbon polymers by, silver nitrate chemical ionization for, 4:453
 - matrix-assisted, structural characterization of normal and modified oligonucleotides by, 2:402
 - poor high mass performance in, a mechanism for, 3:460
- Fourier transform-ion cyclotron glow discharge, external ion injection in, ion guide for, an improved, 4:387
- Fourier transform-ion cyclotron resonance
- capillary electrophoresis, with sustained off-resonance irradiation for the characterization of protein and peptide mixtures, 5:894
 - excitation modes for, 4:433
 - high-frequency, 4:177
 - hyperbolic ion trap for, experimental evaluation of, 3:188
 - improved, of large biomolecules, 4:190
 - probing trapped ion energies via ion-molecule reaction kinetics, 3:727
 - quadrupolar excitation in, simplified application of, 6:448
- and fullerenes, 3:268
- gas chromatography
- high resolution, application to the determination of trace monobromopolychlorodibenzo-p-dioxins in environmental samples, 3:248
 - selective ion monitoring mode in, direct chemical analysis of UV laser ablation products of organic polymers by using, 5:106
- glow discharge
- analysis of solution residues by, 4:47
 - theoretical calculations of relative ion yields for, 3:79
- glow-discharge-Fourier transform ion cyclotron, external ion injection in, ion guide for, 4:387
- of helical unfolding in peptides, 5:425
- heterogeneity of the monomer subunit of *Lumbricus terrestris* hemoglobin, 8:352
- high-performance liquid chromatography atmospheric pressure chemical ionization, quantitative analysis by, determination of the renin inhibitor CP-80,794 in human serum, 2:164
- high resolution, 10:661
- high-resolution, of large molecules in a linear time-of-flight mass spectrometer, 4:2
- high throughput open access, automation and, 10:1174
- of histone modifications in leukemia, 15:77
- hydrocarbon, in situ silver cationization for, 16:437
- of hydrogen/deuterium exchange of proteins
- continuous *versus* pulsed labeling, 10:675
 - in solution, 10:671, 10:672
- hydrogen exchange and, historical perspective, 17:1481
- of hydroxyl radical adducts, 12:1214
- improving sensitivity using high-pressure electrodynamic ion funnel interface, 17:1299
- inductively coupled plasma, 8:1230
- inductively coupled plasma (ICP-MS), 12:1226
- inorganic, 11:822
- interference in, in polypropylene tubes, 16:417, 16:802 (erratum)
- internal glow discharge-Fourier transform ion cyclotron resonance, 7:923
- ion sources, space-charge-dominated, modeling and sensitivity, 2:1
- ion spray, studies on heme binding in myoglobin, hemoglobin, and cytochrome c by, 4:631
- ion spray-tandem, of supramolecular coordination complexes, 5:748
- ion trap
- quadrupole
 - probing trapped ion energies via ion-molecule reaction kinetics, 3:716
 - resonance ejection mode of, mass resolution in, extended theoretical considerations for, 5:676
 - selective chemical ionization of nitrogen and sulfur heterocycles in petroleum fractions by, 4:322
- ion trap/gas chromatography, alternating RF/DC isolations for quantitation with coeluting internal standards in, 3:85
- ionization
- aerosol matrix-assisted laser desorption, 5:1
 - matrix-assisted desorption laser and electrospray, characterization of mouse switch variant antibodies by, 7:707
 - ionization electrospray, gas-phase molecular dications formed from neutral organics in solution via chemical electron-transfer reactions by using, observations of, 5:689
 - ionization Fourier transform ion cyclotron resonance, in-cell matrix-assisted laser desorption, 7:1026
 - ionization tandem reflectron time-of-flight/matrix-assisted laser desorption of fullerenes, 7:590
 - ionspray, high accuracy molecular weight determination and variation characterization of proteins up to 80 ku by, 2:387
 - of iron chelation, 8:1070
 - laser desorption
 - ion trap, of self-assembled monolayers, 4:968
 - negative-ion matrix-assisted, analysis of oligodeoxynucleotides by, 4:955

- laser desorption-ionization, matrix-assisted, matrices used for, effects of heavy-atom substituents on, 5:800
- laser desorption-ionization Fourier transform ion cyclotron resonance, collision-induced dissociation in, improved efficiency of, selective parent ion axialization for, 5:900
- laser desorption time-of-flight, analysis of hydrocarbon dimers by, 5:731
- laser-induced thermal-desorption/Fourier transform, for surface analysis, deflection of laser-produced ions in, 5:583
- laser microprobe, of quaternary phosphonium salts, direct *versus* matrix-assisted laser desorption, 4:798
- LC-TSP and LC-ESI, spin-trapped radicals determination by, 2:413
- liquid chromatography
- for the analysis of N-methyl carbamate pesticides, an evaluation of atmospheric pressure ionization techniques for, 3:378
 - particle beam
 - mass spectra of chlorinated phenoxyacid herbicides obtained by, evidence for thermal decomposition contributions to, 3:823
 - mass transport and calibration in, 3:662
- liquid chromatography-electrospray ionization matrix-assisted laser desorption ionization time-of-flight, rapid analysis of antibiotic-containing mixtures from fermentation broths by using, 7:1227
- liquid flow-injection isotope ratio, direct analysis of carbon isotope variability in albumins by, 7:605
- liquid secondary ion
- alkali chloride cluster formation in, ion-ion recombination for, role of, 4:410
 - analysis of permethylated, n-hexylamine derivatized oligosaccharides, application to baculovirus expressed mouse interleukin-3, 3:345
 - characterization of the opioid peptide-containing protein and of bovine alpha-lactalbumin by, 5:377
- liquid secondary ion/fast-atom bombardment by tandem mass spectrometry, beam-induced reactions occurring in, 4:312
- liquid secondary ionization
- beam-induced reaction between metanitrobenzyl alcohol and dipyrrocyanine dyes in, 5:1017
 - direct desorption of crown ether-metal ion complexes in, 5:638
- of *Lumbricus terrestris* hemoglobin, 8:352
- matching algorithms and databases for, 9:92
- matrix-assisted laser desorption, of biotransformation products of dynorphin A in vitro, 5:10
- matrix-assisted laser desorption ionization
- with 2-(4-hydroxyphenylazo)benzoic acid matrix, 5:399
 - application to low mass poly(methylmethacrylate) polymers, fundamentals of, 7:287
 - charge state distribution shifting of protein ions observed in, 6:1183
 - for polymer characterization, evaluation of, 7:11
 - quantitative assessment of cysteine and cystine in peptides and proteins following organomercurial derivatization and analysis by, 5:359
 - time-of-flight, application of, for molecular weight assignment of peptides in complex mixtures, 4:670
- matrix-assisted laser desorption/ionization time-of-flight of peptides, improved mass accuracy in, 5:955
- methods in, fitness for purpose of, 14:528
- of methylated fullerenes, in-beam electron impact technique in, 16:1311
- micro-electrospray, ultra-high sensitivity analysis of peptides and proteins, 5:605
- in mid-1980s, chemist intrigue with, 12:975, 12:1228
- minimum thickness gate valve design for, 16:441
- modeling data from, in protein-ligand interactions, 15:388
- molecular secondary ion, studies of biological soft tissues, a model matrix for, doped gelatin films as, 4:419
- MS/MS spectra, electron ionization, calibration point for, measured with multiquadrupole mass spectrometers, 5:576
- multi imaging, of ¹³C-oleate transport in adipocytes, 15:1572
- multiple stage
- kinetic method combined with, SiCl₃⁺ and SiCl⁺ affinities for pyridines determined by using, agostic effects in the gas-phase, 7:198
 - in a pentaquadrupole, polar [4+2+] Diels-Alder cycloaddition to, gas stage ions and, 6:1
- multiple-stage, in structural characterization of organophosphorus compounds, 4:125
- multiple stage pentaquadrupole, for generation and characterization of gas-phase ionic species, case of the PyC₂H₅⁺ isomers, 7:1126
- negative ion fast-atom bombardment, of polyanionic analytes, enhancing sensitivity in, a class of cationic spectrometric additives for, N-alkylnicotinium halides, 5:935
- neutralization-reionization
- hydrogen bonding in transient bifunctional hypervalent radicals by, 6:1004
 - neutral counterpart generation by, structure of C₅H₅RFe⁺ (R=F, Cl, Br, I, O, OH, OCH₃, H₆H₅, H) ions in the gas phase, 6:1143
 - organic neutralization agents for, 3:108
- new products in, 8:573
- nitrogen trifluoride in, chemical reaction interface, for detection of phosphorus, deuterium, chlorine, and sulfur, 6:421
- noncovalent protein-ligand complexes, confirmation of binding sites and changes in tertiary structure, 14:460
- oligonucleotide sequencing by, SOS program in, 13:200
- on-line laser, of airborne particles, 10:648
- of *O,O*-dialkyl *N,N*-dialkyl phosphoramides under electron impact conditions, 15:547
- open access atmospheric pressure chemical ionization, for routine sample analysis, 6:387
- particle beam-high-performance liquid chromatography, determination of polycyclic aromatic hydrocarbons by, 4:898
- of peptides, 8:296
- photodissociation-photoionization, unimolecular photochemistry of n-alkenes studied by, 4:65
- plasma
- inductively coupled
 - with an electrically floating sampling interface, 4:733
 - with an enlarged sampling orifice and offset ion lens, ion trajectories and detector performance, 4:16
 - with an enlarged sampling orifice and offset ion lens, polyatomic ion interferences and matrix effects, 4:28
 - interference in, selection of solvent load and first-stage pressure to reduce, 3:128
- mixed-gas helium-argon inductively coupled matrix-induced interferences in, 2:355
- space-charge effects in, time-resolved measurements of individual ion cloud signals to investigate, 7:362
- plasma source ion trap, enhanced abundance sensitivity by resonant ejection of atomic ions, 7:1161
- of platelet-activating factor after isolation by solid phase extraction and direct derivatization with pentafluorobenzoic anhydride, 11:176
- of polymers, 2004 ASMS Fall workshop on, review of, 16:291
- process, in simultaneous analysis of butene isomer mixtures, 11:1079

- Professor Al Nier and his influence on, 3:685
- profiling of whole bacteria by, electron and chemical ionization modes in, 10:747
- of protein
- improved sample processing time and peptide recovery for, 15:784
 - inherent electrochemistry of electrospray for, mass tag generation by, 15:1767
- in protein and proteome research, 14:685
- of protein complex quaternary structure, 12:222
- protein identification by, 12:215
- protein structure information from, selective titration of arginine residues by sulfonates in, 12:810
- putting in hands of end user, 6:394
- putting in the hands of the end user, 6:394
- quadrupole (*See* Quadrupole mass spectrometry)
- quadrupole ion trap, negative chemical ionization in, effects of applied voltages and reaction times, 5:757
- quantitative proteomic analysis by, 12:1238
- with RPLC and CE, of peptides, 8:495
- of some protonated and lithiated 2,5-disubstituted 1,3,4-oxadiazoles, 14:289
- static secondary
- detection of cyclohexamine on soil surfaces exposed to laboratory air, 7:168
 - limit of, organic ion imaging beyond, 6:1047
- structural tool, use of guided ion beam MS/MS to determine thermochemistry, 13:419
- study of the solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes, synthesis and characterization of molecular complexes, 4:255
- tandem
- of acetone and acetone/water cluster ions, 2:113
 - analysis of long-chain fatty acyl coenzyme A thioesters by, 5:416
 - analysis of peptides, optimization of capillary zone electrophoresis/electrospray ionization parameters for, 3:289
 - in characterization of human oxidized apoprotein B-100, 6:242
 - charge inversion, isomer differentiation by, an investigation into the structure of the ionic products from an Sn(AN-RORC) reaction, 3:467
 - collision-induced dissociation, high- and low-energy, in the analysis of glycoalkaloids and their aglycons, comparison of, 7:173
 - curie-point pyrolysis, an investigation of dipeptides containing polar and nonpolar side groups by, 3:750
 - desorption chemical ionization, of polyprenyl and dolichyl phosphates, 7:958
 - of diastereomeric cobalt-glucosyl-glucose disaccharide complexes, differentiation of stereochemistry of glycosidic bond configuration, 7:953
 - differentiation of isomeric C8- and N²-deoxyguanosine adducts of 2-acetylaminofluorene by, 5:58
 - of doubly protonated angiotensin II formed by electrospray ionization, charge-separation reactions in, experimental considerations and structural implications, 7:30
 - electron ionization, of glycosphingolipids, identification of compound-specific sequence ions in the collision-induced dissociation spectra of the immonium ions of two isomeric hexaglycosylceramides, 3:353
 - electrospray, structure analysis of, phenylglycidyl ether adducts of 29-deoxycytidine and 29-deoxyadenosine, 7:682
 - electrospray ionization analysis of peptides derived by enzymatic digestion of oxidized globin subunits, improved method to determine amino acid substitution in the hemoglobin "core," 7:1040
 - electrospray liquid chromatography, quantitation of SR 27417 in human plasma using, a study of ion suppression, 7:1099
- FAB-CAD-MS/MS, characterization of diacylglycerolphosphocholine molecular species by, a general method not sensitive to the nature of the fatty acyl groups, 3:71
- fast atom bombardment
- employing ion-molecule reactions for the differentiation of phospholipid classes, 2:470
 - isomer differentiation in 7,12-dimethylbenz[a]anthracenepyridine adducts by, 2:256
- fast-atom bombardment
- analysis of phosphatidylcholine isolated from heart mitochondrial fractions by, evidence of incorporation of monohydroxylated fatty acyl moieties, 7:50
 - of [¹³C]arachidonic acid labeled phospholipid molecular species, 5:144
 - determination of the anomeric configuration of glycosyl esters of nucleoside pyrophosphates and polyisoprenyl phosphates by, 7:541
- and fast atom bombardment, of macrolide antibiotics, 5:151
- fast-atom bombardment by/liquid secondary ion mass spectrometry, beam-induced reactions occurring in, characterization of, 4:312
- gas-phase labeling combined with, ion-dipole complex formation from deprotonated phenol fatty acid esters evidenced by, 4:343
- high-energy collision-induced dissociation in, low-mass ions produced from peptides by, 4:882
- high-resolution, of large molecules, Fourier-transform electrospray instrumentation for, 4:557
- identification of components in waste streams by, 4:604
- ion trap
- applied to small multiply charged oligonucleotides with a modified base, 5:740
 - quantitative analysis of crude plasma extracts by, 3:372
- linked-scan, product ion characterization by, trimethylsilyl group migration in the mass spectra of trimethylsilyl ethers of cholesterol oxidation products, 4:327
- of lithium-attachment ions from polyglycols, 3:225
- low- and high-energy collision-induced dissociation, use of, in the identification of an unusual amino acid in a semi-synthetic polypeptide, 7:1034
- low energy, and electrospray ionization of polyhydroxy unsaturated fatty acids, 7:140
- low-energy, of the molecular ion derived from fatty acid methyl esters, a novel method for analysis of branched-chain fatty acids, 4:223
- molecular species analysis of arachidonate containing glycerophosphocholines by, 2:45
- negative ion, of leukotriene E₄ and LTE₄ metabolites, identification of LTE₄ in human urine, 2:314
- negative ion fast-atom bombardment, to determine sulfate and linkage position in glycosaminoglycan-derived disaccharides, 3:797
- of poly(ethylene glycol) lithium-attachment ions, 5:1072
- protonated O, O-dimethyl O-aryl phosphorothionates in, fragmentation of, 6:488
- reactivity of collisionally activated dichlorocarbene dications studied by, 6:1019
- secondary ion, imaging large insulating samples by, charge compensation for, 5:37
- of small, multiply charged oligonucleotides, 3:60
- study of C-phenyl-N-tert-butyl nitron spin adducts from *in vitro* rat liver microsomal metabolism of bromotrachelo-methane and carbon, 6:847

- and theoretical calculations, mechanism of gas-phase aldose-ketose isomerization, 4:943
- thermal degradation, structural studies on alkylisocyanate polymers by, 2:130
- tandem-electrospray ionization, positive and negative mode, identification of phosphorylation sites in phosphopeptides by, 7:243
- tandem high-resolution, thiaminase I (42 kDa) heterogeneity and sequence refinement, active site location and, 6:981
- tandem ion trap, and selected ejection chemical ionization, application of nonresonance excitation to, 7:668
- tandem/liquid chromatography analysis of peptide mixtures, data-controlled automation of, 7:532
- tandem magnetic sector/reflectron time-of-flight, detection of femtomole and sub-femtomole levels of peptides by, 2:91
- tandem mass spectrometry of taxol, 3:672
- terms relating to, standard definitions of, report from the Committee on Measurements and Standards of the American Society for Mass Spectrometry, 2:336
- thermal ionization cavity source for, 10:1008
- thermospray, ion formation of N-methyl carbamate pesticides in, effects of additives to the liquid chromatographic eluent and of the vaporizer temperature, 5:913
- time-of-flight
- gridless ion acceleration systems for, 4:782
 - improved energy compensation for, 5:779
 - ion trap source, segmented ring, cylindrical, 7:1009
 - matrix-assisted laser desorption ionization, of underivatized and permethylated gangliosides, 3:785
 - of tirapazamine and its metabolites, insights into mechanism of metabolic transformations and characterization of reaction intermediates, 14:881
 - underwater, for chemical analysis of hydrosphere, 12:677
 - in U.S. space program, 12:656
- Mass spectrometry and good laboratory practices, 7:211
- Mass spectrometry and molecular orbital calculations on pyrrolo[1,2-a][1,4]benzodiazepin-4-one, -6-one, and -4,6-dione, characterization and differentiation of heterocyclic isomers, 7:653
- Mass spectrometry capabilities of a pentaquadrupole mass spectrometer, case study, ion-molecule reactions and collision-activated dissociation of C₄H₄⁺ isomers, 3:518
- Mass spectrometry/chemical reaction interface/gas chromatography, selective detection of sulfur-containing compounds by, 2:250
- Mass spectrometry/electrospray ionization
- characterization of large, heterogeneous proteins by, 7:1219
 - of nucleobases and nucleosides, processes that affect, 7:1106
- Mass spectrometry/electrospray liquid chromatography, selective detection of phosphopeptides in complex mixtures by, 4:710
- Mass spectrometry/fast-atom bombardment, multisample probe for, 4:504
- Mass spectrometry/gas chromatography
- in acute poisoning, scope and limitations of in general unknown screening, 6:995
 - early, 4:367
 - overlapping, spectral deconvolution for, 3:558
 - with a polarizable stationary phase, negative ion chemical ionization and mass-resolved chromatography, high-resolution triglycerol mixture analysis using, 7:350
 - purge-and-trap, measuring volatile organic compounds in human blood by, interferences in, importance of enhanced mass resolution in, 3:831
- Mass spectrometry/gas chromatography/and particle beam liquid chromatography, poly(amic acid) and polyimide characterization using, 5:299
- Mass spectrometry interface
- atmospheric pressure chemical ionization liquid chromatography, optimization of, 7:69
 - particle beam liquid chromatography, performance of, effect of a dc glow discharge on, investigation of, 5:908
- Mass spectrometry/isotope dilution gas chromatography, for platinum determination in urine, 2:85
- Mass spectrometry/isotope-ratio-monitoring gas chromatography/, nitrogen isotope analyses by, 5:387
- Mass spectrometry/liquid chromatography
- with collision induced dissociation, the determination of glycopeptides by, 3:804
 - mass spectrometric analysis of nitrogen- and phosphorus-containing pesticides by, 5:655
 - normal phase, by using coaxial continuous flow fast atom bombardment, 3:575
- Mass spectrometry liquid secondary ionization, beam-induced dehalogenation of organic compounds, secondary electron capture in, evidence for a mechanism that involves, 7:1109
- Mass spectrometry methods, combined, for the characterization of a new variant of human hemoglobin, the double mutant hemoglobin Villeparisis β77(EF1) His → Tyr, β80 (EF₄) Asn → Ser, 7:163
- Mass spectrometry/on-line isotopic dilution cryofocusing gas chromatography, automated high-speed analysis of selected organic compounds in urban air by, 5:1001
- Mass spectrometry on-line monitoring and MS₂ product characterization of TiO₂/UV photocatalytic degradation of chlorinated volatile organic compounds, 9:1321
- Mass spectrometry-particle beam liquid chromatography, with massive cluster impact, 7:293
- Mass spectrometry PittCon® 2006, review, 17:873
- Mass Spectrometry PittCon® 2009, review, R20:3 (June)
- Mass spectrometry PittCon'99 review, 10:457
- Mass Spectrometry School, The Massachusetts Institute of Technology, 5:332
- Mass spectrometry study, tandem, of the gas phase RSH₆H₅Cr⁺L (R=CCH₅; L⁺arene) and H₆H₅SSH₆H₅Cr⁺ ions, 5:928
- Mass spectrometry technique, chemical reaction interface, implementation on a Hewlett-Packard mass-selective detector, 5:765
- Mass spectroscopy
- Fourier transform ion cyclotron resonance, remeasurement at high resolving power, 6:564
 - high energy collision-induced dissociation tandem, charged derivatives, 6:428
 - IgG1 mAb photo-induced methionine oxidation, 20:525
 - low energy tandem, of deoxynucleoside adducts of polycyclic aromatic hydrocarbon dihydrodiol-epoxides, 6:248
 - negative ion chemical ionization, anthraquinone-2-carbonyl chloride as an alcohol derivatization reagent for, 6:148
- Mass spectrum, identifying compound from, tests for, 14:262, 14:264
- Mass spectrum that today is rarely seen, 10:76
- Mass-tag technology, novel assay system using, tumors, 18:106
- Mass tags, by inherent electrochemistry of ESI, 15:1767
- Mass-to-charge-ratio ions, observation and implications of, from electrospray ionization mass spectrometry, 4:536
- Mass-to-charge ratio scale, 5:120
- Mass transport and calibration in liquid chromatography particle beam mass spectrometry, 3:662
- Mass transport calculation, electrospray emitter analyte electrolysis in, 11:951
- Mass uncertainty, in complex MS/MS spectra, 16:183

- Massachusetts Institute of Technology Mass Spectrometry School, 5:332
- Massive cluster impact, liquid chromatography particle beam-mass spectrometry with, 7:293
- Massive cluster impacts, shock wave model for sputtering biomolecules using, 3:311
- Mastoparans, dominulin A and B in cuticle and venom of social paper wasp analyzed by MALDI-TOF, MALDI-TOF/TOF and ESI-ion trap, 17:376
- Material-enhanced laser desorption/ionization time-of-flight mass spectrometry, protein profiling tool with specific carrier materials, 17:1203
- Mathematical study of sample modulation at a membrane inlet mass spectrometer-potential application in analysis of mixtures, A, 7:93
- MATLAB, artificial neural networks for ppb mass accuracy in FT-ICR/MS, 20:1303
- Matrices used for matrix-assisted laser desorption-ionization mass spectrometry, effects of heavy-atom substituents on, 5:800
- Matrilysin, metal binding stoichiometry of by electrospray ionization mass spectrometry, noncovalent enzyme-inhibitor complexes and, 6:1105
- Matrix, nonpolar, in analysis of nonpolar synthetic polymers, MALDI-TOFMS in, 11:731
- Matrix additives, class of cationic, for enhancing the sensitivity in negative ion fast-atom bombardment mass spectrometry of polyanionic analytes, N-alkylnicotinium halides, 5:935
- Matrix amount, dependence on, in MALDI-FTMS, 16:1772
- Matrix application, sublimation method of, MS imaging, 18:1646
- Matrix-assisted filament desorption/ionization mass spectrometry, 9:1226
- Matrix-assisted laser desorption
 - direct *versus*, laser microprobe mass spectrometry of quaternary phosphonium salts, 4:798
 - Fourier transform mass spectrometry, structural characterization of normal and modified oligonucleotides by, 2:402
 - infrared, and ionization by using a tunable mid-infrared free-electron laser, 7:1187
 - neutrals generated by, gas-phase cationization and protonation of, 4:393
 - observed fast fragmentation of peptides in, factors that influence, 7:225
- Matrix-assisted laser desorption ionization, of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, high resolution end group determination of low molecular weight polymers by, 7:449
- Matrix-assisted laser desorption/ionization
 - 193 nm photodissociation characteristics of singly protonated peptides generated by, 17:1643
 - acid hydrolysis of proteins, 20:2106
 - with FTMS aerosol
 - of biomolecules, matrix solutions for, 9:166
 - with collision-induced dissociation, of poly(alkylmethacrylate), 8:1206
 - in detection of pharmaceutical compounds in tissue, 10:1315
 - differentiation of isomeric photomodified oligodeoxynucleotides by fragmentation of ions produced by, 10:329
 - of DNA secondary structure, 10:521
 - effect of poly(ethylene terephthalate) oligomers cationized by alkali ions on, 10:329
 - with FTICR, and capacitive-coupling gated trapping, 8:195
 - with FTMS
 - of hydrocarbon polymers, 8:225
 - of ionized ethylated acetone and deuterated analogs, 8:327
 - initial velocity of ions generated by, TOS-MS effects of, 8:209
 - laser-induced azido photochemistry in, fingerprint patterns from, 10:1337
 - measuring flight time and energy of, 8:1094
 - mercaptobenzothiazoles in, 8:116
 - near-UV-induced, as function of wavelength, 9:885
 - of peptide conformational changes, 10:719
 - of peptides and proteins, 10:91
 - of polymers, 8:225
 - time lag focusing and cation attachment in, 8:132
 - protein modification by, 8:8
 - proton transfer reactions of, 10:1111
 - quaternary protein ensemble analysis by, 8:1046
 - sample preparation by, TOFMS of, 10:104
 - of synthetic polymers, 8:76
 - with TOF-MS, for polydiene analysis, 8:1220
 - ultraviolet
 - of DNA fragment ions, 10:423
 - of oligonucleotides, dihydroxybenzoic acid in, 8:218
 - vitamin B12 fragmentation in, 8:140
 - ions produced by, inspected with Orbitrap LTQ, 20:1451
 - irradiation effects in, 2,5-dihydroxybenzoic acid monocrystals in, 14:893
 - for ruthenium and osmium trisbipyridine complexes, FTMS evaluation of, 14:393
 - solid state NMR exploring solvent-free samples, 20:1906
 - time-resolved photodissociation of peptides by, 20:1522
- Matrix-assisted laser desorption/ionization-collision-induced dissociation, of poly(styrene), 9:269
- Matrix-assisted laser desorption ionization employing a tandem double focusing magnetic-orthogonal acceleration time-of-flight mass spectrometer, peptide sequence determination by, 7:1
- Matrix-assisted laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry, in-cell, 7:1026
- Matrix-assisted laser desorption/ionization Fourier transform mass spectrometry (MALDI-FTMS)
 - in characterization of oligosaccharides labeled with 9-aminofluorene, 12:1254
 - glycosidase and, 12:877
 - herbicide rearrangements in, 15:1820
 - lipid and phospholipid chemistry elucidation by, strategy and data analysis techniques for, 15:1665
- Matrix-assisted laser desorption/ionization Fourier transform ion cyclotron resonance mass spectrometry, peptide imaging in rat brain, 18:145
- Matrix assisted laser desorption/ionization Fourier transform ion cyclotron resonance mass spectrometry, HPLC, shotgun analysis of proteome by, 17:205
- Matrix-assisted laser desorption ionization Fourier transform mass spectrometry
 - acetylcholinesterase inhibitor screening by, 19:1850
 - improved mass accuracy for peptides using SWIFT excitation for, 19:76
- Matrix-assisted laser desorption/ionization Fourier transform mass spectrometry, noncovalent DNA complexes with basic polypeptides or polyamines, 18:1977
- Matrix-assisted laser desorption/ionization-Fourier transform mass spectrometry
 - of iron-oligonucleotide complex, 10:941
 - of LHRH-metal ion complexes, 10:19
- Matrix-assisted laser desorption ionization-Fourier transform mass spectrometry (MALDI-FTMS), high pressure, ganglioside analysis by, 13:402
- Matrix-assisted laser desorption/ionization-FTICR-MS, in peptide mass fingerprint analysis, 14:553

- Matrix-assisted laser desorption/ionization-ion trap MS, cationized phosphotyrosine fragmentation by, 14:581
- Matrix-assisted laser desorption/ionization-ion trap/reflectron time-of-flight mass spectrometry, multistage, LC-ESI-IT-MS and, moenomycin characterization by, 17:1081
- Matrix-assisted laser desorption ionization (MALDI)
 of alkali cation attachment to derivatized fullerene, 13:1448
 alkali metal ion effects on polyether cationization in, 11:639
 atmospheric pressure. (*See* Atmospheric pressure matrix-assisted laser desorption/ionization)
 of bimetallic silver-gold clusters, 15:1455
 biopolymer ions generated by, photodissociation of, tandem TOF MS for, 15:1248
 bioreactive probes in, transthyretin variant detection using, 11:172
 detection limits of, 12:1055
 detection of femtomole and sub-femtomole levels of peptides by, 2:91
 for FTMS to accommodate large targets, 15:128
 ion mobility, of non-covalent complexes, 13:167
 ion pairs and anion adduct generation during, 13:1218
 matrix design for, sensitive determination of PAH-DNA adducts, 5:1021
 oligonucleotide fragmentation in, modified nucleic acids in, 12:744
 for peptide mass fingerprinting, scoring methods in, 13:22
 peptide-peptide interaction study by, 12:88
 photomodified oligodeoxynucleotide determination by, 12:1127
 in polyaniline oligomers analysis, 15:893
 protein identification by, 12:215
 of protonated cross-linked peptides, 13:709
 rapid switchability with ESI quadrupole TOFMS, for protein identification, 11:493
 ultraviolet, of oligodeoxynucleotide metastable decay, 12:180
- Matrix-assisted laser desorption/ionization (MALDI) Fourier transform (FT) mass spectrometry, matrix amount in, dependence on, 16:1772
- Matrix-assisted laser desorption/ionization (MALDI) Fourier transform ion cyclotron resonance (FTICR) mass spectrometry
 of hydrocarbon polymers, 16:2009
 of ionic liquid matrix-induced stabilization of phospholipids and metastable decay of peptides and oligonucleotides, 16:2000
- Matrix-assisted laser desorption/ionization (MALDI) ion source, quadrupole TOF mass spectrometer fitted with, glycosphingolipid analysis with, 12:1220
- Matrix-assisted laser desorption/ionization (MALDI) mass spectrometry
 aminophenylboronic acid immobilization on magnetic beads by, 16:1456
 in analysis of photolabile peptides, 15:1156
 AP and vacuum, on QqLIT instrument, 16:176
 carbon nanotubes as adsorbent of solid phase extraction in, 16:263
 carbon nanotubes as affinity tubes for peptides and proteins in, 15:1629
 of derivatized oligosaccharides, 16:683
 extending solvent-free sample preparation method in, 16:91
 of features smaller than laser beam, 16:1654
 isobutylene/ μ -methylstyrene copolymer compositional analysis by, 15:621
 on-line single droplet deposition for, 15:1471
 peptide ion detection by, esterification and, 16:1269
 polymer, sample deposition for, 15:168
 of proteins in brain structures, 16:1093
 sample preparation for, sodium dodecyl sulfate micelles in, 16:1438
 simple organic molecule cationization in, 15:879
 solvent-free, of biological samples via mini-ball mill approach, 16:542
 of surfactant-treated bacterial cell proteins, 16:1422
- Matrix-assisted laser desorption/ionization (MALDI) matrices
 liquid based UV
 detection limits of proteins from, 15:486
 proton affinity values of, under ESI conditions with kinetic method, 15:431
- Matrix-assisted laser desorption/ionization (MALDI) tandem mass spectrometry (MS/MS)
 of intact protein, 16:482
 microwave-assisted acid hydrolysis of proteins with, 16:471
 of phosphatidylcholines in brains, 16:2052
- Matrix-assisted laser desorption/ionization (MALDI) time-of-flight (TOF) mass spectrometry
 of α -cyano-4-hydroxycinnamic acid cocrystallization with peptides in levitated droplets, 16:254
 automated, of polyethylene glycol, 16:670
 bacterial analysis by, inter-laboratory comparison of, 16:456
 in brain tissue phospholipid analysis, 16:133
 in chemical cross-linking of protein complexes, 16:1921
 immobilized carbon nanotubes for, 16:333
 of intact proteins, 16:482
 oxidized carbon nanotubes for analysis by, 16:883
 phospholipid analysis by, ionic-liquid matrices for, 16:679
 polymer quantitation using, 16:100
 of salt-containing proteins, porous anodic alumina membrane as sample support for, 16:1488
 of urinary nucleosides, 16:940
- Matrix-assisted laser desorption ionization mass spectrometry with 2-(4-hydroxyphenylazo)benzoic acid matrix, 4:399
- aerosol, 5:1
 analysis using fluorescent dyes, 19:1054
 charge state distribution shifting of protein ions observed in, 6:1183
 construction of high precision ambient ionization source for, 19:1527
 direct analysis of proteins immobilized on nylon-based membranes, 5:230
 efficient ion generation in bioaerosol mass spectrometry, 19:315
 imaging solvent-free prepared samples, 19:874
 improved mass accuracy for peptides using SWIFT excitation for MALDI-FTICR MS, 19:76
 matrix dry-coating for phospholipid imaging, 19:882
 N-linked oligosaccharides with uncommon glycoform, 19:671
 ocular integral membrane protein tissue profiling by, 19:814
 for polymer characterization, 7:11
 protected synthetic peptide sample preparation for, 19:614
 quantitative assessment of cysteine and cystine in peptides and proteins following organomercurial derivatization and analysis by, 5:359
 randid skin secreted peptides, oxidation *versus* carboxamidomethylation of disulfide bonds, 19:479
 sensitivity enhancement for, proteins after treatment of tissue sections, 19:1069
 signal intensity and sensitivity improvement using nanoliter spotting by IBF, 19:1086
 UV, pressure probe and, plant single-cell carbohydrate extracts, 19:1842
 UV-, analysis of NBD-tagged glycosphingolipids, 19:923
- Matrix-assisted laser desorption-ionization mass spectrometry
 3-layer sample preparation nanomolar protein samples, 17:780

- amino acid ester fragmentation induced by triphenylcorrole-
(CuIII) complexes, 18:791
- analysis of cardiolipin from rat organ sections, 18:567
- analysis of disulfide-containing proteins by, 20:2284
- analysis of glycosylated globins, nephropathic patients, 18:
1018
- analysis of noncovalent protein complexes, 18:1286
- analysis of peptides and proteins bound to iron oxide nano-
particles, 18:910
- application to low mass poly(methylmethacrylate) polymers,
fundamentals of, 7:287
- of bacteria, internal control for, 10:1131
- ballpoint dye photofading on paper analyzed by LDI and, 17:
297
- calmodulin interactions with peptides investigated by, 20:576
- characterization of complexes between guanidinium deriva-
tives and ssDNA, 17:283
- crustacean hemolymph neuropeptides, measurement by, 20:
708
- DFT study of peptide ions generated by, 20:1327
- enhanced ionization efficiency at metal-metal interface, 17:737
- gas phase scrambling of disulfide bonds, 20:1603
- high speed quantitative, design considerations, 17:1129
- ILM for peptide, protein and carbohydrate analysis by, 20:
1790
- imaging distribution of β -peptides in whole body mice sec-
tions, 18:1921
- impulse-driven droplet deposition system interface combining
LC with MS/MS and, 17:325
- in-source decay, top-down determination of PEGylation sites
using, 20:326
- incoherent ion production reactions, 20:1078
- induction-based fluidics, real-time dispense event and volume
verification, 20:1064
- matrices used for, effects of heavy-atom substituents on, 5:800
- mini-ball mill and, β -amyloid peptide analysis via, 18:1533
- multisample preparation methods of synthetic polymers, 18:
377
- negative-ion, oligosaccharide analysis by, 18:714
- new copper containing matrix for, 20:1263
- peptide trace analysis using 2,5-dihydroxybenzoyl hydrazine
derivative CNTs, 17:1023
- poly(methacrylic acid) analysis by, 18:1507
- protonated deoxyoligonucleotides, sodium controlled reactiv-
ity in gas phase, 20:689
- ribosome topography analyzed by proteolysis and, 18:1304
- solution-phase zinc finger peptide complexation by, 10:27
- solvent-free, improvements in analysis of synthetic polymers
and giant organic molecules, 17:661
- sulfated glycan analysis by sequential double-permethylation,
20:1660
- SUPREX and accuracy of protein unfolding free energy deter-
mination, 17:1535
- ultraviolet, of MRP8 and MRP14 calcium-induced tetramers,
10:1124
- using pyrene methylguanidine, polysulfated oligosaccharides,
20:131
- UV, liquid matrix deposition on conductive hydrophobic sur-
faces, 18:693
- vortex method for preparing solvent-free samples, 20:1115
- Matrix-assisted laser desorption/ionization mass spectrometry
and electrospray ionization mass spectrometry, character-
ization of mouse switch variant antibodies by, 7:707
- Matrix-assisted laser desorption/ionization mass spectrometry
(MALDI-MS)
- activated carbon substrate surface for, 11:644
- of adenovirus proteins extracted by SDS-PAGE, 11:356
- antibody-peptide complex preservation and detection by, 11:
356
- calixarene-amino acid non-covalent complexes formed by, 13:
964
- chemical noise in, 13:129
- to determine composition of *N*-linked carbohydrates from
ovalbumin and co-purified glycoproteins, 11:564
- to determine postsource decay fragmentation of *N*-linked car-
bohydrates from ovalbumin and related glycoproteins,
11:572
- dihydroxybenzoic acids used in, gas phase acidities/basicities
of, 11:544
- fragmentation studies of oligonucleotides in, by charge tag-
ging and H/D exchange, 13:1407
- of IgG nonenzymatic glycation, 11:153
- infrared
- laser pulse duration role in, 13:978
- mean initial ion velocities in, 13:209
- of in-source H/D exchange and ion molecule reactions, 13:308
- of laser microdissected cells, 13:1292
- in locating abasic sites in oligodeoxynucleotides, 13:1418
- of nonpolar polymers, silver cluster interferences in, 12:732
- for oligodeoxynucleotides, improvements in matrix, detection
limits, quantification, and sequencing, 11:854
- of oligonucleotides, 12:5-methoxysalicylic acid and spermine
as matrix for, 12:456
- of peptides and proteins containing nitrotyrosine, 12:438
- of pesticides, 12:590
- of phosphopeptides affinity bound to immobilized metal ion
beads, 11:273
- of protein, 13:107
- surface peptide retention determination by, 11:62
- ultraviolet, stabilizing double-stranded DNA in, 13:1129
- Matrix-assisted laser desorption/ionization-mass spectrometry
of acidic glycoconjugates facilitated by the use of spermi-
ne as co-matrix, 9:1293
- buffer tolerance in, 9:805
- of human immunodeficiency virus-gp120 epitopes recognized
by a limited polyclonal antibody, 9:157
- of oligomer molecular weight, 9:282, 9:651
- phosphopeptide detection by, 10:35
- of proteins, 9:1, 9:88
- Matrix-assisted laser desorption/ionization-MS. *See also* Gel per-
meation chromatography-MALDI-TOFMS
- of bis(imidazole-1-carboxylate) end-functionalized polymers,
14:117
- Cu(II) reduction in, 14:42
- of end-functionalized polystyrenes, 14:648
- of H/D exchange on ATPase supramolecular protein com-
plex, 14:471
- hexamethylcyclotrisiloxane polymerization studied by, 14:95
- matrix mixture for, in peptide mass mapping and protein
analysis, 14:992
- in spatial profiling invertebrate ganglia, 14:752
- Matrix-assisted laser desorption/ionization-MS/MS direct
of peptides affinity-bound to antibody beads, 14:1076
- in trypsin-catalyzed ^{16}O -to- ^{18}O exchange, for comparative pro-
teomics, 14:704
- Matrix-assisted laser desorption/ionization orthogonal quadru-
pole time-of-flight mass spectrometry (MALDI-quadru-
pole TOF-MS), pseudo-MS3 in, 13:1034
- Matrix assisted laser desorption/ionization orthogonal-time-of-
flight mass spectrometry, human and elephant milk oli-
gosaccharide analysis using TLC and, 17:139
- Matrix-assisted laser desorption/ionization-post-source decay
charge-remote fragmentations under, 10:217
- of peptides and proteins, 10:91

- Matrix-assisted laser desorption/ionization post-source decay (MALDI-PSD), of CHCA clusters, 15:336
- Matrix-assisted laser desorption/ionization postsource decay spectra (MALDI-PSD spectra), of charge derivatized peptides, examples of tris(trimethoxyphenyl)phosphoniumtagged proteolytic digestion products of phosphoenolpyruvate carboxykinase, 11:145
- Matrix-assisted laser desorption/ionization-quadrupole ion trap mass spectrometry, atmospheric pressure, ethoxylated surfactant fragmentation by, 17:453
- Matrix-assisted laser desorption/ionization-quadrupole ion trap MS, for de novo peptide sequencing, 14:1012
- Matrix-assisted laser desorption/ionization tandem mass spectrometry
- Conus virgo* conotoxin pyroglutamic acid identification by, 18:1396
 - ILPR binding sites in IGF-2 and insulin, 20:1974
 - In situ characterization of sulfatides and glycerophospholipids, brain, 18:17
- Matrix-assisted laser desorption/ionization tandem mass spectrometry post-source decay (MALDI-MS/MS-PSD), of phenylhydrazone-oligosaccharides from ovalbumin, 15:725
- Matrix-assisted laser desorption/ionization tandem reflectron time-of-flight mass spectrometry of fullerenes, 7:590
- Matrix-assisted laser desorption/ionization tandem time-of-flight (MALDI-TOF/TOF) mass spectrometry, of N-sulfonated peptides, 15:1838
- Matrix-assisted laser desorption ionization/time-of-flight and electrospray ionization mass spectrometry, analysis of serum protein precipitated with antiserum, 6:854
- Matrix-assisted laser desorption/ionization-time-of-flight instrument, automated MS imaging with, 10:67
- Matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry
- of bacterial spores, 15:1444
 - improved PSD and CID on, 15:522
 - noncovalent peptide-DNA interaction detected with, 15:28
 - quantitative analysis by, ionic-liquid matrices for, 15:1833
 - in stress-related fingerprint identification from *Bifidobacterium lactis*, 15:1222
- Matrix-assisted laser desorption ionization-time of flight mass measurement accuracy, by using delayed extraction, 7:995
- Matrix-assisted laser desorption ionization time-of-flight mass spectrometry
- ATP binding to growth factors detected by ESI-MS, 19:91
 - automated digestion and droplet deposition microfluidic chip for, 19:964
 - cryodetection, analysis of polystyrenes, 19:219
 - HPLC, wheat gluten proteins, 19:1542
 - oligosaccharides, post-source decay analysis, 19:1119
 - sialylated glycans and oligosaccharides, analysis using DHB/An and DHB/DMA matrices, 19:1138
- Matrix-assisted laser desorption ionization-time-of-flight mass spectrometry, liquid chromatography-electrospray ionization, rapid analysis of antibiotic-containing mixtures from fermentation broths by using, 7:1227
- Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry
- accurate mass capability, radical ion species, 17:672
 - butyrylcholinesterase inhibition by OP and CB pesticides, MALDI-TOF-MS, 18:698
 - comparison between vacuum sublimed matrices and dried droplet preparation, 20:1104
 - direct analysis of oligonucleotides synthesized on montmorillonite, 17:1265
 - dominulin A and B in cuticle and venom of social paper wasp analyzed by MALDI-TOF/TOF, ESI-ion trap and, 17:376
 - effect of diammonium citrate to CHCA matrix for detection of phosphorylated peptides, 18:1925
 - effect of Tween80 on signal intensity of intact proteins, 18:102
 - honeybee venom detection in envenomed tissues, 20:112
 - 3-hydroxycoumarin as new matrix for, 17:1665
 - identification of mammalian cell lines using LC-ESI-MS/MS, 17:490
 - identification of pigments and dyes, 18:2001
 - improved imaging of proteins, in tissues, 20:89
 - label-free approach to rapid inhibitor screening, 17:815
 - M. bovis* BCG PIMS characterization by ESI MS and, 18:466, 18:479
 - MELDI protein profiling tool with specific carrier materials, 17:1203
 - negative ion mode, neutral lactooligosaccharide fragmentation in UV-MALDI-TOF and UV-MALDI ion trap/TOF MS, 17:67
 - noncovalent interactions, exploring intensity fading using, 18:359
 - oligosaccharides labeled with multifunctioning tag, analysis by ESI-MS and, 17:194
 - oxidized carbon nanotube matrix, monitoring enzyme reactions and screening inhibitors with, 17:1616
 - parallel approach to post source decay, 17:60
 - parameters in optimization of synthetic polymers, 17:246
 - pentacene clustering in, 17:222
 - phosphopeptide enrichment using PAA-membrane, 18:1385
 - S. mliariius barboursi* snake venom, proteomic profiling using, 18:600
 - SBA-15, small molecule analysis by, 20:2167
 - solid state NMR and, analysis of polylactides, 20:67
 - solvent accessibility of protein surfaces, 17:1490
 - titanium oxide characterization of nanoclusters, LDI-TOF-MS and, 18:517
 - triacylglycerols, CID MALDI-TOF/RTOF, 20:1037
 - use of β -cyclodextrin in determining MWs of low-molecular weight explosives, 17:189
 - using spiral ion trajectory, 18:1318
 - UV, protein desolvation in, 18:1880
- Matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry, of polymers
- buffer tolerance in, 9:805
 - of gel-separated proteins, 10:512, 10:784
 - negative, post-source decay calibration of, using fibrinopeptide B, 9:540
 - oligonucleotide ion stability in, co-matrix proton affinity and, 9:668
 - in photodissociation of substance P, 10:1038
 - of polymers
 - accuracy of, 9:275
 - solvent effect on, 9:1303
 - resolution and mass accuracy in, 9:892
 - of silsesquioxane molecular structure, 10:224
 - of triacylglycerols and whole oils, 10:983
- Matrix-assisted laser desorption ionization time-of-flight mass spectrometry and electrospray mass spectrometry, application of, for molecular weight assignment of peptides in complex mixtures, 4:670
- Matrix-assisted laser desorption ionization time-of-flight mass spectrometry (MALDI-TOF-MS)
- in analysis of nonpolar synthetic polymers, 11:731
 - biomedical applications of, 13:1015
 - of complex polyol mixtures, 12:55
 - of diastereomeric benzo[ghi]fluoranthene tetrols, 12:376

- of dihydroxybenzoic acid noncovalent association with cyclic and linear oligosaccharides, 11:228
- of dilute peptide mixtures, matrix-cluster peaks in, 11:88
- in-source decay
- of hyperbranched polyesteramides in, 11:218
 - peptide sequence by, 11:1000
- of in-source peptide decay, 12:420
- mass-correlated pulsed extraction in, 11:841
- in mass measurement of low molecular weight compounds, 11:458
- of novel ions from protonated peptides, 11:345
- of peptides, 13:772, 13:784
- of polyethylene, 12:1186
- of polystyrene, postsorce decay in, 11:1095
- positive-ion, in study of adenine-thymine binary-base oligonucleotide dissociations, 13:1052
- quantitative analysis of technical polymer mixtures by, 13:914
- reflectron, mass correlated acceleration in, 13:135
- of solubilization of phosphopeptides retained on ferric NTA IMAC beads, 13:1042
- substance P molecule detection by, 12:1055
- Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOFMS). *See also* Delayed extraction matrix-assisted laser desorption/ionization time-of-flight mass spectrometry
- Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry of peptides, improved mass accuracy in, 5:955
- Matrix-assisted laser desorption ionization time-of-flight mass spectrometry of underivatized and permethylated gangliosides, 3:785
- Matrix-assisted laser desorption/ionization time-of-flight tandem mass spectrometry
- amino acid analysis by, 18:279
 - gold nanoparticles as assisted matrices for biomolecules, 20:875
 - nano-LC, bacteriotoxin E9:Im9 complex analysis using lysine acetylation and, 17:983
 - PE chloramines detected using 4-chloro- α -cyanocinnamic acid and, 20:867
- Matrix-assisted laser desorption ionization time-of-flight/time-of-flight mass spectrometry
- in analysis of whole cell bacteria, 14:1306
 - Aurum Dataset for protein standards, 18:850
 - H/D-exchange, RNA fragmentation studied by, 17:1353
 - HPLC/ESI-FTICR MS *vs.*, comparison using HIV envelope glycoproteins, 19:1209
 - hydrogen atom scrambling, anionic peptides, collisional activation by, 19:1719
 - in identification of two-dimensional gel proteins, 14:956
 - laser fluence, leucine enkephalin fragmentation with, 18:607
 - multiply-charged intactprotein fragmentation using, 19:231
 - P0 characterization by, bovine myelin sheath, 17:117
 - picolinamidination of phosphopeptides for sequencing by, 20:1751
 - in phosphatidylcholine structural analysis, 14:373
 - from whole bacteria cells, quality and reproducibility of, 14:342
- Matrix-assisted laser desorption mass spectrometry, negative-ion, analysis of oligodeoxynucleotides by, 4:955
- Matrix-assisted laser desorption mass spectrometry of bio-transformation products of dynorphin A in vitro, 5:10
- Matrix-cluster peaks, in MALDI-TOFMS of dilute peptide mixtures, 11:88
- Matrix design for matrix-assisted laser desorption ionization, sensitive determination of PAH-DNA adducts, 5:1021
- Matrix effect, evaluation of, 16:1757
- Matrix effect in bioanalysis of illicit drugs with LC-MS/MS, influence of ionization type, sample preparation, and biofluid, 14:1290
- Matrix effects
- comparison of HPLC-MS/MS and UPLC-MS/PM analyses, pharmaceuticals in surface waters, 19:713
 - polyatomic ion interferences and, inductively coupled plasma mass spectrometry with an enlarged sampling orifice and offset ion lens, 4:28
- Matrix-enhanced secondary ion mass spectrometry (MESIMS), to analyze polymer MALDI matrix solubility, 16:1172
- Matrix-enhanced surface-assisted laser desorption/ionization mass spectrometry
- ionic matrix for, 20:2229
 - metabolite imaging using, 20:80
- Matrix-induced interferences in mixed-gas helium-argon inductively coupled plasma mass spectrometry, investigation of, 2:355
- Matrix metalloproteinase 7, cysteine sulfenic acid in, MS/MS, 18:1544
- Matrix metalloproteinase 12, catalytic domain of, activity anchored on Au(111), 18:961
- Matrix methods, to calculate stability diagrams in quadrupole MS, 13:597
- Matrix protein 1, avian influenza virus (H5N1), AA substitutions mapped by nanoelectrospray MS, 20:312
- Matrix-shimmed ion cyclotron resonance ion trap simultaneously optimized for excitation, detection, quadrupolar axialization, and trapping, 10:759
- Matrix solubility, MESIMS of, 16:1172
- Matrix to analyte proton transfer, in clusters, 12:726
- Matrixes
- influence on analyte fragmentation, atmospheric pressure MALDI, 17:1005
 - oxidized carbon nanotube, monitoring enzyme reactions and screening inhibitors, MALDI-TOF-MS with, 17:1616
- McLafferty or ion-neutral complex rearrangements
- APCI low energy collision-induced dissociation fragmentation of protonated ortho silicates, 5:544
 - bacillus stearothermophilus, glucokinase from, ESI-FTMS-IRMPD of, 9:1222
 - bis(alkylamine)mercury cations, from mercury nitrate surfaces, ion trap-secondary ion MS of, 9:35
 - m/z 69 ion, from β -diketones, 9:29
 - muscidifurax, cuticular hydrocarbons of, GC-MS of, 9:320
 - trans-2-hexenal, peptides modified with, MS/MS of, 10:613
- McLafferty rearrangement, 15:951, 15:956
- binding energies of water to lithiated valine, 15:1014
 - within a complex, 15:966
 - condensation reactions in protonated dipeptides, 15:1025
 - construction of hybrid quadrupole FTICR-MS, for MS/MS, 15:1099
 - decomposition of protonated formic acid, 15:982
 - ECD distinguishing single d-amino acid in, 15:1087
 - Meisenheimer complexes bonded in gas phase, 15:998
 - modeling deoxyribose radicals by neutralization-reionization MS, 15:1055, 15:1068
 - MS/MS of methylbenz(is)oxazole isomers, 15:1005
 - MSn characterization of protonated cyclic peptides and metal complexes, 15:1039
 - orbital symmetry restraint circumvention in, 15:972
 - photoionization of hexyl alcohol, 15:989
 - proton affinity of nucleic acid molecules, kinetic method in, 15:1080
 - in secondary ion images of developing brain, 15:1116
 - solution equivalent of, 15:959
- MCO. *See* Metal-catalyzed oxidation reactions

- MCS. *See* Monte Carlo simulations
- MCS1-molecular ion formation, during sputtering, 11:650
- MDS. *See* Molecular dynamics simulation
- ME-SALDI-MS. *See* Matrix-enhanced surface-assisted laser desorption/ionization mass spectrometry
- Measurement errors, in FTICR-MS, 13:99
- Measurement of collision-induced dissociation rates for tantalum oxide ions in a quadrupole ion trap, 11:1072
- Measurement of high-molecular-weight polycyclic aromatic hydrocarbons in soils by particle beam high-performance liquid chromatography-mass spectrometry, 6:597
- Measurements of mean initial velocities of analyte and matrix ions in infrared MALDI-MS, 13:209
- Mechanism for the loss of 60 u from peptides containing an arginine residue at the C-terminus, 8:253
- Mechanism of gas-phase aldose-ketose isomerization, a study using tandem mass spectrometry and theoretical calculations, 4:943
- Mechanism of matrix to analyte proton transfer in clusters of 2,5-dihydroxybenzoic acid and the tripeptide VPL, 12:726
- Mechanism of poor high mass performance in Fourier transform mass spectrometry, 3:460
- Mechanisms, dynamics, and applications, 14:680
- Mechanisms of single-electron capture by the dichlorocarbene dication, 7:266
- Mechanistic considerations of the protonation and fragmentation of highly functionalized molecules in fast atom bombardment, high resolution mass spectrometry and tandem mass spectrometry analysis of the ions formed by fast atom bombardment of digoxin and related cardiac glycosides, 1:455
- Mechanistic studies
of ionization suppression in ESI, 11:942
of multipole storage assisted dissociation, 11:210
- Mechanistic study of ES-MS: charge gradients within electrospray droplets and their influence on ion response, 12:206
- Mechanistic study of hydrogen/deuterium exchange between [m-1] ions of chlorinated benzenes and D₂O or ND₃, 5:282
- ¹³C, IRMS detection of, 10:153
- Medicinal herbs, saponins from, ESI-MS/MS structural analysis of, 15:133
- Medium pressure laser ionization, ion source for, 15:1885
- Meerwein reaction, of phosphonium ions with epoxides and thioepoxides in gas phase, 15:398
- Meisenheimer complexes, bonded in gas phase, 15:998
- Melanin, from epidermal melanocytes, analysis by pyrolytic GC/MS, 20:464
- MELDI. *See* Material-enhanced laser desorption/ionization
- Melittin
bound to phospholipid micelles, H/D exchange and ESI-FTICR-MS of, 12:1247
calcium-induced association of calmodulin with, 8:809
gas-phase conformation probed by CID, 17:786
- Melphalan-modified 2'-oligodeoxynucleotides, miniaturized LC-ES MS/MS of, 15:568
- Melting points, ion calorimetry using MS to measure, 18:74
- Membrane inlet mass spectrometer, sample modulation at, mathematical study of-potential application in analysis of mixtures, 7:93
- Membrane inlet mass spectrometry (MIMS), cyclic sudden sampling introduction mode in, 15:823
- Membrane introduction mass spectrometry
Candida rugosa lipase immobilized on PDMS membrane, 18:973
radical sampling by, 9:234
- Membrane preconcentration
enhanced sample loading by, 8:15
protein modification by, 8
- Membranes
nylon-based, proteins immobilized on, direct matrix-assisted laser desorption ionization mass spectrometric analysis of, 5:230
proteins, analysis by LILBID MS, 18:1429
- Mercaptobenzothiazoles, new class of matrices for laser desorption/ionization mass spectrometry, 8:116
- Mercapturic acids, LC-MS/MS detection of, 16:1976
- Mercury ions, interactions among polythymine oligonucleotides, ethidium bromide and, ESI MS, 20:1834
- Mercury nitrate surface, bis(alkylamine)mercury cations from, ion trap-secondary ion MS of, 9:35
- [Me₂Sφ0 Σ 77 ε₂], *στρυχτυρε ανδ φραγμαεντατιον οφ*, 8:605
- Meshes, influence of material and mesh characteristics on TM-DESI, 20:584
- MESI-MS. *See* Microelectrospray ionization mass spectrometry
- meso*-Tetrakis(heptafluoropropyl)prophyrin, ESI-MS/MS of, 16:1915
- meso*-Tetraphenylporphyrins, β-substituted, fragmentation induced by high energy collisional activation, 13:1427
- Meso*-tris-*N*-methylpyridinium-4-yl porphyrins, reduction in positive mode ESI-MS, 18:762
- Meta-S-SIMS. *See* Metal-assisted static secondary ion mass spectrometry
- Metabolism, microsomal, of bromotrichloromethane and carbon, tandem mass spectrometry study of C-phenyl-*N*-tert-butyl nitron spin adducts, in vitro rat liver and, 6:847
- Metabolites, nitrogen-containing intracellular, HPLC-MS quantitation, *S. enterica*, 17:37
- Metabolomics, EC-MS in, 15:1717
- Metal
attachment of to polystyrenes, in MALDI, 9:724
changes in vitamin D receptor DNA-binding domain induced by, 9:8
pharmaceutical compound complexation with, ESI vs, 9:463
polyether complex with, in quadrupole ion trap, 8:620
transition, oligosaccharides of, electrospray ionization study of, 8:32
- Metal-assisted static secondary ion mass spectrometry, in situ temperature-time effect, 18:382
- Metal-binding properties, of centrin-2, determined by μESI-MS and UV spectroscopy, 17:1158
- Metal binding stoichiometry of matrilysin by electrospray ionization mass spectrometry, noncovalent enzyme-inhibitor complexes and, 6:1105
- Metal carbonyl anions, insertion reactions of, with methyl formate in the gas phase as revealed by ¹³C- and D-labeling, 5:525
- Metal-catalyzed oxidation reactions, Cu-binding protein AA residues identified using MS and, 17:1552
- Metal cation, alkali-valinomycin complexes, gas-phase studies of, attachment rates and cation affinities, 5:632
- Metal cation-trimethylphosphate, interactions with nucleic acids, 19:305
- Metal cationized peptides, ions from, CID fragmentation pathway for, 16:1305
- Metal chelation, of dinucleotide analogs, in the gas phase by fast-atom bombardment mass spectrometry, 7:42
- Metal cluster ions
group-11, photodissociation of, 14:614
transition, reactive and dissociative behavior of, in the gas phase, characterization of, 2:212
- Metal clusters, molecule interactions with, MALDI-MS of, 15:879

- Metal complexation
in flavonoid glycoside isomer differentiation, 14:1437
flavonoid glycosylation site determination by, 15:1287
- Metal complexes
CAD of, flavonoid glycoside characterization by, 12:537
of crowned spirobenzopyran derivatives, evaluating photoinduced changes by ESI-MS, 14:1110
of di-*o*-benzoyl-tartaric acid dibutyl ester and L-tryptophan, in gas phase, chiral recognition characteristics of, 14:571
dissociation of, metal ion and ligand effects on, 14:152
of flavonoids, transition, threshold dissociation and molecular modeling of, 16:139
gas phase coordination chemistry of, 13:813
Group 10, MS behavior of, 16:94
MSⁿ characterization of, 15:1039
- Metal-complexes, formation in continuous-flow ligand-exchange reactors, ESI/MS, 18:707
- Metal dications, preferential solvation in water/methanol clusters, 19:520
- Metal halide ions, reactivity and binding energy with benzene, 13:477
- Metal hydrides, deuterated, MS of, 18:337
- Metal-induced conformational mutations, FTICR and computational analyses of, 12:1272
- Metal-ion affinity chromatography, on-line immobilized, phosphopeptides isolated by, electrospray ionization mass spectrometry of, 4:662
- Metal ion beads, phosphopeptides affinity bound to, MALDI-MS of, 11:273
- Metal-ion binding and limited proteolysis of betabellin 15D, a designed beta-sandwich protein, 10:969
- Metal ion complexes
with polyamidoamine dendrimers, ECD and CID of, 20:674
transition metal complex cations in gas-phase transformation of deprotonated polypeptides, 20:1718
- Metal ion complexes-crown ether complexes, direct desorption of, in liquid secondary ionization mass spectrometry, evidence for, 5:638
- Metal ion plasma, laser-generated, competitive ionization of tetraphenylporphyrin in, 4:159
- Metal ion source, liquid, Li-Mg, cluster ions ejected from, observation of Mg₂²⁺ and Mg₃²⁺, 2:76
- Metal ion source for gas-phase ion molecule reactions, selective, resonant laser ablation as, 7:664
- Metal ion(s)
alkali
in ESI-MS of peptides and proteins, 11:967
gas-phase anionic complexes of, structure and collision activated decompositions, 5:137
poly(ethylene glycol) doubly and singly cationized by, 13:177
structural effects on polyether cationization, in MALDI, 11:639
alkaline earth, reactions with benzene, 12:250
divalent, interactions with peptides probed by ECD of trications, 17:1731
earth, alkaline, fatty acids cationized with, charge remote fragmentation of, 1:320
interactions of, 12:479
alkali-cationized peptides in, C-terminal peptide sequencing of, 12:497
cobalt-coordinated hexosamine monosaccharides in, 12:517
Cu²⁺-bound complexes in, 12:490
Cu⁺-dimethoxyethane complexes in, 12:480
flavonoid glycoside characterization by metal complex CAD, 12:537
metal-ligated sialic acid containing oligosaccharides as, 12:528
oligodeoxynucleotide ions adducted to metal ions in, 12:550
sodium stabilization of iturin A₂, during ESI-MS, 12:505
photomodified oligodeoxynucleotides adducted with, fragmentation in ESI-ITMS, 12:1174
reactions with long chain alkanes, 12:367
transition, and ESI low energy CID-MS/MS of polyglycols, 12:832
zinc, and angiotensin peptides, interaction of, by electrospray ionization mass spectrometry, 5:959
- Metal-ligand complexation, tunable, for flavonoid diglycosides, ESI-MS, 18:422
- Metal mediated heteroatom removal, by reactions of FeL⁺ [L⁺O, C₄H₆, c-C₅H₆, H₆H₆, C₅H₄(=CH₂)] with furan, thiophene and pyrrole in the gas phase, 7:938
- Metal-metal interfaces, enhanced ionization efficiency, MALDI MS, 17:737
- Metal nitrate cluster ions, anionic group II, IR-MPD spectroscopy, 20:772
- Metal nitrate negative ions, produced by ESI, MS/MS of, 14:671
- Metal-oligonucleotide complexes, MS/MS of, 16:370, 16:799 (erratum)
- Metal oxides, surface fabrication by reactive landing of gas phase ions, 20:915
- Metal/polyether/pyridyl ligand complexes, in quadrupole ion trap mass spectrometer, 10:126
- Metal salts, of aqueous solutions, electrospray mass spectrometry, 3:281
- Metal-transition ions, in the gas phase, ion-molecule reaction of, a study on fragmentation mechanism of the adducts, 2:120
- Metal trications, attachment to peptides, 15:406
- Metallized tips, long-lived, for nanoliter electrospray mass spectrometry, 7:1270
- Metallo-β-lactamase
CphA, monitoring zinc affinity by automated nano-ESI-MS, 17:180
ternary inhibitor complexes, studies with ESI-MS, 17:1000
- Metalloprotein
3+ oxidation state iron-sulfur, 12:817
metal atom oxidation state in, 11:120
- Metallothionein, ion mobility and MS combined investigations of, 16:1787
- Metals
imaging on micro and nanoelectronic surfaces by LA-ICP-MS, 20:883
interactions with α-synuclein and calmodulin monitored by SNAPP-MS, 19:1668
- Metaphosphate anion, monomeric, tropylium, chlorine isotopic abundances, and Conestoga wagon theory, 4:761
- Metastable and collision-induced fragmentation studies of all C₄H₁₂Si⁺ isomers, systematic study of structure-reactivity relations, 5:537
- Metastable atom bombardment, to eliminate concentration dependence of MIAMS of methyl palmitate, 12:754
- Metastable atom bombardment ionization mass spectrometry (MAB-MS), for identifying additives in polymers, 15:1315
- Metastable decay of negatively charged oligodeoxynucleotides analyzed with ultraviolet MALDI post-source decay and deuterium exchange, 12:180
- Metastable decomposition
in MALDI-TOFMS, novel ions for protonated peptides observed by, 11:345
slow, of oligosaccharide cations produced in an external source Fourier transform mass spectrometer, 4:210
- Metastable decomposition of peptide [M + H]⁺ ions via rear-

- rangement involving loss of the C-terminal amino acid residue, 1:249
 Metastable ion formation and disparate charge separation in the gas phase dissection of protein assemblies studied by orthogonal TOFMS, 12:329
 Metastable ion study of substituted cyclopentadienylmanganese cations in the gas phase, 4:216
 Metastable ions, half-protons or doubly charged ions as, 15:1261
 Metastable molecular ion of methyl lactate, fragmentation of, formation of oxygen-protonated methanol $[\text{CH}_3\text{OH}_2]^+$ involving double hydrogen atom transfers, 6:202
 Methane
 elimination of, from protonated acetaldehyde, the ab initio transition state, 5:1102
 membrane inlet MS for online and in situ analysis of, aquatic environments, 19:1395
 protonated, on the structure of, 2:453
Methanobacter thermautotrophicus, archaeal tetraether lipids, analysis by APCI LC-MS/MS, 20:51
 Methanol
 gas phase reactions of, 8:1124
 in induction of conformational transitions of beta-lactoglobulin, cytochrome c, and ubiquitin at low pH, study by ES-IMS, 12:317
 proteins in, ESI-MS of, 12:317
 Methanol and aldehyde, protonated, gas-phase reaction between, 1:481
 Methanol and protonated peptides, interaction of, molecular probe for, gas-phase hydrogen/deuterium exchange as, 5:623
 Methanol solutions, suppression of electric discharge with SF₆ gas, electrospray mass spectrometry of, 2:497
Methanosarcina acetivorans, top-down proteomics reveals proteins expressed in, 20:1743
 Methionine
 mechanisms for selective gas-phase fragmentation reactions, 17:1631
 oxidation in calmodulin, quantification of, 16:1470
 Methionine-containing peptides, analysis by charge derivatization and MS/MS, 16:1131
 Methionine enkephalin, endogenous, quantification of, electrospray mass spectrometry for, 2:149
 Methionine oxidation
 in peptides, CID and ECD of, 14:605
 photo-induced, IgG1 mAbs, MS, 20:525
 Method, 3:863
 Method for screening peptide fragment ion mass spectra prior to database searching, 11:422
 Method for the design of broad energy range focusing reflectrons, 7:1002
 Method of increasing the sensitivity for laser desorption in an ion trap mass spectrometer, 4:706
 Methoxy-carbonyl groups, adjacent, in the gas phase, on the stabilization of carbanions by, 5:159
 2-Methoxychalcone, protonated 2-hydroxychalcone and, Nazarov cyclization, 20:805
 Methoxypropene ions, 3:1- and 3-, fragmentation of, another part of the $\text{C}_4\text{H}_8\text{O}^+$ potential energy surface, 3:99
 5-Methoxysalicylic acid and spermine, new matrix for MALDI-MS analysis of oligonucleotides, 12:456
 Methoxytrimethylsilane, metastable ions of, 9:1029
 Methyl 2-methyl butanoate ions, ethene loss kinetics of, 10:200
 Methyl benzoate
 metastable, decarboxylation of, 17:807
 O reactions with, 12:694
 Methyl cation, neutralization at hydrocarbon and fluorocarbon SAMs, 13:1151
 Methyl esters
 fatty acid
 molecular ion derived from, low-energy tandem mass spectrometry of, a novel method for analysis of branched-chain fatty acids, 4:223
 O⁻ and OH⁻ chemical ionization of some, 5:553
 saturated aliphatic, proton affinities of, 11:789
 Methyl ether, ionized phytol, $\text{C}_3\text{H}_9\text{O}^+$ formation from, the structure and mechanisms of, 7:205
 Methyl formate, insertion reactions of metal carbonyl anions in the gas phase with as revealed by ¹³C- and D-labeling, 5:525
 Methyl groups, migration during fragmentation of trimethyllysine-containing peptides, MS/MS, 20:1172
 Methyl guanine isomer distinction by hydrogen/deuterium exchange using a Fourier transform mass spectrometer, 4:296
 Methyl lactate, metastable molecular ion of, fragmentation of, formation of oxygen-protonated methanol $[\text{CH}_3\text{OH}_2]^+$ involving double hydrogen atom transfers, 6:202
 Methyl nitrate, [M-H] anions of, structures of, 1:295
 Methyl nitrates, protonation sites in, by NRMS, 11:380
 Methyl orange, photolytic degradation in water, LC-ESI-MS, 19:997
 Methyl palmitate, MIAMS of, eliminating concentration dependence of using MAB, 12:754
 Methyl propionate radical cation, 7:482
 Methyl substituted analogs, and phenyl n-propyl ether, chemical ionization of, propene loss initiated by competing proton transfer to the oxygen atom and the aromatic ring, 7:639
 Methyl thioglycolate, molecular ions of, metastable decomposition of, 8:801
 Methylamine systems, ion-molecule reactions in, 2:220
 Methylamines and trimethylaluminum mixtures, ion-molecule reactions in, 4:111
 β-Methylamino-L-alanine, ethylchloroformate derivative, characterization by GC/MS, 18:817
 Methylarginine-containing peptides, fragmentation pathways in, ESI-MS/MS of, 15:142
 Methylated lactones, gas-phase, dissociation behavior of, characterization of the, 3:47
 Methylation, gas-phase, and dissociation of acids and esters, effects of functional group interactions on, 7:565
 Methylbenz(is)oxazole isomers, MS/MS of, 15:1005
 Methylene substitution reactions in a quadrupole ion trap, selectivity of ethylene, ethylene oxide, and dimethyl ether reactive ions, 3:39
 Methylene transfer, ionized, from the distonic radical cation +CH₂O-CH₂ to heterocyclic compounds, pentaquadrupole mass spectrometric study, 6:554
 Methylgermane and germane, gas-phase acidities of, Fourier transform-ion cyclotron resonance study of, bond dissociation energy of germane, 4:54
 N-Methylimidazole, complex with β-cyclodextrin, ESI-MS, 17:9
 2-Methylloxazolopyridine, heterocyclic isomers, characterization and differentiation of, 6:962
 Methylphosphonates, replacing phosphates with, in oligodeoxynucleotides fragmentation, 12:580
 N-Methylpyrrole, complex with β-cyclodextrin, ESI-MS, 17:9
 Methylthioacetic acid, determining gas phase acidity of, Cooks' kinetic method in, 16:535
 Mg₂⁺, observation of, cluster ions ejected from an Li-Mg alloy liquid metal ion source, 2:76
 Mg₃⁺, observation of, cluster ions ejected from an Li-Mg alloy liquid metal ion source, 2:76
 MH⁺ ions of peptides, high-energy collision-induced decompo-

- sition of, comparison of helium and argon as collision gases in, 3:502
- MHC-I. *See* Major histocompatibility complex 20:1 molecules
- MIAMS. *See* Mass isotopomer abundance mass spectrometry
- Mice
interleukin-3, baculovirus expressed, liquid secondary ion mass spectrometry analysis of permethylated, n-hexylamine derivatized oligosaccharides, 3:345
isotope ratios of carbon dioxide in expired air from, following the administration of ¹³C-methyl methionine, the quadrupole-based mass spectrometric evaluation of, 3:742
- Michael-Michael-retro Michael addition, catalyzed by 9-amino-9-deoxyepiquinine, ESI-MS study, 18:2074
- Micro-electrospray, zeptomole/attomole per microliter sensitivity for peptides, 5:867
- Micro-electrospray mass spectrometry, ultra-high sensitivity analysis of peptides and proteins, 5:605
- Micro high-performance liquid chromatography-mass spectrometry, analysis of coumarins by, with a particle beam interface, 6:132
- Micro-liquid chromatography/ES-API-CID-MS hybrid scan technique, to identify IRS-1 phosphorylation site, 14:401
- Microarrays, construction, thiolated deoxyoligonucleotide characterization and sequence variation, HPLC-MS, 17:1396
- Microbial spectral databases, defining and using, 13:875
- Microbore high-performance liquid chromatography-tandem mass spectrometry utilizing pneumatically assisted electrospray ionization, quantitative determination of a non-peptide antithrombotic in dog plasma by, 7:1238
- Microcanonical analysis of the kinetic method, meaning of the "apparent entropy," 13:435
- Microcin C7-C51, *E. coli*, CID characterization, 19:1187
- Microcolumns, packed, continuous flow fast atom bombardment with, comparison of precolumn *versus* coaxial matrix delivery, 1:312
- Microcystins
chirality in, 5:724
ESI-MS of, 10:1138
- Microelectrospray ionization mass spectrometry (MESI-MS), of alkali metal binding selectivities of caged aza-crown ether ligands, 11:884
- Microelectrospray ionization mass spectrometry (μ ESI-MS), of complex polyol mixtures, 12:55
- Microfabricated PDMS multichannel emitter for ESI-MS, 12:463
- Microflow rate particle beam interface, electron capture ionization of explosives with, 7:753
- Microfluidic chips
capillary gel, interfacing with infrared LDI MS, 17:469
for millisecond time-scale kinetic studies, ESI-MS, 20:124
- β_2 -Microglobulin, release from MHC-I, analyzed by HDX-ESI-MS, 20:278
- Micrometer particles, ionization of biomarkers from, BAMS in, 15:900
- Microprobe mass spectrometry, laser of quaternary phosphonium salts, direct *versus* matrix-assisted laser desorption, 4:798
- Microprobe two-step laser desorption/laser ionization (μ L²MS), of polycyclic aromatic hydrocarbons in ancient terrestrial rocks, 12:989
- Microsolvation, role in peptide cation survival after electron capture, 17:1675
- Microsomal metabolism of bromotrichloromethane and carbon, tandem mass spectrometry study of C-phenyl-*N-tert*-butyl nitrene spin adducts, in vitro rat liver and, 6:847
- Microspheres, silica, enzyme inhibitors immobilized on, screening by ESI-MS, 19:865
- Microvolume injection, 50-femtomole, complete large-molecule high-resolution mass spectra from, 6:220
- Microwave-assisted acid hydrolysis, of protein, LC MALDI MS/MS with, 16:471
- Microwave-assisted enzyme-catalyzed reactions, in solvent systems, 16:581
- Microwave-enhanced Akabori reaction for peptide analysis, 13:839
- Microwave-induced plasma ionization mass spectrometry, of underivatized amino acids, 12:117
- Microwave-induced plasma ionization source, for molecular mass spectrometry, 9:42
- Mid-infrared, tunable, free-electron laser, infrared matrix-assisted laser desorption and ionization by using, 7:1187
- Milk
human, oligosaccharides in, structural analysis of, 13:1331, 13:1341
human and elephant, oligosaccharide analysis using TLC and MALDI-MS, 17:139
- MIMS. *See* Membrane introduction mass spectrometry
- Mineral surfaces, tributyl phosphate on, static secondary ionization mass spectrometry analysis of, effect of Fe(II), 6:165
- Mini-ball mill, β -amyloid peptide analysis via, MALDI MS, 18:1533
- Mini-ball mill approach, in solvent-free MALDI-MS of biological samples, 16:542
- Miniature quadrupole mass spectrometers, 12:633
- Miniaturized EI/Q/oa TOFMS, 12:641
- Minor groove-binding molecules and duplex DNA, noncovalent complexes formed between, by electrospray ionization-mass spectrometry, 6:1154
- MIPI. *See* High performance liquid chromatography/atmospheric pressure microwave-induced plasma ionization mass spectrometry
- Mitochondria
cardiolipin concentrations, MALDI-MS analysis from rat organ sections, 18:567
FMN-binding complex I subunit, MS profile of intramolecular disulfides, 19:1876
- Mitochondrial fractions, heart, phosphatidylcholine isolated from, analysis by fast-atom bombardment tandem mass spectrometry, evidence of incorporation of monohydroxylated fatty acyl moieties, 7:50
- Mixture analysis by dissociation and neutralization-reionization, characterization of C₃H₆O⁺ on from 2-methoxyethanol and, 6:1030
- Mixtures, potential application in analysis of, a mathematical study of sample modulation at a membrane inlet mass spectrometer, 7:93
- MMA. *See* Mass measurement accuracy
- MMFF, in ion mobility spectroscopy of charge-solvated vs salt bridge structures of protonated trimers, 16:1009
- MN⁺, excited, superelastic collisions of electrons with, 1:192
- Mn(III) and Fe(III) tetraarylporphyrin complexes, solution and gas phase chemistry of
fast-atom bombardment mass spectrometry, generation of molecular signals, 4:249
by fast-atom bombardment mass spectrometry, synthesis and characterization of molecular complexes, 4:255
- MNM. *See* Multiple neutral loss monitoring
- Mobile phase components, high performance liquid chromatography, on sensitivity in negative atmospheric pressure chemical ionization liquid chromatography-mass spectrometry, 7:1059
- Mobile phases
complementary, reversed-phase microbore liquid chromatography/electrospray mass spectrometry, and sequential

- exoglycosidase digestion, characterization of protein N-glycosylation by, 5:350
- VUV lamps and, detection of PAH compounds, APPI-MS, 18:589
- Mobility resolution spectrometer, ammonia and hydrazine complexes with ketones, mobility resolution and mass analysis of ions, 18:940
- Model for energy transfer in inelastic molecular collisions, applicable at steady state or non-steady state and for an arbitrary distribution of collision energies, 14:1348
- Model large polypeptide ions, RRKM calculations on, effect of ion size on rate of dissociation, 4:11
- Model matrix for molecular secondary ion mass spectrometry, biological soft tissue, doped gelatin films, 4:419
- Model pulsed bombardment ionization method, study of time-dependent desorption using, improved detection limits for fast atom bombardment mass spectrometry, 3:637
- Modeling data, in protein-ligand interactions, 15:388
- Modeling deuterium exchange behavior, of ERK2 using pepsin mapping to probe secondary structure, 10:685
- Modeling of glow discharge sources, with flat and pin cathodes and implications for mass spectrometric analysis, 8:1021
- Modeling the maximum charge state, of arginine-containing peptide ions formed by electrospray ionization, 7:972
- Modified base, small oligonucleotides with, ion trap tandem mass spectrometry applied to, 5:740
- Modified neglect of differential overlap bond orders and ab initio and modified neglect of differential overlap energy partitioning, application of, effect of protonation site on bond strengths in simple peptides, 5:704
- Modified neglect of differential overlap energy partitioning and ab initio and modified neglect of differential overlap bond orders, application of, the effect of protonation site on bond strengths in simple peptides, 5:704
- Modified zeolites for mass spectral analysis of tributyl phosphate using in situ silver ion chemical ionization, 8:718
- Moenomycin, characterization by multistage MALDI-IT/RTOF-MS and LC-ESI-IT-MS, 17:1081
- Molecular anions of SF₆ and C₇F₁₄, decomposition reactions of, cluster-assisted, 1:371
- Molecular beams, supersonic, LC/MS with, 11:587
- Molecular complexes, synthesis and characterization of, study of the solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes by fast-atom bombardment mass spectrometry, 4:255
- Molecular depth profiling
comparison of gold and cobalt ion cluster projectiles, SIMS, 18:406
by MS, Langmuir-Blodgett multilayer model for, 19:96
- Molecular dications, gas-phase
formed from neutral organics in solution, via the controlled-current electrolytic process inherent to electrospray, 7:157
formed from neutral organics in solution via chemical electron-transfer reactions by using electrospray ionization mass spectrometry, 5:689
- Molecular dissociations, new tandem mass spectrometer for, 3:487
- Molecular dynamics, fragmentation of oxidized peptides analyzed by RRKM modeling, SID and, 20:1579
- Molecular dynamics simulation, protein interface footprinting quantified by hydroxyl radical oxidation and, 19:1686
- Molecular dynamics simulations
gas-phase stability of G-quadruplex DNA using ESI-MS and, 18:1760
influence of amino acid composition on CID, model peptides, 18:1625
- Molecular energetics, temporal and spatial variations in, millisecond pulsed radio frequency glow discharge-TOF-MS, 15:87
- Molecular ion derived from fatty acid methyl esters, low-energy tandem mass spectrometry of, novel method for analysis of branched-chain fatty acids, 4:223
- Molecular ion fragmentation and its effects on mass isotopomer abundances of fatty acid methyl esters ionized by electron impact, 10:430
- Molecular ionization, volumetric intensity dependence on, in laser focus, 16:82
- Molecular ions, dissociation of, surface-induced, quadrupole ion trap mass spectrometer, 2:487
- Molecular mass analysis, of antibodies by on-line SEC-MS, 19:502
- Molecular mass determination of saturated hydrocarbons: reactivity of n5-cyclopentadienylcobalt ion CpCo^{•+} and linear alkanes up to C-30, 14:51
- Molecular mass distribution, quantitative measurement by MS, 20:1638
- Molecular mass measurement of intact ribonucleic acids via electrospray ionization quadrupole mass spectrometry, 6:27
- Molecular mass spectrometry, AP-MIPI for, 9:42
- Molecular modeling, to study IgG nonenzymic glycation, 11:153
- Molecular orbital, considerations in probing the stereoselective dissociations of cobalt-coordinated hexamine monosaccharides, 12:517
- Molecular orbital calculations and mass spectrometry, pyrrolo[1,2-a][1,4]benzodiazepin-4-one,-6-one, and -4,6-dione, characterization and differentiation of heterocyclic isomers, 7:653
- Molecular parameters, correlations of relative sensitivities in gas chromatography electron ionization mass spectrometry with, 1:397
- Molecular probe, for the interaction of methanol and protonated peptides, gas-phase hydrogen/deuterium exchange as, 5:526
- Molecular scanner, peptide mass spectra on, visualization and analysis of, 13:221
- Molecular secondary ion mass spectrometry studies of biological soft tissue, model matrix for, doped gelatine films as, 4:419
- Molecular signals, study of the solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes by fast-atom bombardment mass spectrometry, 4:249
- Molecular size, probed by vacuum ultraviolet photoionization, ion decompensation *versus*, 2:103
- Molecular species analysis of arachidonate containing glycerophosphocholines by tandem mass spectrometry, 2:45
- Molecular structure of silsesquioxanes determined by MALDI-TOFMS, 10:224
- Molecular weight
assignment of peptides in complex mixtures, application of electrospray mass spectrometry and matrix-assisted laser desorption ionization time-of-flight mass spectrometry for, 4:670
determination of oligomers, MALDI techniques in, 9:282, 9:651
determination of polymers, by MALDI-TOFMS, accuracy of, 9:275
mass spectrometrically measured, determination of oligonucleotide composition from, 4:204
of proteins, up to 80 ku, determination, high accuracy, by ion spray mass spectrometry, 2:387
- Molecular weight determination, use of β -cyclodextrin in, low-molecular weight explosives, MALDI-TOF MS, 17:189
- Molecular weight distributions, poly(dimethyl siloxane) polymers, using TOF-SIMS, 20:1562

- Molecule fragmentation, when sliding on surfaces, velocity and, 10:471
- Molecule-ion chemistry, gas-phase, of borate and boronate esters, 5:169
- Molecule-ion collisions, in the vacuum chamber of an electrospray time-of-flight mass spectrometer, mass spectra of proteins, 7:342
- Molecule-ion equilibria, how and why, 3:1
- Molecule-ion reaction kinetics
 probing trapped ion energies via Fourier transform ion cyclotron resonance mass spectrometry, 3:727
 quadrupole ion trap mass spectrometry, 3:716
- Molecule-ion reaction studies, flowing afterglow technique, early development of, 3:479
- Molecule-ion reaction under chemical ionization, gas-phase halo alkylation of C₆₀-fullerene by, 4:894
- Molecule-ion reactions
 for the differentiation of phospholipid classes, fast atom bombardment mass spectrometry employing, 2:470
 with dimethyl ether, selective detection of the tolyl cation among other isomers by, 2:270
 elucidating the structures of isomeric silylenium ions (SiC₃H⁺9, SiC₄H⁺11, SiC₅H⁺13), 2:278
 functional group-selective, of ethylene glycol and its monomethyl and dimethyl ethers, 4:97
 gas-phase, of the CH₃O⁺ cation, 3:515
 investigation of the gas-phase structure of electrosprayed proteins using, 5:207
 in methylamine and dimethylamine and trimethylamine systems, 2:220
 in mixtures of trimethylaluminum and methylamines, 4:111
 of oxygenated chemical ionization reagents with vincamine, 7:250
 selective, of lactams with dimethyl ether ions, 4:352
- Molecules
 highly functionalized, in fast atom bombardment, protonation and fragmentation of, mechanistic considerations of, high resolution mass spectrometry and tandem mass spectrometry analysis of the ions formed by fast atom bombardment of digoxin and related cardiac glycosides, 1:455
 isotopic fine structures, hierarchical algorithm for calculation, 19:1868
 multiphoton ionization, comparison between femtosecond and nanosecond laser pulse ionization efficiency, 6:1059
 small
 as substrates for adsorption/desorption in ²⁵²Cf plasma desorption mass spectrometry, 2:29
 UTLC-AP-MALDI-MS of, 16:906
- Molybdenites, terrestrial, isotope ratios of molybdenum in, measurement of, high accuracy, 3:10
- Molybdenum, isotope ratios of, in terrestrial molybdenites, measurement of, high accuracy, 3:10
- Molybdenum disulfide, nanoflakes, closed-caged clusters in gaseous and condensed phases, 18:2191
- Moment theory, for quadrupole ion traps, 17:889
- Monitor for displaying real-time images of low energy ion beams, 2:226
- Monitoring, real-time, atmospheric pollutants, ion trap detector used for, performance of, the effects of air and helium on, 4:774
- Monobromopolychlorodibenzo-p-dioxins, trace in environmental samples, application of gas chromatography high-resolution mass spectrometry to, 3:248
- Monoclonal antibodies
 epitope mapping of, by MS, 9:208
 intact, LTQ Orbitrap hybrid analysis of, 20:1415
 reduced, analysis using SEC-MS, 20:2258
 trap for, reversed-phase HPLC and ion trap MS in, 16:307, 16:797 (erratum)
- Monocyte chemoattractant protein 1, effects of sulfate position on heparin octasaccharide binding to, MS/MS, 17:1114
- Monohydroxylated fatty acyl moieties, analysis by fast-atom bombardment tandem mass spectrometry of phosphatidylcholine isolated from heart mitochondrial fractions, 7:50
- Monoisotopic masses and ion populations, for large biomolecules from resolved isotopic distributions, 6:229
- Monolayer surfaces, self-assembled
 fluorinated, chemical modification of by low energy reactive ion bombardment, 6:187
 laser desorption ion trap mass spectrometry of, 4:968
 reactions of hyperthermal ion beams at, XeF⁺, IF⁺, and other unusual ions generated by, 4:938
- Monomeric metaphosphate anion, tropylium, chlorine isotopic abundances, and Conestoga wagon theory, 4:761
- Monomethyl and dimethyl ethers, ethylene glycol, functional group-selective ion-molecule reactions of, 4:97
- Mononucleotides
 5'- and 3'-, gas-phase hydrogen/deuterium exchange, quadrupole ion trap, 18:724
 gas phase proton affinities of, as determined by the kinetic method, 11:24
 positively charged, H/D exchange of, FTICR-MS in, 12:268
- Monophosphates, hexose, analysis by ion-exchange chromatography and ESI-MS, 17:104
- Monosaccharides
 gas phase acidities and chloride affinities, ranking of, 12:1193
 glycoprotein, analysis by LC-MS/MS, 20:1224
 hexosamine, cobalt-coordinated, stereoselective dissociations of, 12:517
 internal residues, loss of, 8:43
 isobaric, distinguishing by ion pairing and MS/MS, 16:1827
 losses, on collision-induced dissociation mass spectra of anthracycline aminodisaccharides, 9:710
- Monte Carlo simulations, macromolecular ionization by nano-ESI, 19:1098
- Montmorillonite, oligonucleotides synthesized on, direct MALDI analysis, 17:1265
- Morphinans, ESI-MS/MS investigations of, 14:1262
- Mosses, *Sphagnum* and *Polytrichum*, carbon stable isotope analysis of organic matter and nitrocellulose, 18:1453
- Mouse switch variant antibodies, characterization of, by matrix-assisted laser desorption ionization mass spectrometry and electrospray ionization mass spectrometry, 7:707
- MoWeD, computer program to rapidly deconvolute low resolution LC/MS runs to determine component molecular weights, 12:599
- mPC. *See* Membrane preconcentration
- MRP8 and MRP14
 Ca²⁺-induced tetramers of, ESI-MS confirmation of, 11:780
 calcium-induced tetramers of, UV MALDI-MS of, 10:1124
- MS. *See* Mass spectrometry
- MS align software, genetic algorithm feature extraction, alignment of LC-MS and LC-MS/MS data sets, 18:1835
- MS/MS. *See* Tandem mass spectrometry
- MSAD. *See* Multipole storage assisted dissociation
- MS2Assign, automated assignment and nomenclature of MS/MS of chemically crosslinked peptides, 14:834
- MS2Grouper, as algorithm for proteomic MS/MS, 16:1250
- MSn, cyclic protonated peptide and metal complex characterization by, 15:1039
- Multi-collector double focusing inductively-coupled plasma mass spectrometry, mass discrimination of IRMM-007 series using, 17:1412

- Multi imaging mass spectrometry, of 13C-oleate transport in adipocytes, 15:1572
- Multinode detection in ESI-TOFMS, 9:1325
- Multichannel plasma-source mass spectrometer, novel, 7:458
- Multidimensional nano-HPLC for analysis of protein complexes, 14:1003
- Multidimensional protein identification technology (MudPIT), of heart tissue, 16:1207
- Multifunctional alcohols, including triacylglycerols, fragmentation of conjugate bases of esters derived from, 6:498
- Multiparticle simulation of ion motion, in the ion trap mass spectrometer, resonant and direct current pulse excitation, 6:57
- Multipass optical arrangement, infrared multiple photon dissociation in the quadrupole ion trap via, 5:886
- Multiphoton dissociation
 - collision-induced and infrared, on M(acetone)₂⁺ (M=Al, Fe, Co, Cu, ScO) in the gas phase, 4:135
 - infrared, in external ion reservoir, 14:1413
- Multiphoton ionization spectroscopy, Fourier transform ion cyclotron resonance detection, 7:1018
- Multiple inhibitor screening and kinetic constant, determinations for yeast hexokinase using MS-based assays, 14:173
- Multiple neutral loss monitoring, method for post-translational protein modification screening, 17:307
- Multiple scan modes, in the hybrid tandem mass spectrometric screening and characterization of the glutathione conjugate of 2-furamide, 2:55
- Multiple stage mass spectrometry
 - analysis of complex oligosaccharide antibiotics (everninomycins) in quadrupole ion trap, 13:1313
 - determination of daughter ion formulas by, 1:166
 - kinetic method combined with, SiCl₃⁺ and SiCl⁺ affinities for pyridines determined by using, agostic effects in the gas-phase, 7:198
 - pentaquadrupole
 - for generation and characterization of gas-phase ionic species. The case of the PyC₂H₅⁺ isomers, 7:1126
 - polar [4+2⁺] Diels-Alder cycloaddition to, gas stage ions and, 6:1
 - in structural characterization of organophosphorus compounds, 4:125
- Multiply charged ions, resolved isotopic peaks from, automated assignment of, 6:52
- Multiply charged oligonucleotides with a modified base, small, ion trap tandem mass spectrometry applied to, 5:740
- Multiply charged polypeptides, dissociation of, high-energy collision-induced, produced by electrospray, 4:652
- Multiply charged proteins, high mass-to-charge ratio quadrupole mass spectrometer, proton transfer reaction studies of, 5:1064
- Multiply protonated ions, high energy collisional activation of, complete sequence confirmation of large peptides by, 6:234
- Multipole ion guides, driving by high voltage RF oscillator, 13:1370
- Multipole storage assisted dissociation (MSAD), 11:1
 - mechanistic studies of, 11:210
- Multiquadrupole mass spectrometers, calibration point for electron ionization MS/MS spectra measured with, 5:576
- Multisample probe for fast-atom bombardment/mass spectrometry, 4:504
- Multispectral identification of alkyl and chloroalkyl phosphates from an industrial effluent, 2:419
- Multistep tandem mass spectrometry for sequencing cyclic peptides in an ion-trap mass spectrometer, 10:732
- Multivariate analysis, in lactate dehydrogenase classification, 14:236
- Mutagenic/carcinogenic potential of environmental contaminants, estimation of, by ion-molecule reactions and tandem mass spectrometry, 1:110
- Mutations
 - double, hemoglobin Villeparisis β77(EF1) His → Tyr, β80 (EF₄) Asn → Ser, use of combined mass spectrometry methods for the characterization of a new variant of human hemoglobin, 7:163
 - point, changes in protein hydrogen exchange rates that result from, mass spectrometric measurement of, 7:515
- Multiplexed multilinear mass spectrometry, co-occurrence of boundary and resonance ejection, ion traps, 17:56
- Mutual storage mode, in hybrid linear ion trap, 16:71
- Mycobacterial glycopeptidolipid antigens, fast atom bombardment mass spectrometry of, structural characterization by charge remote fragmentation, 3:656
- Mycobacterium*, lipoarabinomannans from, ESI-MS characterization of, 16:1109
- Mycobacterium tuberculosis*, secreted proteins from, ECD-MS of, 14:253
- Myelin basic protein, capillary electrophoresis/MS with ion trap/reflection TOFMS detector, 9:701
- Myelin sheath, bovine, characterization of P0 by MALDI-TOF/TOF MS, 17:117
- Myeloid leukemia, acute, histone modifications in, HPLC and MS of, 15:77
- Myoglobin
 - acid-denatured, stepwise refolding of, evidence from electrospray mass spectrometry, 4:638
 - collision cross sections of, with Ne, Ar, and Kr, 8:681
 - denaturation of, in laser spray, 16:860
 - hemoglobin, and cytochrome c, heme binding in, studies on, by ion spray mass spectrometry, 4:631
- Myoglobin ions, gaseous, stored at greater than 300 K, 5:324
- Myoglobin unfolding, pH effect on, ESI-MS of, 11:312
- Myosin VI complexes, noncovalent, ESI-MS of, 16:1367
- Myristoylated alanine-rich C kinase protein, identification of peptide phosphorylation sites, ECD FTICR MS/MS, 18:2137
- N
- N-Acetylhexosamine, diastereomeric, quantification of, MS3 spectra for, 11:1086
- N-alkylnicotinium halides, class of cationic matrix additives for enhancing the sensitivity in negative ion fast-atom bombardment mass spectrometry of polyanionic analytes, 5:935
- N-benzoylpeptides, deprotonated, fragmentation of, 15:446
- N-C₃H₇CNH⁺, ion-neutral complexes in gas-phase dissociations of protonated nitriles, 2:178
- N²-deoxyguanosine and C8 adducts of 2-acetylaminofluorene, isomeric, by fast-atom bombardment and tandem mass spectrometry, 5:58
- N-formylkynurenine, from tryptophan, identification by MS, 14:406
- N-Glycans, multiantennary, sequencing with ESI-quadrupole TOF-MS/MS, 13:1138
- N-glycosylation, protein, by reversed-phase microbore liquid chromatography/electrospray mass spectrometry, complementary mobile phases, and sequential exoglycosidase digestion, 5:350
- N-heterocycle synthesis, in gas phase, in arylnitrenium ion generation, 15:1679
- N-Phenyl-1,4-phenylenediimine, surface-assisted reduction of, in APCI and APPI, 13:109
- N-(substituted benzalamino)phthalimide radical cations, decom-

- positions of, embody ion-neutral complexes and Stevenson's rule, 5:837
- N-terminal glutamine, studies on a prion protein peptide, tandem mass spectrometry of peptides with, 1:258
- N-terminal processing, of phospholipase A₂, LC-MS/MS of, 15:1780
- Nafion-coated bismuth film electrode, NCMFE and, for ASV with ICP-MS, 17:945
- Nafion-coated mercury film electrode, NCBFE and, for anodic stripping voltammetry with ICP-MS, 17:945
- Nano-electrospray emitter, electrochemical processes in, 12:853
- Nano-electrospray ionization quadrupole time-of-flight mass spectrometry fucosylation study, of low abundance species in complex mixtures, 12:964
- Nano-electrospray-MS/MS, nano-LC combined with, to detect estrogen DNA-adducts in breast tumor and healthy tissue, 14:482
- Nano-HPLC, multidimensional, for protein complex analysis, 14:1003
- Nano-liquid chromatography, nano-ES MS/MS combined with, to detect estrogen DNA-adducts in breast tumor and healthy tissue, 14:482
- Nano-liquid chromatography/microelectrospray ionization, FTICR-MS with, in peptide mass fingerprint analysis, 14:553
- Nanobioparticles, analysis using nanoES GEMMA and PDMA, 19:1062
- Nanocalorimetry, hydrated ion, nonergodicity in ECD investigated by, 18:1217
- Nanoelectrospray
chip-based, evaluation of, parameters for protein identification, 18:1714
combined with a quadrupole ion trap for the analysis of peptides and protein digests, 7:150
more than minimized-flow ESI source, 10:300
- Nanoelectrospray/collision-induced dissociation mass spectrometry, sequencing of proteolytic peptides by, 11:673
- Nanoelectrospray ionization
efficiency of, 16:1702
nonspecific protein-carbohydrate complexes produced by, BIRD of, 16:1583
- Nanoelectrospray ionization Fourier transform ion cyclotron resonance (FTICR) mass spectrometry
glycosphingolipid mixtures screening and sequencing by, 16:571
of Shiga toxin B subunit stability in solution and gas phase, 16:1957
- Nanoelectrospray ionization-mass spectrometry, of proteins, 8:483
- Nanoelectrospray ionization tandem mass spectrometry (nano-ESI-MS/MS)
automated, of cerebellar gangliosides, 15:1649
from chromatographic pipette tips, 15:1333
- Nanoelectrospray mass spectrometry
AA substitutions mapped by, avian influenza virus (H5N1) matrix protein 1, 20:312
of carcinogen-peptide adducts, 9:202
current-controlled, 20:1334
electrochemical properties of, 10:309
to monitor protein reactions, 10:112
solution flow rates in, analyte ion signals and, 14:492
in structural analysis of oligosaccharides in human milk, 13:1331, 13:1341
- Nanoelectrospray triple quadrupole mass spectrometry, protein charge state distribution shifts in, 16:409
- Nanoelectrospray with controllable flow rate, for mass analysis of submicroliter protein samples, 11:94
- Nanoflow electrospray ionization-MS, particle discriminator interface for, 14:1236
- Nanoflow liquid chromatography-ion mobility mass spectrometry, of protein mixtures, 15:1341
- Nanoliter electrospray mass spectrometry, long-lived metallized tips for, 7:1270
- NanoMate, in pharmaceutical compound quantitative analysis, 16:363, 16:798 (erratum)
- Nanoparticles, magnetic, zirconium phosphonate-modified, selective phosphopeptide capture by, 19:1176
- Nanoscale packed capillary liquid chromatography (75 m id), to the analysis of macrolide antibiotics, 3:563
- Nanoscale separations combined with electrospray ionization mass spectrometry, sulfonamide determination, 3:139
- Nanosecond laser pulse ionization efficiency, for multiphoton ionization of molecules, 6:1059
- Nanospray, protein surface reactions produced by, 16:1325
- Nanospray emitters, polyaniline coating for, 11:659
- Nanospray tips, microfabricated, nib-like design for, 15:407
- Nanotubes
carbon, 2,5-dihydroxybenzoyl hydrazine derivative, use in trace peptide analysis, MALDI, 17:1023
oxidized carbon, for MALDI-TOF MS analysis, 16:883
oxidized carbon matrixes, monitoring enzyme reactions and screening inhibitors, MALDI-TOF-MS with, 17:1616
- Naphthalene-2,3 dicarboxaldehyde, enkephalin reaction with, for LC/LIF/MS analysis, 18:1706
- Nazarof cyclization, protonated 2-methoxy and 2-hydroxychalcone, 20:805
- N–C_α bond cleavage of peptide backbone via hydrogen abstraction, 12:1044
- NCBFE. *See* Nafion-coated bismuth film electrode
- NCMFE. *See* Nafion-coated mercury film electrode
- ND₃ or D₂O, hydrogen/deuterium exchange between [m–1]⁺ ions of chlorinated benzenes and, 5:282
- ND₃ with protonated aromatic amines 3, H/D exchange reactions, reactive collisions in quadrupole cells and, 6:19
- NDA. *See* Naphthalene-18:2,18:3 dicarboxaldehyde
- Near-UV-induced MALDI as a function of wavelength, 9:885
- Nebulizer, dual oscillating, in AP-MIPI MS of underivatized amino acids, 12:117
- Nebulizer-assisted electrospray ion sources, improvement by atmospheric pressure ion lens, 13:906
- Nebulizer interface, heated, liquid introduction with, atmospheric-pressure chemical ionization mass spectrometry, characterization of flame-generated C₁₀ to C₁₆₀ polycyclic aromatic hydrocarbons by, 7:276
- Nefazodone, liver microsome metabolism, analysis by linear ion trap/orbitrap MS, 17:363
- Negative atmospheric pressure chemical ionization liquid chromatography-mass spectrometry, sensitivity in, effect of high-performance liquid chromatography mobile phase components on, 7:1059
- Negative chemical ionization in quadrupole ion trap mass spectrometry, effects of applied voltages and reaction times, 5:757
- Negative chemical ionization mass spectrometric fragmentation, electron capture, and calculated internal energies for polychlorinated biphenyls, correlation between, 5:44
- Negative chemical ionization with oxygen, dibenzothiophene subjected to, oxidation reactions of, 4:949
- Negative ion-atmospheric pressure ionization, NI-DART and, ionization mechanisms of, 20:42
- Negative ion atmospheric pressure photoionization mass spectrometry (APPI-MS), 15:203
- Negative ion chemical ionization, polarizable stationary phase, and mass-resolved chromatography, high-temperature

- gas chromatography/mass spectrometry with, high-resolution triglycerol mixture analysis using, 7:350
- Negative ion chemical ionization mass spectroscopy, anthraquinone-2-carbonyl chloride as an alcohol derivatization reagent for, characterization of, 6:148
- Negative ion chemical ionization (NICI), in study of O- reactions with methyl benzoate, 12:694
- Negative ion-direct analysis in real time, NI-APPI and, ionization mechanisms of, 20:42
- Negative ion electrospray mass spectrometry
analysis of lipid hydroperoxides and long-chain conjugated keto acids by, 6:1190
analyte charge state, signal intensity, and stability in, solvent effect on, implications for the mechanism of negative ion formation, 4:546
nucleotides, ionization from water solution with SF₆ discharge suppression, 4:289
- Negative ion electrospray of bromo- and chloroacetic acids and an evaluation of exact mass measurements with benchtop TOFMS, 11:809
- Negative ion electrospray tandem mass spectrometric structural characterization of leukotriene B₄ (LTB₄) and LTB₄-derived metabolites, 7:129
- Negative ion fast-atom bombardment mass spectrometry
analysis of long-chain fatty acyl coenzyme A thioesters by, 5:416
of polyanionic analytes, class of cationic matrix additives for, N-alkylnicotinium halides, 5:935
- Negative ion fast-atom bombardment tandem mass spectrometry, to determine sulfate and linkage position in glycosaminoglycan-derived disaccharides, 3:797
- Negative ion formation
in electrospray mass spectrometry, 4:578
mechanism of, solvent effect on analyte charge state, signal intensity, and stability in negative ion electrospray mass spectrometry, 4:546
- Negative-ion formation in the explosives RDX, PETN, and TNT by using the reversal electron attachment detection technique, 3:243
- Negative ion laser desorption ionization, time-of-flight mass spectrometry of nitrated polycyclic aromatic hydrocarbons, 8:630
- Negative ion liquid chromatography-electrospray mass spectrometry, neutral O-linked oligosaccharide alditol structural determination by, 15:659
- Negative ion MALDI-TOF, post-source decay calibration by using fibrinopeptide B, 9:540
- Negative ion mass spectra, electron capture, not reproducible, 4:270
- Negative ion mass spectrometry, cyclodepsipeptide identification using, 20:2221
- Negative-ion matrix-assisted laser desorption mass spectrometry, analysis of oligodeoxynucleotides by, 4:955
- Negative ion mode, neutral lactooligosaccharide fragmentation in UV-MALDI-TOF and UV-MALDI ion trap/TOF MS, 17:67
- Negative ion tandem mass spectrometry, of leukotriene E₄ and LTE₄ metabolites, identification of LTE₄ in human urine, 2:314
- Negative ionization, electron capture, in the detection of chemically oxidized dexamethasone in human plasma, comparison of tandem and conventional mass spectrometry using, 1:341
- Negative signature mass algorithm, 14:1032
- Neglect, modified, of differential overlap bond orders and ab initio and modified neglect of differential overlap energy partitioning, 5:704
- Neighboring group participation, in the electrospray ionization tandem mass spectra of polyamine toxins of spiders, alpha,omega-diaminoalkane compounds, 6:634
- Neisseria gonorrhoeae* lipooligosaccharides, FTICR-MS in characterization of, 13:571
- Nemorosone, degradation profile by ESI-MS, 20:1688
- Neoglycolipids, synthetic, ESI MS/MS analysis of, 16:565
- Neon, collision cross sections of myoglobin and cytochrome c with, 8:681
- Neosaitoxin, protonated, gas phase dissociation reactions of, 15:462
- Nephelopsis oscura*, hemoglobin from, subunit structure of, ESI-MS of, 15:22
- NEREUS, lake water testing of chemistry with portable and AUV-based MS, 19:1403
- Nerve agents, chemical, APPIS applications to detection of, 20:2093
- Nerve growth factor, ATP binding to, detected by MALDI-TOF and ESI-MS, 19:91
- Networking mass spectrometer data systems, for improved productivity and electronic archiving of data, 4:742
- NeuAc. *See* N-Acetyl neuraminic acid oligomers
- Neural network models, for bacterial classification, 13:10
- Neural networks, artificial, for ppb mass accuracy in FT-ICR/MS, 20:1303
- Neuromelanin, pyrolysis products of, from substantia nigra, GC-MS analysis of, 15:920
- Neuromuscular blocking agents, quaternary ammonium, an analytical strategy using LC-MS/MS, 17:1456
- Neuropeptide research, MS and hyphenated techniques in, 14:81, 14:287
- Neuropeptide Y, gas-phase conformation probed by CID, 17:786
- Neuropeptides
crustacean hemolymph, measurement by MALDI MS, 20:708
mapping of lipids and, by 3D MSI, crustacean brain, 20:1068
- Neutral chemical ionization gas, reaction of analyte ions with, 3:549
- Neutral counterpart generation by neutralization-reionization mass spectrometry, structure of C₅H₅RFe⁺ (R=F, Cl, Br, I, O, OH, OCH₃, H₆H₅, H) ions in the gas phase, 6:1143
- Neutral desorption enclosures, sampling explosives on skin, using EESI-MS and, 20:719
- Neutral loss, constant, oligopeptide fragmentation viewed as, 1:336
- Neutral organic molecules including fullerenes, applications of electrospray ionization mass spectrometry to, 4:596
- Neutral organics in solution, gas-phase molecular dications formed from
via chemical electron-transfer reactions by using electrospray ionization mass spectrometry, 5:689
via the controlled-current electrolytic process inherent to electrospray, 7:157
- Neutral partner, polarity of, effect on complex-mediated reactions, isomeric ion-neutral complexes generated from ionized 2-methylpropanol and n-butanol, 3:409
- Neutralization, of methyl cation via chemical reactions in low energy ion-surface collisions with fluorocarbon and hydrocarbon self-assembled monolayer films, 13:1151
- Neutralization agents, organic, for neutralization-reionization mass spectrometry, 3:108
- Neutralization-reionization
of alkenylammonium cations, intramolecular NNH⁺ · · C=C interactions in cations and hypervalent ammonium radicals, 8:1111
distonic ion CH₂CH₂CH⁺OH, keto ion CH₃CH₂CH₂O⁺, enol ion CH₃CH=CHOH⁺, and related C₃H₆O⁺ radical cations, 7:573

- internal energy distributions of tungsten hexacarbonyl ions after, 5:1093
- Neutralization-reionization mass spectrometry
 angle-resolved, 2:492
 hydrogen bonding in transient bifunctional hypervalent radicals by, 6:1004
 hypervalent pyrrolidinium radicals by, 9:242
 modeling deoxyribose radicals by, 15:1055, 15:1068
 neutral counterpart generation by, structure of $C_5H_5RFe^+$ (R=F, Cl, Br, I, O, OH, OCH_3 , H_6H_5 , H) ions in the gas phase, 6:1143
 organic neutralization agents for, 3:108
 of protonation sites in methyl nitrates, 11:380
- Neutralization-reionization mixture, characterization of $C_3H_6O^+$ on from 2-methoxyethanol and, 6:1030
- Neutralization-reionization studies, novel tandem quadrupole-acceleration-deceleration mass spectrometer for, 3:493
- Neutrals, generated by matrix-assisted laser desorption, gas-phase cationization and protonation of, 4:393
- New mass spectrograph, 4:372
- NI-APPI. *See* Negative ion-atmospheric pressure ionization
- NI-DART. *See* Negative ion-direct analysis in real time
- NICI. *See* Negative ion chemical ionization
- Nickel complex ions, reactions with acetonitrile in gas phase, 15:1128
- Nickel complexes, MS behavior of, 16:94
- Nier, Al, Professor, and his influence on mass spectrometry, 3:685
- NIST Polymer Mass Spectrometry Workshop, fourth annual, report on, review, 17:280
- Nitramines, anion fragmentation, GC/MS, 18:835
- Nitrate adducts, of carbohydrates, 16:622
- Nitrate-anion complex, monopositive, intrinsic hydration of, 15:777
- Nitric oxide-assisted atmospheric pressure corona discharge ionization for the analysis of automobile hydrocarbon emission species, 5:1107
- Nitric oxide synthase oxygenase domain, collisional cooling within, 15:629
- Nitrile sulfides, ionized, gas phase interaction with pyridine, 14:241
- Nitriles, protonated, gas-phase dissociations of, ion-neutral complexes in, $n-C_3H_7CNH^+$ and $i-C_3H_7CNH^+$, 2:178
- Nitrilium and immonium ions in the gas stage, applications of multiple stage mass spectrometry in a pentaquadrupole, polar $[4+2^+]$ Diels-Alder cycloaddition to, 6:1
- Nitro groups
 adjacent, in the gas phase, on the stabilization of carbanions by, 5:159
 protonated, as a gas-phase electrophile, 18:2204
- Nitro *meso*-tetraphenylporphyrins, positional isomers of, MS/MS, 12:381
- Nitro-reduction, in arylnitrenium ion generation, 15:1679
- Nitro-substituted polynuclear aromatic hydrocarbons in airborne particulate matter and soil, sub-parts-per-billion determination by electron capture-tandem mass spectrometry, 7:1255
- Nitroaromatic compounds, denitration of, by arylnitrile radical cations, 11:33
- Nitrobenzyl alcohol-mete, and dipyrrocyanine dyes, beam-induced reaction between, in liquid-secondary-ion mass spectrometry, 5:1017
- Nitrocatechol glucuronides, APCI and ESI of, 10:537
- Nitrocellulose, carbon stable isotope analysis, *Sphagnum* and *Polytrichum* mosses, 18:1453
- Nitrogen
 effect on fluorescence yield of trapped gas-phase ions, 17:855
 nucleophilic substitution at, 12:139
- Nitrogen- and oxygen-containing molecules, small, reactions to the phenylium cation with, 7:473
- Nitrogen- and phosphorus-containing pesticides, mass spectrometric analysis of, by liquid chromatography-mass spectrometry, 5:655
- Nitrogen and sulfur heterocycles in petroleum fractions, selective chemical ionization of, by ion trap mass spectrometry, 4:322
- Nitrogen compounds, $CH_3OCH_2^+$ reactions with, 13:1028
- Nitrogen isotopes
 analyses by isotope-ratio-monitoring gas chromatography/mass spectrometry, 5:387
 carbon isotope ratios to, in cells, 15:478
- Nitrogen trifluoride, a new reactant gas in chemical reaction interface mass spectrometry for detection of phosphorus, deuterium, chlorine, and sulfur, 6:421
- Nitromethane, [M-H] anions of, structures of, 1:295
- Nitropyrenes, mutagenic, LC-MS/MS analysis of, 15:1853
- Nitrosation, of thiols and thioesters, in gas phase, 15:344
- Nitrosopeptides, S- and N-, analysis using electrospray MS/MS, 17:1725
- 3-Nitrosine, in biological tissues and fluids, quantification of, 11:578
- S-Nitrosylation sites, identification in proteins, by QTOF-MS, 19:1353
- Nitrotyrosine, peptides and proteins containing, MALDI-MS of, 12:438
- NO_3^+ , formed via glow discharge in atmospheric gases, structures of, 1:217
- N_2O and CO_2 , gas-phase bond strengths of, with the halide ions, 4:58
- Nod factors, nonreducing terminal residue of, 11:301
- Noise, chemical, in MALDI-MS, 13:129
- Noise discrimination
 Signal and noise evaluation in ESI IT MS, 20:1859
 signal filtering method for quantification and, FT-ICR MS, 20:1087
- Noise filtering, techniques for electrospray quadrupole-TOFMS, 14:766
- Noise reduction
 chemical, reagents for, LC-MS, 20:105
 by supplemental IR activation, ESI MS, 20:2206
- Nomenclature system, for labeling cyclic peptide fragments, 10:360
- Non-labile sites, H/D exchange at, 12:1163
- Non-steroidal anti-inflammatory drugs, ESI-MS/MS of, without chromatography, 9:255
- Nonclassical and classical ethyl structures, tandem mass spectrometric investigation of (C_2H_5) X^+ ions (X=Cl, Br, I), 5:814
- Noncovalent adduction in electrospray mass spectrometry, mass and charge state assignment for proteins and peptide mixtures via, 6:1175
- Noncovalent adducts of electrosprayed nucleotide ions, infrared photodissociation of, 7:209
- Noncovalent associations
 of dihydroxybenzoic acid with cyclic and linear oligosaccharides, MALDI-TOFMS and X-ray crystallographic study of, 11:228
 of glutathione S-transferase and ligands, study using ESI-quadrupole-TOFMS, 11:606
- Noncovalent complexes
 allophycocyanine localization in MALDI-preparations by CLSM, 17:1599
 to the avidin tetramer, by electrospray ionization mass spectrometry, 5:201

- to the avidin tetramer by electrospray ionization mass spectrometry, 5:201
- electrostatic interaction site identified by ECD/ETD, 20:176
- formed between minor groove-binding molecules and duplex DNA by electrospray ionization-mass spectrometry, 6:1154
- in glycosidic bond fragmentation in gas phase, 15:715
- between margatoxin and the KV1.3 peptide ligands, model investigation using ion-spray mass spectrometry, 7:1075
- solution phase H/D exchange of, ESI-MS of, 12:795
- weak interactions analyzed by ESI-MS, 20:303
- Noncovalent enzyme-inhibitor complexes, and metal binding stoichiometry of matrixlysin by electrospray ionization mass spectrometry, 6:1105
- Noncovalent FKBP-FK506 and FKBP-rapamycin complexes, detection of by capillary electrophoresis-mass spectrometry and capillary electrophoresis-tandem mass spectrometry, 6:85
- Noncovalent inhibitors, association constants of adenylate kinase with, 14:442
- Noncovalent interactions
- exploring "intensity fading" by MALDI-TOF-MS, 18:359
 - protein-protein and protein-ligand, application of electrospray mass spectrometry in, 4:624
 - solution phase, ESI-MS diffusion measurements of, 14:430
- Noncovalent protein-ligand complexes, confirming binding sites and tertiary structure of, MS for, 14:460
- Noncovalent streptavidin complexes, tetrameric ions of formed by electrospray ionization, dissociation of, 6:459
- Nonergodicity, in ECD, hydrated ion nanocalorimetry, 18:1217
- Nonionic surfactants, collision-induced dissociation mass spectrometry, following direct supercritical fluid injection, 3:150
- Nonlinear ion acceleration, for improved space focusing in TOFMS, 10:1067
- Nonlinear resonance effects, during ion storage in quadrupole ion trap, 4:917
- Nonresonance excitation, application of, to ion trap tandem mass spectrometry and selected ejection chemical ionization, 7:668
- Nonspecific cleavage products, for protein sequence analysis as shown on calyculin isolated from human granulocytes, 12:1180
- Normal phase liquid chromatography/mass spectrometry, application of, by using coaxial continuous flow fast atom bombardment, 3:575
- Novel 3-hydroxylated leukotriene B₄ metabolites, from ethanol-treated rat hepatocytes, 3:762
- Novel electric ion resonance, cell design with high signal-to-noise ratio and low distortion for FTMS, 10:1000
- Novel interface, for variable flow nanoscale LC/MS/MS for improved proteome coverage, 13:760
- Novel [M-35]⁺ fragment ions, in the mass spectra of 2-alkyl-4,6-dinitrophenols, sequential ortho effects, 2:69
- Novel mass spectral fragmentation, heptafluorobutyl derivatives of acyl analogues of platelet-activating factor, 2:476
- Novel multichannel plasma-source mass spectrometer, 7:458
- Novel nonmetallized tip for EMS at nanoliter flow rate, 10:72
- Novel perfluoroacyl derivatives, of corticosteroids and related substances for potential use in quantitative mass chromatography mass spectrometry, 6:264
- Novel precursor ion, discovery method on a hybrid quadrupole orthogonal acceleration TOF-MS for studying protein phosphorylation, 13:792
- Novel rearranged ions, observed for protonated peptides via metastable decomposition in MALDI-TOFMS, 11:345
- Novel sheathless interface, for capillary electrophoresis/electrospray ionization mass spectrometry using an in-capillary electrode, 8:561
- Novel tandem quadrupole-acceleration-deceleration mass spectrometer, for neutralization-reionization studies, 3:493
- Novel tandem quadrupole mass analyzer, 10:1053
- Nozzle-skimmer region, in CID, in biopolymer sequencing, 12:846
- NRMS. *See* Neutralization-reionization mass spectrometry
- NSAIDs. *See* Non-steroidal anti-inflammatory drugs
- Nuclear magnetic resonance
- solid state, exploring solvent-free MALDI samples, 20:1906
 - solid state, MALDI-TOF MS and, analysis of polylactides, 20:67
- Nuclear receptor coactivator 1, noncovalent complexes, weak interactions analyzed by ESI-MS, 20:303
- Nucleic acid bases, protonated, microhydration, 20:1900
- Nucleic acid molecules, proton affinity of, kinetic method determination of, 15:1080
- Nucleic acids
- building a database of bifunctional probes for MS3D investigation of, 17:1570
 - metal cation-trimethylphosphate interactions with, 19:305
 - modified, in oligonucleotide fragmentation, 12:744
 - monoprotonated peptide, gas phase decomposition of, 11:615
 - peptide mono-deprotonated, complete sequencing of by CID, 12:805
 - regioselective backbone fragmentation of, 8:1257
- Nucleobases
- loss of, Arrhenius activation parameters for, 15:55
 - and nucleosides, electrospray ionization-mass spectrometry of, processes that affect, 7:1106
 - protonated, deamination, 18:2040
- Nucleobases in regioselective backbone fragmentation, of nucleic acids: molecular dynamics and MS/MS investigation on a model dinucleotide phosphotriester, 8:1257
- Nucleophiles
- anionic, fluorotrimethylsilane affinities of, 16:697
 - model biological, reactions of polycyclic aromatic hydrocarbon radical cations with, 7:628
 - pyridine, CH₂⁺ transfer to, means of producing alpha-distonic ions, 4:117
- Nucleophilic substitution, at nitrogen, 12:139
- Nucleophilic substitutions, reactions of halonitrobenzenes and dinitrobenzene with carbanions, gas phase, 18:1351
- Nucleoside adducts, separation and identification of, by CEC- μ ESI/MSI, 9:823
- Nucleoside-carcinogen adducts, identification of, fast atom bombardment mass spectral methods for, 3:360
- Nucleoside pyrophosphates
- glycosyl esters of, anomeric configuration, determination of, by fast-atom bombardment tandem mass spectrometry, 6:516
 - and polyisoprenyl phosphates, glycosyl esters of anomeric configuration of, by fast-atom bombardment tandem mass spectrometry, 7:541
- Nucleosides
- cytosine, self-assembling into dimers, evaluation with ESI-MS/MS, X-ray diffractometry, and theoretical calculations, 15:268, 15:436
 - electron-bound dimers of, 12:176
 - and nucleobases, electrospray ionization-mass spectrometry of, processes that affect, 7:1106
 - urinary, MALDI-TOF MS of, 16:940
- Nucleotide ions, electrosprayed, non-covalent adducts of, infrared photodissociation, 7:209
- Nucleotide peptides, *E. coli* microcin C7-C51 characterization by CID, 19:1187

Nucleotides

- ESI-MS detection of ribonuclease-nucleotide complexes, 19:1199
- negative ion electrospray mass spectrometry of, ionization from water solution with SF₆ discharge suppression, 4:289
- Nylon 66 filter disks, cyclic polyamide oligomer contamination from, LC/MS, 17:652
- Nylon-based membranes, proteins immobilized on, direct matrix-assisted laser desorption ionization mass spectrometric, 5:230

O

- O⁻, methyl benzoate reactions with, 12:694
- O₂⁻, gas-phase reactions of, with alkyl and aryl esters of benzenedicarboxylic acids, 5:990
- O⁻ and H₆H₅F, gas-phase reactions between, acidity of fluorobenzene and 1,4-difluorobenzene, 4:73
- O⁻ and OH⁻ chemical ionization, of some fatty acid methyl esters and triacylglycerols, 5:553
- O⁻ chemical ionization, of carbonyl compounds, 2:292
- O⁻ oxide radical anion, comparison of the reactions of, with simple aromatic compounds at atmospheric and at reduced pressure conditions, 5:367
- O-Acetyl esters, of glucosamine, differentiation of, 11:301
- O-Carbamoyl esters, of glucosamine, differentiation of, 11:301
- O-Diglycosides, flavonoid, glucose loss from, 11:136
- O-linked glycopeptides, structural elucidation of, by high energy collision-induced dissociation, 7:319
- ¹⁶O/¹⁸O peptide ion ratio, calculating for proteome relative quantification, 15:437
- O-rings, in vacuum system, plasticizer contamination from, in quadrupole ion trap mass spectrometer, 13:901
- ¹⁶O-to-¹⁸O exchange, trypsin-catalyzed, for comparative proteomics, 14:704
- Obituaries
 - Alexandra F. Dodonov, 17:113
 - Allan Maccoll, 10:782
 - Andrew Gans "Jack" Sharkey, Jr., 15:627
 - Austin Levy Wahrhaftig, 9:452
 - Ben S. Freiser, 10:917
 - Ben Sherman Freiser, 9:565
 - Frederick W. Lampe, 12:123
 - Harry J. Svec, 18:970
 - Isamu Matsumoto, 15:1689
 - Katheryn A.H. Resing, 20:1576
 - Richard A. Schaeffer, 16:796
 - Robert R. Squires, 10:798
 - Sharon G. Lias, 16:979
 - Victor L. Talroze, 15:1517
 - Vladimír Hanuš, 20:2352
 - Wim J. van der Hart, 20:1944
- 5-Oco-7-glutathionyl-8,11,14-eicosatrienoic acid (FOG7), ESI and MS/MS of, 12:763
- Octadecadienoic acids, homoconjugated, collisional activation of, with fast atom bombardment and tandem mass spectrometry, 1:183
- Octaethylporphyrin, collisionally activated decompositions of, 9:767
- Octafluoropentyl chloroformate, derivatization mediated by, ultratrace determination of hydrophilic compounds by, 10:1328
- Octaglycine, protonated, surface-induced dissociation, 20:939
- Octopamine, molecular ions of, CH₃NH₃⁺ and (CH₃)₂NH₂⁺ formation from, the role of distonic ions in, 1:104
- Octopole ion trap
 - linear, ion extraction from, SIMION analysis of, 13:1304
 - of transition metals, electrospray ionization study of, 8:32

Off-resonance

- irradiation sustained, capillary electrophoresis Fourier transform ion cyclotron resonance mass spectrometry with, for the characterization of protein and peptide mixtures, 5:894
- sustained in FTICR, 8:86
- Off-resonance excitation, linear ion traps, 20:443
- Offset ion lens, inductively coupled plasma mass spectrometry with an enlarged sampling orifice
 - ion trajectories and detector performance, 4:16
 - polyatomic ion interferences and matrix effects, 4:28
- OH⁻ chemical ionization, fatty acid methyl esters and triacylglycerols, 5:553
- Oils
 - essential components of, 8:671
 - whole, MALDI-TOFMS of, 10:983
- Olefin, halide ion clusters with, 12:144
- Oleuropein, localization of noncovalent binding site between amyloid- β -peptide and, ESI-FT-ICR MS, 19:1078
- Oleuropein, amyloid- β -peptide interaction with, ESI-MS, 17:568
- Oligoalanines, protonated, fragmentation of, 15:1810
- Oligodeoxynucleotide dications
 - ECD and IRMPD of, 14:23
 - electron capture and infrared multiphoton dissociation, 14:23
- Oligodeoxynucleotides
 - abasic site location in, by MS/MS, 13:1418
 - adducted to metal ions, ESI-MS/MS of, 12:550
 - analysis of, negative-ion matrix-assisted laser desorption mass spectrometry, 4:955
 - charged anions, transition metal complexes with, 19:281
 - differentiating 8-oxo-dG residues in, 13:195
 - double-stranded, gas phase stability of, 11:450
 - fragmentation mechanisms of, 12:580
 - fragmentation with 5-formyluracil or 2-aminoimidazolone, 13:1190
 - initial fragmentation in quadrupole ion trap, 16:1853
 - isomeric photomodified, MALDI and ESI, 10:329
 - MALDI-MS for, 11:854
 - metastable decay of, 12:180
 - photomodified
 - adducted with metal ions, fragmentation in ESI-ITMS, 12:1174
 - determination by exonuclease digestion, MALDI, and PDSMS, 12:1127
 - 2'-Oligodeoxynucleotides, melphalan-modified, miniaturized LC-ES MS/MS of, 15:568
- Oligomer samples, solubility of, MESIMS of, 16:1172
- Oligomeric peptides, intrinsic basicity of, effects of the alkyl side chain on proton transfer reactions, 6:91
- Oligomerization, of insulin analogs and mutants, SIMSTEX application to, 17:1526
- Oligomers
 - molecular weight determination, MALDI techniques in, 9:282, 9:651
 - polyaniline, MALDI analysis of, 15:893
 - protonated, fragmentation in gas phase, 20:60
- Oligonucleotide anions
 - BIRD of, 9:1117
 - deprotonated, loss of nucleobases from, Arrhenius activation parameters for, 15:55
 - multiply charged, proton transfer reactions of protonated pyridine with, gas phase, ion-ion reactions in, 6:529
- Oligonucleotide composition, from mass spectrometrically measured molecular weight, 4:204
- Oligonucleotide duplexes, stability determination of, gas and solution phase, 15:1354

- Oligonucleotide-metal complexes, fragmentation of, CAD of, 16:28
- Oligonucleotides
 adenine-thymine binary-base, positive-ion MALDI-TOF-MS of, 13:1052
 aflatoxin B1-guanine adducts in, identification and positional mapping, 9:676
 arylamine-modified, sequence verification by LC/MS, 19:1147
 charge states of negative ionization electrospray-MS, function of solution ammonium ion concentration, 8:155
 continuous-flow fast atom bombardment mass spectrometry of, 2:157
 DNA sequencing, by MS/MS, 15:510
 ECD study of, 20:213
 effect of cisplatin on gas phase dissociation of, ESI-MS/MS, 20:792
 ESI-MS of, 8:90
 charge state distribution in, 8:155
 fragmentation of
 in MALDI-MS, by charge tagging and H/D exchange, 13:1407
 modified nucleic acids in, 12:744
 studied by H/D exchange and ESI-MS/MS, 12:193
 gas-phase conformers, use of H/D exchange and ion mobility techniques, 19:938
 improved signal intensity, in electrospray ionization mass spectrometry, charge-state reduction with, 7:697
 ion stability of, co-matrix proton affinity and, 9:668
 ionic liquid matrix-induced metastable decay of, 16:2000
 labeled, 9:660
 MALDI-MS of, 12:5-methoxysalicylic acid and spermine as matrix for, 12:456
 mixed-sequence RNA/DNA, electrospray MS/MS of, 13:936
 modified, sequence confirmation, 14:825
 multiply charged
 modified base, ion trap tandem mass spectrometry applied to, 5:740
 tandem mass spectrometry of, 3:60
 multiply-charged T-rich anions, thermal decomposition in gas phase, 17:1229
 normal and modified, by matrix-assisted desorption Fourier transform mass spectrometry, 2:402
 sequencing of, SOS program in, 13:200
 specific and nonspecific, electrospray ionization mass spectrometry of, dimer formation in, 1:336, 6:159
 synthesized on montmorillonite, direct MALDI analysis, 17:1265
 trapped gas phase, fraying and electron autodetachment dynamics, 14:1330
 UV-MALDI of, 8:218
- Oligo(*p*-phenylethynylene)s, cationization mechanism of, 16:1695
- Oligopeptide-binding protein OppA, competitive binding to, FTICR-MS in-trap cleanup, 11:1023
- Oligopeptide mixtures, analysis of, ion/ion reactions in, 9:585
- Oligopeptides
 matrix-diluted, bombarded with slow and fast projectiles, TOF-SIMS of, 11:553
 protonated, MS/MS of, amide bond cleavage mechanism, 15:103
- Oligoribonucleotides
 fragmentation from ECD and EDD, 17:1369
 gas phase dissociation of, ESI-MS/MS of, 16:1262
- Oligosaccharide alditols, neutral O-linked, structural determination by negative ion LC-ESI-MS, 15:659
- Oligosaccharide antibiotics
 complex, quadrupole ion trap multiple-stage MS analysis of, 13:1313
 structure of, by FAB-MS, 8:1134
- Oligosaccharide cations, produced in an external source Fourier transform mass spectrometer, “slow” metastable decomposition of, 4:210
- Oligosaccharides
 9-aminofluorene-labeled, MALDI-FTMS characterization of, 12:1254
 alginate-derived, sequence analysis by negative-ion ES MS/MS, 17:621
 alkali metal cationized and permethylated, collision-induced dissociation of, influence of the collision energy and of the collision gas for the assignment of linkage position, 4:197
 analysis by negative-ion MALDI MS, 18:714
 analysis using nanoESI-DMS-MS, 18:502
 analysis using negative mode ES-MS, 16:60
 branched permethylated, analyzed by CID and ECD, 19:138
 carbon isotope ratio determination, ICP-MS for, 14:295
 characterization with boronic acid derivatization using IRMPD and QIT, 18:2094
 chloride adducts of, negative ESI-generated, linkage specificity in CID of, 12:1193
 chondroitin sulfate, MS/MS of, 14:1270
 derivatized, ESI and MALDI MS of, 16:683
 GONE sequencing and structural analysis by RP-HPLC-MS, 20:1821
 human and elephant milk, analysis using TLC and MALDI-MS, 17:139
 in human milk, structural analysis of, 13:1331, 13:1341
 labeled with multifunctioning oligosaccharide tag, MALDI-TOF and ESI-MS analysis, 17:194
 linear and branched, linkages of, 13:1322
 metal-ligated, containing sialic acid, stabilization and linkage analysis of, 12:528
 mobile protons in negative ion CID, 18:952
- N-linked
 linkage and branched determination using ion trap MS, 18:248
 with uncommon glycoform, serum of mouse with head and neck tumors, MALDI MS, 19:671
- neutral and sialylated, electron detachment dissociation, 18:2162
- neutral O-linked, exoglycosidase digestion and, 12:877
- from ovalbumin, phenylhydrazones of, MALDI-MS/MS-PSD, 15:725
- permethylated
 glycomics platform for analysis of, 18:1799
 n-hexylamine derivatized, liquid secondary ion mass spectrometry analysis of, 3:345
- polysulfated, MALDI-MS using pyrenemethylguanidine, 20:131
- post-source decay analysis, MALDI linear-field reflectron TOF, 19:1119
- sialylated glycans and, analysis using DHB/An and DHB/DMA matrices, MALDI-TOF MS, 19:1138
- sulfated, structural analysis by positive mode nano-ESI-FTICR-MS and MS/MS by SORI-CID, 17:96
- synthetic mannose-6-phosphate, high-energy collision-induced dissociation mass spectrometry of, 7:182
- underivatized, branching pattern and sequence analysis of, 13:671
- unmodified and methylphosphonate-modified, fragmentation mechanisms of, 18:984
- On-line acquisition, e-mailing of high-resolution exact-mass elec-

- tron impact/chemical ionization mass spectrometry data acquired using an automated direct probe, 10:1212
- On-line capillary liquid chromatography MS/MS, IT/reTOFMS using the sequence tag database search approach for peptide sequencing and protein identification, 11:127
- On-line capillary separations/tandem mass spectrometry, for protein digest analysis by using an ion trap storage/reflectron time-of-flight mass detector, 8:1237
- On-line desalting of physiologically derived fluids, in conjunction with capillary isoelectric focusing-mass spectrometry, 8:743
- On-line immobilized metal-ion affinity chromatography, phosphopeptides isolated by, electrospray ionization mass spectrometry of, 4:662
- On-line isotopic dilution cryofocusing gas chromatography/mass spectrometry, automated high-speed analysis of selected organic compounds in urban air, 5:1001
- On-line single droplet deposition, for MALDI-MS, 15:1471
- On-line strong cation exchange, μ -HPLC-ESI-MS/MS for protein identification and process optimization, 14:719
- One-step microelectroelution concentration, method for efficient coupling of sodium dodecyl sulfate-polyacrylamide gel electrophoresis and MALDI-TOFMS for protein analysis, 9:88
- Onium reactions, H-transfers and, characterization in closed *versus* open shell ions, 18:270
- O,O-dialkyl N,N-dialkyl phosphoramides, MS of, under electron impact conditions, 15:547
- O,O-diethyl O-aryl phosphorothionates protonated, fragmentation of, in tandem mass spectral analysis, 7:189
- Ophiopogon japonicus*, homoisoflavonoids from, HPLC-DAD-ESI MS analysis of, 16:234
- Opioid peptide-containing protein, characterization of, by electrospray ionization and liquid secondary ion mass spectrometry, 5:377
- Opioid peptides, analysis of, electrospray mass spectrometry for, 2:149
- Optical arrangement, multipass, infrared multiple photon dissociation in the quadrupole ion trap via, 5:886
- Optical fibers coiled with platinum wire, electrospray from, 11:464
- Orbital symmetry restraint circumvention, by 1,3-H shifts of enolic radical cations, 15:972
- Orbitrap mass spectrometry
 comparison with LTQ MS for peptide and protein identification, 20:1405
 data acquisition using benchtop model, 20:1441
 drug residue analysis, veterinary, hormone residue analysis, UPLC-Orbitrap MS, 20:451
 dynamics of ions of intact proteins, 20:1486
 HCD, phosphopeptide analysis using, 20:1425
 high-field, performance evaluation, 20:1391
 LTQ hybrid, analysis of intact monoclonal antibodies, 20:1415
 LTQ hybrid, MALDI produced ions inspected with, 20:1451
 mass precision and accuracy for proteome measurements, 20:1477
 methylated arginine quantified with nanoflow LS-ETD-Orbitrap MS, 20:507
 multiresidue screening in food and feed matrices, 20:1464
 nefazodone liver microsome metabolism analysis by, 17:363
 phase-enhanced selective ion ejection, 18:980
 post-acquisition ETD spectral processing for identification of peptides, 20:1435
 rephasing ion packets to improve resolution and peak shape, 20:1397
 spectral accuracy of molecular ions in, 20:2058
- Organic acid modifiers, electrospray use of, iron carboxylate oxygen-centered-triangle complexes detected during, 6:1226
- Organic compounds
 ambient air, BETX analysis using APCI-MS/MS, 20:829
 analysis by ES/VUV-SPI MS, 20:430
 influence of organic layer thickness in TOF-SI-MS, 20:2294
 liquid secondary ionization mass spectrometry beam-induced dehalogenation of, mechanism that involves secondary electron capture in, 7:1109
 mass spectrometric analysis of, characterization of a glow discharge ion source for, 5:305
 responsiveness to negative ion ESI MS, acidity and polarity in, 16:446
 soft ionization of, PERCI for, 15:424
 surface-induced dissociation analyzed using SPALDI and LDI MS, 20:957
 in urban air, automated high-speed analysis of, by on-line isotopic dilution cryofocusing gas chromatography/mass spectrometry, 5:1001
 volatile
 in air, PIT-MS detection and identification of, 16:1316
 using purge-and-trap gas chromatography/mass spectrometry, interferences, removing of, importance of enhanced mass resolution in, 3:831
- Organic ion imaging
 beyond the limit of static secondary ion mass spectrometry, 6:1047
 wide-angled secondary ion probe for, 2:362
- Organic ions, relative collision cross sections of organic ions, 6:803
- Organic material, carbon-13 labeling for improved profiling of, secondary ion MS, 17:1142
- Organic mixtures, complex, suspended trapping gas chromatography/Fourier transform mass spectrometry for, 1:431
- Organic molecules
 containing two functional groups, formation and fragmentation of gas-phase ion-molecule complexes of transition-metal ions with, 3:451
 neutral, including fullerenes, applications of electrospray ionization mass spectrometry to, 4:596
 simple, cationization of, in MALDI-MS, 15:879
- Organic neutralization agents, for neutralization-reionization mass spectrometry, 3:108
- Organic pollutants, halogenated persistent, analyzed by pLVI-GC/ECNI-qMS, 18:1375
- Organic polymers, UV ablation products of, direct chemical analysis, by using selective ion monitoring mode in gas chromatography/mass spectrometry, 5:106
- Organics, neutral, in solution, gas-phase molecular dications formed from
 via chemical electron-transfer reactions by using electrospray ionization mass spectrometry, 5:689
 via the controlled-current electrolytic process inherent to electrospray, 7:157
- Organolithium, ring-opening hexamethylcyclotrisiloxane polymerization initiated by, MALDI-MS study of, 14:95
- Organomercurial derivatization, cysteine and cystine in peptides and proteins following, by matrix-assisted laser desorption ionization mass spectrometry, 5:359
- Organometallic complexes, MS behavior of, 16:94
- Organopalladium complexes, ArPd(PPh₃)₂I, gas phase fragmentation reactions, ESI-FTICR-MS, 17:1582
- Organophosphorus compounds, structural characterization of, multiple-stage mass spectrometry in, 4:125
- Organoruthenium anticancer complexes, analysis using IM-MS, 20:1119

- Organosulfur compounds, fragmentation of, upon electron impact, 8:801
- Organotrifluoroborates, mass determination using ESI-MS, 18:404
- Orifice potential, sampling, characterization of a glow discharge plasma as a function of, 1:448
- Ornithine lipids, glutamine lipids and, from *R. sphaeroides* cell membranes, MS/MS, 20:198
- Ortho-cyclopropylphenyl benzamides, cyclization in gas and liquid phases, 12:956
- Ortho effects, sequential, characterization of novel [M-35]⁺ fragment ions in the mass spectra of 2-alkyl-4,6-dinitrophenols, 2:69
- Ortho silicates, protonated, APCI low energy collision-induced dissociation fragmentation of, McLafferty or ion-neutral complex rearrangements, 5:544
- Orthogonal acceleration
 spontaneous and deflected drift-trajectories in, 5:588
 time-of-flight secondary-ionization, ion intensity in, enhancement of, 7:467
 TOF (See Time-of-flight mass spectrometry of tributyl phosphate, zeolites in, 8:718)
 triple quadrupole, as applied to flame diagnostics, study of the C₂H₄/N₂O/Ar flame, 4:152
 unit resolution, accuracy of, 8:380
 utility of N-peracetylation of proteins for their structure determination by, 4:87
 UV-matrix-assisted laser desorption Fourier transform, for peptides, investigation of, 2:22
- Orthogonal-acceleration time-of-flight mass spectrometer, linear, detector pulse width and jitter in, resolution limitations from, 5:772
- Orthogonal acceleration time-of-flight mass spectrometry, spontaneous and deflected drift-trajectories in, 5:588
- Orthogonal control system, automated, for ESI, 15:1201
- Orthogonal-magnetic acceleration time-of-flight mass spectrometer, tandem double focusing, matrix-assisted laser desorption ionization employing, peptide sequence determination by, 7:1
- Orthogonal time-of-flight mass spectrometry, background reduction in, 8:125
- Ostreopsis ovata*, palytoxin and ovatoxin-a analyzed by LC-MS, 19:111
- Ovalbumin
 assigning N-linked glycan structural isomers from, IMS-MS, 19:1711
 carbohydrates from
 composition of, 11:564
 postsource decay fragmentation of, 11:572
 phenylhydrazone-oligosaccharides from, MALDIMS/MS-PSD fragmentation study of, 15:725
- Ovatoxin-a, *O. ovata*, analyzed by LC-MS, 19:111
- Overlap bond orders, modified neglect of differential overlap energy partitioning, effect of protonation site on bond strengths in simple peptides, 5:704
- Overlap energy partitioning, modified neglect of differential overlap bond orders, effect of protonation site on bond strengths in simple peptides, 5:704
- Overtone mobility spectrometry
 experimental observations, 20:729
 theoretical considerations of resolving power, 20:738
- 1,3,4-Oxadiazoles, protonated and lithiated, MS study of, 14:289
- β -Oxalylaminoalanine, proton affinity of, 16:1151
- Oxazines, 2:2-N-phenyliminoperhydro-3,1-,cyclohexene-fused, fragmentations of, effects of N-substitution on, 2:125
- Oxazinones, stereospecific fragmentation of, 10:393
- Oxazolone structure
 of b₂ fragment ions from tri-alanine, IRMPD spectroscopy, 20:334
 cyclic, of *b* ions, 16:1941
- Oxazolones, deprotonated, formation of, 15:446
- Oxidation
 of analyte, in electrospray ion source using porous flow-through emitter, 15:1755
 of human apoprotein B-100, cysteine acid residue and, tandem mass spectrometric characterization of, 6:242
- Oxidation of arachidonic acid, free-radical, four regioisomers of F₂-isoprostanes formed by, mass spectrometric analysis of, 7:490
- Oxidation products, of tryptophan, identification by MS, 14:406
- Oxidation reactions of dibenzothiophene subjected to negative chemical ionization with oxygen, 4:949
- Oxidation/reduction processes observed in fast-atom bombardment/liquid secondary ion mass spectrometry, true effect of experimental parameters on, evaluation of, 4:482
- Oxidative damage to double-stranded DNA, gas chromatographic-mass spectrometric method for the assessment of, by quantitation of thymine glycol residues, 4:336
- Oxide radical anion (O⁻), comparison of the reactions of, with simple aromatic compounds at atmospheric and at reduced pressure conditions, 5:367
- Oxidized globin subunits, enzymatic digestion of, electrospray ionization-tandem mass spectrometry analysis of, an improved method to determine amino acid substitution in the hemoglobin "core," 7:1040
- 8-Oxo-dG residues, differentiation of, in oligodeoxynucleotides, 13:195
- Oxofatty acids
 collisional activation of, 9:620
 structural determination of, by charge-remote fragmentations, 9:216
- Oxonium ions, fragmentation of, 10:386
- Oxovanadium complexes, bimetallic, ESI-QTOF-MS/MS analysis, 19:1247
- Oxy-carboxylic acid, TMS derivatives of hydroxy-carboxylic acid and, fragmentation of, EIMS, 19:66
- Oxygen
 benzenedicarboxylic ester molecular ion collision with in quadrupole mass spectrometer, 10:1248
 as CID target gas, 12:23
 Meisenheimer complexes bonded at, 15:998
 negative chemical ionization with, dibenzothiophene subjected to, oxidation reactions of, 4:949
 quantitative analysis of, by quadrupole ion trap MS, 13:1120
- Oxygen- and nitrogen-containing molecules, small, reactions to the phenylium cation with, 7:473
- Oxygen addition-induced dechlorination of PCBs and DDT metabolites in electron capture mass spectrometry, regioselectivity of, 7:66
- Oxygen addition to chloroaromatic compounds, collision-induced, 7:1144
- Oxygen atom and the aromatic ring, competing proton transfer to, propene loss initiated by, chemical ionization of phenyl *n*-propyl ether and methyl substituted analogs, 7:639
- Oxygen-centered radicals, DMPO adducts of, separation and identification of, 14:862
- Oxygen compounds, CH₃OCH₂⁺ reactions with, 13:1028
- ¹⁸Oxygen labeling, on ion trap mass spectrometer with Zoom-Quant software, quantification using, 16:916
- Oxygen-protonated methanol [CH₃OH₂]⁺ involving double hydrogen atom transfers, formation of, and metastable molecular ion of methyl lactate, 6:202

- Oxygenated chemical ionization reagents with vincamine, ion-molecule reactions of, 7:250
- Oxysterols, analysis by ES-MS/MS, 17:341
- Ozone, tetracycline degradation by, in aqueous medium, APCI/MS, 18:679
- Ozone oxidation products, amino acids and small peptides, identification by ESI-MS and MS/MS, 11:526
- Ozonolysis, at peptide tyrosine and histidine residues, 17:1289
- P
- P0, characterization MALDI-TOF/TOF MS, bovine myelin sheath, 17:117
- P13 membrane protein, of *Borrelia burgdorferi*, MS of, 13:295
- PAA membrane. *See* Porous anodic alumina membrane
- Packed capillary, LC-RAM-MS of, 13:82
- Packed microcolumns, continuous flow fast atom bombardment with, comparison of precolumn *versus* coaxial matrix delivery, 1:312
- Packed-volume supercritical fluid chromatography/pneumatically, assisted ESI-MS, development and application of, 9:498
- Paclitaxel, ionization and adduct formation of, 15:585
- PAH. *See* Polyaromatic hydrocarbons
- PAH-DNA adducts, sensitive determination of, matrix design for matrix-assisted laser desorption ionization, 5:1021
- PAHs. *See* Polycyclic aromatic hydrocarbons
- Palladium complexes, MS behavior of, 16:94
- Palytoxin, *O. ovata*, analyzed by LC-MS, 19:111
- Para-substituted benzaldehydes, gas-phase base-catalyzed Claisen-Schmidt reactions of the acetone enolate anion with various, 7:82
- Paracoccus denitrificans*, membrane protein analysis by LILBID MS, 18:1429
- Paralytic shellfish poisons, gas phase dissociation reactions of, 15:462
- Paraquat
capillary electrophoresis-mass spectra of, 7:981
in water, LS-(ESI)MS of, 9:830
- Parent ion axialization, selective, for improved efficiency of collision-induced dissociation in laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry, 5:900
- Parent ions
isomeric, reactions from, in the dissociation of dimethylpyrroles, 7:930
resolution in linked scans for dissociations occurring in the first field-free region of sector mass spectrometers, 8:545
- Particle beam high-performance liquid chromatography-mass spectrometry, polycyclic aromatic hydrocarbons measured in soils by, 6:597
- Particle beam interface
coupling, directly to a quadrupole ion trap mass spectrometer, 4:38
with liquid chromatography-mass spectrometry, micro high-performance, analysis of coumarins by, 6:132
microflow rate, electron capture ionization of explosives with, 7:753
- Particle beam liquid chromatography-mass spectrometry
mass spectra of chlorinated phenoxyacid herbicides obtained by, evidence for thermal decomposition contributions to, 3:823
with massive cluster impact, 7:293
- Particle-beam liquid chromatography-mass spectrometry, optimization of the Hewlett-Packard, interface by statistical experimental design, 6:507
- Particle beam liquid chromatography/mass spectrometry and gas chromatography/mass spectrometry, poly(amic acid) and polyimide characterization using, 5:299
- Particle beam liquid chromatography/mass spectrometry interface, performance of, effect of a dc glow discharge on, investigation of, 5:908
- Particle beam mass spectrometry, liquid chromatography, high-performance, investigation of enhanced ion abundances from a carrier process in, 1:92
- Particle beam-mass spectrometry, high-performance liquid chromatography, determination of polycyclic aromatic hydrocarbons by, 4:898
- Particle discriminator interface, for nanoflow ESI-MS, 14:1236
- Particle-in-cell code, three-dimensional, for realistic simulation of ICR MS, 10:136
- Particulate isotopic standard
of plutonium in an aluminosilicate matrix, A, 2:81
of uranium and plutonium in an aluminosilicate matrix, 5:852
- Partners, H-transfer and C-C bond formation between, a photoionization study of the ion-neutral complexes $[\text{CH}_3\text{CH}^+3\text{-CH}_2\text{CH}_3]$ and $[\text{CH}_3\text{CH}_2\text{CH}^+\text{CH}_3\text{-CH}_3]$ in the gas phase, 7:73
- Pattern-based algorithm for peptide sequencing from tandem high energy collision-induced dissociation mass spectra, 3:326
- Paul trap, gas phase cytochrome *c* ions in, 16:1493
- Paul-type ion traps, single-ion trajectories in, computer simulation of, 4:687
- PBDEs. *See* Polybrominated diphenyl ethers
- PCBs and DDT metabolites, oxygen addition-induced dechlorination of, in electron capture mass spectrometry, regioselectivity of, 7:66
- PCR amplicons, single-stranded, production of, 13:232
- PCR (polymerase chain reaction), in genetic diagnosis of familial adenomatous polyposis, 8:737
- PCR product mass measurements, preparing samples for, ES-IFTICR-MS in, 13:338
- PDMS membrane. *See* Polydimethylsiloxane membrane
- PDMS multichannel emitter. *See* Polydimethylsiloxane multichannel emitter
- PE chloramines. *See* Phosphatidylethanolamine chloramines
- Peak assignment, software for, synthetic polymers, MS/MS, 18:1324
- Peak height precision, in HT-TOF MS, 16:1117
- Peak heights, relative, in Fourier transform-ion cyclotron resonance spectra, space charge effects on, 3:734
- Peak interference in FTICR-MS, 13:387
- Peak shape analysis, for unresolved interferences in mass measurement, 9:789
- Peak splitting, with a quadrupole mass filter operated in the second stability region, 10:1263
- Peak-target Fourier transform, automated charge state determination of isotope-resolved MS by, 19:46
- Peak width analysis, EX1 characterization of kinetics by, H/D exchange MS, 17:1498
- PEGylation sites, top-down determination using MALDI ISD MS, 20:326
- Penning trap. *See also* Ion trap
laser-induced fluorescence for ion tomography in, 9:925
addition of ac potential to, 8:962
cooling of negative ions in, 8:793
tetragonal vs cylindrical, 8:283
- Pentacene, clustering in MALDI-TOF MS, 17:222
- Pentadienes, push-pull, from cycloalkylidene malonic acids, EI-MS of, 14:189
- Pentafluorobenzyl chloroformate derivatization, for enhancement of detection of amino acids or alcohols by electron capture negative ion chemical ionization mass spectrometry, 6:525
- Pentafluorobenzyl ester, trimethylsilyl ether derivatives, hy-

- droxy fatty acids as, analysis of by electron ionization gas chromatography/mass spectrometry, 6:40
- Pentaquadrupole, multiple stage mass spectrometry, polar [4+2+⁻] Diels-Alder cycloaddition to, gas stage ions and, 6:1
- Pentaquadrupole mass spectrometry
borinium cations by, 12:948
MS³ capabilities of, case study in the use of, ion-molecule reactions and collision-activated dissociation of C₄H₄⁺ isomers, 3:518
multiple stage, for generation and characterization of gas-phase ionic species, 7:1126
- Pentasaccharides, cobalt-coordinated, linkage position determination in, ion trap-MS for, 9:1125
- Pepsin mapping, of deuterium exchange behavior of ERK2, 10:685
- Peptide aggregates, gaseous, hidden in mass spectra, 9:743
- Peptide amide hydrogen, migration of, during ESI, 12:410
- Peptide and protein ions, proton transfer reactivity, formed by electrospray ionization, 6:1086
- Peptide and protein mixtures, characterization of, capillary electrophoresis Fourier transform ion cyclotron resonance mass spectrometry with sustained off-resonance irradiation for, 5:894
- Peptide anions, ETD of, 16:880
- Peptide backbone, N—C bond cleavage of, hydrogen abstraction in, 12:1044
- Peptide clusters, multiply charged, gas phase dissociation pathways of, 14:1373
- Peptide-crown ether complexes, photodissociation by UV irradiation, 19:257
- Peptide disulfides, oxidized glutathione and contryphan, fragmentation from negative ion mass spectrometry, 19:358
- Peptide fragmentation obtained from a reflector matrix-assisted laser desorption-ionization time-of-flight and a tandem four sector mass spectrometer, comparison of, 6:822
- Peptide hits technique, guidelines for, 16:1231
- Peptide ion dissociation, energetics of, by laser desorption chemical ionization Fourier transform mass spectrometry, 6:1069
- Peptide ion ratio, ¹⁶O/¹⁸O, calculating for proteome relative quantification, 15:437
- Peptide ions
amidated, fragmentation of, 15:158
arginine-containing, formed by electrospray ionization, modeling the maximum charge state of, 7:972
charge location in, decompositions of cationized heterodimers of amino acids in relation to, 4:461
collision-induced dissociation of, the effect of charge state and the localization of charge on, 5:966
collisionally-activated dissociation and 193-nm photodissociation of, in Fourier transform mass spectrometry, efficiency of, 1:288
dissociation energetics of, 9:175
dissociation of, surface-induced, in Fourier transform mass spectrometry, 1:413
gas-phase, low energy decompositions of, role of the site of protonation in, 7:522
MALDI-MS detection of, esterification and, 16:1269
metal cationized, CID fragmentation pathway for, 16:1305
multiply protonated, ubiquitin, insulin B, and renin substrate, apparent gas-phase acidities of, 7:1211
new method to study the effects of peptide sequence on the dissociation energetics of, 9:175
surface-induced dissociation of, 14:1340
vacuum ultraviolet light-induced fragmentation pathways of, 16:1384
- Peptide ions in electrospray ionization mass spectra, adducts (molecular mass=98 u) attached to, origin and removal of, 1:382
- Peptide ions in the quadrupole ion trap, tandem mass spectra of, effects of heavy gases on, 7:1194
- Peptide-iron(III) complex, MS of, 8:1070
- Peptide ligands, KV1.3, noncovalent complexes of, observation of, a model investigation using ion-spray mass spectrometry, 7:1075
- Peptide-lithium adducts, gas-phase, investigations of, tandem mass spectrometry and semiempirical studies, 1:473
- Peptide [M + H]⁺ ions
dehydration of, in the gas phase, 4:477
metastable decomposition of, via rearrangement involving loss of the C-terminal amino acid residue, 1:249
- Peptide mass fingerprints, algorithm measurements of, 16:13
- Peptide mass mapping, of histone H4 acetylated isoforms from lymphosarcoma, 16:1641
- Peptide metal(II)-ion complexes, decompositions of, formed by electrospray ionization, use of scans at a constant ratio of B/E for studying, 6:608
- Peptide mixtures, liquid chromatography/tandem mass spectrometry analysis of, data-controlled automation of, 7:532
- Peptide nucleic acids, mono-deprotonated, complete sequencing of by CID, 12:805
- Peptide-peptide interactions, IM-MALDI, orthogonal TOF-MS, and ESI-MS of, 13:167
- Peptide precursors, in bacterial cell wall, vancomycin antibiotic interaction with, hydrogen/deuterium exchange in, 9:1255
- Peptide radical dications, generated via low energy CID, 16:1795
- Peptide recovery, in MS of protein digests, 15:784
- Peptide sequencing
de novo, MALDI-quadrupole ion trap MS for, 14:1012
determination by matrix-assisted laser desorption ionization employing a tandem double focusing magnetic-orthogonal acceleration time-of-flight mass spectrometer, 7:1
determination from high-energy collision-induced dissociation spectra using artificial neural networks, 6:947
lysine-specific mass tagging in, in human proteome, 14:1
using a patchwork approach and surface-induced dissociation in sector-TOF and dual quadrupole MS, 14:1387
- Peptide Shifter, enhancing separation reproducibility using correlated expression profiles, 18:1638
- Peptide side chain influences, on the metal ion binding site in metal ion-cationized peptides, participation of aromatic rings in metal chelation, 6:1079
- Peptide spectra, Bn (n greater than or equal to 3) ions in, the structure and fragmentation of, 7:233
- Peptide tags, elution time, in proteome analyses with capillary LC-TOFMS, 14:980
- Peptides
α- and β-alanine-containing, CID spectra of, 19:1743
a- and b-type fragment ions, stable isotope labeling ion mobility/MS/MS, 19:1781
a- and b-type ion variance, IM-MS/MS, 19:609
acetylation, detection and location of acetylated lysine residues, 18:1423
amino acid composition and role in charge inversion, gas-phase ion/ion reactions, 20:180
amphipathic, periodic sequence distribution in ECD, 20:1182
analysis by decoupling ETD and CRCID, 20:567
analysis of complex mixtures using FTMS and differential MS, 18:226
anionic, hydrogen atom scrambling, collisional activation by MALDI TOF/TOF MS, 19:1719
anionic radicals, formation in a vacuo, 17:1249

- arginine containing di- and tri-, H/D exchange in, 20:2049
- arginine-containing ions, hydrogen migration upon electron transfer, 19:1726
- b*₀ ions, cyclization of, 20:2248
- b*-type ions produced in ECD of, 18:2007
- β -peptide imaging distribution in whole body mice sections, MALDI-MS, 18:1921
- Boc-protected, positive and negative ESI MS, 18:651
- bound to iron oxide nanoparticles, analysis using MALDI-MS, 18:910
- calmodulin interactions with, investigated by MALDI-MS, 20:576
- cations, survival after electron capture, 17:1675
- chain length, effect on charge-separation reactions, 20:1890
- charge-tagged, electron capture in, 18:2146
- CHCA-modified Au nanoparticles for analysis, LDI-MS, 20:1530
- collisional energy and ion fragmentation, 20:469
- comparison of Orbitrap MS with LTQ MS for identification of proteins and, 20:1405
- containing oxidized cysteine and methionine, sequencing using ETD, 18:1499
- copper complexes with, electrogeneration in microspray MS, 19:560
- crosslinked, effect of proline and aspartic acid residues on dissociation, 19:344
- cyclization reaction of fragment ions, multistage CAD, 18:663
- cysteine-containing
- adenosylated, acting as bisubstrate analogs of protein kinases, mass spectrometric characterization of a series of, 5:443
 - analysis of
 - nano electrospray combined with a quadrupole ion trap for, 7:150
 - sheathless CZE-MS applied to, 10:1271
 - angiotensin and zinc metal ions, interaction of, by electrospray ionization mass spectrometry, 5:959
 - backbone-modified, structural determination of, tandem mass spectrometry for, 1:174
 - calcium binding sites in, MS/MS determination of, 9:1020
 - chemical ionization of, Fe⁺, 4:106
 - collision induced decomposition of, choice of collision parameters, 7:677
 - collision-induced dissociation spectra of, new fragment ion type in, identification of, formation of a2–16 ions, 4:424
 - conformational changes in, MALDI and H/D studies of, 10:719
 - cyclic, multistep MS/MS for sequencing of, 10:732
 - cysteine and cystine in, quantitative assessment of, following organomercurial derivatization and analysis by matrix-assisted laser desorption ionization mass spectrometry, 5:359
 - cysteine-containing
 - complexes of iron(II) with, in the gas phase, 7:977
 - dehydration of, 9:945
 - Fe²⁺ interaction with, gas phase studies of, 9:1285
 - dissociation of, uninterpreted high-energy collision-induced, search of, with sequence databases, 7:1089
 - femtomole and sub-femtomole levels of, detection of, by tandem magnetic sector/reflector time-of-flight mass spectrometry and matrix-assisted laser desorption ionization, 2:91
 - fragmentation of, fast, observed, in matrix-assisted laser desorption, factors that influence, 7:225
 - frozen aqueous solutions of, infrared SALDI-MS of, 9:912
 - gas-phase anionic complexes of, structure and collision activated decompositions, 5:137
 - helical unfolding, the mass spectrometry of, 5:425
 - identification of, by MALDI and MALDI-post-source decay TOFMS, 10:91
 - ions of, proton affinities of, ESI-MS of, 10:483
 - large, complete sequence confirmation of, by high energy collisional activation of multiply protonated ions, 6:234
 - low-mass ions produced from, by high-energy collision-induced dissociation in tandem mass spectrometry, 4:882
 - mass spectrometry and tandem mass spectrometry analysis of, optimization of capillary zone electrophoresis/electrospray ionization parameters for, 3:289
 - matrix-assisted laser desorption/ionization time-of-flight mass spectrometry of, improved mass accuracy in, 5:955
 - MH⁺ ions of, high energy collision-induced decomposition of, comparison of helium and argon as collision gases in, 3:502
 - modified with trans-2-hexenal, MS/MS of, 10:613
 - opioid, analysis of, electrospray mass spectrometry for, 2:149
 - in plasma, mapping of with EMS, 10:45
 - polycyclic aromatic hydrocarbon diolepoxide-adducted, characteristics of high energy collision-induced dissociation tandem mass spectra of, 5:649
 - primary thermal degradation processes of, characterization of, using the mass spectrometric technique K⁺IDS, K⁺ ionization of desorbed species, 5:564
 - protonated
 - complexation of with saccharides in gas phase, decreases rates of hydrogen/deuterium exchange reactions, 6:1247
 - decomposition of, collision-activated, influence of cysteine to cysteic acid oxidation on, evidence for intraionic interactions, 3:337
 - high-energy fragmentation of, determination of, using a BEqQ hybrid mass spectrometer, 4:470
 - and methanol, interaction of, molecular probe for, gas-phase hydrogen/deuterium exchange as, 5:623
 - multiply, surface-induced dissociation of, 3:859
 - segments of, proton affinities of, 10:1
 - sequencing, pattern-based algorithm for, from tandem high energy collision-induced dissociation mass spectra, 3:326
 - simple, bond strengths in, effect of protonation site on, application of Ab initio and modified neglect of differential overlap bond orders and modified neglect of differential overlap energy partitioning, 5:704
 - small singly charged, aggregates of, 10:347
 - tandem mass spectral data of, an attempt to correlate with amino acid sequences in a protein database, 5:976
 - tandem mass spectrometry of, with N-terminal glutamine studies on a prion protein peptide, 1:258
 - thermal energy and its distribution in, calculation of, 10:323
 - ultra-high sensitivity analysis of, micro-electrospray mass spectrometry, 5:605
 - UV matrix-assisted laser desorption Fourier transform mass spectrometry for, investigation of, 2:22
 - zeptomole/attamole per microliter sensitivity for, micro-electrospray, 5:867
- de novo sequencing using exhaustive enumeration, 17:1041
- deciphering the iodination code, 20:965
- DFT study of ions generated by MALDI, 20:1327
- dications, charge location directs ECD of, 17:1704
- disulfide bond cleavage via salt bridge mechanisms, low energy CID, 18:1109
- disulfides, characterization of lanthionine trisulfides and tetrasulfides from, ESI MS, 20:783
- divalent metal ion interactions with, probed by ECD of trications, 17:1731

- doubly-proteinated tryptic, structure of b_2 ions generated from, 20:618
- ECD product ion abundances correlated with amino acid polarities, 20:2273
- EDC monitoring deamidation during processing of proteins and, 19:855
- fingerprinting on porous silica and alumina, LDI MS, 19:632
- folding and unfolding of helix-turn-helix motifs in gas phase, 18:1239
- fragment ion structures, analyzed by SORI-HDX in FTICR instruments, 19:1771
- fragmentation of, editorial, ?
- fragmentation of photolytically generated radical ions, 20:1378
- free-radical-induced site-specific cleavage in gas phase, CID in ESI/MALDI-MS, 18:807
- gas-phase conformation probed by CID, 17:786
- high measured mass accuracy of fragments, CID spectra, 19:891
- hydrogen rearrangements in ECD, 18:113
- identification using IM-CID-MS, 20:907
- ILMs for proteins, carbohydrate and, analysis by MALDI MS, 20:1790
- imaging in rat brain, MALDI-FTICR MS, 18:145
- improved IRMPD using N-terminal phosphonite derivatization, 20:377
- improved mass accuracy using SWIFT excitation for MALDI-FTICR MS, 19:76
- influence of size on sequence scrambling during CID of b -type ions, 20:2174
- intramolecular cross-linked, collision-induced dissociation, 20:557
- ion charge states of proteins and, FTICR, 17:903
- ion dissociation, effect of histidine oxidation on, MS/MS, 18:553
- ion folding, kinetics measured by IR activated DR ECD, 19:780
- ionization efficiency changes caused by phosphorylation, 18:1973
- ions
- factors affecting VUV photofragmentation of, 18:1439
 - soft landed surface effects on charge reduction and desorption kinetics, 20:901
- ions, IRMPD spectroscopy for studying amide position protons, 20:1841
- ions containing S-alkyl cysteine sulfoxide, gas-phase fragmentation reactions, 18:1690
- IRMPD and CID in ion traps, 20:349
- isocyanate-adducts, analyzed by MS/MS, 20:1567
- large hydrogen deficient, tracking radical migration in, 20:1148
- linear, vibrational frequencies of protein ions and, RRKM rate constant calculation using, 17:1749
- MALDI-formed $[M+H]^+$ ions, formation of ($b_{n-1}+H_2O$) by collisional activation, QqTOF MS, 18:1024
- MARCKS, phosphorylation sites identified by ECD FTICR MS/MS, 18:2137
- mass-based classification of, 19:1808
- mass mapping, determination of unique AA substitutions, FT-ICR MS, 17:508
- metalated with alkaline-earth metal ions, EDC, 17:757
- method for prediction of isotopic distribution, 19:703
- mono-oxidized thio-ether bonds, evaluation by CID and ECD, 18:493
- multiply-charged proteins and
- detection by MALDSI coupled to FT-ICR MS, 17:1712
 - generation by RADIO for MS, 20:597
- a_n and a_n^* fragments, isotope labeling, MS/MS and density functional theory calculations, 19:1788
- with N-terminal dimethylation and imine/methylol adduction, MS/MS, 17:859
- neutral loss of isocyanic acid in CID, 20:723
- nucleotide, *E. coli* microcin C7-C51 characterization by CID, 19:1187
- O-glycosylated and O-phosphorylated, sequencing by ECD and fixed charge derivitization, 18:1405
- opioid, protein-containing, characterization of, by electrospray ionization and liquid secondary ion mass spectrometry, 5:377
- oxidized, fragmentation analyzed by SID, RRKM modeling and MD, 20:1579
- ozonolysis at tyrosine and histidine residues, 17:1289
- phospholipids, leucine enkephalin-alka(e)nal-glycerophosphatidylcholine adduct identification by MS/MS, 17:657
- phosphorylated
- biotinylated probe for selective enrichment, 19:741
 - effect of diammonium citrate to CHCA matrix for detection, MALDI-TOF MS, 18:1925
 - enrichment using poly-histidine tag, 18:1007
 - residues in peptide-peptide formation, ESI MS, 19:1535
- post-acquisition ETD spectral processing for identification of, 20:1435
- prion protein, tandem mass spectrometry of peptides with N-terminal glutamine studies on, 1:258
- protected synthetic, sample preparation for MALDI-MS, 19:614
- proteins and
- counting sulfhydryl- and disulfide-bonds using mercurial ions, 19:1108
 - quantification using iTRAQ with ETD, 19:1255
- protonated, time-resolved method for PD mechanistic study of, TOF MS/MS, 18:1729
- protonated, ETD and CID in linear ion traps, 20:1672
- protonated polyproline, tandem MS, 18:2198
- radical ions, dissociation of N-C α bond, 19:1799
- ranid skin secretions, oxidation *versus* carboxamidomethylation of S-S bonds in, MALDI-MS, 19:479
- rapid sequencing, with a picomole-scale method and ESI-MS, 17:995
- rapid UV photodissociation using naphthyl-based adducts, 20:385
- sequence information from linear ion trap MS/MS, 19:1813
- sequence scrambling through cyclization of b_5 ions, CID, 19:1590, 19:1776
- sequencing of, A_1 ion enhancement for protein identification and, 20:1214
- sequencing using alternative enzymes for on-line proteolysis, triaxial electrospray probe, 20:1983
- sequencing using EDD MS/MS, 19:1156
- sequencing using LIAD with CID in FT-ICR-MS, 18:525
- shifting unoccupied spectral space in mass spectrum of fragment ions, 20:2124
- SID study of small molecules, non-covalent protein complexes and, 19:190
- simple di-, tri-, and tetra-peptides, sodium ions affinities of, 18:541
- singly-charged α -type peptide ions formed in 157nm nanometer photodissociation, 17:1315
- singly-charged ions, collision cross-section database, 18:1232
- singly protonated
- generated by MALDI, 193 nm photodissociation characteristics of, 17:1643
 - statistical interpretation of time-resolved PD, 19:645
- spin-trap capped, electron capture in, 18:432

- structures of *y*- and *b*- type ions, 157-nm PD, 19:695
 surface-induced dissociation analyzed using SPALDI and LDI MS, 20:957
 time-resolved photodissociation by MALDI, 20:1522
 trace, analysis using 2,5-dihydroxybenzoyl hydrazine derivative CNTs, MALDI, 17:1023
 trimethyllysine-containing, methyl group migration during fragmentation, MS/MS, 20:1172
 tryptic dications, bifurcating fragmentation behavior, CAD, 19:1755
- Peptide(s). *See also* Protein(s)
 affinity-bound to antibody beads, direct MALDI-MS/MS of, 14:1076
 affinity purified cross-linked, MS detection of, 15:832
 aggregates of, formation during ESI, 12:1020
 analysis of, microwave-enhanced Akabori reaction for, 13:839
 antibiotic linked to, released by ECD-FTICR-MS, 14:302, 14:682
 argentinated, CID of, product ions from, 12:163
 assigning to tandem mass spectra, SILVER in, 15:910
 Ca₂⁺ affinity of, 11:770
 carbon nanotubes as affinity tubes for, in MALDI MS, 15:1629
 charge-derivatized
 fragmentation of, 12:399
 MALDI-PSD spectra of, 11:145
 charge remote fragmentation of, MALDI of, 8:501
 composition-based sequencing of, mass determination by FTICR-MS, 15:703
 conformational differences of, in quadrupole ion trap, 12:1331
 containing arginine, 8:60 u loss from, 8:253
 containing nitrotyrosine, MALDI-MS of, 12:438
 crosslinked, fragmentation and assignment of, 14:834
 α-cyano-4-hydroxycinnamic acid cocrystallization with, promotion within levitated droplets, 16:254
 cyclic, IM-MS of, 15:870
 cyclic protonated, MSn characterization of, 15:1039
 cysteic acid-containing, low energy collision-induced dissociation of, from d-type ion, 8:25
 cysteine-containing
 iron(II) complexes with, 8:199
 protonated ions of, fragmentation of, 15:697
 cysteine-containing oxidized, fragmentation of, 12:228
 deprotonated, fragmentation of, 12:1
 phenylalanine and, 13:1242
 dilute mixtures of, MALDI-TOFMS of, matrix-cluster peaks in, 11:88
 disulfide-linked, SO₂⁻ electron transfer ion/ion reactions with, 16:1020
 DTSSP unexpected reactivity with, 15:736
 ECD of, amino acid side chain loss in, 16:1523
 ECD of, Fourier transform MS, 13:1396
 enhanced sample loading of, 8:15
 ES-generated copper(II) complex ions of, CID of, in formation of molecular radical cations of enkephalin derivatives, 12:1114
 ESI-MS of, alkali-metal adduct ion formation in, 11:967
 fragmentation of, ESI MS/MS for, 16:505
 gas phase H/D exchange of, acidic residues and sodium ion-adduction in, 14:1477
 G1cNAcylation sites of, identification by Q-TOFMS, 12:1106
 H/D exchange mass spectra of, automated analysis of, 12:1153
 in-source decay of, 12:420
 interaction of, MALDI study of, 12:88
 internal residue loss of, 16:607
 ionic liquid matrix-induced metastable decay of, 16:2000
 lariat crown ether lysine probing in, 13:493
 mass fingerprinting of
 analysis with FTICR-MS, nanoLC/microelectrospray ionization, and MALDI, 14:553
 MALDI in, scoring methods in, 13:22
 mass assignment from, 14:943
 matrix mixture for MALDI-MS in, 14:992
 negative signature mass algorithm in, 14:1032
 origins of, 14:931
 mass spectra of
 b₂ ions in, structure and fragmentation of, 11:427
 on molecular scanner, visualization and analysis of, 13:221
 screening method for, before database search, 11:422
 measuring abundance of, increased proteome coverage with DREAMS, 13:954
 metal trication attachment to, 15:406
 methionine-containing, analysis by charge derivatization and MS/MS, 11:31
 methionine oxidation in, CID and ECD of, 14:605
 methylarginine-containing, fragmentation pathways in, ES-IMS/MS of, 15:142
 mixtures of, HPLC, MALDI, and ESI-MS of, 12:215
 model, gas phase H/D exchange reactions of, 12:1272
 modified with photolabile arylazido groups, MALDI-MS analysis of, 15:1156
 monoprotonated nucleic acids of, gas phase decomposition of, 11:615
 MS of, 8:296, 12:125
 MS/MS identification of, increasing through automatic function switching optimization, 16:1818
 multiply protonated, fragmentation of, charge derivatization and, 13:47
 N-sulfonated, MALDI-TOF/TOF-MS of, 15:1838
 noncovalent interaction with DNA, MALDI-TOF detection of, 15:28
 noncovalent interactions of polyether compounds with, CAD evaluation of, 14:1148
 ozone oxidation products of, identification by ESI-MS and MS/MS, 11:526
 phenylisothiocyanate-derivatized electrosprayed tryptic, protein identification from, 12:288
 polymeric substrate retention by, MALDI-MS of, 11:62
 predicting mobilities of, 12:885
 protonated
 condensation reactions in, 15:1025
 derivatization of, in gas phase with acetone, 11:244
 sequence information of, derived by Pronase digestion and ammonium sulfate in-source decay MALDI-TOFMS, 11:1000
 sequencing of
 by LC/ion trap/TOFMS, 11:127
 MALDI-quadrupole TOF-MS in, 13:772
 MALDI TOF-TOF in, 13:784
 sodiated, formation in gas phase, 15:607
 sodium stabilization of, during ESI-MS, 12:505
 tryptic, field strength in separation by IM MS, 16:158
 tris(trimethoxyphenyl)phosphonium acetyl charged derivatives of, ESI-MS and MS/MS of, 11:107
 tryptic, analysis with ESI-FAIMS-MS, 13:1282
- β-Peptides, imaging distribution in whole body mice sections, MALDI-MS, 18:1921
- Peptides derived by enzymatic digestion of oxidized globin subunits, electrospray ionization-tandem mass spectrometry analysis of, an improved method to determine amino acid substitution in the hemoglobin “core,” 7:1040
- Peptides in complex mixtures, molecular weight assignment of, application of electrospray mass spectrometry and ma-

- trix-assisted laser desorption ionization time-of-flight mass spectrometry for, 4:670
- Peptidoglycan, attachment sites to by Braun's lipoprotein, characterization of by ^{242}Cf -PD and MALDI time-of-flight mass spectrometry, 6:892
- Peptidoglycans, from vancomycin-susceptible *E. faecium*, structural variation analyzed by, LC-MS, 19:1467
- Peracetylation-N, of proteins, for their structure determination by mass spectrometry, utility of, 4:87
- Peralkylated disaccharides, low energy CID of, 14:63
- Perfluorocarbon anions, poly(ethylene glycol) cation reactions with, 9:957
- Perfluorohexane, CNF bond cleavage in, 9:1158
- Performance/simplicity trade-offs, understanding ion deceleration lenses, 2:322
- Peripheral mononuclear blood cells, of HIV-infected patients, zidovudine, lamivudine, and stavudine metabolites in, HPLC-MS/MS of, 11:1134
- Permanent magnet, API-FTICR MS, 18:1552
- Permethylated glycosphingolipids, high-energy collision-induced dissociation of ceramide ions from, 5:558
- Permethylated oligosaccharides, collision-induced dissociation of, influence of the collision energy and of the collision gas for the assignment of linkage position, 4:197
- Per(onio)-substituted benzene, electrospray of, 9:15
- Perrhenate systems, ES characterization of, 12:1137
- Perturbation molecular orbital theory of gas phase dissociative electron transfer rates, 8:510
- Perylene diimide, benzannulated perylene diimide ligands and, binding to DNA, ESI-MS, 17:593
- PES. *See* Photoelectron spectroscopy
- Pesticides
 - in aqueous samples, determination of solid-phase microextraction in-line coupled to gas chromatography-mass spectrometry, 6:1119
 - carbamate, thermospray interfaces with, influence of the ion source geometry, 6:656
 - metabolism simulation using EC/LC/MS, 20:138
 - multiresidue screening in food and feed matrices, Oribtrap MS, 20:1464
 - N-methyl carbamate
 - analysis of, by liquid chromatography mass spectrometry, an evaluation of atmospheric pressure ionization techniques for, 3:378
 - ion formation of, in thermospray mass spectrometry, the effects of additives to the liquid chromatographic eluent and of the vaporizer temperature, 5:913
 - nitrogen- and phosphorus-containing, mass spectrometric analysis of, by liquid chromatography-mass spectrometry, 5:655
 - OP and CB, butyrylcholinesterase inhibition by, MALDI-TOF-MS, 18:698
 - studies of using CID, PSD, and MALDI-MS, 12:590
- PETN, negative-ion formation in, by using the reversal electron attachment detection technique, 3:243
- Petrobactin, *B. anthracis*, CID, IRMPD and EC dissociation of, 18:842
- Petroleum, automated analysis by ESI-FTICR-MS, 20:263
- Petroleum fractions, nitrogen and sulfur heterocycles in, selective chemical ionization of, by ion trap mass spectrometry, 4:322
- pH
 - α -synuclein dimer formation dependence on, monomeric structures and, 15:1435
 - myoglobin unfolding and, ESI-MS of, 11:312
 - titration of heterodimeric protein by, ESI-MS in, 11:54
- Pharmaceutical compound detection in tissue, MALDI and LD/CI in, 10:1315
- Pharmaceutical compounds, quantitative analysis of, automated nanospray of, chip-based emitters in, 16:363, 16:798 (erratum)
- Pharmaceutical industry
 - cassette dosing practices in, survey results on, 15:1374
 - chemical synthesis and structural elucidation in, bench marking survey in, 16:599
- Pharmaceuticals
 - in environment, 12:1067
 - in surface waters, comparison of matrix effects in HPLC-MS/MS and UPLC-MS/PM analyses, 19:713
- Phase angle, excitation frequency and, effect on dynamic CID in quadrupole ion trap, 18:749
- Phenetole cations, dissociation dynamics of, by photoelectron photoion coincidence, 2:464
- Phenol, reactivity of rare earth cations with, 9:1035
- Phenol-cyclohexadienone, ionized, isomeric species recognition by ion/molecule reactions, 19:126
- Phenol fatty acid esters, deprotonated, ion-dipole complex formation from, evidenced by using gas-phase labeling combined with tandem mass spectrometry, 4:343
- Phenolic compounds, in rhubarbs, analysis using LC-ESI-MS, 18:82
- Phenols
 - steric and substituent effects in, evaluation of competitive reactions of dimethyl ether ions in a quadrupole ion trap, 6:627
 - sterically hindered, MS fragmentation of, 16:814
- Phenoxyacid herbicides, chlorinated, mass spectra of, obtained by particle beam liquid chromatography/mass spectrometry, evidence for thermal decomposition contributions to, 3:823
- Phenyl allenylmethyl ethers, electron-impact-induced 3,3-sigmatropic rearrangement and cyclization in, 6:195
- Phenyl groups adjacent, in the gas phase, on the stabilization of carbanions by, 5:159
- Phenyl n-propyl ether and methyl substituted analogs, chemical ionization of, propene loss initiated by competing proton transfer to the oxygen atom and the aromatic ring, 7:639
- Phenyl radical/biradical, charged, laser photolysis synthesis of, in FTICR-MS, 9:1135
- Phenyl radical selectivity, in hydrogen atom abstraction from ethanol, 15:913
- Phenylalanine
 - complexation with NaI and KI in gas phase, cation- π effects in, 11:1037
 - effect deprotonated peptide fragmentation, 13:1242
- Phenylglycidyl ether adducts of 29-deoxycytidine and 29-deoxyadenosine, stability in solution and structure analysis by electrospray tandem mass spectrometry, 7:682
- Phenylhydrazone-oligosaccharides, from ovalbumin, MALDIMS/MS-PSD fragmentation study of, 15:725
- Phenylimino derivatization, for double bond location in polyunsaturated long chain alkenones, 15:1161
- Phenylum cation, reactions of, with small oxygen- and nitrogen-containing molecules, 7:473
- Phenylthiocarbonyl-phenylalanine, deprotonated, ion chemistry of, 8:1078
- Phenylthiohydantoin amino acid derivatives, ESI-MS of, 8:1165
- Phillips model catalyst, activation processes and polyethylene formation on, analyzed by LA-MS, LDI-MS and s-SIMS, 17:406
- Phoenix Mars Mission, thermal evolved gas analyzer, 19:1377
- Phosphates
 - alkyl and chloroalkyl, from an industrial effluent, identification of, multispectral, 2:419

- polyisoprenyl, and nucleoside pyrophosphates, glycosyl esters of, anomeric configuration of, determination of, by fast atom bombardment tandem mass spectrometry, 7:541
- polyprenyl and dolichyl, desorption chemical ionization tandem mass spectrometry of, 7:958
- replacing with methylphosphonates and thymines, in oligodeoxynucleotide fragmentation, 12:580
- trimethyl borate gas phase reactions with, 13:1088
- Phosphatidyl-*myo*-inositol mannosides, *M. bovis* BCG, characterization by MALDI-TOF MS, 18:466, 18:479
- Phosphatidylcholine isolated from heart mitochondrial fractions, analysis by fast-atom bombardment tandem mass spectrometry, evidence of incorporation of monohydroxylated fatty acyl moieties, 7:50
- Phosphatidylcholine phospholipids, characterization of, constant neutral loss scanning for, 9:1189
- Phosphatidylcholines
in brain, MALDI-MS/MS of, 16:2052
structural analysis by PSD MALDI-TOFMS, 14:373
tandem quadrupole MS of, fragmentation processes in, 14:352
- Phosphatidylethanolamine chloramines, detection using 4-chloro- α -cyanocinnamic acid and MALDI-TOF-MS, 20:867
- Phosphatidylglycerol
A. thaliana, analysis by ESI QIT and MS/MS, 18:783
triple quadrupole MS/MS studies of, 12:1036
- Phosphatidylinositol, and its phosphates, ESI-MS/MS of, 11:986
- Phosphatidylserine, MS/MS of, ESI with, 16:1510
- Phosphonium ions, stereoisomeric steroid differentiation by reactions with, 13:362
- Phosphoenolpyruvate carboxykinase, tris(trimethoxyphenyl) phosphonium acetyl charged derivatives of, MALDIPSD spectra of, 11:145
- Phospholipase A₂, N-terminal processing of, LC-MS/MS of, 15:1780
- Phospholipid chemistry, MALDI-FTMS of, strategy and data analysis techniques for, 15:1665
- Phospholipid micelles, melittin bound to, H/D exchange and ESI-FTICR-MS of, 12:1247
- Phospholipid molecular species, [¹³C]Arachidonic acid labeled, fast-atom bombardment tandem mass spectrometry of, 5:144
- Phospholipids
analysis by ion mobility TOF MS, 19:1660
in brain tissue, MALDI-TOF and MALDI-IM-TOF MS of, 16:133
headgroups of, gas phase activities of, 16:926
MALDI-TOF MS of, ionic-liquid matrices for, 16:679
matrix dry-coating for imaging of, MALDI MS, 19:882
picomole amounts of, structural determination of via electrospray ionization tandem mass spectrometry, 6:1202
sphingolipids and, DESI/MS and MS/MS of, 19:531
unsaturated acidic, negative ESI-MS structural characterization of, 16:46
- Phospholipid(s), glycerophosphoethanolamine plasmalogen, ESI-CID-MS/MS of, 15:1499
- Phosphonium ions
ketalization by 1,4-dioxane, 14:182
Meerwein reaction with epoxides and thioepoxides, in gas phase, 15:398
- Phosphonium salts, quaternary, laser microprobe mass spectrometry of, direct *versus* matrix-assisted laser desorption, 4:798
- Phosphopeptide fragmentation, in ion trap MS, 9:1175
- Phosphopeptide-Src SH₂ complexes, ESI-MS of, 8:234
- Phosphopeptides
affinity bound to immobilized metal ion beads, MALDI-MS of, 11:273
- analysis by precursor ion scanning, in positive ion mode, derivatization strategy for, 13:996
- analysis by precursor ion scanning in positive ion mode, erratum in, 14:83
- analysis using Orbitrap HCD, 20:1425
- aromatic vs aliphatic, dissociation of, ESI-FTICR-MS and IRMPD of, 15:121
- capture by zirconium phosphonate-modified magnetic nanoparticles, 19:1176
- in complex mixtures, by electrospray liquid chromatography/mass spectrometry, 4:710
- effect on ECD, 19:1263
- enrichment using PAA-membrane, MALDI-TOF-MS, 18:1385
- fragmentation of by AP MALDI and ESI coupled to ITMS, 13:274
- fragmentation pattern manipulation in, gas phase boron derivatization in, 16:1905
- IRMPD of, 15:1581
- isolated by on-line immobilized metal-ion affinity chromatography, electrospray ionization mass spectrometry of, 4:662
- MALDI-MS detection of, 10:35
- new derivatization strategy for analysis of, by precursor ion scanning in positive ion mode, 13:996, 14:83
- in phosphorylation sites, by positive and negative mode electrospray ionization-tandem mass spectrometry, 7:243
- picolinamidation for MALDI-TOF-TOF-MS sequencing, 20:1751
- product ions generated by 157 nm photodissociation of, 20:2334
- quantification with affinity chromatography and stable isotope labeling, 15:363
- retained on ferric NTA IMAC beads, solubilization of, MALDI-TOF-MS of, 13:1042
- in situ enrichment via metal oxide surfaces, 20:915
- sulfopeptides and, atmospheric pressure thermal dissociation, 19:1898
- Phosphorothionates, protonated O,O-diethyl O-aryl, fragmentation of, in tandem mass spectral analysis, 7:189
- Phosphorus, chemical reaction interface mass spectrometry in detection of, nitrogen trifluoride in, 6:421
- Phosphorus- and nitrogen-containing pesticides, mass spectrometric analysis of, by liquid chromatography-mass spectrometry, 5:655
- Phosphorylated carbohydrates, isomeric differences of, probing with ion/molecule reactions and FTICR-MS, 14:323
- Phosphorylation
peptide ionization efficiency changes caused by, 18:1973
protein, stoichiometry determined by MS, 20:2211
sites of, identified in MARCKS peptides by ECD FTICR MS/MS, 18:2137
sulfation distinguished from, glycoproteins and carbohydrates, ion-pairing and MS, 17:1282
- Phosphorylation sites in phosphopeptides, identification of, by positive and negative mode electrospray ionization-tandem mass spectrometry, 7:243
- Phosphotyrosine, cationized, fragmentation by atmospheric pressure MALDI/ion trap spectrometry, 14:581
- Photo-induced dissociation of precursor ions, high efficiency, tandem time-of-flight mass spectrometer, 4:751
- Photochemistry, unimolecular, of n-alkenes studied by photodissociation-photoionization mass spectrometry, 4:65
- Photodissociation
193 nm
characteristics of singly protonated peptides generated by MALDI, 17:1643
compared with TOF-TOFMS, bradykinin analogues, 17:721

- 193-nm
of ions, from saturated and unsaturated aliphatic molecules, 7:114
laser, for isomer differentiation in Fourier-transform mass spectrometry, 1:361
of peptide ions, in Fourier transform mass spectrometry, 1:288
of biopolymer ions generated by MALDI, tandem TOF MS for, 15:1248
bond-selective, of aliphatic disulfides, 6:872
of high molecular weight peptides and proteins in a two-stage linear time-of-flight mass spectrometer, 6:578
infrared, of non-covalent adducts of electrosprayed nucleotide ions, 7:209
laser, determination of positions, velocities, and kinetic energies of resonantly elicited ions in the quadrupole ion trap mass spectrometer by, 4:792
157nm, structures of *y*- and *b*- type peptide ions, 19:695
157nm nanometer, singly-charged α -type peptide ions formed in, 17:1315
of peptide-crown ether complexes by UV irradiation, 19:257
peptide fragmentation of photolytically generated radical ions, 20:1378
product ions generated by 157 nm photodissociation of phosphopeptides, 20:2334
rapid UV, of peptides using naphthyl-based adducts, 20:385
of small group-11 metal cluster ions, fragmentation pathways and photoabsorption cross sections, 14:614
time-resolved
of peptides by MALDI, 20:1522
statistical interpretation of, singly protonated peptides, 19:645
time-resolved method for mechanistic study of protonated peptides, TOF MS/MS, 18:1729
tunable UV-type, LIT in, heparin-derived disaccharides, 20:1645
Photodissociation-photoionization mass spectrometry, unimolecular photochemistry of *n*-alkenes studied by, 4:65
Photodissociation spectroscopy of gas-phase ferrocene cation, 2:97
Photoelectron photoion coincidence, dissociation dynamics of phenetole cations by, 2:464
Photoelectron resonance capture ionization (PERCI), for soft ionization of organic compounds, 15:424
Photoelectron spectroscopy
of CH_3I^- anion, 10:810
of the trimethylenemethane negative ion, 10:800
Photofading, ballpoint dyes on paper, LDI and MALDI MS analysis of, 17:297
Photofragmentation
157nm, analysis of LTC₄ isomers using CID and, 19:14
VUV, of peptide ions, 18:1439
Photoinduced changes, in metal-ion complexes of spirobenzopyran derivatives, evaluating by ESI-MS, 14:1110
Photoion coincidence, photoelectron, dissociation dynamics of phenetole cations by, 2:464
Photoionization
atmospheric pressure, reduction in, 13:109
ion-neutral complexes [$\text{CH}_3\text{CH}^+\text{CH}_3$, CH_2CH_3] and [$\text{CH}_3\text{CH}_2\text{CH}^+\text{CH}_3$, CH_3] in the gas phase, H-transfer and C-C bond formation between partners, and channeling of energy into dissociation, 7:73
of large molecules, 6:883
of hexyl alcohol, 15:989
vacuum ultraviolet, ion decompensation *versus* molecular size probed by, 2:103
Photoionization 193 nm, laser, for isomer differentiation in Fourier-transform mass spectrometry, 1:361
Photoionization mass spectrometry
10.5 eV, of aliphatic compounds, 1:419
mechanism of $[\text{M} + \text{H}]^+$ formation in, 15:1521
Photoionization of large molecules, 6:883
Photolabile arylazido groups, peptides modified with, MALDI-MS analysis of, 15:1156
Photomodified oligodeoxynucleotides
adducted with metal ions, fragmentation in ESI-ITMS, 12:1174
determination by exonuclease digestion, MALDI, and PDSMS, 12:1127
Photon dissociation, infrared multiple, in the quadrupole ion trap via a multipass optical arrangement, 5:886
Photon-induced dissociation, with a four-sector tandem mass spectrometer, 1:107
Photosystem II, structure and dynamics by high resolution FTICR MS, 19:1004
Phthalimide radical cations, N-(substituted benzalamino), decompositions of, embody ion-neutral complexes and Stevenson's rule, 5:837
Phthalocyanine, copper, heterolytic dissociation of negative ions into counter ions, 19:375
Physical/chemical separations, break-up of highly charged droplets from electrosprays, 12:343
Phytochemicals, carbohydrates and, profiling in tulip tissues and bulbs, PESI MS, 20:2304
Phytyl methyl ether, ionized, $\text{C}_3\text{H}_9\text{O}^+$ formation from, structure and mechanisms of, 7:205
Picolinamidation, phosphopeptides, for MALDI-TOF-TOF-MS sequencing, 20:1751
Picolinyl esters
of bacterial fatty esters, 14:58
studies on fragmentation mechanisms, 17:1620
Picomole amounts of phospholipids, structural determination of via electrospray ionization tandem mass spectrometry, 6:1202
Pictograms, for experimental parameters in mass spectrometry, 8:756
PID. *See* Precursor ion discovery
PIF. *See* Precursor ion fingerprinting
Pigments, dyes and, MALDI TOF MS identification, 18:2001
Pilocarpine, reaction sites of, 10:209
 α -Pinene, camphene and, bimolecular and unimolecular contributions to SCI of isomers, 18:2026
Pinus spp., isotope ratios of carbon and oxygen in whole foliage and cellulose, 19:1330
Piperidine, DNA duplex destabilization induced by, in production of single-stranded PCR amplicons, 13:232
Pipette tips, chromatographic, nano-ESI-MS/MS from, 15:1333
PittCon 2003 review, 14:542
Pivaloyl cation, carbocationic rearrangement of, 15:959
Plant virus, identification from infected plants, investigative proteomics in, 14:736
Plants, biotic/abiotic boundary of, probing with LDI-TOF MS, 16:107
Plasma
atorvastatin and active metabolites in, HPLC-MS/MS of, 10:55
desorption, Californium-252, high resolution and tandem Fourier-transform mass spectrometry with, 1:427
dog, nonpeptide antithrombotic in, quantitative determination of, by microbore high-performance liquid chromatography-tandem mass spectrometry utilizing pneumatically assisted electrospray ionization, 7:1238
glow discharge, characterization of, as a function of sampling orifice potential, 1:448
HT2a antagonist and metabolite in, LC-MS/MS of, 8:371

- human
 chemically oxidized dexamethasone in, electron capture negative ionization in, comparison of tandem and conventional mass spectrometry using, 1:341
 SR 27417 in, using electrospray liquid chromatography-tandem mass spectrometry, study of ion suppression, 7:1099
 inductively coupled MS of, 8:1230
 metal ion, laser-generated, competitive ionization of tetraphenylporphyrin in, 4:159
 pulse glow discharge, afterpeak time regime of, ion formation processes in, 4:727
 reagent thermospray, from ammonium acetate, controversial ions (m/z 59, 5:60,5:77, and 119) in, 5:186
- Plasma desorption mass spectrometry, ^{252}Cf
 adsorption/desorption in, small molecules as, 2:29
 fragmentation analysis of bradykinin by, 2:379
- Plasma desorption mass spectrometry- ^{252}Cf , background reduction in, a pulsed ion deflection system for, 3:706
- Plasma mass spectrometry
 inductively coupled
 with an electrically floating sampling interface, 4:733
 with an enlarged sampling orifice and offset ion lens ion trajectories and detector performance, 4:16
 poly atomic ion interferences and matrix effects, 4:28
 determination of chlorine isotope ratios by, alleviation of polyatomic ion interferences for, 1:284
 mixed-gas helium-argon inductively coupled, matrix-induced interferences in, 2:355
 space-charge effects in, 10:159
 time-resolved measurements of individual ion cloud signals to investigate, 7:362
- Plasma-mass spectrometry, inductively coupled, interference effects in, selection of solvent load and first-stage pressure to reduce, 3:128
- Plasma peptides, LC-MS mapping of, 10:45
- Plasma profiling, of arecoline, in treated Alzheimer patients, an automatic reaction control chemical ionization technique in ion trap detector for, 2:168
- Plasma source ion trap mass spectrometry, enhanced abundance sensitivity by resonant ejection of atomic ions, 7:1161
- Plasma-source mass spectrograph, design and performance of, 8:307
- Plasma-source mass spectrometer, multichannel, novel, 7:458
- Plasma source mass spectrometry, 14:171
- Plasma-surface interface reactions, ion-molecule reactions to, 13:518
- Plasma-time-of-flight mass spectrometer, an inductively coupled, for elemental analysis, optimization and characteristics, 5:1008
- Plasticizer contamination, from vacuum system O-rings in a quadrupole ion trap mass spectrometer, 13:901
- Platelet-activating factor
 acyl analogues of, heptafluorobutyl derivatives of, mass spectral fragmentation of, novel, 2:476
 GC/MS of, 11:176
- Platinum, in urine, isotope dilution gas chromatography/mass spectrometry for, 2:85
- Platinum complexes, MS behavior of, 16:94
- Platinum wire, optical fibers coiled with, electrospray from, 11:464
- PLDPI-MS. *See* Laser desorption postionization mass spectrometry
- Plicatamide, fragmentation of, gas phase intramolecular rearrangement in, 12:470
- Plutonium
 240/239 ratios in Lake Ontario sediments, 2:240
 in an aluminosilicate matrix, particulate isotopic standard, 2:81
 particulate isotopic standard in an aluminosilicate matrix, 5:852
- Pneumatic splitter electrospray ionization, nanoliter flow analysis of protein digests using, 8:703
- Pneumatically assisted electrospray ionization, microbore high-performance liquid chromatography-tandem mass spectrometry utilizing, quantitative determination of a non-peptide antithrombotic in dog plasma by, 7:1238
- $\text{P}(\text{OH})_2+/\bar{-}$, ionic and neutral in gas phase, MS/MS and computational chemistry of, 13:250
- Point mutations, changes in protein hydrogen exchange rates that result from, mass spectrometric measurement of, 7:515
- Poisoning, acute, gas chromatography-mass spectrometry in, scope and limitations of in general unknown screening, 6:995
- Polar $[4+2^+]$ Diels-Alder, cycloaddition to nitrilium and immonium ions in the gas stage, applications of multiple stage mass spectrometry in a pentaquadrupole, 6:1
- Polar side groups, dipeptides containing, by Curie-point pyrolysis tandem mass spectrometry, 3:750
- Polarity, and organic molecule responsiveness to negative ion ESI MS, 16:446
- Polarizability and inductive effect contributions to solvent-cation binding observed in ESI-MS, 10:254
- Polarizable ions, collision rate constants for, 14:501
- Polarizable stationary phase, negative ion chemical ionization, and mass-resolved chromatography, high-temperature gas chromatography/mass spectrometry with, high-resolution triglycerol mixture analysis using, 7:350
- Policies, new, for JASMS, 17:1
- Polistes dominulus*, dominulin A and B in cuticle and venom of, analyzed by MALDI-TOF, MALDI-TOF/TOF and ESI-ion trap, 17:376
- Pollutants
 atmospheric, real time monitoring of, ion trap detector used for, effects of air and helium, 4:774
 emerging, in environment, 12:1067
- Poly-histidine tags, enrichment of phosphorylated peptides using, 18:1007
- Polyacrylamide gel electrophoresis, of CD4/Ic-copurifying proteins, 15:558
- Poly(alkylmethacrylate), MALDI-CID of, 8:1206
- Poly(amic acid) and polyimide characterization, using gas chromatography/mass spectrometry and particle beam liquid chromatography/mass spectrometry, 5:299
- Polyamides, evaluation of binding to DNA, by ESI-MS, 17:1742
- Polyamidoamine dendrimers
 investigation of silver binding, ESI-MS/MS, 17:676
 observation of b^* , y cleavages, ECD-MS, 17:536
 structural analysis by ECD, EDD and CID, 19:1312
- Polyamindoamine dendrimers, ECD and CID of metal ion complexes with, 20:674
- Polyamine complexes, gas phase H/D exchange reactions of, 11:711
- Polyamine toxins, of spiders, electrospray ionization tandem mass spectrometry of, 6:634
- Polyamines
 gas phase basicities of, 9:1043
 noncovalent DNA complexes with basic polypeptides or, MALDI-TOF MS, 18:1977
- Polyammonium macrotricyclic ligands, dicarboxylic acid complexes with, ESI-MS of, quantum calculations and, 12:304
- Polyaniline, as conductive polymer coating for durable nanospray emitters, 11:659
- Polyaniline oligomers, MALDI analysis of, 15:893
- Polyanionic analytes, negative ion fast-atom bombardment mass spectrometry of, class of cationic matrix additives for, N-alkylnicotinium halides, 5:935

- Polyaromatic hydrocarbon-deoxynucleoside adducts, identification of, by capillary zone electrophoresis-continuous flow-fast atom bombardment mass spectrometry, 3:757
- Polyaromatic hydrocarbons, detection of, mobile phases and VUV lamps in, APPI-MS, 18:589
- Polyatomic fragment losses, atomic fragment losses slower than, 14:136
- Polyatomic ion excited electron states, detection by charge exchange ionization in collision cells, 12:1120
- Polyatomic ion interferences
alleviation of, for determination of chlorine isotope by inductively coupled plasma mass spectrometry, 1:284
and matrix effects, inductively coupled plasma mass spectrometry with an enlarged sampling orifice and offset ion lens, 4:28
- Polyatomic ions
collision-activated dissociation of, at low energy, a dynamics study of simple C-C cleavage in, fundamentals of tandem mass spectrometry, 1:6
collisions with surfaces, 14:1360
- Polybrominated diphenyl ethers, analysis in soil, dust and human serum, HF-LPME with GC-ICP-MS, 18:1740
- Polychlorinated biphenyls, correlation between electron capture negative chemical ionization mass spectrometric fragmentation and calculated internal energies for, 5:44
- Polycyano push-pull pentadienes, from cycloalkylidene malonic acids, EI-MS of, 14:189
- Polycyclic aromatic compounds, analysis with APCI-MS, charge exchange and proton transfer in, 15:301
- Polycyclic aromatic hydrocarbon
C₁₀ to C₁₆₀, flame-generated, by atmospheric-pressure chemical ionization mass spectrometry with liquid introduction via heated nebulizer interface, 7:276
ion trap MS/MS of, 8:183
nitrated, LDI TOF-MS of, 8:630
- Polycyclic aromatic hydrocarbon diolepoxide adducted peptides, characteristics of high energy collision-induced dissociation tandem mass spectra of, 5:649
- Polycyclic aromatic hydrocarbon (PAH) dihydriol-epoxides, deoxynucleoside adducts, low energy tandem mass spectrometry of, 6:248
- Polycyclic aromatic hydrocarbon radical cations, reactions with model biological nucleophiles, 7:628
- Polycyclic aromatic hydrocarbons
comparison of dopants for charge exchange ionization, RP-LC-APPI-MS, 20:73
fast atom bombardment combined with mass spectrometry for, 1:405
by high-performance liquid chromatography-particle beam-mass spectrometry, 4:898
in solid matrices, 14:562
- Polycyclic aromatic hydrocarbons in ancient terrestrial rocks, μ L2MS vs GC/MS of, 12:989
- Polycyclic aromatic hydrocarbons (PAHs), high-molecular-weight, measurement in soils by particle beam high-performance liquid chromatography-mass spectrometry, 6:597
- Polycyclic polyisoprenylated benzophenones, nemorosone degradation profile by ESI-MS, 20:1688
- Polycyclic polyisoprenylated acylphloroglucinols, *Garcinia* spp., screening using PID and UPLC-ESI-Q-TOF, 20:1846
- Polydeuterated surfactants, isotopic purity in, charge-remote fragmentation in a hybrid (BEqQ) mass spectrometer, 1:85
- Polydiene, MALDI-TOF-MS for, 8:1220
- Poly(dimethyl siloxane) polymers, MWD using TOF-SIMS, 20:1562
- Poly(dimethyl siloxanes), comparative studies of, 14:195
- Polydimethylsiloxane membrane, *Candida rugosa* lipase immobilized on, use in MIMS, 18:973
- Polydimethylsiloxane multichannel emitter, microfabricated, for ESI-MS, 12:463
- Polyelectrolytes, as matrices for secondary ion mass spectrometry, 16:1575
- Polyesteramides, hyperbranched
ECD and CAD-MS of, 14:332
in-source decay of, MALDI-TOFMS of, 11:218
- Polyether, transition metal ion complex with, quadrupole ion trap, 8:620
- Polyether/ammonium ion complexes, gas-phase, hydrogen-bonding interactions in, 5:260
- Polyether cationization, alkali metal ion effects, in MALDI, 11:639
- Polyether ionophore antibiotics, alkali metal complexes of, gas phase stabilities of, by CID, 16:779
- Polyethers
gas-phase basicities and ammonium ion affinities of, kinetic method and ligand exchange technique, 4:718
gas-phase complexation of, with halide ions, 4:242
- Polyethylene
evaluation of field desorption mass spectrometry for, 7:1070
formation on Phillips model catalyst, activation processes and, analyzed by LA-MS, LDI-MS and s-SIMS, 17:406
LDI and MALDI-TOFMS of, 12:1186
- Polyethylene glycol
automated MALDI-TOF MS of, 16:670
as shift reagent in lamivudine analysis, IMS, 20:1
- Poly(ethylene glycol), doubly and singly cationized by different alkali metal ions, relative cation affinities and cation-dependent resolution in a quadrupole ion trap mass spectrometer, 13:177
- Polyethylene glycol-conjugated superoxide dismutase, fingerprinting proteins coupled with polymers by mass spectrometry and, 6:478
- Poly(ethylene glycol) esters, low energy CID ESI-MS of, 13:888
- Poly(ethylene oxide)/polystyrene block copolymer, microstructural study by ESI MS/MS, 19:1163
- Poly(ethylene terephthalate) oligomers, cationized by alkali ions: structures, energetics, effect on mass spectra and the MALDI process, 10:883
- Poly(ethylene)glycol, external ion accumulation of, by ESI-FTMS, 10:1298
- Polyfluorinated chemicals, ionization pathways of, environmental atmospheric pressure and, 14:1177
- Polyglycine, carbon-13 and nitrogen-14 labeled, CN⁻ secondary ion formation, 17:1181
- Polyglycols
ESI low energy CID MS/MS of, 12:832
linear and cyclic, [M + X]⁺ ions (X=H, Li, Na, K) from, dissociation characteristics of, 5:1081
lithium-attachment ions from, tandem mass spectrometry of, 3:225
- Polyhydroxy unsaturated fatty acids, electrospray ionization and low energy tandem mass spectrometry of, 7:140
- Polyimide characterization, using gas chromatography/mass spectrometry and particle beam liquid chromatography/mass spectrometry, 5:299
- Polyiodide ions, thermochemistry of, 8:688
- Polyisobutylene, derivatives of, dopant-assisted APPI-MS, 20:2342
- Polyisobutylene derivatives, APPI-MS of, 19:656
- Polyisoprenyl phosphates and nucleoside pyrophosphates, glycosyl esters of, fast-atom bombardment tandem mass spectrometry, 7:541

- Polyactides, solid state NMR and MALDI-TOF MS analysis, 20:67
- Polyactosamine repeats, in lipooligosaccharides from *Haemophilus ducreyi*, 13:724
- Polymer
 identifying additives in, TPPy/MAB-MS for, 15:1315
 MALDI-MS of, sample deposition for, 15:168
- Polymer distributions, FTMS of, isotope beating effects in, 10:1074
- Polymer mixtures
 DE-MALDI-TOFMS of, isobaric interferences in, 12:911
 quantitative analysis by MALDI-TOF-MS, 13:914
- Polymerase chain reaction, genetic diagnosis of familial adenomatous polyposis, 8:737
- Polymerase chain reaction-mass spectrometry-based analyses, *Thermococcus kodakaraensis* (KOD) DNA polymerase for, 14:601
- Polymerase chain reaction products, ESI-FTICR of, 11:333
- Polymeric electrodes, conductive carbon-filled, as ion optical elements for TOF-MS, 13:1170
- Polymeric substrates, peptide retention of, MALDI-MS of, 11:62
- Polymerization, ethylene catalyzed by Ti⁺ in the gas phase, 3:464
- Polymers
 color bodies on surface of, LC-MS/MS of, 16:295
 fourth annual NIST Polymer Mass Spectrometry Workshop, review, 17:280
 MS of, 2004 ASMS Fall workshop on, review of, 16:291
 nonpolar, MALDI-MS of, silver cluster interferences in, 12:732
 quantitation by MALDI-TOF MS, 16:100
 synthetic
 end group determination of, by ESI-FTICR, 11:536
 multisample preparation methods of, MALDI-MS, 18:377
 parameters in optimization of MALDI-TOF MS for, 17:246
 software for peak assignment from MS/MS, 18:1324
- Polymer(s)
 biodegradable, quantitative TOF-SIMS of, 13:1108
 bis(imidazole-1-carboxylate) end-functionalized, MALDI-MS study of, 14:117
 characterization with SFC-MS and image analysis-based quantitation, 13:1195
 MALDI-FTMS of, 8:225
 MALDI-MS of, time lag focusing and cation attachment in, 8:132
 sample holder for, 8:1214
 synthetic, LSI-MS/MS of, 8:76
- Polymers, MALDI-TOFMS analysis of
 fingerprinting proteins coupled with, by mass spectrometry, investigation of polyethylene glycol-conjugated superoxide dismutase, 6:478
 of high molecular weight, electrospray of, 9:299
 hydrocarbon, laser desorption Fourier transform mass spectrometry, silver nitrate chemical ionization for, 4:453
 low molecular weight, high resolution end group determination of, by matrix-assisted laser desorption ionization of an external ion source Fourier transform ion cyclotron resonance mass spectrometer, 7:449
 MALDI-TOFMS analysis of
 accuracy of, 9:275
 solvent effect in, 9:1303
 organic, UV ablation products of, direct chemical analysis of, by using selective ion monitoring mode in gas chromatography/mass spectrometry, 5:106
 poly(methylmethacrylate), low mass, application of matrix-assisted laser desorption-mass spectrometry to, fundamentals of, 7:287
 synthetic, 9:267
- Poly(methacrylic acid), analysis by MALDI-TOF MS, 18:1507
- Poly(methyl methacrylate) polymers, low mass
 application of matrix-assisted laser desorption-mass spectrometry to, 7:287
 electrospray ionization mass spectrometry and, 6:906
- Poly(methylacrylic acid), oligomers produced by ESI, MS/MS analysis, 20:25
- Polymorphisms, single nucleotide, semi-quantitative genotyping of, LC-MS for, 15:1897
- Polynuclear aromatic hydrocarbons, nitro-substituted, in airborne particulate matter and soil, sub-parts-per-billion determination by electron capture-tandem mass spectrometry, 7:1255
- Polyol mixtures, complex, MS analysis of, 12:55
- Polyoxyethylene dodecylethers, alkali cation affinities of, ESI MS, 18:1914
- Polypeptide, semisynthetic, unusual amino acid in, use of high- and low-energy collision-induced dissociation tandem mass spectrometry in the identification of, 7:1034
- Polypeptide ions, model large, RRKM calculations on, effect of ion size on rate of dissociation, 4:11
- Polypeptides
 deprotonated, transition metal complex cations in gas-phase transformation, 20:1718
 ECD of, in three-dimensional quadrupole ion trap, 16:22
 fragments of, LC-MS analysis of, 16:38
 isotope cluster calculation by probability grouping, 20:295
 matrix-diluted, bombarded with slow and fast projectiles, TOF-SIMS of, 11:553
 methylated, electron transfer dissociation, 20:1349
 multiply charged
 dissociation of, high-energy collision-induced, produced by electrospray, 4:652
 large, produced by electrospray ionization, collision activation and collision-activated dissociation of, 1:53
 noncovalent DNA complexes with basic polyamines or, MALDI-TOF MS, 18:1977
- Polyphosphoesters, characterization by FT-ICR-MS, 20:2238
- Polyposis, familial adenomatous, diagnostic PCR and ESI-MS in, 8:737
- Polyprenyl and dolichyl phosphates, desorption chemical ionization tandem mass spectrometry of, 7:958
- Polypropylenamine dendrimers, dissociation of, 10:414
- Polypropylene glycol, multiple charging by binary mixtures of cations, in electrospray, 16:152
- Polypropylene tubes, MS interferences from, identification of, 16:417, 16:802 (erratum)
- Polypyryl/adenosine triphosphate films, adenosine triphosphate release from, EC/ESMS, 18:919
- Polysaccharides, alkalicationed, MS/MS of, 8:987
- Polysiloxanes, TOFMS of, 9:292
- Polystyrene
 influence of organic layer thickness in TOF-SI-MS, 20:2294
 MALDI-TOFMS of, postsource decay in, 11:1095
- Poly(styrene), MALDI-CID, 9:269
- Poly(styrene) oligomers, cationized, gas phase conformations of, 13:499
- Polystyrenes
 analysis by cryodetection MALDI-TOF MS, 19:219
 end-functionalized, MALDI-MS of, 14:648
 metal attachment to, in MALDI, 9:724
- Polythiophene film, electrodeposited, for LDITMS, 8:1203
- Polythymine oligonucleotides, interactions among ethidium bromide, mercury ions and, ESI MS, 20:1834
- Polyunsaturated fatty acid methyl esters, CI-MS/MS of, 10:1253
- Porous anodic alumina membrane
 phosphopeptide enrichment using, MALDI-TOF-MS, 18:1385

- as sample support for MALDI-TOF MS of salt-containing proteins, 16:1488
- Porous silicon, surface area drives desorption of intact molecules, 18:1945
- Porphorinoids, phenyl-fused, observations using ESI MS, 17:1306
- Porphyrins
double-charged, unimolecular and collision induced reactions of, 3:235
high-performance liquid chromatography-mass spectrometry, using an atmospheric pressure interface, 7:965
meso-tris-*N*-methylpyridinium-4-yl, reduction in positive mode ESI-MS, 18:762
vinylpyridinium substituted, characterization by ESI-MS, 18:218
- Portable systems
3D gas concentration mapping using, 19:1411
DART and DESI MS implementation on fieldable MS, 19:1419
development of a palm portable MS, 19:1442
hand portable GC-TMS for hazardous compound detection, 19:1425
lake water testing of chemistry with AUV-based MS and, 19:1403
- Positive and negative ion fast-atom bombardment high-resolution mass spectrometry, reference compound for, Ultramark 1621 as, 3:842
- Positive ion electrospray ionization mass spectrometry of oligonucleotides, 8:90
- Positive ion electrospray ionization-tandem mass spectrometry, of cleavage reactions of complex ions, 15:689
- Positive ion formation in the ultraviolet matrix-assisted laser desorption/ionization analysis of oligonucleotides by using 2,5-dihydroxybenzoic acid, 8:218
- Positive ion mass spectra, from APCI-MS gases and solvents used in LC and direct liquid injection, 15:311
- Post decay source (PSD) analysis, of protonated cross-linked peptides, 13:709
- Post-ionization separations, FAIMS in, modeling resolution and sensitivity of, 15:1487
- Post source decay, parallel approach to, MALDI-TOF analysis, 17:60
- Post-source decay, analysis of oligosaccharides, MALDI linear-field reflectron TOF, 19:1119
- Post-source decay in analysis of polystyrene by MALDI-TOFMS, 11:1095
- Post-source decay MALDI-TOFMS, phosphatidylcholine structural analysis by, 14:373
- Post-source decay mass spectrometry (PSDMS), photomodified oligodeoxynucleotide determination by, 12:1127
- Post-source decay (PSD)
of diastereomeric benzo[ghi]fluoranthene tetrols, 12:376
improved, on MALDI-TOF, 15:522
of pesticides, 12:590
- Post-translational analysis, qQq-FTICR mass spectrometer for, 16:1985
- Postcolumn addition of propionic acid and isopropanol, signal enhancement for gradient reverse-phase high-performance liquid chromatography-electrospray ionization mass spectrometry analysis, trifluoroacetic and other strong acid modifiers in, postcolumn addition of propionic acid and isopropanol, 6:1221
- Postcolumn derivatization liquid chromatography-mass spectrometry for detection of chemical-weapons-related compounds, 10:440
- Postsource decay fragmentation, of *N*-linked carbohydrates from ovalbumin and related glycoproteins, 11:572
- Potassium, complexation with aromatic acids in gas phase, cation- π effects in, 11:1037
- Potassium halide adducts, as reagent ions in infrared laser desorption/ionization Fourier transform ion cyclotron resonance mass spectrometry, 3:301
- Potential build-up and gradient mapping, in differential electrospray emitter, 15:1737
- PP/ATP films. *See* Polypyrrol/adenosine triphosphate films
- PPAPs. *See* Polycyclic-polyprenylated acylphloroglucinols
- Precipitation, serum protein, with antiserum by matrix-assisted laser desorption ionization/time-of-flight and electrospray ionization mass spectrometry as a clinical laboratory, 6:854
- Precise peptide sequencing, and protein quantification, in the human proteome through in vivo lysine-specific mass tagging, 14:1
- Precolumn delivery, *versus* coaxial matrix, continuous flow fast atom bombardment with packed microcolumns, 1:312
- Preconstructor, GC-FTMS coupled to, 15:1191
- Precursor ion discovery, *Garcinia* spp. PPAP screening using UPLC-ESI-Q-TOF and, 20:1846
- Precursor ion fingerprinting, determination of ion structures using, 20:370
- Precursor ion-independent algorithm, for top-down shotgun proteomics, 20:2154
- Precursor ion scanning
for detection and structural characterization of heterogeneous glycopeptide mixtures, 13:1065
in positive ion mode, for phosphopeptide analysis, derivatization strategy for, 13:996
- Precursor ions, photo-induced dissociation of, high efficiency, in a tandem time-of-flight mass spectrometer, 4:751
- Prediction of peptide ion mobilities, via a priori calculations from intrinsic size parameters of amino acid residues, 12:885
- Preference for an ion-neutral complex-mediated pathway over a five-membered ring H shift in the isomerization of $\text{CH}_3 + \text{HCH}_2\text{CH}_2$ to $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}^+$ by ab initio theory, 6:1037
- Preparation and photoelectron spectrum of the CH_3I^- anion: rare gas cluster-mediated synthesis of an ion-radical complex, 10:810
- Preparative scale mass spectrometry, brief history of calutron, 8:943
- Pressure conditions, reduced, comparison of the reactions of the oxide radical anion (O^-) with simple aromatic compounds at, 5:367
- Pressure limited sustained off-resonance irradiation, for collision-activated dissociation in FTMS, 10:15
- Primaquine, mimetic derivatives of, ESI-MS, 19:1476
- Primary products, nature of, effects of the class and chain length of alcohols on, gas phase reactions of Sc^+ , Y^+ and Lu^+ with alcohols, 7:1157
- Primary thermal degradation processes of peptides, characterization of, using the mass spectrometric technique K^+IDS , K^+ ionization of desorbed species, 5:564
- Prion protein peptide, tandem mass spectrometry of peptides with *N*-terminal glutamine studies on, 1:258
- Prion proteins, detection of attomoles by nanoLC MS/MS, 18:1070
- PRISM. *See* Protein interactions from the spectra of masses
- Probabilities of correct identification, estimating of, from results of mass spectral library searches, 5:316
- Probability grouping, isotope cluster calculation by, polypeptides, 20:295
- Probe
molecular, for the interaction of methanol and protonated

- peptides, gas-phase hydrogen/deuterium exchange as, 5:526
- multisample, for fast-atom bombardment/mass spectrometry, 4:504
- Probe electrospray ionization mass spectrometry, phytochemical and carbohydrate profiling in tulip tissues and bulbs, 20:2304
- Probe insert designs, thermospray, that allow clogged probes to be rapidly replaced, 4:182
- Probes
- clogged, two thermospray probe insert designs that allow to be rapidly replaced, 4:182
 - triaxial, for proteolysis, coupled with HDX-MS, 18:208
- Probing high order structure of proteins, by fast-atom bombardment mass spectrometry, 5:19
- Probing isomeric differences, of phosphorylated carbohydrates through the use of ion/molecule reactions and FTICRMS, 14:323
- Probing the helical content, of growth hormone-releasing factor analogs using electrospray ionization mass spectrometry, 4:646
- Probing trapped ion energies, via ion-molecule reaction kinetics Fourier transform ion cyclotron resonance mass spectrometry, 3:727
- quadrupole ion trap mass spectrometry, 3:716
- Process mass spectrometry, in simultaneous analysis of butene isomer mixtures, 11:1079
- Prodiginines, odd-electron fragments, positive ion ES MS/MS, 19:1857
- Product ion characterization, by linked-scan tandem mass spectrometry, trimethylsilyl group migration in the mass spectra of trimethylsilyl ethers of cholesterol oxidation products, 4:327
- Product ion spectra, dipeptides, study of $b_1 + H_2O$ and b_1 -ions in, 18:1414
- Product ion studies, of diastereomeric benzo[ghi]fluoranthene tetrols by MALDI-TOFMS and PSD, 12:376
- Productivity, improved, networking mass spectrometer data systems for, 4:742
- Proferrioxamines
- collision-activated fragmentation of, evidence for a succinimide-mediated mechanism, 6:836
 - on the collision-activated fragmentation of, evidence for a succinimide-mediated mechanism, 6:836
- Profile generation model, of mass peak, 8:170
- Projectiles, matrix-diluted oligo- and polypeptides bombarded with, TOF-SIMS of, 11:553
- Proline
- b_2 ion containing, gas phase ion structure of, 17:20
 - residues of aspartic acid and, effect on peptide dissociation, 19:344
- Proline analogs, proton affinity of, 13:72
- Pronase digestion, peptide sequence by, 11:1000
- Propanediol ions, solvation by water molecules in gas phase, 15:1123
- Propenal, Ion reactions with ethylene oxide and, selected ion flow tubes, 18:445
- Propene, loss of, from $[M + D]^+$ ions, 9:121
- Propene loss
- initiated by competing proton transfer to the oxygen atom and the aromatic ring, chemical ionization of phenyl n-propyl ether and methyl substituted analogs, 7:639
 - from propolypyridines, deuterium transfer initiation of, 14:1166
- Propionic acid and isopropanol, postcolumn addition of, signal enhancement for gradient reverse-phase high-performance liquid chromatography-electrospray ionization mass spectrometry analysis, trifluoroacetic and other strong acid modifiers in, 6:1221
- Propionitrile, identification in acetonitrile, APPI MS, 19:1579
- Propofol, MS fragmentation of, 16:814
- Propolypyridines, propene loss from, deuterium transfer initiation of, 14:1166
- Propyl propanoate, ionized, formation of the 3-pentanone ion from, through ion-neutral complexes, 3:183
- Prostaglandins
- fast atom bombardment and collision-induced dissociation of, some examples of charge remote fragmentation, 1:325
 - structurally similar, E2, F2alpha, D₂ and DHKF2alpha, low-energy collision-activated dissociation of carboxylate anions, 1:389
- Protease digests, gas phase separations of, 9:1213
- Proteasome, 20S, ESI-MS and ion mobility analysis of, 16:998
- Protectin D1, resolvin D1, and related derived products of DHA, analysis via ES-MS/MS, 18:128
- Protein
- antibiotic linked to, released by ECD-FTICR-MS, 14:302, 14:682
 - carbon isotope ratio determination in, ICP-MS for, 14:295
 - opioid peptide-containing, characterization of, by electrospray ionization and liquid secondary ion mass spectrometry, 5:377
- Protein analyses, applying electrospray ionization-mass spectrometry to, charge discrimination with, 6:1098
- Protein and peptide mixtures, capillary electrophoresis Fourier transform ion cyclotron resonance mass spectrometry with sustained off-resonance irradiation for, 5:894
- Protein and proteome research, MS journey into, 14:685
- Protein antigens, in cell lysates, 9:208
- Protein-carbohydrate complexes, nonspecific, produced by nanoESI, BIRD of, 16:1583
- Protein complexes
- chemical cross-linking of, ESI-FTICR-MS and ESI-TOF-MS in, 16:1921
 - Coulombic repulsion effect on dissociation pathways in gas phase, 18:617
 - effects of single amino acid substitution on dissociation in gas phase, 18:688
 - ESI mechanism features affecting, 15:1424
 - LC-MALDI-TOF/TOF analysis of, 15:803
 - noncovalent, analysis by MALDI MS, 18:1286
 - quaternary structure of, MS analysis of, 12:222
 - sizing by ESI and IMS, 18:1206
 - stability in solution and gas phase, nanoESI-FTICR MS of, 16:1957
- Protein database, amino acid sequences in, approach to correlate tandem mass spectral data of peptides with, 5:976
- Protein digests
- MS of, improved sample processing time and peptide recovery for, 15:784
 - nano electrospray combined with a quadrupole ion trap for, 7:150
 - nanoliter flow analysis of, 8:703
- Protein disulfide isomerase, redox sensitive thiols identified using ICAT MS, 18:260
- Protein-DNA binding affinity, evaluation by ESI-MS, 16:116
- Protein/DNA interactions, SAX for studying, H/D exchange MS, 19:887
- Protein fluorescence, measurements within electrospray droplets, 12:716
- Protein folding, hemoglobin assembly and, monitored by ESI-MS, 18:8
- Protein glycosylation, analysis with Q-trap mass spectrometer, 15:413

- Protein homodimers, asymmetric charge partitioning in, 15:1408
- Protein hydrogen exchange rates, changes in, that result from point mutations, mass spectrometric measurement of, 7:515
- Protein identification, origins of peptide mass fingerprinting, 14:931
- Protein interactions from the spectra of masses, algorithm for identification of, 18:563
- Protein ions
aerodynamic drag model for, 5:17
charge state distribution shifting of, observed in matrix-assisted laser desorption ionization mass spectrometry, 6:1183
collision cross sections for, 4:616
conformer selection of, by ion mobility in a triple quadrupole mass spectrometer, 5:127
declustering and fragmentation of, 8:1053
in electrospray ionization mass spectra, adducts (molecular mass=98 u) attached to, origin and removal of, 1:382
proton transfer reactivity, formed by electrospray ionization, 6:1086
- Protein kinase-2, MAPK-activated, limited proteolysis determined by LC/MS, 19:841
- Protein kinase G, ESI-TOF-MS of, 15:1392
- Protein kinases
bisubstrate analogs of, series of adenosylated peptides acting as, mass spectrometric characterization of, 5:443
novel assay system using mass-tag technology, tumors, 18:106
- Protein-ligand and protein-protein noncovalent interactions, probing, application of electrospray mass spectrometry in, 4:624
- Protein-ligand binding, hydrogen/deuterium exchange-MS study of, 16:869
- Protein-ligand complexes
detection by ESI-precursor ion scan, MS/MS, 20:227
noncovalent, confirming binding sites and tertiary structure of, MS for, 14:460
- Protein-ligand interactions
comparison of ESI, nanoESI and ESSI for dissociation constant determination, 19:332
modeling data in, 15:388
- Protein mass spectrometry, inherent electrochemistry of electrospray for, mass tag generation by, 15:1767
- Protein-metal binding, indirect detection of, ESI-MS in, 15:1658
- Protein mixtures, nanoflow LC-IMS-MS and LC-IMS-CID MS of, 15:1341
- Protein modification, by 4-hydroxy-2-nonenal and other short chain aldehydes analyzed by ESI-MS/MS, 14:215
- Protein N-glycosylation, by reversed-phase microbore liquid chromatography/electrospray mass spectrometry, complementary mobile phases, and sequential exoglycosidase digestion, 5:350
- Protein peptide, priori, tandem mass spectrometry of peptides with N-terminal glutamine studies on, 1:258
- Protein/peptide complexes, isotope-labeled cross-linkers and FTIR MS analysis of, 17:1100
- Protein phosphorylation, stoichiometry of, 15:647
- Protein-protein and protein-ligand noncovalent interactions, probing, application of electrospray mass spectrometry in, 4:624
- Protein-protein complexes, MS/MS of, 13:59
- Protein-protein interaction, allosteric inhibition of, MS of, 14:8
- Protein/protein interactions, bacteriotoxin E9:Im9 complex, analysis using lysine acetylation and nano-LC-MALDI-MS/MS, 17:983
- Protein quantification
in human proteome, lysine-specific mass tagging in, 14:1
lysine-specific mass tagging in, in human proteome, 14:1
- Protein sequence tags, generating using CID in quadrupole TOF MS, 15:1478
- Protein tryptic digests, LC/MS and LC/MS/MS determination of, 1:158
- Protein tyrosine phosphatase 1B, characterization of cysteine oxidation products in active site, 20:1540
- ProteinChip[®] array, SELDI-TOFMS to identify thymosin β -4, differentially secreted protein from lymphoblastoid cell lines, 14:760
- Proteins
acid hydrolysis, MALDI, 20:2106
acid-induced conformational changes and noncovalent complexes, CSI-MS, 20:845
analysis by gCF-MS, 19:1132
automated in-line gel filtration for native state MS, 19:239
binding affinities of mutant DNA complexes with, laser spray MS, 17:611
bound to iron oxide nanoparticles, analysis using MALDI-MS, 18:910
charge state assignment of native ES spectra of complexes, 20:435
charged complexes, theoretical investigations of dissociation in gas phase, 18:2242
collision-induced unfolding analyzed by ion mobility MS, 20:1851
collisional activation of complexes, review, 20:341
comparison of IMAC and TiO₂ surfaces in phosphorylation studies, 18:1932
comparison of Orbitrap MS with LTQ MS for identification of peptides and, 20:1405
conformation using T-WAVE ion mobility MS, 20:1699
conformations probed using ETD in top-down HDX MS, 20:1514
covalent labeling strategies, 20:1193
covalent labeling strategies (reprint, critical insight), 20:1193 in July 20:2009
Cu-binding, AA residues identified using MCO reactions and MS, 17:1552
deaminated, quantification of isoaspartyl residues in, by ECD, 18:48
desolvation in UV MALDI-TOF MS, 18:1880
differential expression analysis, using isotope labeling and PDQ-MS, 20:1287
disulfide-containing, MALDI MS, 20:2284
EDC monitoring deamidation during processing of peptides and, 19:855
effect of stabilization on CSD, ESI MS, 18:1605
efficiency of MS-based screening assay for detection of ligand binding, 19:1303
ESI-MS probing structure of, using site-specific 18-crown-6 attachment, 17:1209
evaluation of chip-based nanoelectrospray parameters for identification, 18:1714
expression by *E. coli* and *Shigella* species monitored by LC/QTOF-MS, 19:1626
folding, monitoring copopulated states during, ESI-IMS-MS, 18:2180
fragment intermediate lifetimes measured by DR ECD, 17:1605
gas-phase ion conformations from different solution conformations, 19:1907
gas phase noncovalent complexes retaining solution binding properties, tandem MS, 18:64
globular and denatured, deprotonation reactions at atmospheric pressure, ESSI-MS, 19:455
hydrogen/deuterium exchange of
bombarded by, implications for the conformation of gas-phase electrosprayed ions, 7:329

- continuous *versus* pulsed labeling/MS of, 10:675
- cysteine and cystine in, following organomercurial derivatization and analysis by matrix-assisted laser desorption ionization mass spectrometry, 5:359
- electroeluted from gels, electrospray MS from, 10:453
- electrospray mass spectra of, Are they related to their aqueous solution chemistry?, 3:216
- electrosprayed, gas-phase structure of, investigation of using ion-molecule reactions, 5:207
- frozen aqueous solutions of, infrared SALDI-MS of, 9:912
- gaseous ions of, hydroiodic acid attachment kinetics of, 10:552
- gel-separated, MALDI-TOFMS of, 10:512, 10:784
- heterogenous characterization of, by electrospray ionization-mass spectrometry, 7:1219
- infrared MALDI-MS of, 9:1, 9:879
- mass spectra of, effect of ion-molecule collisions in the vacuum chamber of an electrospray time-of-flight mass spectrometer on, 7:342
- microelectroelution concentrations of, 9:88
- multiply charged
 - in a high mass-to-charge ratio quadrupole mass spectrometer, proton transfer reaction studies of, 5:1064
 - produced by electrospray ionization, collisional activation and collision-activated dissociation of, 1:53
- N-peracetylation of, for their structure determination by mass spectrometry, utility of, 4:87
- probing by fast-atom bombardment mass spectrometry, 5:19
- rapid identification of, by LC/MS, 9:194
- reactions of, nanoelectrospray MS monitoring of, 10:112
- in solution, 10:671, 10:672
- TOF-MS of, 9:88, 9:879
- ultra-high sensitivity analysis of, micro-electrospray mass spectrometry, 5:605
- unfolding of, ESI-MS monitoring of, 9:1248
- up to 80 ku, molecular weight determination and variation characterization of, high accuracy, by ion spray mass spectrometry, 2:387
- hydroxyl radical footprinting, aliphatic peptidyl hydroperoxides as source of secondary oxidation, 20:1123
- identification by LC-ECD-MS/MS or LC-CID-MS/MS, 18:891
- identification in complex mixtures, DASER-MMF, 19:1915
- identification of, A₁ ion enhancement for peptide sequencing and, 20:1214
- identification of S-nitrosylation sites by QTOF-MS, 19:1353
- identification via SID in FT-ICR/MS, 17:700
- ILMs for peptides, carbohydrate and, analysis by MALDI MS, 20:1790
- intact
 - effect of Tween80 on signal intensity of, MALDI-TOF-MS, 18:102
 - ion dynamics, Orbitrap MS, 20:1486
- intact, MMA, ESI and Q-FT-ICR, 18:1
- interface footprinting quantified by hydroxyl radical oxidation and MDS, 19:1686
- ion charge states of peptides and, FTICR, 17:903
- ions, vibrational frequencies of linear peptides and, RRKM rate constant calculation using, 17:1749
- labeling of free and disulfide-bound thiol functions, 19:1
- local hydrogen exchange data from HDX CAD MS using deconvolution, 17:1543
- material-enhanced laser desorption/ionization profiling tool with specific carrier materials, MALDI-TOF MS, 17:1203
- membrane, analysis by LILBID MS, 18:1429
- modifying charge state distribution in ESI MS, 20:1617
- multiply-charged intact, fragmentation using MALDI-TOF-TOF MS, 19:231
- multiply-charged peptides and, generation by RADIO for MS, 20:597
- nanomolar samples, 3-layer sample preparation, MALDI MS analysis, 17:780
- non-covalent complexes, SID study of small molecules, peptides and, 19:190
- noncovalent complexes
 - allophycocyanine localization in MALDI-preparations by CLSM, 17:1599
 - analysis by ESI-MS, 20:593
- ocular integral membrane, tissue profiling by MALDI, 19:814
- optimization of AI ECD for, 20:763
- peptides and
 - counting sulfhydryl- and disulfide-bonds using mercurial ions, 19:1108
 - detection of multiply charged, by MALDSI coupled to FT-ICR MS, 17:1712
 - quantification using iTRAQ with ETD, 19:1255
- post-translational modification, MNM scening method for, 17:307
- production and clearance rates, quantified using SILT, 18:997
- quantification of isotopic pairs, FT-ICR MS, 19:973
- RNA crosslinks with, detection by nanoLC-ESI-MS/MS, 18:869
- sensitivity enhancement for MALDI IMS after treatment of tissue sections, 19:1069
- SILAC-labeled, quantitation and characterization, 18:2058
- solvent accessibility of surfaces, MALDI-TOF MS, 17:1490
- standards, Aurum Dataset, MALDI-TOF/TOF MS for, 18:850
- stoichiometry of phosphorylation determined by MS, 20:2211
- in tissues, improved MALDI-TOF imaging, 20:89
- ubiquitination analyzed by CID or ETD-based MS/MS fragmentation, 20:1652
- unfolding of, free energy determination, accuracy of MALDI-MS and SUPREX, 17:1535
- wheat gluten, characterization by HPLC/MALDI-TOF-MS, 19:1542
- Protein(s)
 - in brain structure, MALDI-MS of, 16:1093
 - charge reduction ESI-MS of, 16:1876
 - charge state distribution shifts of, in nanoESI triple quadrupole MS, 16:409
 - crystal structure of, and charge states of gaseous proteins in ESI, 16:1409
 - intact, MALDI-TOF-TOF of, 16:482
 - microsecond photochemical oxidation of, laser flash photolysis of hydrogen peroxide in, 16:2057
 - microwave-assisted acid hydrolysis of, LC MALDI MS/MS with, 16:471
 - modified, automated top-down LC-MS of, 16:2027
 - multiple forms of, kinetic characterization by top-down MS, 16:1052
 - post-translational analysis and top-down MS/MS of, qQq-FTICR mass spectrometer for, 16:1985
 - salt-containing, MALDI-TOF-MS of, porous anodic alumina membrane as sample support for, 16:1488
 - surface of, nanospray-related reactions at, 16:1325
- Protein(s). *See also* Peptide(s)
- 1-anilino-8-naphthalene-sulfonate binding of, ESI-MS of, 12:428
- analyzing sequence of, calyculin nonspecific cleavage for, 12:1180
- carbon nanotubes as affinity tubes for, in MALDI MS, 15:1629
- CD4/lc-copurifying, PAGE and MS of, 15:558
- chemical cross-linking of, 15:1604
- complex digest mixtures of, variable LC-MS/MS of, 8:1059
- containing nitrotyrosine, MALDI-MS of, 12:438

- digest analysis of, capillary separations NMS/MS of, 8:1237
 electrochemical oxidation and cleavage of, 15:1707
 enhanced sample loading of, 8:15
 ESI-MS of
 alkali-metal adduct ion formation in, 11:967
 ion-impact energy measurement in, 12:826
 mass error in, water cluster calibration and, 8:1158
 H/D exchange mass spectra of, automatic analysis of, 12:1153
 heterodimeric, pH titration of, by ESI-MS, 11:54
 identification of, from phenylisothiocyanate-derivatized electrosprayed tryptic peptides, 12:288
 by in-gel digestion, HPLC, and MS, peptide analysis by complementary ionization techniques, 12:215
 intact, cumulative distribution functions in MS of, 13:40
 interpreting mass spectra of, 13:107
 ions of, charge states, solvent effects on, 11:976
 from liquid based UV-MALDI matrices, detection limits of, 15:486
 modification of, by mPC-CE-MS/MS, 8
 MS of, 8:296, 12:125
 noncovalent, CAD MS/MS of, 12:329
 phosphorylation of, precursor discovery for, 13:792
 proteolysis and NanoESI-MS/MS of, 8:483
 quantitative proteomic analysis by MS in, 12:1238
 quaternary structure of, enzyme inhibitors and, 12:97
 rapid switchability of MALDI and ESI quadrupole TOFMS for, 11:493
 single d-amino acid distinguished in, by ECD, 15:1087
 structure information of, from MS, selective titration of arginine residues by sulfonates in, 12:810
 submicroliter samples of, nanoelectrospray with controllable flow rate in mass analysis of, 11:94
 in vivo labeling for, 13:804
 Proteins immobilized on nylon-based membranes, direct matrix-assisted laser desorption ionization mass spectrometric analysis of, 5:230
 Proteolysis
 and NanoESI-MS, of proteins, 8:483
 on-line, peptide sequencing using alternative enzymes for, triaxial electrospray probe, 20:1983
 time-limited, MAPK-activated PK-2, determined by LC/MS, 19:841
 triaxial probe for, coupled with HDX-MS, 18:208
 Proteolytic digests, liquid chromatography with MALDI or nano-ESI-MS for, 12:1205
 Proteolytic peptides, sequencing of, by C-terminal derivatization and nano-ES-CID-MS, 11:673
 Proteome
 human, peptide sequencing and protein quantification in, 14:1
 shotgun analysis by HPLC-MALDI FT-ICR MS, 17:205
 Proteome analysis, using selective incorporation of isotopically labeled amino acids, 11:78
 Proteome coverage, in peptide abundance measurements, using DREAMS, 13:954
 Proteomes
 human plasma, mapping using SCX-LC-IMS-MS, 18:1249
 low-mass, top-down MS/MS, 20:2183
 QCAL, standard for assessing instrument conditions for analysis of, 19:1275
 relative quantification of, calculating $^{16}\text{O}/^{18}\text{O}$ peptide ion ratio for, 15:437
 Proteome(s)
 complex, accurate mass and LC elution time utility for, 16:1239
 low molecular weight, quantitative analysis in serum, 16:1221
 Proteomics
 analysis of trypanosome flagellar proteome using ETD/ETD/CAD, 20:167
 analyzing stable isotope labeling data in, 16:1181
 bovine myelin sheath, characterization of P0 by MALDI-TOF/TOF MS, 17:117
 evaluation of chip-based nanoelectrospray parameters for protein identification, 18:1714
 in heart tissue investigation, 16:1207
 high-throughput comparative, MS/MS using nanoLC-FTICR-MS, 18:1332
 MMA of intact proteins and product ions, ESI and Q-FT-ICR, 18:1
 MS/MS in, MS2Grouper in, 16:1250
 Orbitrap mass precision and accuracy for proteome measurements, 20:1477
 phosphopeptide capture by zirconium phosphonate-modified magnetic nanoparticles, 19:1176
 photosystem II structure and dynamics by high resolution FTICR MS, 19:1004
 probability-based shotgun cross-linking site analysis, 20:1896
 profiling *S. miliaryus barbouri* venom using MALDI-TOF MS, 18:600
 qQq Fourier transform ion cyclotron resonance (qQqFTICR) mass spectrometer, for post-translational analysis and top-down MS/MS of whole proteins, 16:1985
 quantitative, reductive alkylation by acetone, 20:1366
 shotgun, comparison of Orbitrap with LTQ MS, 20:1405
 top-down, reveals proteins expressed in *M. acetivorans*, 20:1743
 top-down shotgun, precursor ion-independent algorithm for, 20:2154
 tryptic peptides dications, bifurcating fragmentation behavior, CAD, 19:1755
 Proteomics. *See also* Industrial proteomics
 in analysis of chromatin-associated factors, 14:687
 in analysis of *Pseudomonas aeruginosa* grown under magnesium limitation, 14:742
 analysis using accurate mass and elution time peptide tags with capillary LC TOFMS, 14:980
 comparative, trypsin-catalyzed ^{16}O -to- ^{18}O exchange for, 14:704
 complementary nature of LC-MALDI-MS/MS and LC-ESI-IMS/MS in, 14:971
 complex component analysis in, 14:728
 de novo peptide sequencing in, MALDI-quadrupole ion trap MS for, 14:1012
 high performance LC-FTICR mass spectrometer for, 15:212
 investigative, of unknown plant virus from infected plants, 14:736
 ion trap data-dependent MS/MS in, 15:150
 ligand-induced site-specific phosphorylation of EGF receptor in, 14:1022
 negative signature mass algorithm in, 14:1032
 on-line strong cation exchange μ -HPLC-ESI-MS/MS, protein identification and process optimization in, 14:719
 peptide mass fingerprinting in
 mass assignment from, 14:943
 origins of, 14:931
 peptide mass mapping and protein analysis in, matrix mixture for MALDI-MS in, 14:992
 protein complex analysis in, multidimensional nano-HPLC for, 14:1003
 research in, MS in, 14:685
 two-dimensional gel proteins in, identification by MALDI-TOF or μ -LC-ESI MS/MS, 14:956
 Proton affinities
 of aza-18-crown-6-ether, 13:493
 of benzamides and proton-bound complexes, 5:826
 of bromo- and iodoacetonitrile, using kinetic method with full entroply analysis, 11:601

- computed, reevaluation of, α -amino acids, 20:2116
 ESI-MS determination of, 10:483
 mononucleotide gas phase, as determined by the kinetic method, 11:24
 of the N- and C-terminal segments arising upon the dissociation of the amide bond in protonated peptides, 10:1
 of nucleic acid molecules, kinetic method determination of, 15:1080
 of polyglycines assessed by using, 3:863
 of polyglycines assessed by using the kinetic method, 3:863
 of proline analogs using kinetic method with full entropy analysis, 13:72
 of saturated aliphatic methyl esters, 11:789
 of SO₃, 10:856
 of uracil, computational study of protonation sites, 11:1065
 Proton affinity, of β -oxalylaminoalanine, 16:1151
 Proton affinity scale, revising of, α -amino acids, 17:1275
 Proton affinity values, of MALDI matrices under ESI conditions with kinetic method, 15:431
 Proton-bound complexes and proton affinity of benzamides, dissociation of, 5:826
 Proton-bound dimers, reactions of, 5:695
 Proton-bound homodimers and heterodimers, of amides and amines in gas phase, equilibrium studies by FTICR-MS, 13:1273
 Proton complexes, of di-*o*-benzoyl-tartaric acid dibutyl ester and L-tryptophan, in gas phase, chiral recognition characteristics of, 14:571
 Proton transfer
 in APCI-MS, in analyzing polycyclic aromatic compounds, 15:301
 competing, to the oxygen atom and the aromatic ring, propene loss initiated by, chemical ionization of phenyl n-propyl ether and methyl substituted analogs, 7:639
 intracluster, to biomolecular analytes, 9:1060
 light-induced, from gas phase cations to reference bases, 12:359
 matrix to analyte, in clusters, 12:726
 observed in mass spectra, 13:1209
 in radical-cation clusters of nucleosides, 12:176
 reactivity of peptide and protein ions formed by electrospray ionization, 6:1086
 ternary copper (II) complexes with histidine, macrocyclic ligand effects on electron transfers and, 20:972
 Proton-transfer ion trap mass spectrometry, of volatile organic compounds in air, 16:1316
 Proton transfer reaction time of flight mass spectrometry, two-stage, isomeric species differentiation by, 19:325
 Proton transfer reactions
 of derivatized fullerene trications, 9:114
 gas phase, ion-ion reactions in, proton transfer reactions of protonated pyridine with multiply charged oligonucleotide anions, 6:529
 gas-phase, involving multiply charged cytochrome c ions and water under thermal conditions, 3:624
 ion/ion, in linear ion traps, 18:882
 of matrix-assisted laser desorption/ionization matrix monomers and dimers, 10:1111
 onium reactions and, characterization in closed *versus* open shell ions, 18:270
 studies of multiply charged proteins in a high mass-to-charge ratio quadrupole mass spectrometer, 5:1064
 of ubiquitin ions, 8:954
 Proton transport catalysis, protonated 2-methoxy and 2-hydroxychalcone, Nazarov cyclization, 20:805
 Protonated acetaldehyde, elimination of methane from, ab initio transition state, 5:1102
 Protonated acetaldehyde and methanol, gas-phase reaction between, 1:481
 Protonated alkyl dipeptides, in gas phase, bimolecular hydrogen-deuterium exchange behavior of, 6:466
 Protonated amides, fragmentation of, 9:454
 Protonated amine/polyether complexes, in quadrupole ion trap, threshold dissociation energies of, 14:383
 Protonated amines, deamination of, 9:1146
 Protonated and methylated lactones, gas-phase, dissociation behavior of, characterization of the, 3:47
 Protonated angiotensin II, doubly, formed by electrospray ionization, charge-separation reactions in tandem mass spectrometry of, experimental considerations and structural implications, 7:30
 Protonated aromatic amines with ND₃, 3, H/D exchange reactions, relative collisions in quadrupole cells and, 6:19
 Protonated aromatic carbonyl compounds, remote fragmentations of, via internal reactions in intermediary ion-neutral complexes, 3:417
 Protonated clusters
 structure of water-alcohol and ammonia-alcohol, 4:507
 water-alcohol and ammonia-alcohol, on the structure of, 4:507
 Protonated cysteine, fragmentation of, sulfhydryl group in, 9:1275
 Protonated dipeptides, condensation reactions in, 15:1025
 Protonated ethanol, and its neutral counterparts, 2:459
 Protonated formic acid, decomposition of, 15:982
 Protonated H-Gly-Gly-Sar-OH and H-Gly-Sar-Sar-OH, fragmentation pathways of, 14:1454
 Protonated ions, of cysteine-containing peptides, fragmentation of, 15:697
 Protonated methane, on the structure of, 2:453
 Protonated nitriles, gas-phase dissociations of, ion-neutral complexes in, n-C₃H₇CNH⁺ and i-C₃H₇CNH⁺, 2:178
 Protonated O
 O-diethyl O-aryl phosphorothionates, fragmentation of, in tandem mass spectral analysis, 7:189
 O-dimethyl O-aryl phosphorothionates in tandem mass spectral analysis, fragmentation of, 6:488
 Protonated ortho silicates, APCI low energy collision-induced dissociation fragmentation of, McLafferty or ion-neutral complex rearrangements, 5:544
 Protonated peptide ions, multiply, ubiquitin, insulin B, and renin substrate, apparent gas-phase acidities of, 7:1211
 Protonated peptides
 complexation of with saccharides in gas phase, decreased rates of hydrogen/deuterium exchange reactions, 6:1247
 cyclic, MS_n characterization of, 15:1039
 decomposition of, collision-activated, influence of cysteine to cysteic acid oxidation on, evidence for intraionic interactions, 3:337
 high-energy fragmentation of, determination of, using a BEqQ hybrid mass spectrometer, 4:470
 multiply, surface-induced dissociation of, 3:859
 Protonated peptides and methanol, interaction of, molecular probe for, gas-phase hydrogen/deuterium exchange as, 5:623
 Protonated species, protonation of benzonitrile hydrogen interchange in, site of, 1:225
 Protonated trimers, charge-solvated vs salt bridge structures of, ion mobility spectroscopy of, 16:1009
 Protonated ubiquitin, copper adducts of, electrospray ionization mass spectrometry of, 6:521
 Protonation
 in ESI-MS, 11:961
 in the low energy decompositions of gas-phase peptide ions, 7:522

- of peptides, novel ions for, MALDI-TOFMS of, 11:345
- of serine and its derivatives, fragmentation reactions of, in gas phase, 11:1047
- sites for, in methyl nitrate and the formation of transient CH₄NO₃ radicals, neutralization-reionization mass spectrometric and computational study of, 11:380
- of uracil, 11:1065
- Protonation and fragmentation of highly functionalized molecules in fast atom bombardment, mechanistic consideration of, high resolution mass spectrometry and tandem mass spectrometry analysis of the ions formed by fast atom bombardment of digoxin and related cardiac glycosides, 1:455
- Protonation and gas-phase cationization, of neutrals generated by matrix-assisted laser desorption, 4:393
- Protonation of benzonitrile hydrogen interchange, in the protonated species, 1:225
- Protonation of ferrocene, in the gas phase, 5:274
- Protonation reactions, deprotonation and, gas phase reactions, fluorinated 1,2,4-triazines, ESI MS, 19:686
- Protonation site on bond strengths, in simple peptides, application of Ab initio and modified neglect of differential overlap bond orders and modified neglect of differential overlap energy partitioning, 5:704
- Protons, half or doubly charged, metastable ions as, 15:1261
- Protoporphyrin IX ions, induction by ECD, 19:809
- PSD. *See* Post decay source analysis; Post-source decay entries
- Pseudo-MS3, in a MALDI orthogonal quadrupole-TOF-MS, 13:1034
- Pseudomonas aeruginosa*
- 4-hydroxy-2-alkylquinolines from, ESILC-MS of, 15:862
- grown under magnesium limitation, proteomic analysis of, 14:742
- PTFT. *See* Peak-target Fourier transform
- Pulse excitation, direct and resonant current, and multiparticle simulation of ion motion in ion trap mass spectrometer, 6:57
- Pulse glow discharge plasmas, afterpeak time regime of, ion formation processes in, 4:727
- Pulsed bombardment ionization method, model, study of time-dependent desorption using, improved detection limits for fast atom bombardment mass spectrometry, 3:637
- Pulsed extraction, mass-correlated, in MALDI-TOFMS, 11:841
- Pulsed fluorescence measurements, of trapped molecular ions with zero background detection, 13:696
- Pulsed gas introduction into quadrupole ion traps, 1:308
- Pulsed glow discharge, as an elemental ion source, 1:138
- Pulsed ion deflection system, for background reduction in ²⁵²Cf-plasma desorption mass spectrometry, 3:706
- Pulsed-leak valve, for use with ion trapping mass spectrometers, 7:118
- Pulsed Q dissociation mass spectrometry, differential expression analysis of proteins using isotope labeling and PQD-MS, 20:1287
- Pulsed-sample-inlet system, quadrupole ion trap equipped with, carbon addition reactions in, information on the location of carbon-carbon double bonds in H₆-C₂₃ linear alkenes from, 2:305
- Pump, using a fast direct current pulse as, determination of ion frequencies in a quadrupole ion trap by, 5:29
- Pumps, high dead volume, delay time reduction by short-term low-ratio split flow, LC/MS, 18:245
- Purge-and-trap gas chromatography/mass spectrometry, measuring volatile organic compounds in human blood by, importance of enhanced mass resolution in, 3:831
- PyC₂H₅⁺ isomers, multiple stage pentaquadrupole mass spectrometry for generation and characterization of gas-phase ionic species, 7:1126
- Pyrenemethylguanidine, polysulfated oligosaccharides, MALDI-MS using, 20:131
- Pyridine
- gas phase interaction with ionized nitrile sulfides, 14:241
- gas phase reactions of, 8:1124
- sulfur trifluoride cation affinities of, 8:68
- Pyridine-7,12-dimethylbenz[a]anthracene-pyridine adducts, isomer differentiation in, by fast atom bombardment tandem mass spectrometry, 2:256
- Pyridine alkaloids, tobacco, determination by PFTBA ions/analyte molecule reaction IT-MS, 18:1774
- Pyridine-containing Schiff base derivatives, for the structural determination of long-chain aldehydes by gas chromatography combined with mass spectrometry, 2:245
- Pyridine nucleophiles, CH₂⁺ transfer to, a means of producing alpha-distonic ions, 4:117
- Pyridines, SiCl₃⁺ and SiCl⁺ affinities for, determined by using the kinetic method with multiple stage mass spectrometry, agostic effects in the gas-phase, 7:198
- Pyridinium ions, homolytic cleavages in, 14:790
- Pyrimidin-4(3H)-ones, 5:2-substituted, electron impact ionization mass spectrometry and intramolecular cyclization in, 5:113
- Pyrimidines, antiviral agents, CID mechanisms of ion formation, ESI-MS/MS, 18:1477
- 2-Pyrimidinyl-*N*-arylbenzyl amines, chemical transformations predicted by tandem mass spectrometry, 17:253
- Pyroglutamic acid, *Conus virgo* conotoxin, identification by MALDI-MS/MS, 18:1396
- Pyrolysis mass spectrometry, cell culture media investigation by, 10:502
- Pyrophosphates, nucleoside, and polyisoprenyl phosphates, glycosyl esters of, anomeric configuration, by fast-atom bombardment tandem mass spectrometry, 7:541
- Pyrrole, transition-metal mediated heteroatom removal by reactions of FeL⁺ [L⁺O, C₄H₆, c-C₅H₆, H₆H₆, C₅H₄(=CH₂)] with, in the gas phase, 7:938
- Pyrrolidinium radicals, hypervalent, by NRMS, 9:242
- Pyrrolo[1,2-a][1,4]benzodiazepin-4-one,-6-one, and -4,6-dione, mass spectrometry and molecular orbital calculations, characterization and differentiation of heterocyclic isomers, 7:653
- Pyrrolo[2,1-b][1,3]oxazin-6-one derivatives, isomers of, stereospecific fragmentation of, 10:393
- Q
- Q-TOFMS. *See* Quadrupole time-of-flight mass spectrometry
- Q-trap mass spectrometer, in glycoprotein analysis, 15:413
- QCAL, standard for assessing instrument conditions for analysis of proteomes, 19:1275
- QIT. *See* Quadrupole ion-trap mass spectrometry
- QqLIT instrument, AP and vacuum MALDI on, 16:176
- Qscore, an algorithm for evaluating SEQUEST database search results, 13:378
- Quadrupolar excitation, in Fourier transform ion cyclotron resonance mass spectrometry, simplified application of, 6:448
- Quadrupole. *See also* Tandem quadrupole-MS
- radiofrequency, collisional focusing effects in, 3:398
- transmission of, ion source emittance and, 9:1074
- Quadrupole-acceleration-deceleration mass spectrometer, tandem, for neutralization-reionization studies, 3:493
- Quadrupole-based mass spectrometric, evaluation of isotope ratios of carbon dioxide, expired air from mice and men following the administration of 13-C-methyl methionine, 3:742
- Quadrupole cells, reactive collisions in
- H/D exchange reactions of emolate ions with CH₃OD and C₂H₅OD, 3:853

- H/D exchange reactions of protonated aromatic amines with ND_3 , 6:19
- Quadrupole collision cell
 high pressure, for low-energy collision-induced dissociation, 5:1042
 high-pressure radio frequency-only, 9:775
- Quadrupole excitation
 frequencies in a linear ion trap, 11:1016
 in linear quadrupole traps, 19:510
 resonance shifts in, 13:577
- Quadrupole excitation frequencies, higher order, in linear ion trap, 11:1016
- Quadrupole field, ions confined in, excitation frequencies of, 11:10
- Quadrupole/Fourier transform ion cyclotron resonance mass spectrometer, construction of, for MS/MS, 15:1099
- Quadrupole Fourier transform ion cyclotron resonance mass spectrometry, ESI and, MMA, intact proteins and product ions, 18:1
- Quadrupole Fourier transform mass spectrometry, hybrid, solvated ion accumulation in accumulation/collision hexapole, 20:411
- Quadrupole instrument, in CAD, of glycosphingolipids as lithiated adducts, 12:61
- Quadrupole ion guide, improving ion beam quality exiting, 17:335
- Quadrupole ion trap. *See* Ion trap, quadrupole
 anomericity of glycosidic bond by, 10:269
 applications of external customized waveforms to, 10:355
 CAD in, cyclic protonated peptide and metal complex characterization by, 15:1039
 competitive reactions of dimethyl ether ions in, steric and substituent effects in phenols evaluated by, 6:627
 in detection of pharmaceutical compounds in tissue, 10:1315
 determination of binding selectivities by, 9:845, 9:1049
 determination of cooling rates, 17:932
 dynamic collision-induced dissociation in, 18:749
 an empirical approach to estimation of critical energies by using, 7:1116
 equilibrium space charge in, 5:64
 equipped with a pulsed sample-inlet system, carbon addition reactions in, information on the location of carbon-carbon $\text{H}_6\text{-C}_{23}$ linear alkenes from, 2:305
 ESI identification of tetracycline antibiotics in, 9:1089
 field-modulated selective ion storage in, 6:928
 H/D exchange of gas phase bradykinin ions in, 14:85
 heavy gases in, using static pressures of, 14:1099
 improving IRMPD efficiency in, 20:1127
 infrared multiple photon dissociation in, via a multipass optical arrangement, 5:886
 initial fragmentation of oligodeoxynucleotides in, 16:1853
 ion frequencies in, by using a fast direct current pulse as pump and a laser probe, 5:29
 ion isolation in, evidence of isomerization during, 3:680
 ion storage in, nonlinear resonance effects during, 4:917
 linear, axial ejection from, 14:1130
 linear, ion excitation in, 16:835
 measurement of CID of tantalum oxide ions in, 11:1072
 methylene substitution reactions in, selectivity of ethylene, ethylene oxide, and dimethyl ether reactive ions, 3:39
 5'- and 3'-mononucleotides, gas-phase hydrogen/deuterium exchange, 18:724
 oligosaccharide characterization with boronic acid derivatization using IRMPD and, 18:2094
 nano electrospray combined with, for the analysis of peptides and protein digests, 7:150
 peptide ions in, conformational differences of, 12:1331
 peptide ions in, tandem mass spectra of, effects of heavy gases on, 7:1194
 peptides in, MS/MS spectra of, 9:341
 of phosphatidylserine, ESI with, 16:1510
 prediction of collective characteristics for ion ensembles, 17:889
 product ions in the MS/MS spectra of peptides in a, 9:341
 protonated amine/polyether complexes in, threshold dissociation energies of, 14:383
 pulsed gas introduction into, 1:308
 resonance excitation in, competition between resonance ejection and ion dissociation during, 5:1031
 rhodamine 575 fluorescence and photodissociation of cations, 20:96
 selective ion isolation/rejection over a broad mass range in, 2:11
 three-dimensional, ECD of polypeptides in, 16:22
- Quadrupole ion trap mass spectrometer
 coupling a particle beam interface directly to, 4:38
 high resolution on, 2:198
 internal pulsed valve sample introduction on, 6:976
 isomerization of linear C_3H_3^+ in its reaction with acetylene, and collisional stabilization of the $[\text{C}_5\text{H}_5^+]$ * collision complex in, 3:631
 positions, velocities, and kinetic energies of resonantly elicited ions in, by laser photodissociation, 4:792
 surface-induced dissociation of molecular ions in, 2:487
- Quadrupole ion trap mass spectrometry. *See also* Ion trap mass spectrometry
 of alkali metal binding selectivities of caged aza-crown ether ligands, 11:884
 dissociation of doubly charged transition metal/polyether/pyridyl ligand complexes in, 10:126
 distinction among isomeric unsaturated fatty acids as lithiated adducts by ESI-MS using low energy CAD on, 10:600
 dynamic CID in, with two-frequency excitation waveform, 18:2017
 ESI with, sulfatide studies using, 15:536
 HASTE-CID, low mass product ion detection, 17:81
 multiple stage, *E. coli* cardiolipin characterization by ESI with, 17:420
 negative chemical ionization in, effects of applied voltages and reaction times, 5:757
 probing trapped ion energies via ion-molecule reaction kinetics, 3:716
 resonance ejection mode of, mass resolution in, extended theoretical considerations for, 5:676
 tandem, of cardiolipin, 16:491
 time scale of internal energy relaxation, by AP-MALDI and nanoESI in, 20:1801
 transition metal/polyether/pyridyl ligand complexes in, 10:126
- Quadrupole ion trap-MS
 fast excitation CID in, 14:785
 ion motion in, analytical description of, 14:818
- Quadrupole ion trap/trapping
 calculating stability diagrams in, matrix methods for, 13:597
 IRMPD in, erythromycin analog characterization by, 13:630
 in multiple-stage MS analysis of complex oligosaccharide antibiotics, 13:1313
 plasticizer contamination from vacuum system O-rings in, 13:901
 poly(ethylene glycol)-cation interactions in, 13:177
 for proton affinity of proline analogs, 13:72
 two-dimensional, 13:659
- Quadrupole mass analyzer, 10:1053
 analytical approach to, 9:50

- multiple pass, mass resolution of 11,000–22,000 with, 11:407
 single series peptide fragment ion spectra generated by CID in, 9:606
 transmission and resolution of, 9:70
- Quadrupole mass filter
 inductively coupled plasma MS with, 8:1230
 modulating voltages applied to, 16:379, 16:800 (erratum)
 peak splitting with, 10:1263
- Quadrupole mass filters, microengineered, ion coupling strategies for, 20:146
- Quadrupole mass spectrometer
 benzenedicarboxylic ester molecule ion collision with oxygen in quadrupole mass spectrometer, 10:1248
 double, effects of target gas in collision-induced dissociation using, 5:845
 high-mass-to-charge ratio, proton transfer reaction studies of multiply charged proteins in, 5:1064
 triple
 collisional activation in, alkenamide fragmentation by, 15:1462
 ion mobility in, conformer selection of protein ions by, 5:127
- Quadrupole mass spectrometers, development using rapid prototyping technology, 20:1359
- Quadrupole mass spectrometry
 collision dynamics in, 9:101, 9:606
 ion energy in, 15:48
 pentaquadrupole, borinium cations by, 12:948
 triple
 as applied to flame diagnostics, study of the $C_2H_4/N_2O/Ar$ flame, 4:152
 biopolymer sequencing in, 12:846
 water clusters in, 9:814
- Quadrupole MS, dual, peptide sequencing using, 14:1387
- Quadrupole reaction cell, inductively coupled plasma mass spectrometer with axial field in, 13:1176
- Quadrupole tandem mass spectrometry, triple, phosphatidylglycerol studies with, 12:1036
- Quadrupole tandem mass spectrometry (MS/MS), triple, in molecular characterization of thiols, 15:288
- Quadrupole time-of-flight mass spectrometer
 CID in, generating protein sequence tags by, 15:1478
 fitted with MALDI ion source, glycosphingolipid analysis with, 12:1220
 MALDI orthogonal, pseudo-MS3 in, 13:1034
 precursor discovery for protein phosphorylation on, 13:792
- Quadrupole time-of-flight mass spectrometry
 identification of S-nitrosylation sites in proteins, 19:1353
 for structural elucidation of drug metabolites, 10:1305
 Trp-cage conformation studied by, 18:195
- Quadrupole time-of-flight mass spectrometry (Q-TOFMS)
 in determining CAD of substituted indole, 12:1145
 in fucosylation study of low abundance species in complex mixtures, 12:964
 in identifying G1cNAcylation sites of peptides and α -crystallin, 12:1106
- Quadrupole time-of-flight (Q-TOF) instrument, stable gradient nanoflow LC-MS on, 16:1545
- Quadrupole time-of-flight (Q-TOF) mass spectrometry, target compound verification by, mass measurement for, 16:406
- Quadrupole time-of-flight tandem mass spectrometry, *B. stolpii* UKi2, identification of SPNL molecular species, 18:394
- Quadrupole TOFMS
 of diazinon photocatalytic degradation products, 14:803
 high resolution, of glycopeptides, 14:777
 noise filtering techniques for, 14:766
- Quadrupoles
 linear, with added octopole fields
 quadrupole ion excitation, 19:510
 selective ejection of axial ions from, 19:270
 RF-only, collision-activated dissociation in, instrument-independent measurements of, a kinetics-based protocol for, precision and accuracy in tandem mass spectrometry measurements, 1:272
- Qualitative assessment, monomer ratios, in putative ionic terpolymer samples by ESI-MS with CID, 13:921
- Qualitative comparison between the quantum calculations and ESI-MS of complexes of polyammonium macrotricyclic ligands with dicarboxylic acids, 12:304
- Qualitative selected ion flow tube MS, influence of ionic diffusion and mass discrimination in, 12:863
- Quality control
 QCAL standard for assessing instrument conditions for analysis of proteomes, 19:1275
 risk of false positive identification using multiple ion monitoring in qualitative MS, 17:823
- Quantification algorithms, classical, acquisition and quantification of ion images with, 1:37
- Quantification of 3-nitrotyrosine in biological tissues and fluids, generating valid results by eliminating artifactual formation, 11:578
- Quantification of a potent 5-HT_{2a} antagonist and an active metabolite in rat plasma and brain microdialysate by liquid chromatography-tandem mass spectrometry, 8:371
- Quantitation
 of ion abundances in FTICR-MS, 9:1204
 of major elements with secondary ion mass spectrometry by using M₂⁺ molecular ions, 9:638
- Quantitation of SR 27417 in human plasma using electrospray liquid chromatography-tandem mass spectrometry, study of ion suppression, 7:1099
- Quantitative analysis by high-performance liquid chromatography atmospheric pressure chemical ionization mass spectrometry, the determination of the renin inhibitor CP-80,794 in human serum, 2:164
- Quantitative analysis of crude plasma extracts by ion trap tandem mass spectrometry, 3:372
- Quantitative analysis of technical polymer mixtures by MALDI-TOF-MS, 13:914
- Quantitative assessment, of cysteine and cystine in peptides and proteins, following organomercurial derivatization and analysis by matrix-assisted laser desorption ionization mass spectrometry, 5:359
- Quantitative determination of a nonpeptide antithrombotic in dog plasma by microbore high-performance liquid chromatography-tandem mass spectrometry utilizing pneumatically assisted electrospray ionization, 7:1238
- Quantitative determination of peptide retention of polymeric substrates using MALDI-MS, 11:62
- Quantitative evaluation of noncovalent chorismate mutase inhibitor binding by ESI-MS, 14:1470
- Quantitative mass chromatography mass spectrometry, novel perfluoroacyl derivatives of corticosteroids and related substances for potential use in, 6:264
- Quantitative plasma profiling, of arecoline in treated Alzheimer patients, an automatic reaction control chemical ionization technique in ion trap detector for, 2:168
- Quantitative proteomic analysis
 of chromatin-associated factors, 14:696
 by MS, 12:1238
- Quantitative secondary ion MS, depth profiling of surface layers of cubic silver halide microcrystals, 10:1016
- Quantitative TOF-SIMS analysis, of oligomeric degradation

- products at the surface of biodegradable polymers, 13:1108
- Quaternary ensembles of, by MALDI, 8:1046
- Quaternary phosphonium salts, laser microprobe mass spectrometry of, direct *versus* matrix-assisted laser desorption, 4:798
- Quaternary protein ensembles, MALDI analysis of, 8:1046
- Quaternary structure, of protein complexes, MS analysis of, 12:222
- Quaternary structure of protein complexes, new crosslinker for MS analysis of, 12:222
- R
- RABA. *See* Reductive alkylation by acetone
- Radiative stabilization, of trimethylsilyl adduct ions, 1:144
- Radical cation
- to the benzene structure, 1,3-hexadien-5-yne, and to nonclassical ion structures, *Ab initio* calculations of, 7:731
 - methyl propionate, 7:482
- Radical-cation clusters of nucleosides, electron transfer vs proton transfer in, 12:176
- Radical cations, conventional and distonic, distinguishing, by using dimethyl diselenide, 7:1245
- Radical ejection in a cubic trap, effect of ion initial axial position on, simulated ion trajectory and induced signal in ion cyclotron resonance ion traps, 5:807
- Radical ion species, acute mass capability, MALDI-TOF-MS, 17:672
- Radical ions
- long-lived cations, gas phase fragmentation of, cysteine, 20:985
 - peptide fragmentation of photolytically generated, 20:1378
 - tracking migration in large hydrogen deficient peptides, 20:1148
 - triglycine cations, tautomerization and dissociation of, 20:996
- Radical trap moieties, effect on ECD spectra of substance P, 17:1428
- Radicals, spin-trapped, determination, by LC-TSP-MS and LC-ESI-MS, 2:413
- RADIO. *See* Radiofrequency acoustic desorption and ionization
- Radio frequency glow discharge-time of flight mass spectrometry, millisecond pulsed, temporal and spatial variations in molecular energetics, 15:87
- Radio frequency ion traps and guides, collisional activation in, 15:1616
- Radio frequency multipoles, charge state selection in, 14:1229
- Radiochemical detection for packed capillary LC-MS, 13:82
- Radiofrequency acoustic desorption and ionization, multiply-charged proteins and peptides for MS generated by, 20:597
- Radiofrequency capacitive coupling, with efficient gated trapping in internal matrix-assisted laser desorption ionization-Fourier transform ion cyclotron resonance, 8:195
- Radiofrequency glow discharge, effects of target gas in collision-induced dissociation using, 5:845
- Radiofrequency glow discharge mass spectrometry, cooled polymer sample holder from, 8:1214
- Radiofrequency-only mode, ion cloud manipulation using, improvement for high mass detection in Fourier transform mass spectrometry, 4:186
- Radiofrequency-only-mode event, high pressure trapping in Fourier transform mass spectrometry, 3:590
- Radiofrequency-only quadrupole collision cell, for measuring ion collision cross section, 8:697
- Radiofrequency quadrupoles, collisional focusing effects in, 3:398
- Ramsperger statistical theory, 8:103
- Rana* spp., ranid skin secreted peptides, oxidation *versus* carboxamidomethylation of S-S bonds in, MALDI-MS, 19:479
- Ranking of gas phase acidities and chloride affinities of monosaccharides and linkage specificity in CID of negative ESI-generated chloride adducts of oligosaccharides, 12:1193
- Rapamycin, new metabolite of, identification by HPLC and electrospray MS/MS, 11:516
- Rapid analysis, of antibiotic-containing mixtures from fermentation broths, using liquid chromatography-electrospray ionization-mass spectrometry matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry, 7:1227
- Rapid and sensitive differentiation of anomers, linkage, and position isomers of disaccharides using FAIMS, 14:265
- Rapid and sensitive identification, of epitope-containing peptides by direct MALDI MS/MS of peptides affinity-bound to antibody beads, 14:1076
- Rapid confirmation/quantitation of cocaine and benzoylecgonine in urine utilizing HPLC and MS/MS, 11:257
- Rapid protein identification, using a microscale electrospray LC/MS system, an ion trap mass spectrometer, 9:194
- Rapid prototyping technology, quadrupole mass spectrometer development using, 20:1359
- Rapidly switchable MALDI and ESI quadrupole TOFMS for protein identification, 11:493
- Rare earth cations, reactivity of, with phenol, 9:1035
- Rare earth elements, trace amounts of, determination using MA-PTFE fibers as sorbent, ICP-MS, 17:1258
- Rat hepatocytes, ethanol-treated, novel 3-hydroxylated leukotriene B₄ metabolites from, 3:762
- Rat liver, in vitro, and microsomal metabolism of bromotrichloromethane and carbon, tandem mass spectrometry study of C-phenyl-N-tert-butyl nitrene spin adducts, 6:847
- Rate constants for the reactions of Ar⁺ with CO₂ and SO₂ as a function of temperature (300–1500 K), 10:815
- R=C₆H₅L⁺arene, gas phase RSH₆H₅Cr⁺L and H₆H₅SSH₆H₅Cr⁺ ions, a tandem mass spectrometry study, 5:928
- RDX, negative-ion formation in, by using the reversal electron attachment detection technique, 3:243
- Reaction from isomeric parent ions in the dissociation of dimethylpyrroles, 7:930
- Reaction of analyte ions with neutral chemical ionization gas, 3:549
- Reaction times and applied voltages, negative chemical ionization in quadrupole ion trap mass spectrometry, 5:757
- Reactions of atomic transition metal ions with long chain alkanes, 12:367
- Reactions of CH₃OCH₂⁺ with hydrocarbons and O, N, and S compounds, applications for chemical ionization in selected ion flow tube studies, 13:1015
- Reactions of ground state Cu⁺ and Fe⁺ with the 20 common amino acids, 7:722
- Reactions of O⁻ with methyl benzoate, NICI and FTICR study of, 12:694
- Reactions of polycyclic aromatic hydrocarbon radical cations, with model biological nucleophiles, 7:628
- Reactions of poly(ethylene glycol) cations, with iodide and perfluorocarbon anions, 9:957
- Reactions of proton-bound dimers, 5:695
- Reactions of strong bases with vinyl fluoride formation and characterization of 1-fluorovinyl anion and the fluoride-acetylene hydrogen-bonded complex, 3:91
- Reactions of the phenylium cation with small oxygen- and nitrogen-containing molecules, 7:473
- Reactive behavior of transition metal oxide cluster ions in the gas phase, characterization of, 2:212

- Reactive collisions in quadrupole cells
H/D exchange reactions of emolate ions with CH₃OD and C₂H₅OD, 3:853
H/D exchange reactions of protonated aromatic amines with ND₃, 6:19
- Reactive ion scattering spectrometry, small molecules, peptides and non-covalent protein complexes, 19:190
- Reactivity and binding energies of transition metal halide ions with benzene, 13:477
- Reactivity of CINP+NCl toward cyclic organic ethers, 10:379
- Reactivity of collisionally activated dichlorocarbene dications studied by tandem mass spectrometry, 6:1019
- Reactivity-structure relations, systematic study of, metastable and collision-induced fragmentation studies of all C₄H₁₂Si⁺ isomers, 5:537
- Reagent gas, isotopically labeled, chemical ionization mass spectrometry using, 5:596
- Reagent ions
in infrared laser desorption/ionization Fourier transform ion cyclotron resonance mass spectrometry, potassium halide adducts as, 3:301
injection of, into the seldedge region in fast-atom bombardment mass spectrometry, 4:259
- Reagent plasma, thermospray, from ammonium acetate, controversial ions (m/z 59, 60, 77, and 119) in, on the origin of some, 5:186
- Real-time images of low energy beams, monitor for displaying, 2:226
- Real-time monitoring of some atmospheric pollutants, ion trap detector used for, effects of air and helium on, 4:774
- Real-time quantitative analysis, of H₂, He, O₂, and Ar by quadrupole ion trap MS, 13:1120
- Realistic simulation of ICR-MS using a distributed three-dimensional particle-in-cell code, 10:136
- Rearrangement involving loss of the C-terminal amino acid residue, metastable decomposition of peptide [M + M]⁺ ions via, 1:249
- Recognition and quantification of binary and ternary mixtures of isomeric peptides by the kinetic method: metal ion and ligand effects on dissociation of metal-bound complexes, 14:152
- Recombinant human macrophage colony stimulating factor, disulfide linkage and disulfide bond scrambling in, by fast-atom bombardment mass spectrometry of enzymatic digests, 6:638
- Recorder, transient, dynamic range of, increasing the, by using two analog-to-digital converters, 7:107
- Redox processes by ruthenium(II) polypyridine complexes induced by fast-atom bombardment mass spectrometry, 4:306
- Reduced pressure conditions, comparison of the reactions of the oxide radical anion (O^{•-}) with simple aromatic compounds at, 5:367
- Reduction of copper(II) complexes by electron capture in an electrospray ionization source, 9:1217
- Reduction of Cu(II) in MALDI-MS, 14:42
- Reduction/oxidation processes observed in fast-atom bombardment/liquid secondary ion mass spectrometry, true effect of experimental parameters on, 4:482
- Reductive alkylation by acetone, quantitative proteomics, 20:1366
- Reference compound for positive and negative ion fast-atom bombardment high-resolution mass spectrometry, Ulmark 1621 as, 3:842
- Reflector matrix-assisted laser desorption-ionization time-of-flight and tandem four sector mass spectrometer, peptide fragmentation obtained from, 6:822
- Reflectron MALDI-TOF mass spectrometer, mass correlated acceleration in, 13:135
- Reflectron time-of-flight mass spectrometry, detection of femtomole and sub-femtomole levels of peptides by, 2:91
- Reflectron time-of-flight mass spectrometry (reTOFMS), for peptide sequencing and protein identification, 11:127
- Reflectron time-of-flight (TOF) mass spectrometry, of hydrocarbon polymers, 16:2009
- Reflectrons, broad energy range focusing, design of, 7:1002
- Refolding, stepwise, of acid-denatured myoglobin, evidence from electrospray mass spectrometry, 4:638
- Regioisomers of F2-isoprostanes, four, formed by free radical oxidation of arachidonic acid, mass spectrometric analysis of, 7:490
- Regional stability changes, in oxidized and reduced cytochrome c located by hydrogen exchange and mass spectrometry, 8:1039
- Regioselectivity of the oxygen addition-induced dechlorination, of PCBs and DDT metabolites in electron capture mass spectrometry, 7:66
- Reionization-neutralization, internal energy distributions of tungsten hexacarbonyl ions after, 5:1093
- Reionization-neutralization and dissociation, stabilities and isomerization proclivities studied by, distonic ion CH₂CH₂CH⁺OH, keto ion CH₃CH₂CH₃O⁺, enol ion CH₂CH=CHOH⁺, and related C₃H₆O⁺- radical cations, 7:573
- Reionization-neutralization mass spectrometry
angle resolved, 2:492
organic neutralization agents for, 3:108
- Reionization-neutralization studies, novel tandem quadrupole-acceleration-deceleration mass spectrometer for, 3:493
- Relative collision cross sections of organic ions, 6:803
- Relative dissociation energy measurements using ion trap collisional activation, 5:250
- Relative ion yields for glow discharge mass spectrometry, theoretical calculations of, 3:79
- Relative ion yields measured with a high-resolution glow discharge mass spectrometer operated with an argon/hydrogen mixture, 4:278
- Relative peak heights in Fourier transform-ion cyclotron resonance spectra, space charge effects on, 3:734
- Relative silver(I) ion binding energies of α -amino acids, determination by means of the kinetic method, 9:760
- Relaxation, collisional, metastable electronic states of Fe⁺, 6:543
- Relaxation of vibrationally excited gas-phase Cr(CO)₅⁻, 10:950
- Remeasurement at high resolving power in Fourier transform ion cyclotron resonance mass spectroscopy, 6:564
- Remote fragmentations of protonated aromatic carbonyl compounds via internal reactions in intermediary ion-neutral complexes, 3:417
- Remote sampling, remote MS sampling using air ejector, ESI and DESI, 18:1844
- Renin inhibitor, CP-80,794, in human serum, quantitative analysis by high-performance liquid chromatography atmospheric pressure chemical ionization mass spectrometry, 2:164
- Renin substrate, apparent gas-phase acidity of, 7:1211
- Reporter molecules, nonspecific ligand binding identification using, ESI-MS, 20:1242
- Reproducibility, evaluation of, complex spectra, applied to tubulin, 19:367
- Research, philosophy of, building mass spectrometers and, 1:119
- Residence time, ion source, estimating, 4:787
- Residue, cysteic acid, in oxidized human apoprotein B-100, tandem mass spectrometric characterization of, 6:242
- Residue determination of compounds, by LC-MS/MS, 16:1619

- Residue mass plot and abundance plot, in detection of isobaric interferences in DE-MALDI-TOFMS of complex polymer mixtures, 12:911
- Resin, Egyptian mummy, by mass spectrometry, 3:582
- Resing, Kathryn A.H., obituary, 20:1576
- Resolution and mass accuracy in MALDI-TOF, 9:892
- Resolution equations, for high-field ion mobility, 15:1320
- Resolution limitations from detector pulse width and jitter in a linear orthogonal-acceleration time-of-flight mass spectrometer, 5:772
- Resolved isotopic distributions, monoisotopic masses and ion populations for large biomolecules from, 6:229
- Resolved isotopic peaks for multiply charged ions, charge states from, automated assignment of, 6:52
- Resolvin D1, protectin D1, and related derived products of DHA, analysis via ES-MS/MS, 18:128
- Resolving power
attainable from collisionally activated processes in sector instruments, influence of length and position of collision cell on, 6:212
in complex MS/MS spectra, 16:183
- Resonance, Fourier transform ion cyclotron, multiphoton ionization spectroscopy detection of, 7:1018
- Resonance effects, nonlinear, during ion storage in a quadrupole ion trap, 4:917
- Resonance ejection, co-occurrence of boundary ejection and, ion traps, multiplexed multilinear MS, 17:56
- Resonance ejection and ion dissociation, competition between, during resonant excitation in a quadrupole ion trap, 5:1031
- Resonance ejection mode of quadrupole ion trap mass spectrometry, mass resolution in, extended theoretical considerations for, 5:676
- Resonance shifts in the excitation of the $n = 0$, $K = 1$ to 6 quadrupole resonances for ions confined in a linear ion trap, 13:577
- Resonant current pulse excitation, and multiparticle simulation of ion motion in ion trap mass spectrometer, 6:57
- Resonant ejection of atomic ions, enhanced abundance sensitivity by, plasma source ion trap mass spectrometry, 7:1161
- Resonant excitation in a low pressure linear ion trap, 14:622
- Resonant excitation in a quadrupole ion trap, competition between resonance ejection and ion dissociation during, 5:1031
- Resonant laser ablation as a selective metal ion source for gas-phase ion molecule reactions, 7:664
- Resonantly elicited ions, positions, velocities, and kinetic energies of, in the quadrupole ion trap mass spectrometer, by laser photodissociation, 4:792
- Resorcarene capsule complexes, alkali metal-mediated, 13:851
- Resorcarene capsules, ammonium ion-mediated, ESI-FRIRMS study of, 14:143
- Retinoic acid-binding protein-1, cellular, hydrogen/deuterium exchange MS study of, 16:869
- Retinoid X receptor α ligand binding domain, ligands with, ES MS of, 16:1631
- Retinoid X receptors, ligand interactions with, H/D exchange and MS, 17:1510
- reTOFMS. *See* Reflectron time-of-flight mass spectrometry
- Retro-Michael fragmentation reactions, in gas phase, of 2-hydroxybenzyl-*N*-pyrimidinylamine derivatives, 16:1552
- Reversal electron attachment detection technique, negative-ion formation in the explosives RDX, PETN, and TNT by using, 3:243
- Reverse phase-high performance liquid chromatography mass spectrometry, oligosaccharides, GONE sequencing and structural analysis by, 20:1821
- Reverse phase-liquid chromatography-atmospheric pressure ionization mass spectrometry, comparison of dopants for charge exchange ionization of PAHs, 20:73
- Reversed phase liquid chromatography, of peptides, 8:495
- Reversed-phase microbore liquid chromatography/electrospray mass spectrometry, complementary mobile phases, and sequential exoglycosidase digestion, characterization of protein N-glycosylation by, 5:350
- Reviewers, plea to, editorial, i (April)
- Reviews
1996 ASMS Fall Workshop, 8:1180
of Mass spectrometry PittCon'99, 10:457
of PittCon 2003, 14:542
of 11th Sanibel Conference on Mass Spectrometry: Mass Spectrometry in Clinical Diagnosis of Disease, 10:561
- RF/DC isolations, alternating, for quantitation with coeluting internal standards in gas chromatography/ion trap mass spectrometry, 3:85
- RF-only quadrupoles, collision-activated dissociation in, instrument-independent measurements of, kinetics-based protocol for, precision and accuracy in tandem mass spectrometry measurements, 1:272
- RF oscillator, high voltage, for driving multipole ion guides, 13:1370
- Rhenium compounds, transformation of, by ESI-MS, 8:191
- Rhenium oxide surface, high-efficiency-surface-induced dissociation on, 4:869
- Rhodamine 575, fluorescence and photodissociation of cations, quadrupole ion trap, 20:96
- Rhodamine 6G
detection of attamole amounts by DSI-MS and fluorescence spectroscopy, 18:1093
effect of buffer gas on fluorescence yield, 17:855
- Rhodobacter capsulatus*, cytochrome c_2 stability probed by solution-phase H/D exchange and MS, 17:1518
- Rhodobacter sphaeroides*, characterization of ornithine and glutamine lipids from cell membranes, MS/MS, 20:198
- Rhubarbs, analysis of phenolic compounds, LC-ESI-MS, 18:82
- Ribonuclease B, bovine, high mannose glycans characterized by ion trap MS, 20:539
- Ribonuclease S, thermal denaturation of, by electrospray ionization mass spectrometry, 5:614
- Ribonucleases, ESI-MS detection of ribonuclease-nucleotide complexes, 19:1199
- Ribonucleic acids, intact, molecular mass measurement of, via electrospray ionization quadrupole mass spectrometry, 6:27
- Ribonucleosides, 2'-*O*- and 3'-*O*-methylated, differentiation by MS/MS, 17:1096
- Riboses, reactivity of aromatic σ, σ -biradicals toward, 17:1325
- Ribosomes, topography analyzed by proteolysis and MALDI-MS, 18:1304
- Rice-Ramsperger-Kassel-Marcus rate constant, calculation using vibrational frequencies of linear peptide and protein ions, 17:1749
- Rice-Ramsperger-Kassel-Marcus unimolecular reaction rate constants, SB algorithm modification for calculation of, 18:1063
- Rice-Ramsperger-Kassel-Markus quasiequilibrium theory, 8:103
- Ring-opening H-transfer in ionized cyclopentanone, Ab initio characterization of, similarity to ion-neutral complex-mediated alkane eliminations, 7:1251
- RNA
binding to, high throughput screening for, 15:884
crosslinks with proteins, detection by nanoLC-ESI-MS/MS, 18:869

- fragmentation studied by H/D-exchange, MALDI-TOF-TOF, 17:1353
- hairpins, probing with cobalt(III)-hexamine and ESI-MS, 17:1376
- multiply deprotonated, ion trap collision-induced dissociation, 19:1832
- normal and chemically modified, AP MALDI-FTMS of, 16:199
- RNA/DNA oligonucleotides, mixed-sequence, electrospray MS/MS of, 13:936
- Robertson, Carol, Critical Insights feature, 19:627
- Robinson, Carol V., Biemann Medal awardee, 15:1379
- Rocks, ancient terrestrial, polycyclic aromatic hydrocarbons in, μ L2MS vs GC/MS of, 12:989
- Role of acidic residues and sodium ion adduction on gas phase H/D exchange of peptides and peptide dimers, 14:1477
- Role of ion-ion recombination for alkali chloride cluster formation in liquid secondary ion mass spectrometry, 4:410
- Role of the site of protonation in the low-energy decompositions of gas-phase peptide ions, 7:522
- Role of the sulfhydryl group on the gas phase fragmentation reactions of protonated cysteine and cysteine-containing peptides, 9:1275
- Ron Hites Award, editorial, i (Nov)
- Route 66 method, insulin disulfide bond mapping with, 20:157
- Routine sample analysis, open access atmospheric pressure chemical ionization mass spectrometry for, 6:387
- RP-LC-APPI-MS. *See* Reverse phase-liquid chromatography-atmospheric pressure ionization mass spectrometry
- RPLC (reversed phase liquid chromatography), of peptides, 8:495
- RRKM. *See* Rice-Ramsperger-Kassel-Marcus rate constant; Rice-Ramsperger-Kassel-Marcus unimolecular reaction rate constants
- RRKM calculations on model large polypeptide ions, effect of ion size on rate of dissociation, 4:11
- RRKM modeling, fragmentation of oxidized peptides analyzed by SID, MD and, 20:1579
- RRKM/QET, 8:103
- RSH₆H₅Cr⁺L (R=CcH₅;L⁺arene), gas phase, and H₆H₅SSH₆H₅Cr⁺ ions, tandem mass spectrometry study of, 5:928
- Ru⁺, activation of C₂H₆, C₃H₈, HC(CH₃)₃, and c-C₃H₆ by, 10:821
- Ruthenium, organoruthenium anticancer complexes, analysis using IM-MS, 20:1119
- Ruthenium arene complexes, reactions with glutathione, nanoLC-FTICR MS, 19:544
- Ruthenium carbonyl clusters, gas-phase reactivity, ED-ESI-MS, 20:658
- Ruthenium(II) complexes, photo-oxidized, 8:713
- Ruthenium(II) polypyridine complexes, redox processes of, induced by fast-atom bombardment mass spectrometry, 4:306
- S
- S100 protein CP10, dimeric complexes of, ESI-MS of, 9:533
- S-SIM. *See* Static secondary ion mass spectrometry
- Saccharides
- complexation of with protonated peptides in gas phase, decreases rates of hydrogen/deuterium exchange reactions, 6:1247
- derivatized, ESI and MALDI MS of, 16:683
- sodiated isobaric, separation using ESI-APIM-TOF MS, 16:660
- Vibrio cholerae* 0:1, ESI IT-MS fragmentation of analogs of, 17:749
- Saccharomicins, structural validation of, by FT-ICR MS, 10:1285
- Saddle field fast-atom bombardment gun, flux from, 5:100
- SALDI-MS. *See* Surface-assisted laser desorption/ionization-mass spectrometry
- SALDI TOF MS. *See* Surface-assisted laser desorption/ionization time-of-flight mass spectrometry
- Salen complex analysis, by FAB and APCI, 8:996
- Salmonella enterica*, nitrogen-containing intracellular metabolites, HPLC-tandem MS quantitation, 17:37
- Salmonella typhimurium*, acylphosphatidylglycerols from, ESI-MS/MS of, 15:1
- Salt
- anionic, in arginine molecular recognition, 15:616
- tetraalkylammonium, molecular ions sputtered from a gelatin matrix, formation and emission of, 4:930
- Salt bridge mechanisms, disulfide bond cleavage via salt bridge mechanisms, peptides, low energy CID, 18:1109
- Salt cluster formation, in ESI, 12:1077
- Salt complexes, in gas phase, CAD of, stereoselectivity in, 15:1509
- Salt concentration, analyte response and, ESI-MS of, 10:625
- Salt-containing proteins, MALDI-TOF-MS of, porous anodic alumina membrane as sample support for, 16:1488
- Salt tolerance, of DESI, 18:2218
- Salts, quaternary phosphonium, laser microprobe mass spectrometry of, direct *versus* matrix-assisted laser desorption, 4:798
- Sample deposition, for polymer MALDI-MS, 15:168
- Sample holder, cryogenically cooled, radiofrequency glow discharge MS in, 8:1214
- Sample modulation at a membrane inlet mass spectrometer, mathematical study of, potential application in analysis of mixtures, A, 7:93
- Sample preparation
- 3-layer, nanomolar protein samples, MALDI MS analysis, 17:780
- comparison between vacuum sublimed matrices and dried droplet preparation, MALDI-TOF MS, 20:1104
- multiple, of synthetic polymers, MALDI-MS, 18:377
- vortex method for preparing solvent-free samples, MALDI MS, 20:1115
- Sample processing time, in MS of protein digests, 15:784
- Sampling, remote, using air ejector, ESI and DESI, 18:1844
- Sampling interface, electrically floating, inductively coupled plasma mass spectrometry with, 4:733
- Sampling of aryldiazonium, anilino, and aryl radicals by membrane introduction mass spectrometry: thermolysis of aromatic diazoamino compounds, 9:234
- Sampling orifice, enlarged, and offset ion lens, inductively coupled plasma mass spectrometry with
- ion trajectories and detector performance, 4:16
- polyatomic ion interferences and matrix effects, 4:28
- Sampling orifice potential, characterization of a glow discharge plasma as a function of, 1:448
- SAMs. *See* Self-assembled monolayer surfaces
- Saponins, from medicinal herbs, ESI-MS/MS structural analysis of, 15:133
- Sarcosine oxidase, ESI-MS of, 9:510
- Saturated and unsaturated aliphatic molecules, 193-nm photodissociation of ions from, 7:114
- Saturn's moon (Titan), organic environment on, simulation by FT-ICR and ion trap MS, 16:850
- SAX. *See* Strong anion exchange
- Saxitonix, protonated, gas phase dissociation reactions of, 15:462
- SBA-15, small molecule analysis by MALDI-TOF-MS, 20:2167
- Sc⁺, Y⁺, and Lu⁺, gas-phase reactions with alcohols, effects of the class and chain length of alcohols on the nature of primary products, 7:1157
- Scan modes, multiple, in the hybrid tandem mass spectrometric screening and characterization of the glutathione conjugate of 2-furamide, 2:55

- Scanning, linked
 artifact peaks in, 8:554
 fast, of quadrupole mass spectrometer, 8:539
 parent ion resolution in, 8:545
- Scanning microprobe matrix-assisted laser desorption ionization (SMALDI) mass spectrometry: instrumentation for submicrometer resolved LDI and MALDI surface analysis, 13:735
- Scans at a constant ratio of B/E for studying decompositions of peptide metal(II)-ion complexes formed by electrospray ionization, 6:608
- Schiff base derivatives, pyridine-containing, for the structural determination of long-chain aldehydes by gas chromatography combined with mass spectrometry, 2:245
- School, Mass Spectrometry, The Massachusetts Institute of Technology, 5:332
- Schweppe dye collection, pigments and dyes, MALDI TOF MS identification, 18:2001
- Sco, Al, Fe, Co, Cu, M(acetone)₂⁺, collision-induced and infrared multiphoton dissociation studies on, in the gas phase, 4:135
- Scope and limitations of a general unknown screening by gas chromatography-mass spectrometry in acute poisoning, 6:995
- Scoring methods in MALDI peptide mass fingerprinting, ChemScore and the ChemAppex program in, 13:22
- Scrambling
 carbon, via isomerization to fulvene structure, 1,2-hydrogen shifts in, benzene radical cation, Ab initio calculations on, 6:513
 disulfide linkage and disulfide bond, in recombinant human macrophage colony stimulating factor by fast-atom bombardment mass spectrometry of enzymatic digests, 6:638
- Screening, multiresidue, food and feed matrices, Orbitrap, 20:1464
- Screening for drugs in serum by ESI/CID and library searching, 10:1028
- SDS-PAGE. *See* Sodium dodecyl sulfate polyacrylamide gel electrophoresis
- SDZ RAD, new metabolite of, identification by HPLC and electrospray MS/MS, 11:516
- SDZ RAD metabolism, ESI-MS of, 9:1267
- Search of sequence databases with uninterpreted high-energy collision-induced dissociation spectra of peptides, 7:1089
- SEC-MS. *See* Size exclusion mode mass spectrometry
- Secondary electron capture in the liquid secondary ionization mass spectrometry beam-induced dehalogenation of organic compounds, evidence for a mechanism that involves, 7:1109
- Secondary electrospray ionization, explosives, ambient vapors detected by, 20:287
- Secondary electrospray ionization mass spectrometry, vapors on human skin detected by, 20:1060
- Secondary ion emission and solution chemistry from aminoglycerol solutions, 2:33
- Secondary ion images, of rodent brain, 11:191
- Secondary ion mass spectrometry
 carbon-13 labeling for improved organic material profiling, 17:1142
 comparison of gold and cobalt ion cluster projectiles, molecular depth profiling, 18:406
 investigation of secondary ion yield enhancement using, 19:33
 liquid, characterization of the opioid peptide-containing protein and of bovine alpha-lactalbumin by, 5:377
 of matrix diluted oligo- and polypeptides bombarded with slow and fast projectiles, 11:553
 MCs⁺-molecular ion formation in, during sputtering, 11:650
 polyelectrolytes as matrices for, 16:1575
 in situ analysis and reactions of surfaces prepared by SL of ions, IT MS, 20:949
 static, organic ion imaging beyond, 6:1047
 tandem, imaging large insulating samples by, charge compensation for, 5:37
 under silver bombardment, 16:733
 use for resolution and imaging of biological molecules, 17:1195
- Secondary-ion mass spectrometry, liquid, beam-induced reaction between meta-nitrobenzyl alcohol and dipyrrocyanine dyes in, 5:1017
- Secondary ion microprobe ion trap MS, 13:1099
- Secondary ion source, external, for Fourier transform mass spectrometry, 1:208
- Secondary ionization mass spectrometry
 liquid, direct desorption of crown ether-metal ion complexes in, 5:638
 static, detection of cyclohexamine on soil surfaces exposed to laboratory air, 7:168
- Secondary-ionization mass spectrometry, time-of-flight, ion intensity in, enhancement of, 7:467
- Secondary ionization mass spectrometry analysis, static, of tributyl phosphate on mineral surfaces, effect of Fe(II), 6:165
- Sector instruments, collisionally activated processes in, resolving power attainable from, influence of length and position of collision cell on, 6:212
- SED. *See* Simultaneous excitation/detection
- Sediments, Lake Ontario, plutonium 240/239 ratios in, 2:240
- Segmented radiofrequency-only quadrupole collision cell for measurements of ion collision cross section on a triple quadrupole mass spectrometer, 8:697
- Segmented ring, cylindrical ion trap source for time-of-flight mass spectrometry, 7:1009
- SELDI-TOF MS. *See* Surface enhanced laser desorption/ionization-TOF MS
- Selected ion flow tube mass spectrometry (SIFT-MS)
 ionic diffusion and mass discrimination in, 12:863
 of stable isotopes in water: isotopic composition of H₃O⁺ and H₃O⁺(H₂O)³ ions in exchange reactions with water vapor, 11:866
- Selected ion flow tubes, ion reactions with ethylene oxide and propenal, 18:445
- Selection of solvent load and first-stage pressure to reduce interference effects in inductively coupled plasma-mass spectrometry, 3:128
- Selective chemical ionization of nitrogen and sulfur heterocycles in petroleum fractions by ion trap mass spectrometry, 4:322
- Selective detection of phosphopeptides in complex mixtures by electrospray liquid chromatography/mass spectrometry, 4:710
- Selective detection of sulfur-containing compounds by gas chromatography/chemical reaction interface mass spectrometry, 2:250
- Selective detection of the tolyl cation among other [C₇H₇]⁺ isomers by ion/molecule reaction with dimethyl ether, 2:270
- Selective generation of charge-dependent/independent ion energy distribution from a heated capillary electrospray source, 5:221
- Selective ion isolation/rejection over a broad mass range in the quadrupole ion trap, 2:11
- Selective ion/molecule reactions of lactams with dimethyl ether ions, 4:352
- Selective noncovalent adduct protein probing mass spectrometry

- try, metal interactions with calmodulin and α -synuclein monitored by, 19:1668
- Selective parent ion axialization for improved efficiency of collision-induced dissociation in laser desorption-ionization Fourier transform ion cyclotron resonance mass spectrometry, 5:900
- Selectivity, role of basicity in determination of, ESI MS, 19:719
- Selenium, loss from selenoproteins, 14:593
- Selenocysteine, conversion to dehydroalanine, 14:593
- Selenosulfonate, mass spectra of, 15:1325
- Self-assembled monolayer (SAM) surfaces, of hydrocarbon and fluorocarbon, methyl cation neutralization at, 13:1151
- Self-assembled monolayer surfaces
fluorinated, chemical modification of by low energy reactive ion bombardment, 6:187
reactions of hyperthermal ion beams at, XeF^+ , IF^+ , and other unusual ions generated by, 4:938
- Self-assembled monolayers, laser desorption ion trap mass spectrometry of, 4:968
- Self-chemical ionization, bimolecular and unimolecular contributions to, α -pinene and camphene isomers, 18:2026
- Self-complementary deoxydinucleotides, complex ions derived from, cleavage reactions of, positive ion ESI-MS/MS of, 15:689
- Selvedge region, injection of reagent ions into in fast-atom bombardment mass spectrometry, 4:259
- Semiconductor nanoclusters, ESMS of, 11:338
- Semiempirical studies, investigations of gas-phase lithium-peptide adducts, 1:473
- Semivolatile organic compounds, aerosols, analysis by DESI-MS, 19:450
- Sensitivity
enhancement for MALDI IMS of proteins after treatment of tissue sections, 19:1069
improved using high-pressure electrodynamic ion funnel interface, MS, 17:1299
signal intensity and sensitivity improvement using nanoliter spotting by IBF, MALDI MS, 19:1086
- Sensitivity analysis, ultra-high, of peptides and proteins, microelectrospray mass spectrometry, 5:605
- Sensitivity for laser desorption in an ion trap mass spectrometer, method for increasing, 4:706
- Sensitivity in negative atmospheric pressure chemical ionization liquid chromatography-mass spectrometry, effect of high-performance liquid chromatography mobile phase components on, 7:1059
- Sensitivity in negative ion fast-atom bombardment mass spectrometry of polyanionic analytes, class of cationic matrix additives for enhancing, N-alkylnicotinium halides, 5:935
- Separation and detection of the α - and β -chains of hemoglobin of a single intact red blood cell using CE/ESI TOFMS, 10:184
- Separation and identification of DMPO adducts of oxygen-centered radicals formed from organic hydroperoxides by HPLC-ESR, ESI-MS, and MS/MS, 14:862
- Separation and identification of positively charged and neutral nucleoside adducts by CEC- μ ESI/MS, 9:823
- Separation-charge reactions in tandem mass spectrometry of doubly protonated angiotensin II formed by electrospray ionization, experimental considerations and structural implications, 7:30
- Separation of leucine and isoleucine by ESI-FAIMS-MS, 10:1279
- Sequence and side-chain specific photofragment (193 nm) ions from protonated substance P by MALDI-TOFMS, 10:1038
- Sequence confirmation, of large peptides by high energy collisional activation of multiply protonated ions, 6:234
- Sequence confirmation of modified oligonucleotides using IRMPD in external ion reservoir of ESI-FTICR spectrometer, 14:825
- Sequence databases, with uninterpreted high-energy collision-induced dissociation spectra of peptides, 7:1089
- Sequence-dependent fragmentation of peptides generated by MALDI quadrupole time-of-flight mass spectrometry and its implications for protein identification, 13:772
- Sequence ions, compound specific, in the collision-induced dissociation spectra of the immonium ions of two isomeric hexaglycosylceramides, electron-ionization-tandem mass spectrometry of glycosphingolipids, 3:353
- Sequence length, sequence frequency, and data acquisition rate effects on performance of a Hadamard transform TOFMS, 12:1302
- Sequence refinement, and active site location from high-resolution tandem mass spectrometry, thiaminase I (42 kDa) heterogeneity, 6:981
- Sequence-specific fragmentation, of deprotonated peptides containing H or alkyl side chains, 12:1
- Sequence-specific stabilization, of linear analog of the antifungal lipopeptide iturin A2 by sodium during low energy ESI-MS conditions, 12:505
- Sequence tag database search, for peptide sequencing and protein identification, MS/MS on IT/reTOF using, 11:127
- Sequencing, rapid, of peptides, with a picomole-scale method and ESI-MS, 17:995
- Sequencing of tri- and tetraantennary N-glycans containing sialic acid by negative mode ESI-quantitative TOF-MS/MS, 13:1138
- Sequential exoglycosidase digestion, reversed-phase microbore liquid chromatography/electrospray mass spectrometry, complementary mobile phases, characterization of protein N-glycosylation by, 5:350
- Sequential ortho effects, characterization of novel $[\text{M}-35]^+$ fragment ions in the mass spectra of 2-alkyl-4,6-dinitrophenols, 2:69
- Sequential product ion scanning, of vertebrate type steroid hormones, 8:365
- SEQUEST searches, Qscore algorithm for, 13:378
- Serene, dipeptides of, gas-phase basicities of, 5:718
- Serine
alkaline metal-cationized clusters, SSI-MS/MS, 18:856
chiral enrichment of, via ion soft-landing, 15:1360
gas-phase basicities of, 5:718
protonated, and derivatives, gas phase fragmentation reactions of, 11:1047
- Serum, human, CP-80,794 in, quantitative analysis by high-performance liquid chromatography atmospheric pressure chemical ionization mass spectrometry, 2:164
- SESI. *See* Secondary electrospray ionization
- SF6, molecular anions of, decomposition reactions of, cluster-assisted, 1:372
- SF6 discharge suppression, ionization from water solution with, negative ion electrospray mass spectrometry of nucleotides, 4:289
- SF6 gas, suppression of electric discharge with, electrospray mass spectrometry of methanol and water solutions, 2:497
- SFC-MS. *See* Supercritical fluid chromatography-mass spectrometry.
- SFCMS. *See* Supercritical fluid chromatography-mass spectrometry
- Sheathless preconcentration-capillary zone electrophoresis-mass spectrometry applied to peptide analysis, 10:1271
- Shell ions, closed *versus* open, onium reaction and H-transfer characterization in, 18:270
- Shift reagents, use in IMS, 20:1

- Shiga toxin β subunits, stability in solution and gas phase, nanoESI-FTICR MS of, 16:1957
- Shiga toxins, Coulombic repulsion effect on dissociation pathways of protein complexes in gas phase, 18:617
- Shigella*, *E. coli* species and, protein expression monitored by LC/QTOF-MS, 19:1626
- Shock wave model for sputtering biomolecules using massive cluster impacts, 3:311
- Shrimp, vertebrate type steroid hormones in, MS/MS and sequential product ion scanning of, 8:365
- Sialic acid, metal-ligated oligosaccharides containing, stabilization and linkage analysis of, 12:528
- Sialic acid synthase, ESI-MS characterization of, 16:324
- Sialyl anomers, distinction on ESI- and FAB-MS/MS, 20:394
- Sialylation, glycan, effect on negative ion dissociation, 17:844
- SiC_3H^+9 , structures of, by using specific ion/molecule reactions, 2:278
- SiC_4H^+11 , structures of, by using specific ion/molecule reactions, 2:278
- SiC_5H^+13 , structures of, by using specific ion/molecule reactions, 2:278
- SiCl_3^+ and SiCl^+ affinities for pyridines determined by using the kinetic method with multiple stage mass spectrometry, agostic effects in the gas-phase, 7:198
- Side chain effects, aspartic acid, experimental and theoretical insights, 18:121
- Siderophores
B. anthracis petrobactin, CID, IRMPD and EC dissociation, 18:842
 enterobactin, Fe(III) complex with, IRMPD, 18:632
- SiF_4 , basicity and proton affinity of, 10:848
- SIFT ion-molecule study, of reactions in Titan/Es atmosphere, 15:1148
- SIFT-MS. *See* Selected ion flow tube mass spectrometry
- σ , σ -Biradicals, aromatic, reactivity toward riboses, 17:1325
- Signal, induced
 in a cubic trap, effect of ion initial axial position on, simulated ion trajectory and induced signal in ion cyclotron resonance ion traps, 5:807
 in ion cyclotron resonance ion traps, 5:238
 and simulated ion trajectory in ion cyclotron resonance ion traps, effect of ion initial axial position on ion coherence, induced signal, and radial or z ejection in a cubic trap, 5:807
- Signal enhancement, boundary activated, 13:650
- Signal enhancement for gradient reverse-phase high-performance liquid chromatography-electrospray ionization mass spectrometry analysis with trifluoroacetic and other strong acid modifiers by postcolumn addition of propionic acid and isopropanol, 6:1221
- Signal filtering, method for quantification and noise discrimination, FT-ICR MS, 20:1087
- Signal intensity, analyte charge state, and stability in negative ion electrospray mass spectrometry, solvent effect on, implications for the mechanism of negative ion formation, 4:546
- Signal intensity of oligonucleotides in electrospray ionization mass spectrometry, improved, charge-state reduction with, 7:697
- Silica
 porous alumina and, peptide fingerprinting on, LDI MS, 19:632
 SBA-15, small molecule analysis by MALDI-TOF-MS, 20:2167
- Silica microspheres, enzyme inhibitors immobilized on, screening by ESI-MS, 19:865
- Silicates, ortho, protonated, APCI low energy collision-induced dissociation fragmentation of, McLafferty or ion-neutral complex rearrangements, 5:544
- Silicon, porous, surface area drives desorption of intact molecules, 18:1945
- Silicone nanoparticle assisted laser-desorption/ionization, surface-induced dissociation analyzed using LDI MS and, peptides, 20:957
- Siloxane bond cleavage, endgroup-assisted, in gas phase, 14:1039
- Silsesquioxanes
 low molecular weight components of, ESI-FTMS of, 16:524
 molecular structure of, MALDI-TOFMS of, 10:224
- SILT. *See* Stable isotope labeling tandem mass spectrometry
- SILVER, in assigning peptides to tandem mass spectra using intensity-based scoring, 15:910
- Silver
 binding by polyamidoamine dendrimers, investigation by ESI-MS/MS, 17:676
 bombardment of
 by energetic carbon and gallium ions, 16:1677
 SIMS under, 16:733
 cationization of, for hydrocarbon MS, 16:437
 chemical reaction interface mass spectrometry in detection of, nitrogen trifluoride in, 6:421
- Silver binding energy, of amino acids, 9:760
- Silver CIS-MS, in analysis of unsaturated compounds, 14:872
- Silver cluster interferences in MALDI-MS of nonpolar polymers, 12:732
- Silver-gold clusters, bimetallic, MALDI of, 15:1455
- Silver halide microcrystals, depth profiling of, MS in, 10:1016
- Silver ion chemical ionization, in MS analysis of tributyl phosphate, 8:718
- Silver nitrate chemical ionization for analysis of hydrocarbon polymers by laser desorption Fourier transform mass spectrometry, 4:453
- Silver(I) complex, produced in ESI-MS, 8:781
- Silylenium ions, isomeric (SiC_3H^+9 , SiC_4H^+11 , SiC_5H^+13), structures of, by using specific ion/molecule reactions, 2:278
- Similarity between condensed phase and gas phase chemistry: fragmentation of peptides containing oxidized cysteine residues and its implications for, proteomics, 12:228
- Similarity index, in spectral comparison, 13:85
- SIMION analysis, of linear octopole from linear octopole ion trap, 13:1304
- SIMION simulations, of gap-tripole ion trap, 19:1367
- Simple approach for coupling liquid chromatography and electron ionization mass spectrometry, 13:265
- Simple cylindrical ion mirror with three elements, 11:759
- Simple geometry gridless ion mirror, 11:764
- Simple method for estimating effective ion source residence time, 4:787
- Simplicity/performance trade-offs, understanding ion deceleration lenses, 2:322
- Simplified application of quadrupolar excitation in Fourier transform ion cyclotron resonance mass spectrometry, 6:448
- SIMS. *See* Secondary ion mass spectrometry
- SIMSTEX, application to oligomerization of insulin analogs and mutants, 17:1526
- Simulated ion trajectory and induced signal in ion cyclotron resonance ion traps, 5:238
 effect of ion initial axial position on ion coherence, induced signal, and radial or z ejection in a cubic trap, 5:807
- Simultaneous analysis of 14 non-steroidal anti-inflammatory drugs in human serum by ESI-MS/MS without chromatography, 9:255

- Simultaneous analysis of butene isomer mixtures using process MS, 11:1079
- Simultaneous excitation/detection (SED), FTICR-MS with, real-time excitation profiles by, 11:1009
- Simultaneous measurement of flight time and energy of large MALDI desorption ionization ions with a superconducting tunnel junction detector, 8:1094
- Simultaneous or scanning data acquisition: a theoretical comparison relevant to ICP sector-field mass spectrometers, 14:227
- Simultaneous quantitation of the 5'-triphosphate metabolites of zidovudine, lamivudine, and stavudine in peripheral mononuclear blood cells of HIV-infected patients by HPLC MS/MS, 11:1134
- Sinapic acids, deprotonated monomeric and dimeric, gas phase basicities of, 10:1111
- Sinapinic acid
 guanosine/deoxyguanosine clusters with, electron transfer vs proton transfer in, 12:176
 proton transfer from, to biomolecular analytes, 9:1060
- Single cells
 plant carbohydrate extracts, pressure probe and UV-MALDI-TOF MS, 19:1842
 preparation for imaging/profiling MS, 19:1230
- Single-ion trajectories in Paul-type ion traps, computer simulation of, 4:687
- Single isolated droplets with a net charge as a source of ions, 11:393
- Single nucleotide polymorphisms, *ApoE* indels and, genotyping of by MS, 17:3
- Single photon ionization time-of-flight mass spectrometry, tunable, laser-produced plasma as VUV light source, 20:2192
- Single series peptide fragment ion spectra generated by two-stage CID in a triple quadrupole, 9:606
- Single trap electrode, high performance Fourier transform ion cyclotron resonance mass spectrometry via, 6:812
- Sistrurus mliarius barbouri*, snake venom, proteomic profiling using MALDI-TOF MS, 18:600
- Site of protonation of benzonitrile hydrogen interchange in the protonated species, 1:225
- Site-selective gas-phase reactions of amino alcohols with dimethyl ether ions, 4:230
- Sites of reaction of pilocarpine, 10:209
- Sixty u, peptide loss of, 8:253
- Size effects in ion-neutral complex-mediated alkane eliminations from ionized aliphatic ethers, 2:261
- Size exclusion chromatography mass spectrometry, reduced monoclonal antibodies, analysis using, 20:2258
- Size exclusion chromatography-microelectrospray ionization mass spectrometry, in study of enzyme inhibitor effect on protein structure, 12:97
- Size exclusion mode mass spectrometry, online, molecular mass analysis of antibodies by, 19:502
- Skeleton, troponin C in, calcium binding site and, 11:770
- Skin
 human, vapors detected by SESI-MS, 20:1060
 sampling explosives on, using neutral desorption and EESI-MS, 20:719
- SLCC. *See* Spontaneous loss of coherence catastrophe
- Slit interfaces, improving sensitivity of FAIMS with, 20:1768
- Slower atomic than polyatomic fragment losses, 14:136
- Sludge, nonylphenol carboxylates in, LC-ES-MS/MS of, 14:516
- SMALDI. *See* Scanning microprobe matrix-assisted laser desorption ionization
- Small-bore electrospray ionization source, construction and characterization of, 4:964
- Small molecules as substrates for adsorption/desorption in ²⁵²Cf plasma desorption mass spectrometry, 2:29
- Smiles rearrangement reactions, gas phase, 2-(4,6-dimethoxy-pyrimidin-2-ylsulfanyl)-N-phenylbenzamide, ESI-MS, 18:1813
- Snake venom, *S. mliarius barbouri*, proteomic profiling using MALDI-TOF MS, 18:600
- Sn(ANRORC) reaction, ionic products from, isomer differentiation by charge inversion tandem mass spectrometry, 3:467
- SNAPP-MS. *See* Selective noncovalent adduct protein probing mass spectrometry
- SNPs. *See* Single nucleotide polymorphisms
- SO₃, proton affinity of, 10:856
- SO₂•electron transfer ion/ion reactions, with disulfide-linked peptides, 16:1020
- Social paper wasps, dominulin A and B in cuticle and venom of, analyzed by MALDI-TOF, MALDI-TOF/TOF and ESI-ion trap, 17:376
- Sodiated isobaric saccharides, separation using ESI-APIM-TOF MS, 16:660
- Sodium
 complexation with aromatic acids in gas phase, cation- π effects in, 11:1037
 iturin A₂ stabilization by, during ESI-MS, 12:505
- Sodium cationized peptides, formation in gas phase, 15:607
- Sodium dodecyl sulfate micelles, in MALDI-MS sample preparations, 16:1438
- Sodium dodecyl sulfate polyacrylamide gel electrophoresis, adenovirus proteins extracted by, MALDI-MS of, 11:356
- Sodium gel-induced protein modifications, in dodecylsulfate polyacrylamide gel electrophoresis, accurate molecular weight determination of gel-separated proteins by MALDI-TOFMS, 10:512, 10:784
- Sodium ion adduction, in gas phase H/D exchange of peptides and peptide dimers, 14:1477
- Sodium ions, affinities of simple di-, tri-, and tetra-peptides, 18:541
- Sodium trifluoroacetate as a tune/calibration compound for positive- and negative-ion ESI-MS in the mass range of 100–400 Da, 9:977
- Soft tissue, biological, molecular secondary ion mass spectrometry studies of, model matrix for, doped gelatin films as, 4:419
- Software
 HD desktop platform for analysis of H/D exchange data, 20:601
 LipidQA, for identifying and quantifying complex lipid mixtures, 18:1848
 MATLAB, artificial neural networks for ppb mass accuracy in FT-ICR/MS, 20:1303
 MS align, genetic algorithm feature extraction, alignment of LC-MS and LC-MS/MS data sets, 18:1835
 Peptide Shifter, enhancing separation reproducibility using correlated expression profiles, 18:1638
- Software-based mass spectral enhancement to remove interferences from spectra of unknowns, 7:598
- Soil, nitro-substituted polynuclear aromatic hydrocarbons in, sub-parts-per-billion determination of, by electron capture-tandem mass spectrometry, 7:1255
- Soil surfaces exposed to laboratory air, cyclohexamine on, static secondary ionization mass spectrometry detection of, 7:168
- Soils, MS of, 8:200
- Solid phase extraction
 carbon nanotubes as adsorbent of, in MALDI MS, 16:263
 to detect steroids in human urine, 10:1322

- Solid-phase microextraction in-line coupled to gas chromatography-mass spectrometry, pesticides in aqueous samples determined by, 6:1119
- Solid phase peptide synthesis, S-SIM in monitoring of, 12:1099
- Solution
 amide anions in, ESI-MS of, 11:1061
 gas-phase molecular dications formed from neutral organics in, via chemical electron-transfer reactions by using electrospray ionization mass spectrometry, 5:689
 hydrogen/deuterium exchange of proteins in, 10:671, 10:672
 neutral organics in, gas-phase molecular dications formed from, via the controlled-current electrolytic process inherent to electrospray, 7:157
 stability in, phenylglycidyl ether adducts of 29-deoxycytidine and 29-deoxyadenosine, 7:682
- Solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes by fast-atom bombardment mass spectrometry
 generation of molecular signals, 4:249
 synthesis and characterization of molecular complexes, 4:255
- Solution chemistry and secondary ion emission from amine-glycerol solutions, 2:33
- Solution composition, and organic molecule responsiveness to negative ion ESI MS, 16:446
- Solution composition and thermal denaturation for the production of single-stranded PCR amplicons: piperidine-induced destabilization of the DNA duplex, 13:232
- Solution flow rates, in nano-ESI-MS, analyte ion signals and, 14:492
- Solution phase
 noncovalent interactions in, ESI-MS diffusion measurements of, 14:430
 oligonucleotide duplexes in, stability determination of, 15:1354
 structure of, water binding energies to lithiated valine in, 15:1014
- Solution phase chemistry, ion structural analog formation by, 16:607
- Solution phase H/D exchange, of non-covalent complexes, ESI-MS of, 12:795
- Solution-phase zinc finger peptide complexation, MALDI-MS of, 10:27
- Solution residues, analysis of, by glow discharge mass spectrometry, 4:47
- Solvated ions, accumulation in accumulation/collision hexapole, hybrid QFTMS, 20:411
- Solvation of copper ions by acetone, structures and sequential binding energies of Cu^+ (acetone)_x, x = 1–4 from CID and theoretical studies, 13:453
- Solvation studies of electrospray ions: method and early results, 9:1241
- Solvent, effects on charge states of protein ions produced by ESI-MS, 11:976
- Solvent and counterion, on ion pairing and observed charge states of diquatery ammonium salts in electrospray ionization mass spectrometry, 7:1050
- Solvent-cation binding, in ESI-MS, 10:254
- Solvent effect, in MALDI-TOFMS analysis of polymers, 9:1303
- Solvent effect on analyte charge state, signal intensity, and stability in negative ion electrospray mass spectrometry, implications for the mechanism of negative ion formation, 4:546
- Solvent load to reduce interference effects in inductively coupled plasma-mass spectrometry, 3:128
- Solvent systems, microwave-assisted enzyme-catalyzed reactions in, 16:581
- Solvents
 disintegration of solvent droplets during ESI, 20:496
 evaporated, analysis of residues from, TM-DSI, 19:1617
 surface effects and, effects on analyte ionization, DIOS MS, 19:8
- Sonic spray ionization, for drug analysis, 13:204
- Sonic spray ionization tandem mass spectrometry, alkaline metal-cationized serine clusters, 18:856
- Soot formation, calcote mechanism for, 7:559
- SORI. *See* Sustained off resonance irradiation; Sustained off-resonance irradiation excitation
- SORI-CAD. *See* Sustained off-resonance irradiation collision-activated dissociation; Sustained off resonance irradiation for collision-activated dissociation
- SORI-CID. *See* Sustained off resonance irradiation-collision induced dissociation
- SOS, simple interactive program for ab initio oligonucleotide sequencing by mass spectrometry, 13:200
- Space-charge, effects of on FTICR, experimental observations and three-dimensional trajectory simulations, 8:319
- Space-charge-dominated mass spectrometry ion sources, modeling and sensitivity, 2:1
- Space-charge effects, in plasma mass spectrometry, 10:159
- Space-charge effects in plasma mass spectrometry, time-resolved measurements of individual ion cloud signals to investigate, 7:362
- Space charge effects on relative peak heights in Fourier transform-ion cyclotron resonance spectra, 3:734
- Space focusing, in TOFMS, 10:1067
- Space program, of U.S., MS in, 12:656
- SPALDI. *See* Silicone nanoparticle assisted laser-desorption/ionization
- Spatial profiling invertebrate ganglia using MALDI-MS, 14:752
- Specific and nonspecific dimer formation in the electrospray ionization mass spectrometry of oligonucleotides, 6:159
- Specificity of cyclodextrin complexes detected by electrospray MS, 13:946
- Spectra, similarity index and contrast angle in, 13:85
- Spectral accuracy, molecular ions in Orbitrap MS, 20:2058
- Spectral deconvolution, automated ESI-MS, 9:225
- Spectral deconvolution for overlapping GC/MS components, 3:558
- Spectral quality, influence of controllable parameters on, improving fast atom bombardment mass spectra, 1:149
- Spectroscopic characterization of the isomers of the allyl bromide molecular ion, 2:372
- Spectroscopy
 multiphoton ionization, Fourier transform ion cyclotron resonance detection of, 7:1018
 photodissociation, of gas-phase ferrocene cation, 2:97
- Spectrum extraction and compound identification, GC/MS for, 10:770
- Spermine, as co-matrix, in MALDI-TOF of acidic glycoconjugates, 9:1293
- Spermine and 5-methoxysalicylic acid, as new matrix for MALDI-MS analysis of oligonucleotides, 12:456
- Sphingolipids, phospholipids and, DESI/MS and MS/MS of, 19:531
- Sphingomyelin, structural determination of, by MS/MS with ESI, 11:437
- Sphingophosphonolipids, *B. stolpii* UKi2, identification of SPNL molecular species, qTOF-II MS, 18:394
- Spider venom acylpolyamides, MS/MS of, 15:1636
- Spin adducts, C-phenyl-N-tert-butyl nitron, from in vitro rat liver, and microsomal metabolism of bromotrchloromethane and carbon, tandem mass spectrometry study of, 6:847
- Spin-trapped radicals, determination by LC-TSP-MS and LC-ESI-MS, 2:413

- Spiral ion trajectory
MALDI-TOF-MS utilizing, 18:1318
in new TOF mass spectrometer, 16:1969
- Spirobenzopyran derivatives, metal-ion complexes of, evaluating photoinduced changes by ESI-MS, 14:1110
- Spleen, tyrosine kinase of, conformational dynamics upon diphosphorylated ITAM binding to, 16:1039
- $S=P(OH)^{2+/\bullet}$, ionic and neutral, generation and characterization in gas phase, 16:1353
- Spontaneous and deflected drift-trajectories in orthogonal acceleration time-of-flight mass spectrometry, 5:588
- Spontaneous loss of coherence catastrophe, FTICR-MS, 20:247
- Spores, bacterial, MALDI-TOF-MS of, 15:1444
- Sputtering, biomolecules, shock wave model for, using massive cluster impacts, 3:311
- SR 27417 in human plasma, quantitation of, using electrospray liquid chromatography-tandem mass spectrometry, study of ion suppression, 7:1099
- Src SH₂-phosphopeptide complexes, ESI-MS of, 8:234
- SSI. *See* Sonic spray ionization
- sSIMS. *See* Static secondary ion mass spectrometry
- Stabilities and isomerization proclivities studied by dissociation and neutralization-reionization, distonic ion CH₂CH₂CH⁺OH, keto ion CH₃CH₂CH₂O⁺, enol ion CH₃CH=CHOH⁺, and related C₃H₆O⁺ radical cations, 7:573
- Stability, analyte charge state, and signal intensity, in negative ion electrospray mass spectrometry, implications for the mechanism of negative ion formation, 4:546
- Stability conditions for multiply reflecting electrostatic ion traps, 10:241
- Stability diagrams, in quadrupole MS, calculation with matrix methods, 13:597
- Stabilization and linkage analysis of metal-ligated sialic acid containing oligosaccharides, 12:528
- Stabilization of carbanions by adjacent phenyl, cyano, methoxy-carbonyl, and nitro groups in the gas phase, 5:159
- Stable association complex ES-MS, for determination of cyanuric acid, 12:1085
- Stable association complex ESI, for cyanuric acid determination, 12:1085
- Stable isotope incorporation triples the upper mass limit for determination of elemental composition by accurate mass measurements, 11:835
- Stable isotope labeling by amino acids in cell culture, proteins labeled by, quantitation and characterization, 18:2058
- Stable isotope labeling tandem mass spectrometry, protein production and clearance rates quantified using, 18:997
- Stable species, ionic, in peptide spectra?, 6:1165
- Stacked-ring electrostatic ion guide, 7:101
- π -Stacking interactions, in crown ether complex self-assembly detection by ESI-MS, 16:1162
- Stages of mass spectrometry, multiple, determination of daughter ion formulas by, 1:166
- Stainless steel, surface-induced dissociation and chemical reactions on carbon, diamonds and, 20:927
- Standard definitions of terms relating to mass spectrometry, report from the Committee on Measurements and Standards of the American Society for Mass Spectrometry, 2:336
- Standard of atomic weight, ¹²C₆₀ as, unsuitability of, 6:1243
- Standards
calibrant ion delivery for MS, 18:991
protein, Aurum Dataset, MALDI-TOF/TOF MS for, 18:850
- STAT6, noncovalent complexes, weak interactions analyzed by ESI-MS, 20:303
- Static pressure, of heavy gases in quadrupole ion trap, 14:1099
- Static secondary ion mass spectrometry
activation processes and polyethylene formation on Phillips model catalyst analyzed by LA-MS, LDI-MS and, 17:406
limit of, organic ion imaging beyond, 6:1047
- Static-secondary ion mass spectrometry, ion suppression and enhancement of secondary formation, 18:1559
- Static-secondary ion mass spectrometry (S-SIM), in monitoring solid phase peptide synthesis, 12:1099
- Static secondary ionization mass spectrometry analysis of tributyl phosphate on mineral surfaces, effect of Fe(II), 6:165
- Static secondary ionization mass spectrometry detection of cyclohexamine on soil surfaces exposed to laboratory air, 7:168
- Stationary phase, polarizable, negative ion chemical ionization, and mass-resolved chromatography, high-temperature gas chromatography/mass spectrometry with, high-resolution triglycerol mixture analysis using, 7:350
- Statistical experimental design, optimization of the Hewlett-Packard particle-beam liquid chromatography-mass spectrometry interface by statistical experimental design, 6:507
- Stavudine metabolites, in peripheral mononuclear blood cells of HIV-infected patients, HPLC-MS/MS of, 11:1134
- Stepwise-external calibration, FT-ICR MS, sub part-per-million mass accuracy, 17:1681
- Stepwise refolding of acid-denatured myoglobin, evidence from electrospray mass spectrometry, 4:638
- Stereochemistry of glycosidic bond configuration, differentiation of, tandem mass spectrometry of diastereomeric cobalt-glucosyl-glucose disaccharide complexes, 7:953
- Stereoisomeric steroids, differentiation by reactions with phosphonium ions, 13:362
- Stereoselectivity, in CAD of gas phase salt complexes, 15:1509
- Stereoselectivity and ion response in electrospray mass spectrometry, correlation of, and electrospray ionization-mass spectrometry as tool to predict chemical behavior, 6:207
- Stereospecific fragmentation processes in cycloalkane/cycloalkane-fused isomers of saturated pyrrolo[2,1-b][1,3]oxazin-6-one derivatives, 10:393
- Stereospecific fragmentations in the mass spectra of stereoisomeric isoindoloquinazolines, 14:1049
- Steric and substituent effects in phenols, evaluation of by competitive reactions of dimethyl ether ions in a quadrupole ion trap, 6:627
- Steroid hormones
by LC-MS, 8:1010
vertebrate type, MS/MS and sequential product ion scanning of, 8:365
- Steroids
quantitative signatures using GC-MS, 20:1626
thermally-labile, in supersonic molecular beams, fast, very fast and ultra-fast gas chromatography-mass spectrometry of, 7:737
- Steroids in human urine, solid-phase extraction and LC-MS/MS in detection of, 10:1322
- Stevenson's rule, decompositions of N-(substituted benzal-amino)phthalimide radical cations embody, 5:837
- Stoichiometry, metal binding, of matrilysin by electrospray ionization mass spectrometry, noncovalent enzyme-inhibitor complexes and, 6:1105
- Storage temperature, greater than 300 K, gaseous myoglobin stored at, 5:324
- Stored waveform inverse Fourier transform excitation, MALDI-FTICR MS, improved mass accuracy for peptides using, 19:76
- STR database, as online resource for characterization of PCR products using ESI-MS, 13:87

- Streams, waste, components of, by electrospray and tandem mass spectrometry, 4:604
- Streptavidin, Coulombic repulsion effect on dissociation pathways of protein complexes in gas phase, 18:617
- Streptavidin binding to biotin, electrospray ionization mass spectrometry of, 6:912
- Streptavidin complexes, noncovalent, tetrameric ions of formed by electrospray ionization, 6:459
- Stress-related fingerprint identification, from *Bifidobacterium lactis*, by MALDI-TOF-MS, 15:1222
- Strong anion exchange, for studying protein/DNA interactions, H/D exchange MS, 19:887
- Strong-cation-exchange, human plasma proteome mapping using SCX-LC-IMS-MS, 18:1249
- Strong cation exchange μ -HPLC-ESI-MS/MS, on-line, 14:719
- Structural analysis of chromophore-labeled disaccharides, by CE-MS/MS, using IT-MS, 11:292
- Structural analysis of phosphatidylcholines by PSD MALDI-TOFMS, 14:373
- Structural analysis of photo-oxidized (ethylenediamine)bis(2,2'-bipyridine)ruthenium(II) complexes by using one-line electrospray mass spectrometry of labeled compounds, 8:713
- Structural analysis of underivatized neutral human milk oligosaccharides in negative ion mode by nano-electrospray MS
application to isomeric mixtures, 13:1341
methodology in, 13:1331
- Structural characterization and differentiation of modified isomeric tryptophans, 13:1298
- Structural characterization of normal and modified oligonucleotides by matrix-assisted laser desorption Fourier transform mass spectrometry, 2:402
- Structural characterization of organophosphorus compounds, multiple-stage mass spectrometry in, 4:125
- Structural characterization of triacylglycerols as lithiated adducts by ESI-MS using low-energy CAD on a triple stage quadrupole instrument, 10:587
- Structural determination of picomole amounts of phospholipids via electrospray ionization tandem mass spectrometry, 6:1202
- Structural determination of sphingomyelin by MS/MS with ESI, 11:437
- Structural effects on polyether cationization by alkali metal ions in MALDI, 11:639
- Structural elucidation
by charge-remote fragmentations, of oxofatty acids, 9:216
by ESI-MS, in vitro metabolism of the macrolide immunosuppressant SDZ RAD and, 9:1267
generation of molecular signals, 4:249
of O-linked glycopeptides by high energy collision-induced dissociation, 7:319
studies on 14- and 16-membered macrolide aglycones by accurate-mass electrospray sequential mass spectrometry, 13:862
- Structural studies on alkylisocyanate polymers by thermal degradation tandem mass spectrometry, 2:130
- Structural studies on ceramides as lithiated adducts by low energy CAD MS/MS with ESI, 13:680
- Structural validation of saccharomycins by high resolution and high mass accuracy FT-ICR MS and infrared multiphoton dissociation MS/MS, 10:1285
- Structure
of flavonoid glycosides, characterization by CAD of metal complexes, 12:537
glycosphingolipids as lithiated adducts, determination by ESI-MS using low-energy CAD on triple stage quadrupole instrument, 12:61
of ionized 1,5-hexadiene in gas phase, 12:840
of melittin bound to phospholipid micelles studied using H/D exchange and ESI-FTICR-MS, 12:1247
of neutral O-linked oligosaccharides, exoglycosidase digestion and, 12:877
- Structure analysis by electrospray tandem mass spectrometry, phenylglycidyl ether adducts of 2-deoxycytidine and 2-deoxyadenosine, 7:682
- Structure and collision activated decompositions, gas-phase anionic complexes of alkali metal ions and peptides, 5:137
- Structure and fragmentation mechanisms of isomeric T-rich oligodeoxynucleotides, comparison of four MS/MS methods, 9:683
- Structure and fragmentation of b_2 ions in peptide mass spectra, 11:427
- Structure and fragmentation of Bn ($n-3$) ions in peptide spectra, 7:233
- Structure and mechanism of formation of $C_3H_9O^+$ from ionized phytyl methyl ether, 7:205
- Structure characterization of dinucleoside and nucleoside glucopyranosides lipophilic and phosphotriesters by FAB MS/MS, 10:975
- Structure confirmation, accurate masses for, 5:57
- Structure determination, by MS, four decades of, 13:1254
- Structure determination of ceramides and neutral glycosphingolipids by collisional activation of $[M + Li]^+$ ions, 3:260
- Structure elucidation, gas phase hydrogen-deuterium exchange in electrospray ionization mass spectrometry as a practical tool for, 5:434
- Structure elucidation of everninomicin-6, new oligosaccharide antibiotic, by chemical degradation and FAB-MS methods, 8:1134
- Structure of $C_5H_5RFe^+$ ($R=F, Cl, Br, I, O, OH, OCH_3, H_6H_5$) ions in the gas phase and the generation of their neutral counterparts by neutralization-reionization mass spectrometry, 6:1143
- Structure of protonated methane, 2:453
- Structure-reactivity relations, systematic study of, metastable and collision-induced fragmentation studies of all $C_4H_{12}Si^+$ isomers, 5:537
- Structures and fragmentation of $[Me_2S \cdot Sme_2]$ and $[Et_2S \cdot SEt_2]^+$: tandem mass spectrometry metastable and collision-induced dissociation results, 8:605
- Structures of gas-phase $C_7F_7^+$ ions, 4:813
- Studies of pesticides by CID, PSD, and MALDI-MS, 12:590
- Studies of structure and mechanism in acetonitrile chemical ionization tandem mass spectrometry of polyunsaturated fatty acid methyl esters, 10:1253
- Studies on heme binding in myoglobin, hemoglobin, and cytochrome c by ion spray mass spectrometry, 4:631
- Studies on phosphatidylglycerol with triple quadrupole tandem mass spectrometry with electrospray ionization, 12:1036
- Study of adduct ions of meso-phenyl-substituted tetrabenzoporphyrins by fast-atom bombardment-mass spectrometry, 8:62
- Study of diketone/metal ion complexes by ESI-MS, influence of keto-enol tautomerism and chelation, 10:402
- Study of fast and metastable dissociations of adenine-thymine binary-base oligonucleotides by using positive-ion MALDI-TOF-MS, 13:1052
- Study of kynurenine fragmentation, ESI-MS/MS in, 12:786
- Study of light-induced proton transfer from gas phase (radical) cations to reference bases, bracketing of proton transfer from excited ions and associated reaction kinetics, 12:359
- Study of metal chelation of dinucleotide analogs in the gas

- phase by fast-atom bombardment mass spectrometry, 7:42
- Study of noncovalent enzyme-inhibitor complexes and metal binding stoichiometry of matrilysin by electrospray ionization mass spectrometry, 6:1105
- Study of Src SH₂ domain protein-phosphopeptide binding interactions by electrospray ionization mass spectrometry, 8:234
- Study of the solution and gas-phase chemistry of Mn(III) and Fe(III) tetraarylporphyrin complexes by fast-atom bombardment mass spectrometry, synthesis and characterization of molecular complexes, 4:255
- Study of the thermal denaturation of ribonuclease S by electrospray ionization mass spectrometry, 5:614
- Sub-femtomole, levels of, peptide, by tandem magnetic sector/reflectron time-of-flight mass spectrometry and matrix-assisted laser desorption ionization, 2:91
- Sub-miniature ExB sector-field mass spectrometer, 12:619
- Sub-parts-per billion determination of nitro-substituted polynuclear aromatic hydrocarbons in airborne particulate matter and soil by electron capture-tandem mass spectrometry, 7:1255
- Substance P
effect of radical trap moieties on ECD spectra, 17:1428
MALDI-TOFMS detection of molecules in, 12:1055
photodissociation of, 10:1038
- Substantia nigra, neuromelanin from, GC-MS analysis of, 15:920
- Substituents, heavy-atom, on matrices used for matrix-assisted laser desorption-ionization mass spectrometry, 5:800
- Substituted cyclopentadienylmanganese cations in the gas phase, metastable ion study of, 4:216
- Substrates for adsorption/desorption in ²⁵²Cf plasma adsorption/desorption mass spectrometry, small molecules for, 2:29
- Substructure identification rules
automatically generated, evaluation of, from tandem mass spectra, 3:169
generation of, using feature-combinations from tandem mass spectra, 3:159
- Succinimide-mediated mechanism, for collision-activated fragmentation of proferrioxamines, 6:836
- Sudden sampling introduction mode, cyclic, in MIMS, 15:823
- Sugar-sugar complexes, noncovalent, infrared atmospheric pressure MALDI of, 14:1158
- Sulfadiazine metabolites, structural characterization of, by hydrogen/deuterium exchange and MS/MS, 16:1687
- Sulfate position, in glycosaminoglycan-derived disaccharides, negative fast-atom bombardment tandem mass spectrometry to determine, 3:797
- Sulfatides
quadrupole ion trap MS with ESI of, 15:536
in situ characterization of glycerophospholipids and, brain, using MALDI-MS/MS, 18:17
- Sulfation, phosphorylation distinguished from, glycoproteins and carbohydrates, ion-pairing and MS, 17:1282
- Sulfation positions, in chondroitin sulfate oligosaccharides, MS/MS of, 14:1270
- Sulfhydryl bonds, disulfide bonds and, counting using mercurial ions, peptides and proteins, 19:1108
- Sulfhydryl group, in protonated cysteine fragmentation, 9:1275
- Sulfonamide cross-linked peptides, CID and MALDI PSD of, 13:709
- Sulfonamide determination, nanoscale separations combined with electrospray ionization mass spectrometry, 3:139
- Sulfonated peptides, MALDI-TOF/TOF-MS of, 15:1838
- Sulfonates, selective arginine residue titration by, 12:810
- Sulfones and sulfoxides, Li⁺ and H⁺ affinities of, ESI-FTICR of, 8:262
- Sulfopeptides
fragmentation in ECD and ETD, 18:1617
phosphopeptides and, atmospheric pressure thermal dissociation, 19:1898
- Sulfur, Chinese and American drywall materials analyzed by DART with TOF MS, 20:2082
- Sulfur and nitrogen heterocycles in petroleum fractions, selective chemical ionization of, by ion trap mass spectrometry, 4:322
- Sulfur compounds, CH₃OCH₂ reactions with, 13:1028
- Sulfur-containing compounds, selective detection of, by gas chromatography/chemical reaction interface mass spectrometry, 2:250
- Sulfur-containing distonic radical anion, CH₂-S-CH-CN, formation and characterization of, in gas phase, 6:71
- Sulfur mustards, hydrolysis products of, analyzed by APCI-MS, 18:1364
- Sulfur trifluoride cation affinities of pyridines determined by the kinetic method: stereoelectronic effects in the gas phase, 8:68
- Superacid medium, carbocationic rearrangement of pivaloyl cation and protonated pivalaldehyde in, 15:959
- Supercharging, origin from aqueous solutions during ESI, 20:1933
- Supercritical fluid chromatography-mass spectrometry, ion-neutral reactions in the ion source, 18:2173
- Supercritical fluid chromatography-mass spectrometry (SFCMS), long column, polymer characterization with, 13:1195
- Superelastic collisions of electrons with excited MN⁺, 1:192
- Superoxide dismutase, polyethylene glycol-conjugated, fingerprinting proteins coupled with polymers by mass spectrometry and, 6:478
- Supersonic molecular beams
alkanes in, electron impact mass spectrometry of, 6:120
cluster chemical ionization and deuterium exchange mass spectrometry in, 7:550
laser desorption fast GC-MS in, 9:628
LC/MS with, 11:587
thermally labile steroids, carbamates, and drugs in, fast, very fast, and ultra-fast gas chromatography-mass spectrometry of, 7:737
- Suppression of the lower charge state ions in the external accumulation radio frequency multipole with a reduced trapping DC potential, 14:1229
- Supramolecular coordination complexes, ion spray-tandem mass spectrometry of, 5:748
- SUPREX
accuracy of MALDI-MS and, protein unfolding free energy determination, 17:1535
efficiency of MS-based screening assay for detection of ligand binding, 19:1303
- Surface-assisted laser desorption/ionization mass spectrometry
estrogen determination using, 19:1343
silver-coated nanoparticles as a matrix in analysis of aminoglycosides, 20:1925
use of titania nanotube arrays, 19:1014
using two-layered silicon-coated nanostructures, 20:1723
- Surface-assisted laser desorption/ionization-mass spectrometry, infrared, of protein and peptide frozen aqueous solutions, 9:912
- Surface-assisted laser desorption/ionization time-of-flight mass spectrometry, signal enhancement by oxidized GCB nanoparticles, 20:1207
- Surface-assisted reduction of aniline oligomers, N-phenyl-1,4-phenylenediimine and thionine in APCI and APPI, 13:109

- Surface-bound proteins, digestion of, 14:203
- Surface effects, solvents and, effects on analyte ionization, DIOS MS, 19:8
- Surface enhanced laser desorption/ionization-TOFMS, Protein-Chip[®], thymosin β -4 identification by, 14:760
- Surface-induced dissociation
- comparison with CID for large system structural studies, 19:903
 - fragmentation of oxidized peptides analyzed by RRKM modeling, MD and, 20:1579
 - FT-ICR/MS, protein identification via, 17:700
 - high-efficiency, on a rhenium oxide surface, 4:869
 - in ICR-MS/MS, 11:1107
 - N*-protonated diglycine, collision energy effect on, 14:1402
 - peptide sequencing using, 14:1387
 - small molecules, peptides and non-covalent protein complexes, 19:190
- Surface-induced dissociation from a liquid surface, 4:769
- Surface-induced dissociation in tandem quadrupole mass spectrometers, comparison of three designs, 3:27
- Surface-induced dissociation of molecular ions in a quadrupole ion trap mass spectrometer, 2:487
- Surface-induced dissociation of multiply protonated peptides, 3:859
- Surface-induced dissociation of peptide ions: kinetics and dynamics, 14:1340
- Surface-induced dissociation of peptide ions in Fourier-transform mass spectrometry, 1:413
- Surface-induced dissociation of singly and multiply protonated polypropylenamine dendrimers, 10:414
- Surfaces
- dissociation induced by
 - analyzed using SPALDI and LDI MS, peptides, 20:957
 - protonated octaglycine, 20:939 - fabrication by reactive landing of gas phase ions, metal oxides, 20:915
 - peptide ions, soft landed surface effects on charge reduction and desorption kinetics, 20:901
 - in situ SIMS analysis and reactions after SL of ions, IT MS, 20:949
 - surface-induced dissociation and chemical reactions on stainless steel, carbon and diamonds, 20:927
- Surfactant, bacterial cell proteins treated with, MALDI-MS of, 16:1422
- Surfactants
- ethoxylated, fragmentation by AP-MALDI-QIT, 17:453
 - nonionic, collision-induced dissociation mass spectrometry of, following direct supercritical fluid injection, 3:150
 - selectively polydeuterated, isotopic purity in, charge-remote fragmentation in a hybrid (BEqQ) mass spectrometer to determine, 1:85
- Survivor ions
- delayed dissociation spectra of, from high-energy collisional activation, 4:845
 - energy spectra of, angle-resolved translational, high-energy collisional activation studied via, 4:835
- Suspended trapping gas chromatography/Fourier transform mass spectrometry for analysis of complex organic mixtures, 1:431
- Sustained off-resonance collision-induced dissociation, peptide fragment ion structures, analyzed by SORI-HDX in FTICR instruments, 19:1771
- Sustained off-resonance irradiation, capillary electrophoresis Fourier transform ion cyclotron resonance mass spectrometry with, for the characterization of protein and peptide mixtures, 5:894
- Sustained off resonance irradiation, CID induced by, precalculating parameters for, 13:318
- Sustained off-resonance irradiation collision-activated dissociation
- direct dynamic combinatorial library screening using FTICR MS and, 17:1074
 - structural analysis of sulfated oligosaccharides by positive mode nano-ESI-FTICR-MS and MS/MS by, 17:96
- Sustained off-resonance irradiation excitation
- CID, study of isomeric diglycosyl flavonoids, ESI FTICR MS, 18:2127
 - collisional and radiative processes, 18:2119
- Sustained off-resonance irradiation for collision-activated dissociation, in FTMS, 10:15
- Svec, Harry J., in memoriam, 18:970
- SVOCs. *See* Semivolatile organic compounds
- SWIFT. *See* Stored waveform inverse Fourier transform excitation
- Sympathetic cooling of trapped negative ions by self-cooled electrons in FTICR-MS, 8:793
- Synephrine, molecular ions of, CH_3NH_3^+ and $(\text{CH}_3)_2\text{NH}_2^+$ formation from, the role of distonic ions in, 1:104
- Synthesis of charged phenyl radicals and biradicals by laser photolysis in a FTICR-MS, 9:1135
- Synthetic chemistry, automated high resolution MS for, 10:546
- Synthetic polymers
- giant organic molecules and, improvements in analysis by solvent-free MALDI-MS, 17:661
 - nonpolar matrices in analysis of, MALDI-TOFMS in, 11:731
- α -Synuclein, calmodulin and, metal interactions monitored by SNAPP-MS, 19:1668
- α -Synuclein, monomeric structures and pH-dependent dimer formation in, 15:1435
- Systematic errors originating from Coulomb mutual interaction and image charge in Fourier transform ion cyclotron resonance precise mass difference measurements, analysis and estimation of, 4:855
- T
- t*-Boc group, elucidation using hydrogen/deuterium exchange HPLC/ESI/MS, 16:553
- Tailing signal, on the low-energy side of the main beam in mass-analyzed ion kinetic energy spectra, 3:644
- Tailing signal on the low energy side of the main beam in mass-analyzed ion kinetic energy spectra, origin of, 3:644
- Tandem and electrospray mass spectrometry, identification of components in waste streams by, 4:604
- Tandem capillary liquid chromatography-mass spectrometry, low flow high-performance liquid chromatography solvent delivery system designed for, 6:571
- Tandem double focusing magnetic-orthogonal acceleration time-of-flight mass spectrometer, matrix-assisted laser desorption ionization employing, peptide sequence determination by, 7:1
- Tandem four sector and reflector matrix-assisted laser desorption-ionization time-of-flight mass spectrometer, peptide fragmentation obtained from, 6:822
- Tandem Fourier-transform mass spectrometry with Californium-252 plasma desorption, 1:427
- Tandem-ion spray mass spectrometry, of supramolecular coordination complexes, 5:748
- Tandem magnetic sector/reflector time-of-flight mass spectrometry, detection of femtomole and sub-femtomole levels of peptides by, 2:91
- Tandem mass spectra
- assigning peptides to, SILVER in, 15:910
 - feature-combinations from, generation of substructure identification rules using, 3:159

- high energy collision induced, of polycyclic aromatic hydrocarbon diolepoxide adducted peptides, 5:649
- high energy collision-induced dissociation, peptide sequencing from, pattern-based algorithm for, 3:326
- substructure identification rules from, automatically generated, 3:169
- time dependence of, fast atom bombardment-induced condensation of glycerol with ammonium surfactants, 5:92
- Tandem mass spectra of peptide ions in the quadrupole ion trap, effects of heavy gases on, 7:1194
- Tandem mass spectral analysis, fragmentation of protonated O,O-diethyl O-arylphosphorothionates in, 7:189
- Tandem mass spectral data of peptides, an approach to correlate with amino acid sequences of a protein database, 5:976
- Tandem mass spectrometer
- four sector
 - interface region of, enhancement of ion transmission at low collision energies via, 5:460
 - photon-induced dissociation with, 1:107
 - quadrupole, surface-induced dissociation in, comparison of three designs, 3:27
 - quadrupole-acceleration-deceleration, novel, for neutralization-reionization studies, 3:493
 - for the study of molecular dissociations, 3:487
 - time-of-flight, high efficiency photo-induced dissociation of precursor ions in, 4:751
- Tandem mass spectrometric characterization of a specific cysteine acid residue in oxidized human apoprotein B-100, 6:242
- Tandem mass spectrometric investigation of (C₂H₅)X⁺ ions (X=Cl, Br, I), involvement of classical and nonclassical ethyl structures therein, 5:814
- Tandem mass spectrometric screening and characterization, hybrid, of the glutathione conjugate of 2-furamide, multiple scan modes in, 2:55
- Tandem mass spectrometric structural characterization, negative ion electrospray, of leukotriene B₄ (LTB₄) and LTB₄-derived metabolites, 7:129
- Tandem mass spectrometry. *See also* Capillary separation-tandem mass spectrometry; Electrospray ionization-MS/MS; Fast-atom bombardment tandem mass spectrometry; Liquid chromatography-tandem mass spectrometry
- A. thaliana* phosphatidylglycerol analysis by ESI QIT and, 18:783
 - ADP-ribosylated Kemptide, 20:477
 - analysis of long-chain fatty acyl coenzyme A thioesters by, 5:416
 - analysis of peptides, optimization of capillary zone electrophoresis/electrospray parameters for, 3:289
 - analysis of the ions formed by fast atom bombardment of digoxin and related cardiac glycosides, mechanistic considerations of the protonation and fragmentation of highly functionalized molecules in fast atom bombardment, 1:455
 - arsenic-glutathione complex analysis by HPLC/ES MS/MS, 19:1559
 - binding of xylobiose inhibitors to β-1,4 exoglucanase from *C. fimi*, 18:64
 - of C-phenyl-N-tert-butyl nitron spin adducts from in vitro rat liver microsomal metabolism of bromotrichloromethane and carbon, 6:847
 - of calcium binding sites in peptides, 9:1020
 - of carcinogen-peptide adducts, 9:202
 - characterization bovine insulin glycation sites, 20:1319
 - characterization of cysteine oxidation products in PTP1B active site, 20:1540
 - charge inversion, isomer differentiation by, an investigation into the structure of the ionic products from an Sn (AN-RORC) reaction, 3:467
 - chemical ionization with, internal energy deposition in, 14:102
 - collision-induced dissociation, high- and low-energy, in the analysis of glycoalkaloids and their aglycons, 7:173
 - collisional activation of a series of homoconjugated octadecadienoic acids with, 1:183
 - column switching LC/APCI, urinary etheno-DNA adduct quantification by, 17:605
 - for comparative proteomics, trypsin-catalyzed ¹⁶O-to-¹⁸O exchange in, 14:704
 - and conventional mass spectrometry, comparison of, using electron capture negative ionization of chemically oxidized dexamethasone in human plasma, 1:341
 - crosslinked Lys-Lys peptide influence of cross-linker identity on gas-phase dissociation, 17:395
 - Curie-point pyrolysis, an investigation of dipeptides containing polar and nonpolar side groups by, 3:750
 - cysteine sulfenic acid in MMP-7, 18:1544
 - desorption chemical ionization, of polyprenyl and dolichyl phosphates, 7:958
 - for determination of sulfation positions and uronic acid epimerization in chondroitin sulfate oligosaccharides, 14:1270
 - of diastereomeric cobalt-glucosyl-glucose disaccharide complexes, differentiation of stereochemistry of glycosidic bond configuration, 7:953
 - differentiation of isomeric C8- and N²-deoxyguanosine adducts of 2-acetylaminofluorene by, 5:58
 - differentiation of 2'-O- and 3'-O-methylated ribonucleoside, 17:1096
 - distinguishes hyaluronic acid from N-acetylheparosan, 19:82
 - of doubly protonated angiotensin II formed by electrospray ionization, charge-separation reactions in, experimental considerations and structural implications, 7:30
 - effect of histidine oxidation on peptide ion dissociation patterns, 18:553
 - electron capture, sub-parts-per-billion determination of nitro-substituted polynuclear aromatic hydrocarbons in airborne particulate matter and soil by, 7:1255
 - electrospray, structure analysis of, phenylglycidyl ether adducts of 29-deoxycytidine and 29-deoxyadenosine, 7:682
 - electrospray ionization, structural determination of picomole amounts of phospholipids via, 6:1202
 - ESI, fragmentation pathways of anabolic steroids, 17:477
 - ESI MS and, investigation of enantioselective multivalency by, 19:1634
 - estimation of mutagenic/carcinogenic potential of environmental contaminants by, 1:110
 - ETD-based, protein ubiquitination analyzed by CID or fragmentation by, 20:1652
 - F. tularensis novicida* lipid A, environmentally-regulated remodeling, 18:1080
 - and fast atom bombardment, of macrolide antibiotics, 5:151
 - fast atom bombardment
 - employing ion-molecule reactions for the differentiation of phospholipid classes, 2:470
 - isomer differentiation in 7,12-dimethylbenz[a]anthracene-pyridine adducts by, 2:256
 - negative ion, to determine sulfate and linkage position in glycosaminoglycan-derived disaccharides, 3:797 - fast-atom bombardment
 - analysis of phosphatidylcholine isolated from heart mitochondrial fractions by, evidence of incorporation of monohydroxylated fatty acyl moieties, 7:50
 - determination of the anomeric configuration of glycosyl es-

- ters of nucleoside pyrophosphates and polyisoprenyl phosphates by, 7:541
 determination of the anomeric configuration of glycosyl esters of nucleoside pyrophosphates by, 6:516
 of [¹³C]arachidonic acid labeled phospholipid molecular species, 5:144
 fast-atom bombardment by, liquid secondary ion mass spectrometry, beam-induced reactions occurring in, 4:312
 fast atom bombardment combined with, determination of structure and properties of modified chlorophylls using, 1:72
 in fragmentation and assignment of crosslinked peptides, 14:834
 fundamentals of, dynamics study of simple C-C bond cleavage in collision-activated dissociation of polyatomic ions at low energy, 1:6
 gas-phase labeling combined with, ion-dipole complex formation from deprotonated phenol fatty acid esters evidenced by, 4:343
 guided ion beam, mechanism of decomposition and, protonated asparagine, 20:852
 high-energy collision-induced dissociation in, low-mass ions produced from peptides by, 4:882
 high-resolution
 of large molecules, Fourier-transform electrospray instrumentation for, 4:557
 thiaminase I (42 kDa) heterogeneity and sequence refinement, active site location and, 6:981
 HIV-1 packaging signal mapping noncovalent ligand binding to stemloop domains, 17:1401
 hybrid mass spectrometers for, 19:161
 investigations of gas-phase lithium-peptide adducts, 1:473
 impulse-driven droplet deposition system interface combining LC with MALDI MS and, 17:325
 ion mobility, variance of a- and b-type peptide fragment ions, 19:609
 ion trap
 APCI of fluorinated phenols in, 10:1157
 applied to small multiply charged oligonucleotides with a modified base, 5:740
 in detection of pharmaceutical compounds in tissue, 10:1315
 isomeric dibenzopyrene-adenine adduct determination by, 8:977
 of meso-phenyl-substituted tetrabenzoporphyrin adduct ions, 8:62
 of model peptides modified with trans-2-hexenal, product of lipid peroxidation, 10:600
 multistep, for sequencing cyclic peptides, 10:732
 polycyclic aromatic hydrocarbon analysis by, 8:183
 protein modification by, 8
 of proteins, 8:483
 quantitative analysis of crude plasma extracts by, 3:372
 selected ejection chemical ionization, application of non-resonance excitation to, 7:668
 of synthetic polymers, 8:76
 α_3 ions, observation frequency, 19:1764
 ions in ILTs, 20:1714
 of labeled oligonucleotides, 9:660
 LC-ESI, identification of mammalian cell lines using MALDI-TOF and, 17:490
 leucine enkephalin-alka(e)nal-glycerophosphatidylcholine adduct identification, 17:657
 linked-scan, production characterization by, trimethylsilyl group migration in the mass spectra of trimethylsilyl ethers of cholesterol oxidation products, 4:327
 of lithium-attachment ions from polyglycols, 3:225
 loss of CO and ammonia, protonated AGG, 18:1291
 low- and high-energy collision-induced dissociation, in the identification of an unusual amino acid in a semisynthetic polypeptide, 7:1034
 low-energy
 afterglow-guided ion beam in, 8:688
 of alkali-cationized polysaccharides, in quadrupole ion trap, 8:987
 dinucleotide phosphodiester, 8:1257
 with FTICR, of ionized ethylated acetone and deuterated analogs, 8:327
 of HT2a antagonist and metabolite in, 8:371
 of ions containing $S\phi 0 \Sigma \beta\omega\delta\sigma$, 8:605
 of the molecular ion derived from fatty acid methyl esters, novel method for analysis of branched-chain fatty acids, 4:223
 low energy and, electrospray ionization of polyhydroxy unsaturated fatty acids, 7:140
 of metal nitrate negative ions produced by ESI, 14:671
 methyl group migration during fragmentation of trimethyllysine-containing peptides, 20:1172
 and molecular orbital calculations, 3-methylisooxazolo- and 2-methylloxazopyridine, heterocyclic isomers, characterization and differentiation of, 6:962
 molecular species analysis of arachidonate containing glycerophosphocholines by, 2:45
 negative ion, of leukotriene E₄ and LTE₄ metabolites, identification of LTE₄ in human urine, 2:314
 3-O-sulfated disaccharide heparin isomers, 20:652
 ornithine and glutamine lipids from *R. sphaeroides* cell membranes, 20:198
 a_n and *a_n** peptide fragments, isotope labeling, density functional theory calculations and, 19:1788
 peptide isocyanate-adducts analyzed by, 20:1567
 peptides with N-terminal dimethylation and imine/methylol adduction, 17:859
 of peptides with N-terminal glutamine studies on a prion protein peptide, 1:258
 pharmaceuticals in surface waters, matrix effects in HPLC-MS/MS and UPLC-MS/PM analyses, 19:713
 of poly(ethylene glycol) lithium-attachment ions, 5:1072
 poly(methylacrylic acid) oligomers produced by ESI, 20:25
 product ions in, 9:341
 protein-ligand complex detection by ESI-precursor ion scan, 20:227
 protonated polyproline peptides, 18:2198
 protonated O, O-dimethyl O-aryl phosphorothionates in, fragmentation of, 6:488
 2-pyrimidinylxy-N-arylbenzyl amines, chemical transformations predicted, 17:253
 of quadrupole ion trap peptides, 9:341
 reactivity of collisionally activated dichlorocarbene dications studied by, 6:1019
 secondary ion, imaging large insulating samples by, charge compensation for, 5:37
 in separation and identification of oxygen-centered radical DMPO adducts, 14:862
 of small, multiply charged oligonucleotides, 3:60
 software for peak assignment, synthetic polymers, 18:1324
 sphingolipid and phospholipids, DESI/MS and, 19:531
 for the structural determination of backbone-modified peptides, 1:174
 studies of acetone and acetone/water cluster ions, 2:113
 study of the gas phase RSH₆H₅Cr⁺L (R=H₆H₅; I=arene) and H₆H₅SSH₆H₅Cr⁺ ions, 5:928
 of structure and fragmentation mechanisms of isomeric T-rich oligodeoxynucleotides, 9:683

- and theoretical calculations, mechanism of gas-phase aldose-ketose isomerization, 4:943
- thermal degradation, structural studies on alkylisocyanate polymers by, 2:130
- top-down, low-mass proteomes, 20:2183
- tryptophan O₂ (¹Δ_g)-derived oxidation products, 20:188
- UGT1A1 regioselectivity towards flavonoids, metal complexation and, 19:246
- using nanoLC-FTICR-MS, high-throughput comparative proteomics, 18:1332
- Tandem mass spectrometry-electron ionization, of glycosphingolipids, identification of compound-specific sequence ions in the collision-induced dissociation spectra of the immonium ions of two isomeric hexaglycosylceramides, 3:353
- Tandem mass spectrometry-electrospray ionization
- analysis of peptides derived by enzymatic digestion of oxidized globin subunits, an improved method to determine amino acid substitution in the hemoglobin core, 7:1040
 - positive and negative mode, identification of phosphorylation sites in phosphopeptides by, 7:243
- Tandem mass spectrometry electrospray ionization (MS/MSESI), sphingomyelin structural determination by, 11:437
- Tandem mass spectrometry-electrospray liquid chromatography, quantitation of SR 27417 in human plasma using, study of ion suppression, 7:1099
- Tandem mass spectrometry/liquid chromatography, analysis of peptide mixtures, data-controlled automation of, 7:532
- Tandem mass spectrometry measure-merits, precision and accuracy in, kinetics-based protocol for instrument-independent measurements of collision-activated dissociation in RF-only quadrupoles, 1:272
- Tandem mass spectrometry measurements, precision and accuracy in, kinetics-based protocol for instrument-independent measurements of collision-activated dissociation in RF-only quadrupoles, 1:274
- Tandem mass spectrometry-microbore high-performance liquid chromatography, utilizing pneumatically assisted electrospray ionization, quantitative determination of a nonpeptide antithrombotic in dog plasma by, 7:1238
- Tandem mass spectrometry (MS/MS). *See also* Electrospray ionization tandem mass spectrometry; Electrospray tandem mass spectrometry.
- ceramide characterization by, 13:558
 - in combined ion mobility and MS metallothionein investigations, 16:1787
 - complex, mass uncertainty and resolving power in, 16:183
 - of complex polyol mixtures, 12:55
 - in confirmation/quantitation of cocaine and benzoylecgonine in urine, 11:257
 - construction of quadrupole-FTICR-MS for, 15:1099
 - of cytochrome *c*-cytochrome *b*₅ protein complex, 13:59
 - in DNA cross-link determination, 15:1565
 - DNA sequencing by, 15:510
 - flavonoid glycosylation site determination by, 15:1287
 - of gas phase reactivity of isomeric arylglycoside toward amines, 15:244
 - of growth hormone secretagogue: amide bond cleavage and resultant gas phase rearrangement, 13:371
 - guided ion beam, to determine thermochemistry, 13:419
 - of heparin-like glycosaminoglycans, competing fragmentation processes in, 15:1534
 - hydrogen/deuterium exchange with, in sulfadiazine metabolite structural characterization, 16:1687
 - to identify ozone oxidation products of amino acids and small peptides, 11:526
 - improved mass accuracy for, 16:463
 - ion trap data-dependent, in proteomics, 15:150
 - of ionic and neutral P(OH)₂+/- in gas phase, 13:250
 - isobaric monosaccharides distinguished by, 16:1827
 - of lithiated ceramides, 13:680
 - in locating abasic sites in oligodeoxynucleotides, 13:1418
 - of metal-oligonucleotide complexes, 16:370, 16:799 (erratum)
 - methionine-containing peptide analysis by, 16:1131
 - of methylbenz(is)oxazole isomers, 15:1005
 - of nitro *meso*-tetraphenylporphyrins positional isomers, 12:381
 - peptide identification by, increasing through automatic function switching optimization, 16:1818
 - of phosphatidylserine, ESI with, 16:1510
 - of protein, 13:107
 - proteomic, MS2Grouper in, 16:1250
 - of protonated oligopeptides, amide bond cleavage mechanism in, 15:103
 - quadrupole, of cardiolipin, 16:491
 - screening of, for database search, 11:422
 - in sequencing multiantennary N-glycans, 13:1138
 - of singly and doubly charged ions in negative ion ESI MS, 13:671
 - of spider venom acylpolyamides, 15:1636
 - triple quadrupole
 - in molecular characterization of thiols, 15:288
 - phosphatidylglycerol studies with, 12:1036
 - of tris(trimethoxyphenyl)phosphonium acetyl charged derivatives of peptides, 11:107
 - ultrafiltration, of estrogens, 16:271
 - verification of (M + Fe-5H)²⁻ peptide ion composition by, 11:83
- Tandem mobility mass spectrometry, of electrosprayed tetraheptyl ammonium bromide clusters, 16:717
- Tandem quadrupole mass analyzer, 10:1053
- Tandem quadrupole mass spectrometry
- disulfur monoxide-water charged complexes, isotope exchanges, FT-ICR, CI-CAD and, 18:1664
 - of phosphatidylcholines, fragmentation processes in, 14:352
- Tandem quadrupole time-of-flight mass spectrometry, MALDI-formed peptide [M+H] ions, formation of (b_{n-1}+H₂O) by collisional activation, 18:1024
- Tandem reflectron time-of-flight mass spectrometry, ionization/matrix-assisted laser desorption of fullerenes, 7:590
- Tandem time-of-flight experiment, for low energy collision studies, 11:160
- Tandem time-of-flight mass spectrometer, for photodissociation of biopolymer ions generated by MALDI, 15:1248
- Tanshinone IIA, effect on warfarin-HSA interactions, ESI MS, 19:1568
- Tantalum oxide ions, collision-induced dissociation rates of, measurement in ion trap, 11:1072
- Target gas, effects of, in collision-induced dissociation using a double quadrupole mass spectrometer and radiofrequency glow discharge, 5:845
- Targeted use of exoglycosidase digestion for structural elucidation of neutral O-linked oligosaccharides, 12:877
- Tate Gallery, pigments and dyes, MALDI TOF MS identification, 18:2001
- Tau-related peptides, quantification using GLaD with LC-IT-TOF MS, 18:201
- Tauroconjugated bile acid anions, fast-atom bombardment mass spectrometry and low energy collision-induced tandem mass spectrometry of, 6:588
- Tautomerization, effect of, on the fast-atom bombardment tandem mass spectra of azo dyes, 5:407
- Taxol, mass spectrometry of, 3:672
- TCP. *See meso*-Trisphenylcorrole
- Tears, analysis of glucose concentration by ESI MS, 18:332

- Technical chlordane, highly chlorinated dimethanofluorenes in, 1:99
- Temperature
in alkali chloride cluster ion dissociation, 13:1388
low, ECD at, 15:1869
rate constants of Ar⁺ reaction with CO₂ and SO₂ as function of, 10:815
vaporizer, ion formation of N-methyl carbamate pesticides in thermospray mass spectrometry, 5:913
- Temperature-constrained cascade correlational networks, in bacterial classification, 13:10
- Temperature-dependent H/D exchange, of compact and elongated cytochrome c ions in the gas phase, 13:506
- Temperature-programmed pyrolysis/metastable atom bombardment ionization mass spectrometry (TPPy/MAB-MS), for identifying additives in polymers, 15:1315
- Temperature-time effects, in situ, in Meta-S-SIMS, 18:382
- Temporal encoding in time-of-flight mass spectrometry, beam deflection for, 1:440
- Tenax collection with jet-separator enrichment and ion trap mass spectrometric analysis, air analysis using, 7:1172
- Ter-body intermediates in gas phase, ionized reaction of enols with tert-butanol, 12:938
- Termolecular ion-molecule reactions
in Titan/Es atmosphere: search made at up to 1 micron in pure hydrocarbons, 14:900
in Titan/Es atmosphere: structure of the association adducts of HCNH with C₂H₂ and C₂H₄, 12:557
- Terpenoids, plasma determination of flavonoids and, from *G. biloba*, efficiency of ESI, 19:445
- Terrestrial molybdenites, isotope ratios of molybdenum in, measurement of, high accuracy, 3:10
- tert-Butanol, enol ionized reaction with, 12:938
- Tests, for identification of compound from mass spectrum, 14:262, 14:264
- Tetanus toxin C, fragment of, NeuAc oligomers associated with, ESI-MS, 17:967
- Tetraalkylammonium salt molecular ions sputtered from a gelatin matrix, formation and emission of, 4:930
- Tetraarylporphyrin complexes, Mn(III) and Fe(III), solution and gas phase chemistry of
by fast-atom bombardment mass spectrometry. 1, Generation of molecular signals, 4:249
by fast-atom bombardment mass spectrometry. 2, Synthesis and characterization of molecular complexes, 4:255
- Tetrabenzoporphyrin, of meso-phenyl-substituted, FAB-MS of adduct ions of, 8:62
- Tetrachloride anions, group 15, thermochemistry of, 13:469
- Tetracycline, degradation by ozone in aqueous medium, APCI/MS, 18:679
- Tetracycline antibiotics, ESI identification of, in quadrupole ion trap, 9:1089
- Tetracyclines, MS characterization of, 13:543
- Tetraether lipids, archaeal, analysis by LC-MS/MS, 20:51
- Tetraglycine, protonated, study of formation of a₃⁺ ions, 17:1654
- Tetrahedral supramolecular Ti₄L₄ cluster, ESI-ion trap MS of, 9:1099
- Tetraheptyl ammonium bromide clusters, electrosprayed, tandem mobility MS of, 16:717
- meso-Tetrakisphenylporphorin, TCP and, observations using ESI MS, 17:1306
- Tetrameric ions, of noncovalent streptavidin complexes formed by electrospray ionization, 6:459
- Tetraphenylporphyrin, competitive ionization of, in a laser generated metal ion plasma, 4:159
- Tetrasaccharides, GAG, EDD of, 18:234
- Tetronasin, ESI-CID-MS/MS fragmentation studies of, 15:325
- Theoretical calculations
of glycine and alanine gas-phase acidities, 10:318
of relative ion yields for glow discharge mass spectrometry, 3:79
- Theoretical calculations and tandem mass spectrometry, mechanism of gas-phase aldose-ketose isomerization, 4:943
- Theoretical considerations, extended, for mass resolution in the resonance ejection mode of quadrupole ion trap mass spectrometry, 5:676
- Thermal conditions, gas-phase proton transfer reactions involving multiply charged cytochrome c ions and water under, 3:624
- Thermal decomposition, contributions to the mass spectra of chlorinated phenoxyacid herbicides obtained by particle beam liquid chromatography/mass spectrometry, 3:823
- Thermal degradation processes, primary, of peptides, using the mass spectrometric technique K⁺IDS, K⁺ ionization of desorbed species, 5:564
- Thermal degradation tandem mass spectrometry, structural studies on alkylisocyanate polymers by, 2:130
- Thermal denaturation
of cytochrome c, analyzed by ESI-MS, 20:819
for single-stranded PCR amplicon production, 13:232
- Thermal denaturation of ribonuclease S, by electrospray ionization mass spectrometry, 5:614
- Thermal-desorption, laser-induced/Fourier-transform mass spectrometry for surface analysis, deflection of laser-produced ions in, 5:583
- Thermal dissociation of protein homodimer ecotin, in gas phase, 13:1432
- Thermal energy and its distribution, in peptides, calculation of, 10:323
- Thermal energy reactions of mass-selected ions, high pressure drift reaction cell to study, a hybrid double focusing mass spectrometer, 1:197
- Thermal evolved gas analyzer, Phoenix Mars Mission, 19:1377
- Thermal ionization cavity, for MS, 10:1008
- Thermally labile steroids, carbamates, and drugs in supersonic molecular beams, fast, very fast and ultra-fast gas chromatography-mass spectrometry of, 7:737
- Thermochemistry
determination by guided ion beam MS/MS, 13:419
of gaseous ethylsilanes and their radical cations, 9:597
of group 15 tetrachloride anions, 13:469
ion, 13:417
of polyiodide ions, 8:688
- Thermococcus kodakaraensis* (KOD) DNA polymerase, for PCR mass spectrometry based analyses, 14:601
- Thermodynamic properties, thiomethyl anion, 1:366
- Thermodynamics, mechanism of decomposition and, protonated asparagine, GIB-MS/MS, 20:852
- Thermolysis, of aromatic diazoamino compounds, 9:234
- Thermometer ions, strategy based on, ESI MS reproducibility using, 20:398
- Thermospray interfaces with carbamate pesticides, comparative study of different, influence of the ion source geometry, 6:656
- Thermospray mass spectrometry, ion formation of N-methyl carbamate pesticides in, effects of additives to the liquid chromatographic eluent and of the vaporizer temperature, 5:913
- Thermospray probe insert designs, that allow clogged probes to be rapidly replaced, 4:182
- Thermospray reagent plasma from ammonium acetate, controversial ions (m/z 59, 60, 77, and 119), on the origin of some, 5:186

- Thermus thermophilus*, ribosome topography analyzed by proteolysis and MALDI-MS, 18:1304
- Thexyl alcohol, photoionization of, 15:989
- Thiaminase I (42kDa) heterogeneity, sequence refinement, and active site location from high-resolution tandem mass spectrometry, 6:981
- Thiazines, cyclohexene-fused-2-N-phenyliminoperhydro-3,1-oxazines related, fragmentation of, effects of N-substitution on, 2:125
- Thin layer chromatography matrix-assisted laser desorption/ionization orthogonal time-of-flight mass spectrometry, ganglioside analysis by, 16:1552
- Thioacylium ions, iso(thio)cyanate interactions with, 16:1602
- Thioepoxides, Meerwein reaction of phosphonium ions with, in gas phase, 15:398
- Thioesters
 long-chain fatty acyl, coenzyme A, by negative ion fast-atom bombardment mass spectrometry and tandem mass spectrometry, 5:416
 nitrosation in gas phase, 15:344
- Thioether conjugates of acrolein, protonated, fragmentation of, 8:727
- Thiols
 free and disulfide-bound functions, protein labeling, 19:1
 redox-sensitive, protein disulfide isomerase, identified using ICAT MS, 18:260
- Thiol(s)
 molecular characterization of, ESI and triple quadrupole MS/MS in, 15:288
 nitrosation in gas phase, 15:344
- Thiomethyl anion, formation, reactivity, and thermodynamic properties, 1:366
- Thionine, surface-assisted reduction of, in APCI and APPI, 13:109
- Thiophene, transition-metal mediated heteroatom removal by reactions of $\text{FeL}^+ [\text{L}^+ \text{O}, \text{C}_4\text{H}_6, \text{C}-\text{C}_5\text{H}_6, \text{H}_6\text{H}_6, \text{C}_5\text{H}_4(=\text{CH}_2)]$ with, in the gas phase, 7:938
- Thiouracils, gas-phase interactions between lead(II) ions and, 20:359
- Thiourea, substituted N in, cyclization of, in gas phase and solution, 16:1739
- Thomson, J.J., American experiences, 20:1964
- Three rods ion optic, high focusing and mass filtering capabilities, computer simulations, 18:413
- Threshold collision-induced dissociation, with guided ion beam mass spectrometer, ESI source for, 18:1124
- Threshold dissociation energies, of protonated amine/polyether complexes in quadrupole ion trap, 14:383
- Threshold dissociation energy, of flavonoid transition metal complexes, 16:139
- Thromboxanes, fast atom bombardment and collision-induced dissociation of, some examples of charge remote fragmentation, 1:325
- Thymine glycol residues, quantification of, gas chromatographic-mass spectrometric method for the assessment of oxidative damage to double-stranded DNA by, 4:336
- Thymines, replacing phosphates with, in oligodeoxynucleotides fragmentation, 12:580
- Thymosin β -4, identification by ProteinChip[®] SELDI-TOFMS, 14:760
- Ti⁺, polymerization of ethylene catalyzed by, in the gas phase, 3:464
- Ti4L4 cluster, ESI-ion trap MS of, 9:1099
- Time
 FAB-MS dependence on, 8:337
 geometry and energy, ion formation from charged droplets, 4:524
- Time dependence, of mass spectra and tandem mass spectra, Fast atom bombardment-induced condensation of glycerol with ammonium surfactants, 5:92
- Time-dependent desorption, using a model pulsed bombardment ionization method, improved detection limits for fast atom bombardment mass spectrometry, 3:637
- Time-lag focusing, and cation attachment in the analysis of synthetic polymers by matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry, 8:132
- Time-of-flight, ionization/matrix-assisted laser desorption mass spectrometry of peptides, improved mass accuracy in, 5:955
- Time of flight mass measurement, matrix-assisted laser desorption ionization, accuracy, by using delayed extraction, 7:995
- Time-of-flight mass spectrometer
 an inductively coupled plasma, for elemental analysis, An. Part I, optimization and characteristics, 5:1008
 constant-momentum/energy selector, 18:92
 electrospray, vacuum chamber of, ion-molecule collisions in, effect of, on mass spectra of proteins, 7:342
 linear
 large molecules in, high-resolution mass spectrometry of, 4:2
 postsource pulse focusing conditions and mass scale, 6:889
 linear orthogonal acceleration, detector pulse width and jitter in, resolution limitations from, 5:772
 magnetic-orthogonal acceleration, tandem double focusing, matrix-assisted laser desorption ionization employing, peptide sequence determination by, 7:1
 tandem, high efficiency photo-induced dissociation of precursor ions in, 4:751
 two-stage linear, photodissociation of high molecular weight peptides and proteins in, 6:578
- Time-of-flight mass spectrometry. *See also* Matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry
 analysis of serum protein precipitated with antiserum, 6:854
 of bacterial lipooligosaccharides, 8:645
 Chinese and American drywall sulfur materials analyzed by DART with, 20:2082
 FAIMS/FAIMS coupled to, ion trapping in, 12:1320
 GC with, field-portable high speed spectrometer for, 12:648
 gridless ion acceleration systems for, 4:782
 Hadamard transform, sequence length, sequence frequency, and data acquisition rate effects on, 12:1302
 high resolution ion mobility with, of synthetic failures in combinatorial libraries, 11:352
 human urine screening for cocaine and metabolites by, 20:891
 improved energy compensation for, 5:779
 ion association by, study of V-P-O catalysts, 1:129
 ion trap source for, a segmented ring, cylindrical, 7:1009
 ion trap storage and, for protein digest analysis, 8:1237
 ion mobility cell combined with collision cell and, 17:691
 ionization tandem/matrix-assisted laser desorption of fullerenes, 7:590
 laser desorption, analysis of hydrocarbon dendrimers by, 5:731
 laser desorption electron attachment, to detect involatile compounds, 12:1339
 laser ionization orthogonal, multielemental analysis of biological samples by, 20:1355
 linear, development and ion characterization of, 8:1085
 linear ion trap, axially-resonant excitation, duty cycle enhancement, 17:1669
 MALDI, Braun's lipoprotein and determination of its attachment sites to peptidoglycan by, 6:892
 MALDI-generated initial ion velocity and, 8:209
 MALDI with, for polydiene analysis, 8:1220

- MALDI with
 of complex polyol mixtures, 12:55
 of diastereomeric benzo[ghi]fluoranthene tetrols, 12:376
 of in-source peptide decay, 12:420
 for mass measurement of bromo- and chloroacetic acids, 11:809
 matrix-assisted laser desorption ionization
 and electrospray mass spectrometry, application of, for molecular weight assignment of peptides in complex mixtures
 reflectron, CE/MS with, applications of to myelin basic protein, 9:701
 in review of 10th Sanibel conference on mass spectrometry, 9:849
 space focusing in, 10:1067
 of underivatized and permethylated gangliosides, 3:785
 matrix-assisted laser desorption ionization, and electrospray mass spectrometry, application of, for molecular weight assignment of peptides in complex mixtures, 4:670
 of DNA fragment ions, 10:423
 MALDI sample preparation investigation by, 10:104
 of polysiloxanes, 9:292
 protein analysis by, 9:88, 9:1879
 quadrupole, for structural elucidation of drug metabolites, 10:1305
 miniaturized EI/Q/oa spectrometer for, 12:641
 of nitrated polycyclic aromatic hydrocarbons, 8:630
 novel ion trap with orthogonal injection, 20:1342
 orthogonal
 background reduction in, 8:125
 of noncovalent protein assemblies, 12:329
 orthogonal acceleration, spontaneous and deflected drift-trajectories in, 5:588
 protein modification by, 8:8
 quadrupole
 in determining CAD of substituted indole, 12:1145
 in identifying GlcNAcylation sites of peptides and betacrySTALLIN, 12:1106
 reflectron, detection of femtomole and sub-femtomole levels of peptides by, 2:91
 temporal encoding in, beam deflection for, 1:440
 Time-of-flight mass spectrometry. *See also* Matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry.
 Time-of-flight mass spectrometry (TOF-MS). *See also* Matrix-assisted laser desorption ionization time-of-flight mass spectrometry.
 ion optical elements for, conductive carbon-filled polymeric electrodes as, 13:1170
 orthogonal, of peptide-peptide interactions, 13:167
 of peptides, 13:772, 13:784
 quadrupole, in sequencing multiantennary N-glycans, 13:1138
 of α,w -bis(4-hydroxybutyl)poly(dimethylsiloxane), 13:170
 Time-of-flight-matrix-assisted laser desorption ionization mass spectrometry, liquid chromatography-electrospray ionization, rapid analysis of antibiotic-containing mixtures from fermentation broths by using, 7:1227
 Time-of-flight MS. *See also* Gel permeation chromatography-MALDI-TOFMS; Hadamard transform-TOFMS; quadrupole TOFMS.
 electrospray quadrupole, noise filtering techniques for, 14:766
 incorporation with LC-IMS-CID techniques, 14:1242
 Time of flight-secondary ion mass spectrometry, factors affecting mass scale calibration in, 17:514
 Time-of-flight secondary ion mass spectrometry
 influence of organic layer thickness in, 20:2294
 poly(dimethyl siloxane) polymer MWDs using, 20:1562
 Time-of-flight-secondary ion mass spectrometry (TOF-SIMS), quantitative, of biodegradable polymers, 13:1108
 Time-of-flight secondary-ionization mass spectrometry
 ion intensity in, enhancement of, 7:467
 of matrix-diluted oligo- and polypeptides bombarded with slow and fast projectiles, positive and negative matrix and analyte ion yields, background signals, and sample aging, 11:553
 Time-of-flight tandem mass spectrometry, time-resolved method for PD mechanistic study of protonated peptides, 18:1729
 Time-of-flight time-of-flight mass spectrometry
 compared with 193 nm photodissociation, bradykinin analogues, 17:721
 fragmentation reactions, arginine-containing protonated dipeptides, 18:1959
 Time-of-flight (TOF), tandem, as experiment for low energy collision studies, 11:160
 Time-of-flight (TOF) bioaerosol mass spectrometry (BAMS), sensitivity and mass range of, 16:1866
 Time-of-flight (TOF) mass spectrometer, with spiral ion trajectory, 16:1969
 Time-of-flight (TOF) secondary ion mass spectrometry (SIMS), improvement with bismuth cluster ion source, 16:1608
 Time regime, afterpeak, of pulsed glow discharge plasmas, ion formation processes in, 4:727
 Time-resolved electrospray ionization mass spectrometry, of pH effect on myoglobin unfolding, 11:312
 Time-resolved measurements
 of individual ion cloud signals to investigate space-charge effects in plasma mass spectrometry, 7:362
 with single droplet introduction to investigate space-charge effects in plasma MS, 10:159
 Tingle compounds, fragmentation of, by collisional activation in triple quadrupole mass spectrometer, 15:1462
 TiO₂ surface, light-activated, dexamethasone transformation products on, ion trap MS/MS of, 12:1286
 Ti₆O₄(O₂C₄H₅)₈(OCH₂CH₃)₈, ESMS-TOF of, 15:260
 Tips
 metallized, long-lived, for nanoliter electrospray mass spectrometry, 7:1270
 nonmetallized, for EMS, 10:72
 Tirapamazine and metabolites, MS study of, 14:881
 Titan, atmosphere of, study of ion-molecule reactions relevant to, ICR MS, 17:544
 Titania nanotube arrays, surface-assisted laser desorption/ionization mass spectrometry, 19:1014
 Titanium dioxide, comparison of IMAC surfaces and, in protein phosphorylation studies, 18:1932
 Titanium oxide, characterization of nanoclusters, MALDI-TOF-MS, 18:517
 Titan's atmosphere
 pure hydrocarbons, ion-molecule reactions in, 14:900
 SIFT ion-molecule study of reactions in, 15:1148
 termolecular ion-molecule reactions in, 12:557
 Titration, modeling data from, in protein-ligand interactions, 15:388
 TM-DESI. *See* Transmission mode desorption electrospray ionization
 TM-DSI. *See* Transmission mode desorption electrospray ionization
 TNT, negative-ion formation in, by using the reversal electron attachment detection technique, 3:243
 Tobacco, pyridine alkaloid determination by PFTBA ions/analyte molecule reaction IT-MS, 18:1774
 Tobramycin samples, new compound in, CAD of, 11:200
 TOF. *See* Time-of-flight entries
 TOF-MS. *See* Time-of-flight mass spectrometry

- TOF-SIMS. *See* Time-of-flight-secondary ion mass spectrometry
- TOFMS. *See* Time-of-flight MS; Time-of-flight mass spectrometry
- Tolyl cation, selective detection of, among other $[C_7H_7]^+$ isomers by ion/molecule reaction with dimethyl ether, 2:270
- Top down characterization of secreted proteins from *Mycobacterium tuberculosis* by ECD-MS, 14:253
- Top-down liquid chromatography mass spectrometry, automated, of modified proteins, 16:2027
- Top-down mass spectrometry, in kinetic characterization of multiple protein forms, 16:1052
- Top-down tandem mass spectrometry (MS/MS), qQq-FTICR mass spectrometer for, 16:1985
- Toxicological screening, LC-ES-MS vs HPLC-DAD for, 14:14
- TPEPICO studies, ionized *o*-,*m*-,*p*-difluorobenzene, 20:20
- TPP. *See* *meso*-Tetrakisphenylporphyrin
- Trace analysis, MS in, 9:643
- Trace element analysis, laser ablation ICP-MS in, 15:769
- Trade-offs in miniature quadrupole designs, 12:633
- Trans-activation responsive element region, binding sites of TAR and TAR mutants for peptide ligands, LILBID, 19:1605
- Transacetalization, with gaseous carboxonium and carbosulfonium ions, 12:14
- Transferrin, In^{3+} and Bi^{3+} binding to, indirect detection with ESI-MS, 15:1658
- Transferrins, mass spectrometry of, 14:635
- Transformation of neutral rhenium compounds during electrospray ionization-mass spectrometry, 8:191
- Transient bifunctional hypervalent radicals, hydrogen bonding in, by neutralization-reionization mass spectrometry, 6:1004
- Transient intermediate, dissociation kinetics of, 10:119
- Transient recorder, dynamic range of, increasing the, by using two analog-to-digital converters, 7:107
- Transition metal cluster ions, in the gas phase, reactive and dissociative behavior of, characterization of, 2:212
- Transition metal complexes, gas phase reactions of, 8:1124
- Transition metal ions
and ESI low energy CID-MS/MS of polyglycols, 12:832
reactions with long chain alkanes, 12:367
- Transition-metal ions, gas-phase ion-molecule complexes of, formation and fragmentation of, with organic molecules containing two functional groups, 3:451
- Transition-metal ions in the gas phase, ion-molecule reaction of, study on fragmentation mechanism of the adducts, 2:120
- Transition-metal mediated heteroatom, removal by reactions of FeL^+ [L^+O , C_4H_6 , $c-C_5H_6$, H_6H_6 , $C_5H_4(=CH_2)$] with furan, thiophene and pyrrole in the gas phase, 7:938
- Transition metals
10:1-(2-benzothiazolyl)-3,5-diphenyl formazan complex formation with under laser desorption/ionization, 10:339
atomic, as ion mobility source for guided ion beam experiments, 18:1196
complexation by 3-azidopropionitrile, ESI-MS, 18:453
inorganic compounds, cluster ions of, LA-MS, 19:488
metal complexes with oligodeoxynucleotide charged anions, 19:281
polyether/pyridyl ligand complexes with, in quadrupole ion trap mass spectrometer, 10:126
- Translational energy spectra, angle-resolved, of survivor ions, high-energy collisional activation studied via, 4:835
- Transmission mode desorption electrospray ionization analysis of residues from evaporated solvents, 19:1617
influence of material and mesh characteristics on, 20:584
- Transthyretin variant detection, using immunoprecipitation and MALDI bioreactive probes, clinical application of MS, 11:172
- Trapping and detection of ions, generated in a high magnetic field electrospray ionization Fourier transform ion cyclotron resonance mass spectrometer, 3:615
- Trapping electrode, central, in FTICR-MS, 12:30
- Trapping motion in elongated Fourier-transform mass spectrometry trapped ion cells, direct observation of, 1:351
- Trapping ring electrocells, FT-ICR-MS, improved ion trapping and detection capabilities, 20:755
- Traps, electrically-compensated, designed to 8th order for FT-ICR, 19:1281
- Trastuzumab, heterogeneity profile, ESI-Q-IM-TOF MS, 20:2021
- Traveling-wave ion mobility mass spectrometry, hemoglobin structure probed by, 20:625
- Tri-alanine, oxazolone structure of b_2 fragment ions, IRMPD spectroscopy, 20:334
- Triacylglycerol mixture analysis, high-resolution, using high-temperature gas chromatography/mass spectrometry with a polarizable stationary phase, negative ion chemical ionization, and mass-resolved chromatography, 7:350
- Triacylglycerols
CID MALDI-TOF/TOF, 20:1037
ESI-MS of, 16:1498
fragmentation of conjugate bases of esters derived from, 6:498
MALDI-TOFMS of, 10:983
 O^- and OH^- , chemical ionization of some, 5:553
structural characterization of, by ESI-MS, 10:587
- Triaxial electrospray probe, peptide sequencing using alternative enzymes for on-line proteolysis, 20:1983
- Triaxial probes, for proteolysis, coupled with HDX-MS, 18:208
- 1,2,4-Triazines, fluorinated, protonation and deprotonation and gas phase reactivity, ESI MS, 19:686
- Tribenuron methyl, hydrolytic and photodegradation, HPLC-DAD-MS/MS, 19:1221
- Tributyl phosphate
MS analysis of, zeolites in, 8:718
static secondary ionization mass spectrometry analysis of, on mineral surfaces, effect of Fe(II), 6:165
- Tricationic ion-pairing reagents, evaluation in positive mode ESI-MS, 20:529
- Trideoxynucleotide anions, structures, fragmentation and protonation, 19:987
- Triethylantimony, tris(*tert*-butyl)gallium, trimethylgallium and, thermally-induced decomposition of, 19:947
- Trifluoroacetic and other strong acid modifiers, gradient reverse-phase high-performance liquid chromatography-electrospray ionization mass spectrometry analysis with, by postcolumn addition of propionic acid and isopropanol, signal enhancement of, 6:1221
- Triglycine, tautomerization and dissociation of radical cations, 20:996
- Triiodothyronine, ESI-MS/MS detection and quantification of, 16:1781
- Trilinolenoylglycerol, calibration experiments of, chemometric approaches by LC-ion trap MA, 16:388, 16:801 (erratum)
- Trimannosyl, of transition metals, electrospray ionization study of, 8:32
- Trimethyl borate, gas phase reactions with phosphates and non-covalent complexes, 13:1088
- Trimethylaluminum and methylamines mixtures, ion-molecule reactions in, 4:111
- Trimethylamine, mechanism of loss from acetylcholine, density functional theory study, 20:238
- Trimethylamine systems, ion-molecule reactions in, 2:220
- Trimethylenemethane negative ion, PES of, 10:800

- Trimethylgallium, tris(*tert*-butyl)gallium, triethylantimony and, thermally-induced decomposition of, 19:947
- Trimethylphosphate, metal cation, interactions with nucleic acids, 19:305
- Trimethylsilyl adduct ions, radiative stabilization of, 1:144
- Trimethylsilyl cyanide, HCN in situ generation for MS, 17:51
- Trimethylsilyl derivatives, of hydroxy- and oxo-carboxylic acid, fragmentation of, EIMS, 19:66
- Trimethylsilyl ether derivatives, hydroxy fatty acids as, analysis of by electron ionization gas chromatography/mass spectrometry, 6:40
- Trimethylsilyl ethers of cholesterol oxidation products, mass spectra of, trimethylsilyl group migration in, production characterization by linked-scan tandem mass spectrometry, 4:327
- Trimethylsilyl group migration in the mass spectra of trimethylsilyl ethers of cholesterol oxidation products, production characterization by linked-scan tandem mass spectrometry, 4:327
- Trimethylsilylmethanol, metastable ions of, 9:1029
- Trinucleotides, deprotonated, gas phase conformations of, 14:161
- Tripeptide radical cations, facile generation of, 16:763
- Tripeptide VPL, 5-dihydroxybenzoic acid in cluster with, 12:2, 12:726
- Tripeptides
 fragmentation of doubly-protonated, formation of b_2^{2+} ions, ESI MS, 20:2135
 selected, gas-phase basicities of histidine and lysine and their, 7:1203
- Triphenylcorrole(CuIII) complexes, amino acid ester fragmentation induced by, MALDI MS, 18:791
- Triphenylphosphonium-aminoethyl derivatives of fatty acids, FAB-CAD-B/E linked scan mass spectrometry of, charge-remote fragmentation during, 3:769
- Triple quadrupole mass spectrometer
 collisional activation in, alkenamide fragmentation by, 15:1462
 ion mobility in, conformer selection of protein ions by, 5:127
- Triple quadrupole mass spectrometry as applied to flame diagnostics, study of the $C_2H_4/N_2O/Ar$ flame, 4:152
- Triple quadrupole tandem mass spectrometry (MS/MS), in molecular characterization of thiols, 15:288
- TRIPOLE. *See* Three rods ion optic
- Tris (perfluoroalkylethyl)silyl alkyl amines, as calibration standards for ESI-MS in the mass range of 100–3000 Da, 12:1050
- Tris(2,4,6-trimethoxyphenyl) phosphonium, hydrogen/deuterium exchange of, computational investigation of, 16:1067
- Tris (trimethoxyphenyl)phosphonium acetyl charged derivatives of peptides
 ESI-MS and MS/MS of, 11:107
 MALDI-PSD spectra of, 11:145
- Trisaccharides, sodiated isobaric, separation using ESI-APIM-TOF MS, 16:660
- meso*-Trisphenylcorrole, TPP and, observations using ESI MS, 17:1306
- Tris(*tert*-butyl)gallium, trimethylgallium, triethylantimony and, thermally-induced decomposition of, 19:947
- Triterpene saponin mixture, *C. quinoa*, nLC-ESI-MS/MS, 17:795
- Triterpenoids, *Ganoderma lucidum*, analysis using LC-ESI/MS, 18:927
- Trityl cations, benzene loss from, mechanistic study, 17:730
- Troponin C
 cardiac, small Ca^{2+} -induced conformational changes in, 10:703
 skeletal, calcium binding site and, 11:770
- Tropylum, chlorine isotopic abundances, monomeric metaphosphate anion, and Conestoga wagon theory, 4:761
- Trp-cage, conformation studied by Q/TOF MS, 18:195
- Trypanosoma brucei, complex component analyses in, 14:728
- Trypanosomes, analysis of flagellar proteome using ETD/ETD/CAD, 20:167
- Trypsin-catalyzed ^{16}O -to- ^{18}O exchange for comparative proteomics: MS/MS comparison using MALDI-TOF, ES-IQTOF, and ESI-ion trap mass spectrometers, 14:704
- Tryptic digests, protein, LC/MS and LC/MS/MS determination of, 1:158
- Tryptic peptides, field strength and separation of, by IM MS, 16:158
- Tryptophan
 complexation with Na^+ and K^+ in gas phase, cation- π effects in, 11:1037
 metal and proton complexes of, in gas phase, chiral recognition characteristics of, 14:571
 neutral alkaline earth-metal complexes, gas-phase formation, 19:1021
 O_2 ($^1\Delta_g$)-derived oxidation products, MS/MS, 20:188
 oxidation products and free radicals of, identification by MS, 14:406
 protonated, gas phase reactions of, 15:65
- Tryptophans, modified isomeric, structural characterization and differentiation of, 13:1298
- Tubulin, evaluating reproducibility in complex spectra, 19:367
- Tulips, phytochemicals and carbohydrate profiling in tulip tissues and bulbs, PESI MS, 20:2304
- Tumors
 head and neck, mouse, N-linked oligosaccharide glycoforms, MALDI MS, 19:671
 novel assay system using mass-tag technology, 18:106
- Tunable complexes, metal-ligand complexation for flavonoid diglycosides, ESI-MS, 18:422
- Tunable mid-infrared free-electron laser, infrared matrix-assisted laser desorption and ionization by using, 7:1187
- Tungstate anions, detection of gaseous $W_{12}O_{41}^{\bullet}$ and, by LDI-MS, 20:1782
- Tungsten hexacarbonyl dication, ground and excited states of, double-ionization energies to, 2:108
- Tungsten hexacarbonyl ions
 doubly and singly charged, generated by charge stripping, electron impact, and charge exchange, internal energy distributions deposited in, 1:16
 internal energy distributions, after neutralization-reionization, 5:1093
- Tween80, effect on signal intensity of intact proteins, MALDI-TOF-MS, 18:102
- Two-dimensional charge injection device, direct generation of ion beam images with, 6:140
- Two-dimensional Coulomb-induced frequency modulation in Fourier transform ion cyclotron resonance, a mechanism for line broadening at high mass and for large ion populations, 4:909, 5:605
- Two-dimensional electrophoresis, proteins separated by, peptide mass fingerprints from, 14:943
- Two-dimensional gel proteins, identification by MALDI-TOF or μ -LC-ESI MS/MS, 14:956
- Two-stage linear time-of-flight mass spectrometer, photodissociation of high molecular weight peptides and proteins in, 6:578
- Two-step laser desorption/ionization, aerosol TOFMS with, for analysis of individual organic particles, 9:1068
- Two thermospray probe, insert designs that allow clogged probes to be rapidly replaced, 4:182
- Tyrosine
 chemical cross-linking with lysine, 15:1604

- complexation with N^+ and K^+ in gas phase, cation- π effects in, 11:1037
- Tyrosine kinase, of spleen, conformational dynamics upon diphosphorylated ITAM binding to, 16:1039
- Tyrosylprotein sulfotransferases, TPST-1 and TPST-2, analysis by LC/ESI-MS, 19:1459
- U
- Ubiquitin
- apparent gas-phase acidity of, 7:1211
 - conformation of gas-phase ions, 19:1907
 - elongated conformers of, ESI-FAIMS-MS of, 12:894
 - elongated states, resolution and structural transitions, 18:322
 - in methanol, ESI-MS of, 12:317
 - protonated, copper adducts of, electrospray ionization mass spectrometry of, 6:521
 - role of conformation on ECD, 17:1469
- Ubiquitin conformers, multidimensional separations of, in gas phase, 16:1427
- Ubiquitin cross-sections, by FAIMS/energy loss, 11:738
- Ubiquitin ions
- conformations of, dissociation of, 13:719
 - proton transfer reactions of, 8:954
- Ubiquitination, protein, analyzed by ETD-based MS fragmentation or CID, 20:1652
- UDP-Glucuronosyl-transferase 1A1, regioselectivity towards flavonoids, metal complexation and MS/MS, 19:246
- UGT1A1. *See* UDP-Glucuronosyl-transferase 19:1A19:1
- Ultra high performance liquid chromatography, pharmaceuticals in surface waters, matrix effects in HPLC-MS/MS and UPLC-MS/PM analyses, 19:713
- Ultra-high sensitivity analysis of peptides and proteins, micro-electrospray mass spectrometry, 5:605
- Ultra performance liquid chromatography, improves hydrogen/deuterium exchange mass spectrometry, 17:163
- Ultra performance liquid chromatography electrospray ionization Q-time-of-flight, *Garcinia* spp. PPAP screening using PID and, 20:1846
- Ultra thin layer chromatography (UTLC) atmospheric pressure (AP) matrix-assisted laser desorption/ionization (MALDI) mass spectrometry, of small molecules, 16:906
- Ultramark 1621 as a reference compound for positive and negative ion fast-atom bombardment high-resolution mass spectrometry, 3:842
- Ultratrace determination of highly hydrophilic compounds by 2,2,3,3,4,4,5,5-octafluoropentyl chloroformate-mediated derivatization directly in water, 10:1328
- Ultraviolet irradiation
- NBD-tagged glycosphingolipids, UV-MALDI MS analysis, 19:923
 - photodissociation of peptide-crown ether complexes by, 19:257
- Ultraviolet MALDI
- DNA decay under, 9:866
 - of DNA fragment ions, 10:423
 - of protein, 9:1
- Ultraviolet-MALDI matrices, liquid based, detection limits of proteins from, 15:486
- Ultraviolet matrix-assisted laser desorption/ionization
- of oligodeoxynucleotides metastable decay, 12:180
 - of oligonucleotides, dihydroxybenzoic acid in, 8:218
- Ultraviolet matrix-assisted laser desorption ionization mass spectrometry, stabilizing double-stranded DNA in, 13:1129
- Ultraviolet photoionization, vacuum, ion decompensation *versus* molecular size probed by, 2:103
- Ultraviolet spectroscopy, centrin-2 metal-binding properties determined by μ ESI-MS and, 17:1158
- Understanding ion deceleration lenses, what are the simplicity/performance trade offs?, 2:322
- Underwater mass spectrometers for in situ chemical analysis of hydrosphere, 12:677
- Unequivocal determination of metal atom oxidation state in naked heme proteins, Fe(III)myoglobin, Fe(III)cytochrome *c*, Fe(III)cytochrome b5, and Fe(III)cytochrome b5 L47R, 11:120
- Unexpected high energy ions from a chemical ionization source, 8:1262
- Unfolding proteins monitored by electrospray ionization mass spectrometry: a comparison of positive and negative ion modes, 9:1248
- Unimolecular and collision-induced reactions of double charged porphyrins, 3:235
- Unimolecular photochemistry, of n-alkenes studied by photo dissociation-photoionization mass spectrometry, 4:65
- Uninterpreted high-energy collision-induced dissociation spectra of peptides, search of, with sequence databases, 7:1089
- Unit resolution mass spectrometry, of 112-kDa molecules, 3-kDa accuracy of, 8:380
- United States space program, MS in, 12:656
- Universal algorithm for fast and automated charge state deconvolution of electrospray mass-to-charge ratio spectra, 9:225
- Unknown identification using reference mass spectra: quality evaluation of databases, 10:1229
- Unknowns, spectra of, interferences from, software-based mass spectral enhancement to remove, 7:598
- Unsaturated and saturated aliphatic molecules, 193-nm photodissociation of ions from, 7:114
- Unsaturated fatty acids, polyhydroxy, electrospray ionization and low energy tandem mass spectrometry of, 7:140
- Unsuitability of $12H_6O$ as a standard of atomic weight, 6:1243
- Unusual ions, XeF^+ , IF^+ , and other, generated by reactions of hyperthermal ion beams at self-assembled monolayer surfaces, 4:938
- Unusually high proton affinity of aza-18-crown-6 ether: implications for the molecular recognition of lysine in peptides by lariat crown ethers, 13:493
- UPLC. *See* Ultra performance liquid chromatography
- Uracil, protonation sites of, 11:1065
- Uracil and its derivatives, collision-induced dissociation of, 5:339
- Uracil/uracil analogs, acidity in gas phase, 13:985
- Uranium, a particulate isotopic standard in an aluminosilicate matrix, 5:852
- Uranyl cation, CID pathways containing, 14:1205
- Uranyl hydroxide-anion complex, monopositive, intrinsic hydration of, 15:777
- Uranyl nitrate, ions generated from, by ESI, detected by FT-ICR MS, 17:230
- Urban air, selected organic compounds in, automated high-speed analysis of, by on-line isotopic dilution cryofocusing gas chromatography/mass spectrometry, 5:1001
- Urea, substituted *N* in, cyclization of, in gas phase and solution, 16:1739
- Urinary nucleosides, MALDI-TOF MS of, 16:940
- Urine
- ethylglucuronide in, LC-ESI-MS/MS of, 15:188
 - human
 - LTE₄ in, identification of, negative ion tandem mass spectrometry of leukotriene E₄ and LTE₄ metabolites, 2:314
 - modified nucleosides and ribosylated metabolites, ESI-FTICR MS and ESI-IT-MS, 19:1500
 - screening for cocaine and metabolites by TOF-MS, 20:891
 - platinum determination in, isotope dilution gas chromatography/mass spectrometry for, 2:85

- Uronic acid epimerization, in chondroitin sulfate oligosaccharides, MS/MS of, 14:1270
- Uronosyl substitution patterns, in acidic heteroxylans, ES-IMS/MS of, 15:43
- Use of a kinetic energy orifice as a probe of metastable dissociation in FTICR-MS, 11:1118
- Use of an internal control for MALDI-TOFMS of bacteria, 10:1131
- Use of combined mass spectrometry methods for the characterization of a new variant of human hemoglobin, the double mutant hemoglobin Villeparisis $\beta 77(\text{EF1}) \text{His} \rightarrow \text{Tyr}$, $\beta 80(\text{EF}_4) \text{Asn} \rightarrow \text{Ser}$, 7:163
- Use of double-depleted ^{13}C and ^{15}N culture media for analysis of whole cell bacteria by MALDI TOF and FTMS, 14:1306
- Use of heated capillary dissociation and collision-induced dissociation to determine the strength of noncovalent binding interactions in gas phase peptide-cyclodextrin complexes, 8:244
- Use of low- and high-energy collision-induced dissociation tandem mass spectrometry in the identification of an unusual amino acid in a semisynthetic polypeptide, 7:1034
- Use of region II of the a/q stability diagram for fast scanning of a linear quadrupole mass spectrometer, 8:539
- Use of static pressures of heavy gases within a quadrupole ion trap, 14:1099
- Utility of membrane preconcentration-capillary electrophoresis-mass spectrometry in overcoming limited sample loading for analysis of biologically derived drug metabolites, peptides, and proteins, 8:15
- Utility of N-peracetylation of proteins for their structure determination by mass spectrometry, 4:87
- Utility of peak shape analyses in determining unresolved interferences in exact mass measurements at low resolution, 9:789
- Utilization of MS³ spectra for the multicomponent quantification of diastereomeric N-acetylhexosamines, 11:1086
- Utilizing MALDI-CID for generation of structural information from poly (alkylmethacrylate)s, 8:1206
- UV laser ablation products of organic polymers, direct chemical analysis of, by using selective ion monitoring mode in gas chromatography/mass spectrometry, 5:106
- UV-MALDI. *See* Ultraviolet matrix-assisted laser desorption/ionization
- UV-matrix-assisted laser desorption Fourier transform mass spectrometry for peptides, investigation of, 2:22
- V
- V-P-O catalysts, ion association by time-of-flight mass spectrometry, 1:129
- Vaccinia virus, class I HLA-A*0201 peptide from, identification by mass FTICR MS/MS, 16:1812
- in Vacuo, anionic peptide radical formation in, 17:1249
- Vacuum chamber, concentric tube, for high magnetic field, high-pressure ionization in a Fourier transform ion cyclotron resonance mass spectrometer, 4:168
- Vacuum chamber of an electrospray time-of-flight mass spectrometer, ion-molecule collisions in, effect of, on mass spectra of proteins, 7:342
- Vacuum matrix-assisted laser desorption/ionization (MALDI), on QqLIT instrument, 16:176
- Vacuum system O-rings, plasticizer contamination from, in quadrupole ion trap MS, 13:901
- Vacuum ultraviolet light, peptide ion fragmentation pathways induced by, 16:1384
- Vacuum ultraviolet light source
laser-produced plasma as, tunable SPI-TOF-MS, 20:2192
product ions generated by 157 nm photodissociation of phosphopeptides, 20:2334
- Vacuum ultraviolet photofragmentation, of peptide ions, factors affecting, 18:1439
- Vacuum ultraviolet photoionization, ion decompensation *versus* molecular size probed by, 2:103
- Vacuums, ultrahigh, ion fragmentation and fluorescence of in, ESI, 20:1731
- Validation using sensitivity and target transform factor analyses of neutral network models for classifying bacteria from mass spectra, 13:10
- Valine, lithiated, water binding energies to, 15:1014
- Valinomycin-alkali metal cation complexes, gas-phase studies of, attachment rates and cation affinities, 5:632
- Valve, a pulsed-leak, for use with ion trapping mass spectrometers, 7:118
- Van Berkel, Gary J., Biemann Medal 2005, 17:887
- van der Hart, Wim J., obituary, 20:1944
- Vancomycin antibiotics, bacterial cell wall precursor peptide interaction with, hydrogen/deuterium exchange in, 9:1255
- Vaporizer temperature, the effects of, ion formation of N-methylcarbamate pesticides in thermospray mass spectrometry, 5:913
- Variability of calcium binding to EF-hand motifs probed by ES-IMS, 12:1296
- Variable flow liquid chromatography-tandem mass spectrometry and the comprehensive analysis of complex protein digest mixtures, 8:1059
- Variation characterization, of proteins, up to 80 ku, determination of, high accuracy, by ionspray mass spectrometry, 2:387
- Varying counterions, effect of, on the sensitivity of electrohydrodynamic and fast atom bombardment mass spectrometry, 1:233
- Velocities, positions, and kinetic energies of resonantly elicited ions in the quadrupole ion trap mass spectrometer, determination of, by laser photodissociation, 4:792
- Velocity
above thermal and below Bohr, molecule fragmentation at, 10:471
initial, of ions, MALDI-generated, TOF-MS effects of, 8:209
in reflectron TOF-MS, 10:992
- Venom, honeybee, detection in envenomed tissues, MALDI MSI, 20:112
- Vertebrate type steroid hormones, MS/MS and sequential product ion scanning of, 8:365
- Vibrational frequencies, of linear peptide and protein ions, RRKM rate constant calculation using, 17:1749
- Vibrio cholerae* 0:1, ESI-MS fragmentation of analogs of saccharides from, 17:749
- Villeparisis, double mutant hemoglobin, $\beta 77(\text{EF1}) \text{His} \rightarrow \text{Tyr}$, $\beta 80(\text{EF}_4) \text{Asn} \rightarrow \text{Ser}$, use of combined mass spectrometry methods for the characterization of a new variant of human hemoglobin, 7:163
- Vincamine, oxygenated chemical ionization reagents with, ion-molecule reactions of, 7:250
- Vinyl fluoride formation, reactions with strong bases, and characterization of 1-fluorovinyl anion and the fluoride-acetate hydrogen-bonded complex, 3:91
- Vinylchloride, vinylbromide, vinyliodide, long-lived excited electronic states of, 14:110
- Viscous gas flow through capillaries, ion transport by, 5:873
- Visualization and analysis of molecular scanner peptide mass spectra, 13:221
- Vitamin B12, fragmentation of, in aerosol MALDI, 8:140
- Vitamin D receptor, DNA-binding domain of, zinc-induced changes in, 9:8
- Volatile organic compounds
in air, PIT-MS detection and identification of, 16:1316

- chlorinated, degradation of, 9:1321
in human blood, measuring, by using purge-and-trap gas chromatography/mass spectrometry, interferences, removing of, importance of enhanced mass resolution in, 3:831
- Voltage
applied to quadrupole mass filter, modulation of, 16:379, 16:800 (erratum)
compensation, peak shapes of, using domed FAIMS, 16:948
- Voltages, applied, and reaction times, negative chemical ionization in quadrupole ion trap mass spectrometry, 5:757
- VX, on concrete, IT-SIMS of, 11:69
- W
- Warfarin, interactions with HSA, effect of tanshinone IIA on, ESI MS, 19:1568
- Wasps, cuticular hydrocarbons of, GC-MS of, 9:320
- Waste streams, components of, identification of, by electrospray and tandem mass spectrometry, 4:604
- Water
binding energies to lithiated valine, 15:1014
electrochemically-induced reactions of hexafluorophosphate anions with, ES-MS, 17:939
electrospraying in air, at atmospheric pressure, 15:253
elimination from GlyGlyH⁺ carboxyl group, 14:1192
elimination in condensation reactions in protonated dipeptides, 15:1025
gas-phase proton transfer reactions involving, under thermal conditions, 3:624
nonylphenol carboxylates in, LC-ES-MS/MS of, 14:516
surface, pharmaceuticals in, matrix effects in HPLC-MS/MS and UPLC-MS/PM analyses, 19:713
- Water/acetone cluster ions, tandem mass spectrometry studies of, 2:113
- Water-alcohol and ammonia-alcohol protonated clusters, on the structure of, 4:507
- Water cluster calibration, reduces mass error in electrospray ionization mass spectrometry of proteins, 8:1158
- Water clusters, in triple quadrupole MS, 9:814
- Water molecules, propanediol ion solvation by, in gas phase, 15:1123
- Water solutions
ionization from, with SF₆ discharge suppression, negative ion electrospray mass spectrometry of nucleotides, 4:289
suppression of electric discharge with SF₆ gas, electrospray mass spectrometry of, 2:497
- Water-solvated ammonia radical cation, reactions with carbonyl compounds, 15:966
- Water vapor, H₃O⁺ and H₃O⁺(H₂O)³ ion isotopic composition in exchange reactions with, SIFT-MS analysis of, 11:866
- Waveforms
asymmetric, FAIMS for, 16:1447
external customized, application to quadrupole ion trap, 10:355
- Wavelength, near-UV-induced MALDI as function of, 9:885
- Weighted two-band target entropy minimization for the reconstruction of pure component mass spectra: simulation studies and application to real systems, 14:1295
- Why are alkane eliminations from ionized alkanes so abundant?, 9:21
- Why are electron capture negative ion mass spectra not reproducible? An ion source problem, 4:270
- Why are ions stable species in peptide spectra?, 6:1165
- Wide-angle secondary ion probe for organic ion imaging, A, 2:362
- Wire-in-a-capillary bulk-loaded nano-electrospray emitter, electrochemical processes in, 12:853
- Wittig reaction, thermally unstable intermediate in, 9:1168
- X
- X-ray crystallographic study, of 2,5-dihydroxybenzoic acid non-covalent association with cyclic and linear oligosaccharides, 11:228
- X-ray diffractometry, of cytosine nucleosides self-assembling into dimers, 15:268, 15:436
- X=Cl, Br, I (C₂H₅) X⁺ ions, a tandem mass spectrometric investigation of, the involvement of classical and nonclassical ethyl structures therein, 5:814
- XeF⁺, IF⁺, and other unusual ions generated by reactions of hyperthermal ion beams at self-assembled monolayer surfaces, 4:938
- Xylobiose inhibitors, binding to β-1,4 exoglucanase from *C. fimi*, MS/MS, 18:64
- Y
- Y⁺, Sc⁺, and Lu⁺, gas-phase reactions with alcohols, effects of the class and chain length of alcohols on the nature of primary products, 7:1157
- Yates, John, III, 16:1192
- Yeast hexokinase, MS-based assays of, 14:173
- Z
- Z-ionic fragments, hydrogen rearrangements to and from, peptides, ECD, 18:113
- Zeolites, for MS of tributyl phosphate, 8:718
- Zeptomole/attomole per microliter sensitivity for peptides, micro-electrospray, 5:867
- Zidovudine metabolites, in peripheral mononuclear blood cells of HIV-infected patients, HPLC-MS/MS of, 11:1134
- Zinc
affinity to CphA monitored by automated nano-ESI-MS, 17:180
deposition during peptide-zinc complex analysis, ESI MS, 20:2199
- Zinc finger peptide complexation, solution-phase, MALDI-MS of, 10:27
- Zinc fingers, of estrogen receptors, cysteine reactivity in, 16:2017
- Zinc-induced conformational changes in the DNA-binding domain of the vitamin D receptor determined by electrospray ionization mass spectrometry, 9:8
- Zinc-ligand-hexose diastereomers, ion mobility measurement and density functional theory calculations of, 13:284
- Zinc-octaethylporphyrin, collisionally activated decomposition of, 9:767
- Zinc(II) ion complexes, chiral recognition of, by CID, 15:795
- Zirconium phosphonate, magnetic nanoparticles modified by, selective phosphopeptide capture, 19:1176
- ZoomQuant
with ion trap mass spectrometer, ¹⁸O labeling on, quantification using, 16:916
for stable isotope label quantitation, 16:302
- Zubarev, Roman, Biemann Medal recipient, 19:751
- Zwitterion, formation in gas-phase H/D-exchange with CH₃OD, 18:512
- Zwitterion formation
of deprotonated trinucleotides, 14:161
in gas phase cyclodextrin complexes, 16:166
- Zwitterions
formation, gas-phase enantioselectivity and, amino acids, 17:442
sodiated amino acids, gas phase H/D exchange, 17:29