



Original Article

Anatomical characterization of ultra-structures, biominerals and histolocalization of metabolites in leaves of *Genipa americana*



Alex L. Vasconcelos^a, Andréa V. Santos^a, Rafael J.R. Padilha^b, Luiz C. Alves^b, Karina P. Randau^{a,*}

^a Laboratório de Farmacognosia, Departamento de Ciências Farmacêuticas, Universidade Federal de Pernambuco, Recife, PE, Brazil

^b Setor de Microscopia Eletrônica, Laboratório de Imunopatologia Keizo Asami, Universidade Federal de Pernambuco, Recife, PE, Brazil

ARTICLE INFO

Article history:

Received 13 April 2017

Accepted 19 May 2017

Available online 13 June 2017

Keywords:

Crystals morphotype

Histochemistry

Histolocalization

Jenipapo

Leaf anatomy

ABSTRACT

Inserted in one of the largest families among the Angiosperms, *Genipa americana* L., Rubiaceae, can be found in all Brazilian territory, presenting great medicinal importance, where several uses have been attributed. In view of this, this work has the purpose of analyzing the ultrastructural, biomineral, phytochemical and histochemical characteristics of the leaves of this native species from Brazil. For this, light microscopy, polarization and scanning electron microscopy techniques were used with X-ray scattering energy, associated to chromatographic and histochemical tests. The anatomical ultrastructural characteristics of the leaves detailed information about the type and arrangement of the cuticle, trichomes, surface and arrangement of the tissues that determine the botanical identity of this species. The phytochemical tests allowed determining their chromatographic pattern and histochemistry to determine the exact storage site of these substances in the leaf. It was observed that the characterization of the crystalline macro-pattern present in the analyzed species, as well as its exact elemental composition, can be considered an important differential diagnosis factor. The results characterize the leaves of this species in different aspects, being a native species and pharmacologically promising, with different popular uses and proven pharmacological activities, and more in depth studies is needed.

© 2017 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Farmacognosia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Rubiaceae is considered as one of the largest families among Angiosperms, it comprises the fourth largest number of species, with approximately 13,000 species distributed in 550 genera. The *Genipa* genus is characterized by two species: *Genipa americana* L. and *G. infundibuliformis* Zappi & Semir (Delprete et al., 2005). Among the studies carried out on the genus, the researches are concentrated on the *G. americana* species.

Popularly known as jenipapo, this species can be found all over Brazilian territory. It is a native plant of Brazil, of arboreal size, reaching up to 30 m in height (Corrêa, 1984; Delprete et al., 2005; Zappi, 2015). Naturalists, who visited Brazil in the colonial period, have long described the use of *G. americana* as a body dye (Alves, 2010, 2013).

Several medicinal uses have traditionally been attributed to this species. The leaf decoction has an antidiarrheal or antisyphilitic

action, whereas the leaf infusion is used against liver diseases (Corrêa, 1984). Based on ethnobotanical surveys, Souza et al. (2013) point out the traditional use of this species in the treatment of cough, anemia, contusions, dislocations; as depurative and associated with popular beliefs. Among the secondary metabolites found in this species, it highlights the presence of tannins (Revilla, 2001), monoterpenes (Ono et al., 2007), steroids (Conceição et al., 2011) and mostly iridoids which are characteristic of the Rubiaceae family (Jensen, 1983; Ueda and Iwahashi, 1991; Ono et al., 2005).

Different studies have indicated the potential of *G. americana* for use in the treatment of asthma (Deng et al., 2013), ophthalmic diseases (Koriyama et al., 2013; Song et al., 2013), antiviral action (Lin et al., 2013), cardiovascular disorders (Zhang et al., 2013), antidepressant activity (Tian et al., 2010), in the treatment of Alzheimer's disease (Nam et al., 2013; Gao et al., 2014; Wang et al., 2012; Zhang et al., 2012), anti-inflammatory (Fu et al., 2012; Li et al., 2012) (Kim et al., 2012; Dando et al., 2013; Yang et al., 2013).

In view of the recognized popular use of this species and distinct pharmacological studies demonstrating its therapeutic potential, this work aims at analysing the anatomical ultrastructural, biomineral, phytochemical and histochemical characteristics of *G. americana*, native to Brazil.

* Corresponding author.
E-mail: karina.prandau@ufpe.br (K.P. Randau).

Materials and methods

Plant material

Genipa americana L., Rubiaceae, mature leaves were collected from adult species on the campus of the Universidade Federal de Pernambuco (–8,0464168, –34,9451315), Recife, Northeast Brazil, at October 12, 2015, in the fruiting season. The voucher specimen was deposited at the Herbarium Dárdano de Andrade Lima, for botanical legitimation, by the number 89999.

Anatomical characterization

Mature leaves were collected from second or third node, from up to five individuals. The samples were fixed in FAA50 (Johansen, 1940). Cross and paradermic sections of the middle portion of the leaflets were freehand obtained, submitted to double staining process with astra blue and safranin (Johansen, 1940; Kraus and Arduin, 1997) and mounted in glycerine on semi-permanent slides. Analyze of the structures was performed using optical and polarized light microscopy.

Phytochemistry determination

The methanolic extract from fresh leaves was processed by infusion (0.1%) and subsequently analyzed by thin-layer chromatography (TLC) (Kieselgel 60, 0.2 mm, Merck), with different eluents systems and reagents to search alkaloids, mono- and sesquiterpenes, triterpenes and steroids, saponins, flavonoids, cinnamic derivatives and phenylpropanoid glycosides, hydrolyzable and condensed tannins. Phytochemical screening of the extract was carried out using a conventional protocol (Harborne, 1998; Randau et al., 2004).

Histochemical analyses

Freehand cross sections from the middle portion of fresh leaves were subjected to reaction with specific reagents, according to the different groups of metabolites to be investigated: Lugol reagent (Johansen, 1940) for starch, Sudan IV (Kraus and Arduin, 1997) for detection of epicuticle wax, acid phloroglucin (Jensen, 1962) for lignins, potassium dichromate (Gabe, 1968) for phenolic compounds, NADI reagent (David and Carde, 1964) for essential oils and oil-resins, vanillin-sulphuric acid (Wagner and Bladt, 1996) for iridoids, antimony trichloride (Mace and Howell, 1974) for steroids, vanillin-hydrochloric acid (Mace and Howell, 1974) for tannins, Dragendorff reagent (Yoder and Mahlberg, 1976) for alkaloids and hydrochloric acid (Johansen, 1940) for calcium oxalate crystals. For each procedure employed on the samples slides without any treatment (blanks) were also mounted. Images documentation was performed with a light microscope (Alltion) equipped with a digital camera.

Scanning electron microscopy

Cross sections obtained by hand were fixed with 2.5% glutaraldehyde (GA) + 4% formaldehyde buffer. After fixation and washing, the dehydrated samples by increasing ethanolic series (Johansen, 1940) were submitted to CO₂ Critical Point (HCP-2). Afterwards, the samples were mounted on metal brackets and coated with a gold layer metalizer Quorum Q150T ES (Haddad et al., 1998). The images obtained were used in the anatomical characterization of ultra-structures with a QUANTA 200 FEG scanning electron microscope (SEM).

Morphological analyze and elemental composition of crystals

The chemical microanalyses by EDS were done with X-ray detector attached to the Zeiss-EVO-LS15 scanning electron microscope, under the same operational conditions. Absorbance peaks indicating relative elemental chemical concentration $\geq 0.5\%$ were considered significant.

Results

Anatomical characterization

The epidermal cells, which constitute the adaxial face, present polygonal contours containing amorphous inclusions within them that vary in size (Fig. 1A). On the abaxial side the cells are smaller and it can be noticed the presence of paracytic and anomocytic stomata, restricted to this face, characterizing the leaf as hypostomatic. Several inclusions are also observed on the abaxial face of the leaf blade, inside the epidermal cells and on stomatal guard cells (Fig. 1B). The images obtained by Scanning Electron Microscopy (SEM) address that the cuticle presents in the format of crystalline projections, such as an extensive layer of nanofilaments that cover the entire epidermal surface on both faces (Fig. 1C and D), as well as on the stomata (Fig. 1E).

In cross-section the epidermis is covered by a thick cuticular layer. The epidermis is unstratified, formed by larger cells on the adaxial side, more elongated in the anticlinal direction, as opposed to the abaxial epidermal cells, smaller and more elongated in the periclinal direction. The mesophyll has a dorsiventral organization and presents a palisade parenchyma facing the adaxial face, constituted by two-three layers of elongated cells (Fig. 1F). The lacunar parenchyma is multi-stratified composed of loosely organized rounded cells, with small intercellular spaces between them, comprising the abaxial region of the mesophyll. Some secondary vascular bundles can be seen dipped into the mesophyll. The presence of scarce multicellular and glandular tector trichomes (Fig. 1G and H) can be observed on the abaxial surface. Furthermore, the presence of these trichomes was also detected by electron microscopy, revealing small and scarce glandular trichomes on the abaxial side of the blade that are inserted between the stomata and the epicuticular layer (Fig. 1I).

The midrib presents biconvex symmetry, being prominent on the abaxial surface, with vascular bundles of concentric arrangement with no perivascular sclerenchymal fibers (Fig. 1J and K). At the center of the main vascular cylinder, in the medullary parenchyma region, there are accessory vascular caps, of which the presence of druses can be confirmed (Fig. 1L and M). Underlying the epidermal monolayer there are several strata of angular-type collenchyma cells where the presence of crystalline inclusions of druse type is recognized (Fig. 1N). Through polarized light microscopy, the presence of these crystals throughout the leaf blade can be observed, presenting a higher incidence in the cortical parenchyma of the midrib (Fig. 1O).

Phytochemistry determination and histochemical characterization

According to the results, the presence of iridoids, mono and sesquiterpenes and of triterpenes and steroids in the extract obtained from the leaves was observed. The tests also revealed the presence of hydrolyzable tannins, proanthocyanidins, cinnamic derivatives and phenylpropanoglycosides. The analyzes for alkaloids and saponins were negative. The results obtained formed the basis for performing the histochemical tests.

These tests were performed by comparing the sections after reaction with specific reagents under the negative control,




Fig. 1. Anatomical structures of leaves of *Genipa americana*. Leaf blade. (A) Adaxial epidermal cells. (B) Abaxial epidermal cells. (C) Adaxial epidermis covered by tubular cuticle (SEM). (D) Abaxial epidermis covered by tubular cuticle and presence of stomata (SEM). (E) Stoma in the abaxial face (SEM). (F) Dorsiventral mesophyll with druses. (G) Multi-serrated trichome. (H) Glandular trichome. (I) Detail of the glandular trichoma in the abaxial face (SEM). (J) Biconvex midrib. (K) Detail of the midrib (SEM). (L) Vascular cylinder with concentric arrangement. (M) Detail of the vascular cylinder (SEM). (N) Crystalline inclusion of druse type. (O) Druses observed through polarized light microscopy. ad, adaxial surface; ab, abaxial surface; avb, accessory vascular bundles; cp, cortex parenchyma; co, collenchyma; ct, cuticle; dr, druse; ep, epidermis; es, ergastic substance; gtr, glandular trichome; mdr, midrib; mep, medullary parenchyma; mtr, multiseriate trichome; pp, palisade parenchyma; sp, spongy parenchyma; st, stomata; vb, vascular bundle; xy, xylem; ph, phloem. Bars: A, B, F, H, I, K, M, O = 25 μm . C, D, J = 25 μm . E = 10 μm . G, N = 40 μm . L = 200 μm .




Fig. 2. Histolocalization of metabolites present in leaves of *Genipa americana*. Control sample: (A) Midrib. (B) Mesophyll. Histochemical tests: (C) Starch grains in parenchyma cells. (D) Covering cuticular epidermal cells. (E) Presence of lignins in woody vascular bundles. (F) Polyphenols present in epidermal cells and palisade parenchyma. (G) Volatile oils inside abaxial epidermal cells. (H) Iridoids inside the palisade parenchyma cells. (I) Steroids in perivascular idioblasts in the midrib. (J) Proanthocyanidins inside the palisade parenchyma cells. cp, cortex parenchyma; ep, epidermis; ew, epicuticular wax; ir, iridoids; lg, lignin; pa, parenchyma; pcd, proanthocyanidins; ph, phloem; phc, phenolic compounds; pp, palisade parenchyma; sp, spongy parenchyma; st, starch grain; std, steroids; vo, volatile oil; xy, xylem. Bars: A = 100 μm . B, C, D, E, F, G, H, I, J = 25 μm .

cross-sections of the midrib and mesophyll without treatment with the reagents (Fig. 2A and B). The results demonstrate the presence of starch granules present in the mesophyll, both in the palisade parenchyma and in the epidermal cells (Fig. 2C). The intense accumulation of amyloplasts can also be observed in the medullary parenchyma region, in the center of the vascular cylinder.

From the reaction with Sudan III, the presence of a thick epicuticular waxy layer can be observed, covering both sides of the blade. The cuticle presents as an orange layer (Fig. 2D). The presence of lignins can also be verified in the woody vascular bundles, characterized by the pink color conferred by floroglucinol (Fig. 2E).

After reaction with potassium dichromate, the presence of phenolic compounds can be verified, characterized by reddish coloration, distributed throughout the leaf blade region, mostly in the

prismatic cells that make up the palisade parenchyma and epidermal cells of the adaxial face (Fig. 2F). The volatile oils present in the leaves of *G. americana* were evidenced by the reaction with the NADI reagent, conferring blue color to the spherical droplets described above, accumulated in the epidermal and subsidiary cells and abaxial epidermal cells (Fig. 2G).

The presence of iridoids was confirmed by the sulfuric vanillin reaction, showing reddish staining, determining the accumulation site of these substances in the xylem cells and some cells of the palisade parenchyma close to the phloem (Fig. 2H). The presence of steroids is observed in idioblasts located in the perivascular regions, close to the phloem, along the entire midrib, showing a characteristic orange color after reaction with antimony trichloride (Fig. 2I).




Fig. 3. Scanning electron micrograph and elemental composition of the crystals observed in *Genipa americana*. (A) Presence of druse in the midrib of the leaf in cross-section. (B) Detail of the rosette-like druse (whewellite type) covered by the membrane of the idioblast. (C) Analysis of elemental composition of the crystal. (D) Percentage of the chemical constituents of the crystal. Bars: A = 20 μm . B = 2 μm .

The presence of proanthocyanidins was characterized by poor red coloration as well as the results found in phytochemistry. The accumulation of these substances was verified in some cells of the palisade parenchyma, after reaction with sulfuric vanillin (Fig. 2J). The reaction with hydrochloric acid demonstrated the soluble nature of the crystalline inclusions, confirming the chemical composition of calcium oxalate of these crystals.

Morphological analyze and elemental composition of crystals

The dominant crystalline macro-pattern in the leaves of *G. americana* is druse type of crystals in rosette. This type of crystal can be found along the mesophyll, in the parenchymal cortex and in the medullary parenchyma of the midrib (Fig. 3A and B).

The microanalysis of druses, performed by scanning electron microscopy (SEM), by energy dispersive X-ray analyze (SEM), revealed three peaks of absorbance for calcium (0.37, 3.7 and 4.0 keV), a peak for oxygen (0.37 keV) and one for carbon (0.25 keV). According to the results above, we can state that the druse is composed mainly of calcium (0.13%), followed by oxygen (0.06%) and carbon (0.02%) (Fig. 3C and D).

Discussion

Anatomical characterization

Characteristics such as hypostomatic leaf, epidermal cells with straight anticline walls, paracytic and anomocytic stomata, uni and multicellular tector trichomes and dorsiventral mesophyll are common occurrences for the family, among the representatives of Rubiaceae (Metcalf and Chalk, 1979), as verified in *Tocoyena*

formosa (Cham and Schltdl.) K. Schum. (Coelho et al., 2006) and *Psychotria viridis* Ruiz & Pav. (Quinteiro et al., 2006).

The scarce presence of uni and multicellular tector trichomes can be verified, besides glandular trichomes. The presence of glandular trichomes is a characteristic that differs from the results achieved by Santos et al. (2006) and Urbano and Duarte (2010) in *G. americana* leaves. Some studies have indicated that exogenous factors such as herbivory, temperature, luminosity, water availability and seasonality are able to influence the formation and functioning of glandular trichomes in different plant groups (Viljoen et al., 2005; Gonzáles et al., 2008; Martínez-Natarén et al., 2011). The presence of idioblasts is detected in the parenchymatic region, external and internally to the vascular cylinder, which present crystalline inclusions in the form of druses. Superior plants commonly present crystals stored in their tissues, being related to physical protection, removal of oxalate from the metabolic system, storage of calcium, and regulation of light during photosynthesis (Franceschi and Nakata, 2005).

The crystal type is an important diagnostic feature for genus and species of Rubiaceae. The presence of crystals like raphide, druse, styloid, prismatic or crystalline sand can be confirmed in species belonging to this family, being common the presence of druse and crystalline sand for *Genipa* genus (Metcalf and Chalk, 1979). Raphides were noticed in *Palicourea longepedunculata* (Pereira et al., 2003), crystalline sand in some species of the genus *Rondeletia* L. (Kocsis et al., 2004) and druse in *T. formosa* (Coelho et al., 2006). Urbano and Duarte (2010) confirmed the presence of calcium oxalate druse on leaves and stem of *G. americana*. Moraes et al. (2011) report the presence of styloid prismatic and rusty crystals on leaves of some species of the genus *Psychotria*. The birefringence presented by druse in the analyzed species is considered

an anisotropic characteristic of crystals that have a chemical composition of a single element. Crystals that have mixed chemical composition are referred to as non-birefringent, making it impossible to evaluate the different refractive indexes that they possess, since they present alterations in optical properties in relation to pure crystals (Hammond, 2009; He et al., 2012).

Phytochemistry determination and histochemical characterization

The cuticle shape is dependent on the compounds present in the wax. Asymmetric secondary alcohols and β -diketones form wax spaces in the form of nanotubes, whereas primary alcohols and symmetrical secondary alcohols form flat plates (Hallam, 1967; Jeffrey et al., 1975). Differences in cuticle patterns can be taxonomically useful in generic terms and even between species (Metcalf and Chalk, 1979; Cutter, 1986). The fibers are associated with the vascular system, as cited by Metcalf and Chalk (1979) for Rubiaceae.

Analyses revealed phenolic compounds intensely visible throughout the palisade parenchyma, being also observed in the adaxial epidermal cells of less intense form. Studies have also reported the presence of phenolic compounds in different species of the Rubiaceae family, such as *Chomelia obtusa* Cham. & Schltdl. (Barros et al., 2008), *Palicourea rigida* Kunth (Rosa et al., 2010) and *G. americana* (Porto et al., 2014).

Most studies on essential oils in this species are limited to fruits only. Some studies indicate that these metabolites are composed mainly of carboxylic acids, esters and alcohols (Borges and Rezende, 2000; Pino et al., 2005; Pinto et al., 2006). In the analyzed leaves, the essential oils present a greater incidence of distribution in the abaxial face, being found in the form of granules with storage site in the epidermal and subsidiary cells.

The iridoids are present in most of the species of Rubiaceae, constituting an important chemical and pharmacological marker in *G. americana* (Alves and Ming, 2015). According to the analyses, the accumulation site of these substances was observed in the phloem cells and in some palisade cells, nearby these vessels. The presence of these compounds is observed in *Eucommia ulmoides* Oliv. (Hirata et al., 2011; Nam et al., 2013; Zhang et al., 2013), *Bellia trixago* (L.) All. (Venditti et al., 2013), *Gardenia jasminoides* J. Ellis (Zhang et al., 2013), *Castilleja tenuiflora* Benth (Carrillo-Ocampo et al., 2013).

In fruits of *G. americana* were identified the presence of different phyosteroids (Barbosa, 2008). Conceição et al. (2011) report that the presence of steroids in extracts of *G. americana*, being related to anticancer activity. There are few studies that indicate the presence of these metabolites in the leaves. The assays performed in this work revealed that, in the leaves, these substances are stored in idioblasts located in the perivascular regions, close to the phloem.

The phytochemical, results in the search for proanthocyanidins, contributed with the histochemical tests, where little reddish coloration was observed inside some cells of the palisade parenchyma. According to Porto et al. (2014), there is a low accumulation of these substances in this species, corroborating the histochemical results described for the leaves analyzed in this study.

Morphological analyze and elemental composition of crystals

Scanning electron micrographs, coupled with spectrometric and histochemical assays, revealed the presence of crystalline inclusions of the druse type of rosette crystals (whewellite) throughout the leaf blade. From these tests, the chemical composition of calcium oxalate of these crystals was confirmed. The presence of these inclusions, the determination of their chemical nature and their morphology, as well as the location site within the medicinal plant, can help in the characterization of taxonomic groups, becoming

relevant in the identification and purity of plant drugs (Metcalf and Chalk, 1979; Monje and Baran, 2002; Anitha and Sandhiya, 2014).

In the last decades, many plant crystals made of calcium oxalate have not been accurately identified. This biomineral occurs in two hydration states in plants: monohydrate (whewellite) or dihydrate (weddellite) (Franceschi and Horner Junior, 1980; Frey-Wyssling, 1981; Arnott, 1982). Monje and Baran (2002), in their works distinguished the whewellite of the weddellite druse from its star-like forms. Different crystalline morphotypes can be observed in Rubiaceae as styloid crystals, raphide and druses (Metcalf and Chalk, 1979). Zini et al. (2016) report the presence of druses in *Borreria latifolia* K. Schum., *B. orientalis* E.L. Cabral, R.M. Salas & L.M. Miguel, *B. palustris* (Cham. & Schltdl.) Bacigalupo & E.L. Cabral. Styloid crystals are observed in some species of the genus *Psychotria* L. Urbano and Duarte (2010) report in general the presence of druses into leaves and stem of *G. americana*, but details about the morphological analyze, elemental composition and histolocalization of these inclusions in medicinally important leaves are scarce in the literature referring to these species. These druses can be considered relevant structures to the ultrastructural anatomic characterization.

Conclusion

The anatomical characteristics commonly attributed to the representatives of Rubiaceae are not considered relevant in interspecific differentiation, since they are of general occurrence for the family. In response to this, the use of SEM becomes a useful tool in the anatomical characterization of the *G. americana* foliar ultrastructures. Besides the phytochemistry, the histochemistry allowed to obtain more specific data, determining the exact storage site of these substances in the leaf. The characterization of the crystalline macro-pattern, as well as its exact elemental composition, can be considered an important factor of differential diagnosis. The results presented allowed the characterization of this species in different aspects and point to *G. americana* as a promising native species with different popular uses and proven pharmacological activities, and further studies are needed.

Authors' contributions

ALV and AVS assisted in carrying out the laboratory work. RJRP and LCA helped in conducting the scanning electron microscopy (SEM) analyses. ALV wrote the paper and KPR provided a critical reading of the manuscript. ALV and KPR planned the project, collected the plant material and was responsible for its identification. KPR supervised the laboratory work. All the authors have read the final manuscript and approved the submission.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors thanks to Centro de Tecnologias Estratégicas do Nordeste (CETENE) and Laboratório de Imunopatologia Keizo Asami (LIKA) for the valuable work on the electronic microscopy slides preparation. Acknowledgment is also due to Laboratório de Gemologia – UFPE to the polarization microscopy and to the herbarium Dárdano de Andrade Lima for the botanical identification. Thanks to CNPq for research funding.

References

- Alves, L.F., 2010. Plantas medicinais e fitoquímica no Brasil: Uma visão histórica. Editora Pharmabooks, São Paulo.
- Alves, L.F., 2013. Produção de Fitorpéuticos no Brasil: História, Problemas, Perspectivas. *RVQ* 5, pp. 450–513.
- Alves, L.F., Ming, L.C., 2015. Chemistry and pharmacology of some plants mentioned in the letter of Pero Vaz de Caminha. *Ethnobot. Conserv.* 4, 3.
- Anitha, R., Sandhiya, T., 2014. Occurrence of calcium oxalate crystals in the leaves of medicinal plants. *IJP* 1, 389–393.
- Arnott, H.J., 1982. Three systems of biomineralization in plants with comments on the associated organic matrix. In: Nancollas, G.H. (Ed.), *Biological Mineralization and Demineralization*. Springer Verlag, Berlin, pp. 199–218.
- Barbosa, D.A., (Dissertação de Mestrado) 2008. Avaliação fitoquímica e farmacológica de *Genipa americana* L. (Rubiaceae). Programa de Pós-graduação em Ciências Farmacêuticas, Universidade Federal do Rio de Janeiro.
- Barros, M.P., Santin, S.M.O., Costa, W.F., Vidotti, G.J., Sarragiotto, M.H., Souza, M.C., Bersani-Amado, C.A., 2008. Constituintes químicos e avaliação do potencial anti-inflamatório e antioxidante de extratos das folhas de *Chomelia obtusa* Cham. & Schlttdl. (Rubiaceae). *Quim. Nova* 31, 1987–1989.
- Borges, E.S., Rezende, C.M., 2000. Main aroma constituents of genipap (*Genipa americana* L.) and bacuri (*Platonia insignis* M.). *J. Essent. Oil Res.* 12, 71–74.
- Carrillo-Ocampo, D., Bazaldúa-Gómez, S., Bonilla-Barbosa, J.R., Aburto-Amar, R., Rodríguez-López, V., 2013. Anti-inflammatory activity of iridoids and verbascoside from *Castilleja tenuiflora*. *Molecules* 18, 12109–12118.
- Coelho, V.P.M., Agra, M.F., Barbosa, M.R.V., 2006. Estudo farmacobotânico das folhas de *Tocoyena formosa* (Cham. & Schlttdl.) K. Schum. (Rubiaceae). *Rev. Bras. Farmacogn.* 16, 170–177.
- Conceição, A.O., Rosi, M.H., Oliveira, F.F., Takser, L., Lafond, J., 2011. *Genipa americana* (Rubiaceae) fruit extract affects mitogenactivated protein kinase cell pathway in human trophoblast-derived BeWo cells: implication for placental development. *J. Med. Food* 14, 483–494.
- Corrêa, P.M., 1984. Dicionário das plantas úteis do Brasil e das exóticas cultivadas. IBDF, Rio de Janeiro.
- Cutter, E.G., 1986. Anatomia Vegetal. Roca, São Paulo, pp. 304.
- Dando, I., Fiorini, C., Pozza, E.D., Padroni, C., Costanzo, C., Palmieri, M., Donadelli, M., 2013. UCP2 inhibition triggers ROS-dependent nuclear translocation of GAPDH and autophagic cell death in pancreatic adenocarcinoma cells. *Biochim. Biophys. Acta* 1833, 672–679.
- David, R., Carde, J.P., 1964. Coloration différentielle des inclusions lipidique et terpéniques des pseudophylles du pin maritime au moyen du réactif Nadi. *C. R. Acad. Sci. Paris* 258, 1338–1340.
- Delprete, P.G., Smith, L.B., Klein, R.M., 2005. Flora Ilustrada Catarinense: Rubiáceas. *Herbário Barbosa Rodrigues*, Itajaí.
- Deng, Y., Guan, M., Xie, X., Yang, X., Xiang, H., Li, H., Zou, L., Wei, J., Wang, D., Deng, X., 2013. Geniposide inhibits airway inflammation and hyperresponsiveness in a mouse model of asthma. *Int. Immunopharmacol.* 17, 561–567.
- Erbano, M., Duarte, M.R., 2010. Morfoanatomia de folha e caule de *Genipa americana* L. Rubiaceae. *Rev. Bras. Farmacogn.* 20, 825–832.
- Franceschi, V.R., Horner Junior, H.T., 1980. Calcium oxalate crystals in plants. *Bot. Rev.* 46, 361–427.
- Franceschi, V.R., Nakata, P.A., 2005. Calcium oxalate in plants: formation and function. *Annu. Rev. Plant Biol.* 56, 41–71.
- Frey-Wyssling, A., 1981. Crystallography of the two hydrates of crystalline calcium oxalate in plants. *Am. J. Bot.* 68, 130–141.
- Fu, Y., Liu, B., Liu, J., Liu, Z., Li, D., Li, F., Cao, Y., Zhang, X., Zhang, N., Yang, Z., 2012. Geniposide, from *Gardenia jasminoides* Ellis, inhibits the inflammatory response in the primary mouse macrophages and mouse models. *Int. Immunopharmacol.* 14, 792–798.
- Gabe, M., 1968. Techniques histologiques. Masson & Cie, Paris.
- Gao, C., Liu, Y., Jiang, Y., Ding, J., Li, L., 2014. Geniposide ameliorates learning memory deficits, reduces Tau phosphorylation and decreases apoptosis via GSK3b pathway in streptozotocin-induced Alzheimer rat model. *Brain Pathol.* 24, 261–269.
- González, W.L., Negritto, M.A., Suárez, L.H., Gianoli, E., 2008. Induction of glandular and non-glandular trichomes by damage in leaves of *Madia sativa* under contrasting water regimes. *Acta Oecol.* 33, 128–132.
- Haddad, A., Sessos, A., Attias, M., Farina, M., Nazareth, M.M., Silveira, M., Benchimol, M., Soares, M.J., Barth, M.O., Machado, D.R., Souto-Pradón, T., Souza, W., 1998. Técnicas básicas de microscopia eletrônica aplicadas às Ciências Biológicas. Sociedade Brasileira de Microscopia Eletrônica, Rio de Janeiro.
- Hallam, N.D., (PhD thesis) 1967. An electron microscope study of the leaf waxes of the genus *Eucalyptus* L'Heritier. University of Melbourne.
- Hammond, C., 2009. The Basics of Crystallography and Diffraction. Oxford University Press, New York, pp. 421.
- Harborne, J.B., 1998. Phytochemical Methods, 3rd ed. Chapman & Hall, London.
- He, H., Bleby, T.M., Veneklaas, E.J., Lambers, H., Kuo, J., 2012. Morphologies and elemental compositions of calcium crystals in phylloides and branchlets of *Acacia roborum* (Leguminosae: Mimosoideae). *Ann. Bot.* 109, 887–896.
- Hirata, T., Kobayashi, T., Wada, A., Ueda, T., Fujikawa, H., Miyashita, H., Ikeda, T., Tsukamoto, S., Nohara, T., 2011. Anti-obesity compounds in green leaves of *Eucommia ulmoides*. *Bioorg. Med. Chem. Lett.* 21, 1786–1791.
- Jeffrey, C.E., Baker, E.A., Holloway, P.J., 1975. Ultrastructure and recrystallisation of plant epicuticular waxes. *New Phytol.* 75, 539–549.
- Jensen, S.R., 1983. Iridoids in *Rothmannia globosa*. *Phytochemistry* 22, 1761–1765.
- Jensen, W.A., 1962. Botanical Histochemistry: Principles and Practice. W.H. Freeman & Co, San Francisco.
- Johansen, D.A., 1940. *Plant Microtechnique*. McGraw-Hill Book Company, New York.
- Kim, E.S., Jeong, C.S., Moon, A., 2012. Genipin, a constituent of *Gardenia jasminoides* Ellis, induces apoptosis and inhibits invasion in MDA-MB-231 breast cancer cells. *Oncol. Rep.* 27, 567–572.
- Kocsis, M., Darók, J., Borhidi, A., 2004. Comparative leaf anatomy and morphology of some neotropical *Rondeletia* (Rubiaceae) species. *Plant Syst. Evol.* 218, 205–218.
- Koriyama, Y., Takagi, Y., Chiba, K., Yamazaki, M., Sugitani, K., Arai, K., Suzuki, H., Kato, S., 2013. Requirement of retinoic acid receptor b for genipin derivative-induced optic nerve regeneration in adult rat retina. *PLoS ONE* 8, e71252.
- Kraus, J.E., Arduin, M., 1997. Manual básico de métodos em morfologia vegetal. Editora da Universidade Federal Rural do Rio de Janeiro, Seropédica.
- Li, C.C., Hsiang, C.Y., Lo, H.Y., Pai, F.T., Wu, S.L., Ho, T.Y., 2012. Genipin inhibits lipopolysaccharide-induced acute systemic inflammation in mice as evidenced by nuclear factor- κ B bioluminescent imaging guided transcriptomic analysis. *Food Chem. Toxicol.* 50, 2978–2986.
- Lin, Y.J., Lai, C.C., Lai, C.H., Sue, S.C., Lin, C.W., Hung, C.H., Lin, T.H., Hsu, W.Y., Huang, S.M., Hung, Y.L., et al., 2013. Inhibition of enterovirus 71 infections and viral IRES activity by *Fructus gardenia* and geniposide. *Eur. J. Med. Chem.* 62, 206–213.
- Mace, M.E., Howell, C.R., 1974. Histochemistry and identification of condensed tannin precursor in roots of cotton seedlings. *Can. J. Bot.* 52, 2423–2426.
- Martínez-Natarén, D.A., Tabla, V.P., Dzib, G., Irbién, L.M.C., 2011. Morphology and density of glandular trichomes in populations of Mexican oregano (*Lippia graveolens* H.B.K., Verbenaceae), and the relationship between trichome density and climate. *J. Torrey Bot. Soc.* 138, 134–144.
- Metcalfe, C.R., Chalk, L., 1979. Anatomy of the Dicotyledons, 2nd ed. Clarendon Press, Oxford, pp. 276.
- Monje, P.V., Baran, E.J., 2002. Characterization of calcium oxalates generated as biominerals in cacti. *Plant Physiol.* 128, 707–713.
- Moraes, T.M.S., Rabelo, G.R., Alexandrino, C.R., Silva Neto, S.J., Cunha, M., 2011. Comparative leaf anatomy and micromorphology of *Psychotria species* (Rubiaceae) from the Atlantic Rainforest. *Acta Bot. Brasil.* 25, 178–190.
- Nam, J.W., Kim, S.Y., Yoon, T., Lee, Y.J., Kil, Y.S., Lee, Y.S., Seo, E.K., 2013. Heat shock factor 1 inducers from the bark of *Eucommia ulmoides* as cytoprotective agents. *Chem. Biodivers.* 10, 1322–1327.
- Ono, M., Ishimatsu, N., Masuoka, C., Yoshimitsu, H., Tsuchihashi, R., Okawa, M., Kinjo, J., Ikeda, T., Nohara, T., 2007. Three new monoterpenoids from the fruit of *Genipa americana*. *Chem. Pharm. Bull.* 55, 632–634.
- Ono, M., Ueno, M., Masuoka, C., Ikeda, T., Nohara, T., 2005. Iridoid glucosides from the fruit of *Genipa americana*. *Chem. Pharm. Bull.* 53, 1342–1344.
- Pereira, Z.V., Meira, R.M.S.A., Azevedo, A.A., 2003. Morfoanatomia foliar de *Palaquium longepedunculata* Gardiner (Rubiaceae). *Rev. Arvore* 27, 759–767.
- Pino, J., Marbot, R., Vazquez, C., 2005. Volatile constituents of genipap (*Genipa americana* L.) fruit from Cuba. *Flavour Fragr. J.* 21, 488–586.
- Pinto, A.B., Guedes, C.M., Moreira, R.F.A., Maria, C.A.B., 2006. Volatile constituents from headspace and aqueous solution of genipap (*Genipa americana*) fruit isolated by the solid-phase extraction method. *Flavour Fragr. J.* 21, 488–491.
- Porto, R.G.C.L., Cardoso, B.V.S., Barros, N.V.A., Cunha, E.M.F., Araújo, M.A., Araújo, R.S.R.M., 2014. Chemical composition and antioxidant activity of *Genipa americana* L. (Jenipapo) of the Brazilian Cerrado. *J. Agric. Environ. Sci.* 3, 51–61.
- Quinteiro, M.M.C., Teixeira, D.C., Moraes, M.G., Silva, J.G., 2006. Anatomia foliar de *Psychotria viridis* Ruiz & Pav. (Rubiaceae). *Rev. Univ. Rural* 26, 30–41.
- Randau, K.P., Florêncio, D.C., Ferreira, C.P., Xavier, H.S., 2004. Estudo Farmacognóstico de *Croton rhamnifolius* H.B.K. e *Croton rhamnifolioides* Pax & Hoffm. (Euphorbiaceae). *Rev. Bras. Farmacogn.* 14, 89–96.
- Revilla, J., 2001. Plantas da Amazônia. Programa de desenvolvimento empresarial e tecnologia, Manaus.
- Rosa, E.A., Silva, B.C., Silva, F.M., Tanaka, C.M.A., Peralta, R.M., Oliveira, C.M.A., Kato, L., Ferreira, H.D., Silva, C.C., 2010. Flavonoides e atividade antioxidante em *Palaquium rigidum* Kunth, Rubiaceae. *Rev. Bras. Farmacogn.* 20, 484–488.
- Santos, B.F.S., Silva, L.C., Azevedo, A.A., Aguiar, R., 2006. Effects of simulated acid rain on leaf anatomy and micromorphology of *Genipa americana* L. (Rubiaceae). *Braz. Arch. Biol. Technol.* 49, 313–321.
- Song, J., Bi, H., Xie, X., Guo, J., Wang, X., Liu, D., 2013. Natural borneol enhances geniposide ocular absorption in rabbits. *Int. J. Pharm.* 445, 163–170.
- Souza, R.K.D., Mendonça, A.C.A.M., Silva, M.A.P., 2013. Ethnobotanical, phytochemical and pharmacological aspects Rubiaceae species in Brazil. *Rev. Cubana Plant Med.* 18, 140–156.
- Tian, J.S., Cui, Y.L., Hu, L.M., Gao, S., Chi, W., Dong, T.J., Liu, L.P., 2010. Antidepressant-like effect of genipin in mice. *Neurosci. Lett.* 479, 236–239.
- Ueda, S., Iwahashi, Y., 1991. Production of anti-tumor-promoting iridoid glucosides in *Genipa americana* and its cell cultures. *J. Nat. Prod.* 54, 1677–1680.
- Venditti, A., Serrilli, A.M., Bianco, A., 2013. Iridoids from *Bellardia trixago* (L.) all. Nat. Prod. Res. 27, 1413–1416.
- Viljoen, A.M., Subramoney, S., Van Vuuren, S.F., Başer, K.H., Demirci, B., 2005. The composition, geographical variation and antimicrobial activity of *Lippia javanica* (Verbenaceae) leaf essential oils. *J. Ethnopharmacol.* 96, 271–277.
- Wagner, H., Blatt, S., 1996. *Plant Drug Analysis: A Thin Layer Chromatography Atlas*. Spinger-Verlag, Berlin.
- Wang, Q.S., Xiang, Y., Cui, Y.L., Lin, N.K.M., Zhang, X.F., 2012. Dietary blue pigments derived from genipin, attenuate inflammation by inhibition of LPS-induced iNOS and COX-2 expression via NF- κ B inactivation. *PLoS ONE* 7, e34122.
- Yang, X., Yao, J., Luo, Y., Han, Y., Wang, Z., Du, L., 2013. P38 MAP kinase mediates apoptosis after genipin treatment in non-small-cell lung cancer H1299 cells via a mitochondrial apoptotic cascade. *J. Pharmacol. Sci.* 121, 272–281.

- Yoder, L.R., Mahlberg, P.G., 1976. Reactions of alkaloid and histochemical indicators in laticifers and specialized parenchyma cells of *Catharanthus roseus* (Apocynaceae). *Am. J. Bot.* 63, 1167–1173.
- Zappi, D., 2015. Genipa in Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro, Electronic resource: <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB14045> (accessed March 2016).
- Zhang, G., He, J.L., Xie, X.Y., Yu, C., 2012. LPS-induced iNOS expression in N9 microglial cells is suppressed by geniposide via ERK, p38 and nuclear factor- κ B signaling pathways. *Int. J. Mol. Med.* 30, 561–568.
- Zhang, Q., Su, Y.Q., Zhang, J.F., 2013. Seasonal variation in antioxidant capacity and active compounds contents of *Eucommia ulmoides* Oliver leaf. *Molecules* 18, 1868.
- Zini, A.S., Martins, S., Toderke, M.L., Temponi, L.G., 2016. Anatomia foliar de Rubiaceae ocorrentes em fragmento florestal urbano de Mata Atlântica, PR, Brasil. *Hoehnea* 43, 173–182.