

## CORRECTION

In “A randomized trial of exercise treadmill ECG versus stress SPECT myocardial perfusion imaging as an initial diagnostic strategy in stable patients with chest pain and suspected CAD: Cost-analysis” (J Nucl Cardiol 2007;14:174-186), there were errors in Table 1. The table should not have included a listing for Blood pressure mmol/L; this error created the incorrect measurements for all subsequent values. Please see the table below as it should have printed.

**Table 1.** Patient characteristics

<b>Variable</b>	<b>ETT</b>	<b>MPI</b>
No.	207	250
Age (y) (SD)	58.9 (11.4)	59.7 (12.2)
Male gender [% (No.)]	57.5 (119)	55.6 (139)
White [% (No.)]	46.9 (97)	55.6 (139)
Hypertension [% (No.)]	46.3 (96)	53.2 (133)
Nonsmokers [% (No.)]	83.6 (173)	87.2 (218)
Diabetes [% (No.)]	14.5 (30)	19.2 (48)
Family history of CAD [% (No.)]	46.3 (96)	43.2 (108)
Blood pressure (mmHg)	132/80	140/85
Cholesterol (mmol/L) [mean (SD)]	5.81 (1.05)	5.79 (1.79)
Cholesterol (mg/dL) [mean (SD)]	225 (41)	224 (69)
HDL cholesterol (mmol/L) [mean (SD)]	1.14 (0.29)	1.22 (0.33)
HDL cholesterol (mg/dL) [mean (SD)]	44 (11)	47 (13)
Triglycerides (mmol/L) [mean (SD)]	2.11 (1.34)	2.01 (1.12)
Triglycerides (mg/dL) [mean (SD)]	82 (52)	78 (43)
BMI (kg/m <sup>2</sup> ) [mean (SD)]	27.6 (4.6)	26.9 (4.5)
Framingham score [mean (SD)]	17 (11)	17 (9)

*HDL*, High-density lipoprotein cholesterol; *BMI*, body mass index.

## CORRECTION

In *Abstracts of Original Contributions, CT Angiography for the Cardiologist, April 12-14, 2007, Chicago, IL* (J Nucl Cardiol 2007;14:S93), Abstract 20.0 “Costs and clinical outcomes in individuals undergoing coronary computed tomography versus myocardial perfusion imaging,” was not presented, and therefore, should have been withdrawn.