- Rouse B. M. (1966) Phenylalanine deficiency syndrome. *J Paediatr* **69**: 246–249.
- Scriber CR (1988) We mean well: treatment of Mendelian disease. *Acta Paediatr Jpn* **30**: 472–480.
- Smith I, Beasley MG, Ades AE (1990) Intelligence and quality of dietary treatment in phenylketonuria. *Arch Dis Child* **65**: 472–478.
- Van Spronsen FJ, Berger R, Smit GPA, et al (1989) Tyrosinaemia type I: orthotopic liver transplantation as the only definitive answer to a metabolic as well as an oncological problem. *J Inherit Metab Dis* **12**: 339–342.
- Van Wyk KG, Clayton PT (1997) Dietary management in tyrosinaemia type 1. In: International Metabolic Dietitians Group, 2nd Dietitians meeting SSIEM, Gothenburg, Sweden.

Erratum

Skladal D, Grissenauer G, Konstantopoulou V, Felker S, Sperl W (2000) High plasma leucine concentrations without neurological symptoms in a patient with classical maple syrup urine disease. *J Inherit Metab Dis* **23**(5): 513–514.

It has been found that the amino acid analyser used in this study systematically overestimated plasma leucine at high concentrations. The concentration reported as 2249 $\mu\text{mol/L}$ had a true value of 1430 $\mu\text{mol/L}$ and leucine values reported as $>2000 \mu\text{mol/L}$ were approximately 1500 $\mu\text{mol/L}$.