Erratum

Root competition for phosphorus between coconut, multipurpose trees and kacholam (*Kaempferia galanga* L.) in Kerala, India

S. S. KUMAR¹, B. M. KUMAR¹, P. A. WAHID², N. V. KAMALAM³ and R. F. FISHER^{4, *}

¹ College of Forestry, Kerala Agricultural University, Vellanikkara, Thrissur 680 654, India;

² College of Agriculture, Padanakkad, 671 328, Kerala, India; ³ Radiotracer Laboratory, Kerala Agricultural University, Vellanikkara, Thrissur 680 654, India; ⁴ Department of Forest Sciences, Texas A&M University, College Station, TX 77843-2135, USA (*Author for correspondence)

Agroforestry Systems 46: 131–146, 1999.

On pages 134, 136 and 140 respectively of the above article, the correct figures should be as follows.

Figure 1. Experimental units involving coconut palms, multipurpose trees and kacholam (not to scale). A – single hedge and B – double hedge system of multipurpose trees.

Figure 2. Schematic diagram (not to scale) showing a radio-phosphorus applied coconut basin.

Figure 3. Absorption of ^{32}P by three associated multipurpose trees at different lateral distances from the root zones of coconut palms in Kerala, India (cpm = counts per minute per g foliage dry weight).