

ERRATUM

In the article entitled "An Experimental Study of the Neurogenic and the Immunological Contribution to 'Tennis Elbow' in Rats," by Eva Haker, Elvar Theodorsson, and Thomas Lundeberg (*Inflammation*, Volume 21, Number 1, pp. 35–44) Table 1 should read:

Table 1. Concentrations of Substance P-, Neurokinin A-, Calcitonin Gene-Related Peptide- and Neuropeptide Y-LI in Cerebrospinal Fluid, Plasma and Synovial Fluid.

Post injection	CSF	Plasma	Perfusate R	Perfusate L	CSF	Plasma	Perfusate R	Perfusate L
	Neurokinin A-LI							
Saline								
2 h	<0.1	<0.1	<0.1	<0.1	<0.1	11.6 ± 3.1	<0.1	<0.1
6 h	<0.1	<0.1	<0.1	<0.1	<0.1	12.2 ± 3.4	<0.1	<0.1
24 h	<0.1	<0.1	<0.1	<0.1	<0.1	9.1 ± 2.6	<0.1	<0.1
Substance P								
2 h	29.3 ± 6.2 ^a	<0.1	9.2 ± 2.5 ^b	5.5 ± 2.8	59.2 ± 35.2 ^a	<0.1 ^c	5.4 ± 1.2 ^b	5.1 ± 3.4
6 h	35.2 ± 21.3	<0.1	3.1 ± 0.9 ^b	8.1 ± 4.9	65.2 ± 39.7 ^a	<0.1 ^c	5.1 ± 0.9 ^b	4.6 ± 2.9
24 h	12.4 ± 11.3	<0.1	6.9 ± 2.1 ^a	<0.1	95.8 ± 47.6 ^a	<0.1 ^c	4.7 ± 1.1 ^b	<0.1
hr Interleukin-1 α								
2 h	14.4 ± 4.7 ^a	<0.1	9.4 ± 2.2 ^a	4.7 ± 1.6 ^a	15.7 ± 5.8 ^a	<0.1 ^c	18.6 ± 3.1 ^a	5.6 ± 4.4
6 h	13.9 ± 3.2 ^a	<0.1	5.3 ± 1.9 ^a	4.2 ± 1.9	19.6 ± 11.3	<0.1 ^c	5.6 ± 1.4 ^a	3.9 ± 2.2
24 h	5.9 ± 4.8	<0.1	<0.1	<0.1	53.2 ± 29.6	<0.1 ^c	5.5 ± 0.7	6.1 ± 4.7
	Calcitonin gene-related peptide-LI							
Saline								
2 h	7.8 ± 3.6	7.9 ± 2.4	<0.1	<0.1	2231.5 ± 499.2	535.1 ± 49.2	17.2 ± 2.4	14.3 ± 3.6
6 h	12.5 ± 4.7	8.7 ± 3.1	<0.1	<0.1	1935.2 ± 487.6	476.4 ± 93.6	18.6 ± 3.1	15.6 ± 6.7
24 h	11.7 ± 5.1	9.5 ± 3.2	<0.1	<0.1	2214.3 ± 512.9	547.5 ± 102.1	17.5 ± 1.9	12.2 ± 1.9
Substance P								
2 h	48.9 ± 9.6 ^a	22.3 ± 4.9 ^a	4.5 ± 1.2 ^a	4.9 ± 6.1	2982.5 ± 936.5	839.7 ± 212.4	52.2 ± 11.6	45.3 ± 7.3
6 h	35.4 ± 11.2 ^a	11.5 ± 4.7	4.9 ± 0.7 ^b	5.5 ± 3.7	2476.8 ± 537.5	639.5 ± 161.2	47.2 ± 3.5 ^c	46.9 ± 9.8
24 h	13.2 ± 7.5	10.6 ± 5.2	5.3 ± 1.4 ^b	5.3 ± 1.7	2523.4 ± 698.4	529.4 ± 99.4	33.5 ± 3.4 ^c	45.7 ± 16.2
hr Interleukin-1 α								
2 h	29.8 ± 6.1 ^a	19.3 ± 2.2 ^a	4.7 ± 0.7 ^b	4.9 ± 1.2	2437.0 ± 372.8	639.2 ± 96.3	68.7 ± 9.2 ^b	43.6 ± 9.1
6 h	31.5 ± 9.9 ^a	10.7 ± 1.3	4.5 ± 0.9 ^b	5.6 ± 2.4	2743.9 ± 321.5 ^a	1304.5 ± 312.8	39.9 ± 8.6 ^a	29.2 ± 9.7
24 h	15.5 ± 10.1	12.4 ± 3.3	4.9 ± 1.3 ^b	11.3 ± 9.9	2956.6 ± 299.4 ^a	598.6 ± 221.8	35.5 ± 14.2	24.4 ± 6.1

Concentrations are presented in fmol/ml as mean ± SEM. Statistical analysis was carried out by comparing the neuropeptide-LI in the cerebrospinal fluid, plasma and perfusates of rats treated with pro-inflammatory substances intra-ECRB with the respective saline-injected control group. Number of animals in each group 9–10. Significant differences from control groups: ^a $P < 0.05$, ^b $P < 0.01$, ^c $P < 0.001$.