This week in therapeutics

Indication	Target/marker/ pathway	Summary	Licensing status	Publication and contact information
Cancer				
Hepatocellular carcinoma (HCC)	<i>Interleukin-6 signal transducer (IL6ST; GP130); catenin (cadherin-associated protein), β1, 88kDa (CTNNB1; β-catenin)</i>	Transcriptional analysis of human liver tissue suggests that mutations in the <i>IL6ST</i> gene could help predict susceptibility to HCC. In benign inflammatory hepatocellular adenomas capable of converting to malignant HCC, there was greater activation of proinflammatory IL-6 signaling compared with that seen in noncancerous hepatocytes. In 60% of those benign adenomas, mutations in <i>IL6ST</i> , which encodes GP130, were associated with constitutive activation of signal transducer and activator of transcription 3 (STAT3), which can potentially contribute to tumorigenesis. No adenomas transformed into malignant GP130-mutated cells without an accompanying β -catenin mutation. Next steps include validating the association between <i>IL6ST</i> mutations and the inflammatory hepatocellular tumor phenotype in a larger patient cohort. Reata Pharmaceuticals Inc.'s RTA 402, a synthetic triterpenoid that inhibits the transcriptional activity of NF- κ B and STAT3, is in Phase II testing to treat cancer.	Unpatented; licensing status not applicable	Rebouissou, S. <i>et al. Nature</i> ; published online Nov. 19, 2008; doi:10.1038/nature07475 Contact: Jessica Zucman-Rossi, Institut National de la Santé et de la Recherche Médicale (INSERM), Paris, France e-mail: zucman@cephb.fr
SciBX 1(43); doi:10.1038/scibx.2008.1044 Published online Dec. 4, 2008				