The cross-talking hedgehog

By Michael J. Haas, Senior Writer

For about a decade, abnormal activation of the hedgehog (Hh) signaling pathway has been thought to play a role in tumorigenesis, possibly mediated by the tumor suppressor p53, but the molecular mechanisms linking Hh to cancer remained elusive. Newly reported *in vitro* studies in the *Proceedings of the National Academy of Sciences* now have shown a signaling connection between Hh and p53-induced apoptosis, thereby establishing the first molecular pathway between Hh and tumor survival and growth.

Though it remains to be seen how extensive this pathway is in human cancers, the findings could offer new cancer targets and provide markers for predicting whether existing therapies are likely to be effective. Until now, exploiting suggestive evidence about the Hh signaling path has proved unfruitful for drug developers (see **Figure 1**, “The elusive hedgehog”).

In the *PNAS* paper, researchers at **Nippon Medical School**, the **National Cancer Center Research Institute** in Tokyo and the **Weizmann Institute of Science** show that mutations along the Hh signaling pathway could enable tumorigenesis by inhibiting p53-induced apoptosis and accelerating cell proliferation.¹ The team was led by Nobuyuki Tanaka, professor of molecular oncology at Nippon Medical School.

“Not a lot has been known about the relationship between hedgehog signaling and p53,” said Margaret Read, senior director of cancer biology at **Infinity Pharmaceuticals Inc.** “This paper appears to be the first example detailing Hh-p53 cross-talk at the cell-signaling level.”

Hh is an extracellular signaling molecule in a pathway that regulates embryonic development in vertebrates and may also regulate maintenance and regeneration of adult stem cells. Hh interacts with the transmembrane protein Patched, a negative regulator of the signal transducer Smoothened (Smo), resulting in derepression of Smo and activation of its downstream transcriptional program(s).

In mouse fibroblast cells, Tanaka’s team found that introduction of gain-of-function mutant Smo increased the activity of a phosphoinositide 3-kinase/protein kinase B (PI3K/Akt) pathway, which appeared to upregulate a pair of transcription factors—glioma-associated oncogene homologs 1 and 2 (Gli1 and Gli2). These two transcription factors in turn activated unknown MDM2-phosphorylating kinases, which led to degradation of p53 and thus partial inhibition of p53-induced apoptosis.

In mouse embryonic fibroblasts, mutant Smo also accelerated cell proliferation by enhancing the expression of cyclin D1 and cyclin E.

Taken together, the results indicate that Smo simultaneously activates two pathways that enable a nascent tumor to survive and grow (see **Figure 2**, “Hedgehog-mediated suppression of p53-induced apoptosis”).

Hh signaling

The consensus among industry and academic scientists contacted by SciBX is that the results could offer new targets or diagnostic markers, although more work is needed to confirm the findings.

“How the hedgehog pathway initiates cancer has not been clear,” said James Chen, assistant professor of chemical and systems biology at **Stanford University**. “This paper is a good example that shows how a known tumor suppressor [p53] is lost when the Hh pathway is turned on. It provides a molecular explanation for how Hh *might* promote tumorigenesis in multiple cancers.”

But Chen also said Hh signaling is probably not completely responsible for the loss of p53 in all tumors.

“In mouse models, deletion of Patched results in tumors in only about 15% of the mice,” he said. “But with both Patched and p53 deletion, 100% of the mice get tumors. In other words, the genetic loss of p53 is a stronger factor than turning on Hh. This suggests either only partial p53 inhibition by Hh signaling, or that the effect is tumor dependent.”

In either case, he said, “Hh signaling alone is not sufficient for complete inhibition of p53 in all cancers. Other oncogenic factors besides Hh signaling are probably involved.”

Chen also stressed that the findings of Tanaka’s team aren’t conclusive. The study laid out the p53 end of the pathway well, “but how this connects to Hh on the molecular level is not clearly established,” he said.

For example, he noted the researchers showed that cyclopamine—a known Smo inhibitor—reversed the loss of p53 function caused by mutant Smo, but did not show whether cyclopamine had this same effect if molecules downstream from Smo on the Hh-p53 pathway were overexpressed.

He said a good control would have been to induce expression of Gli1 and Gli2 and observe whether cyclopamine could still restore p53 function.

Nevertheless, Chen does think the findings imply that Hh signaling is involved in tumor maintenance, even where it is not required for tumor initiation. “If Hh signaling can downregulate p53 function, then other signaling and damage events that might trigger p53-mediated apoptosis or growth arrest could go unchecked, resulting in tumorigenesis,” he said.

Thus, Chen said, the Hh pathway could present a broad therapeutic target if Hh signaling is required for the maintenance of a wide range of cancers. “There is evidence in the literature that several cancers—of the stomach, pancreas, esophagus, lung, breast, blood and others—may require Hh signaling for their proliferation and maintenance,” he added.

Infinity’s Read noted the findings in *PNAS* could lead to treatments that increase the effectiveness of chemotherapy.

Figure 1. The elusive hedgehog. Since the *hedgehog* gene was discovered more than a quarter-century ago, efforts to exploit hedgehog signaling in a number of therapeutic settings have yet to produce commercially useful results.

"p53 is often sensitive to toxic agents, but it is deregulated in many types of tumors," Read said, explaining that this can decrease the apoptotic response of cancer cells to chemotherapy. She speculated that Hh inhibition might render tumors with nonmutant p53 more sensitive to chemotherapy by restoring normal p53 function. Not much is known about how effective such combination therapy would be, "but it's worth exploring," she said.

Infinity's lead cancer compound, IP-926, is a derivative of cyclopamine that operates upstream of both the Gli1 and Gli2 and the cyclin D1 and cyclin E pathways. The company plans to submit an IND this year.

According to Read, the findings also indicate that "targeting Gli1 and Gli2 for cancer is definitely one approach." She cited recent work by Rolf Toftgård and colleagues at the **Karolinska Institute** that implicated the two transcription factors in tumor development and metastasis.²

However, Read also wanted to see more work establishing the role of Gli1 and Gli2 along the Hh-p53 pathway.

"In their [PNAS] study, forced expression of Gli1 and Gli2 is shown to downregulate p53. But they could have silenced Gli with siRNA to show whether interfering with Gli re-established p53 function," she said.

Read also said the findings did not firmly establish that the path from Smo to p53 ran through Gli1 and Gli2. The researchers showed that PI3K inhibition restored p53 function, but not whether PI3K inhibition did so by downregulating Gli1 and Gli2, she said. This is important, noted Read, because the Hh-p53 pathway proposed in the PNAS paper depends on Gli1- and Gli2-mediated transcription and synthesis of MDM2-phosphorylating kinases.

"It would have been nice to see them close that loop," Read said.

She added: "The main point of interest is the demonstration of a signaling connection between Hh and p53. Whether Hh-mediated downregulation of p53 via MDM2 really 'overrides p53-mediated tumor suppression', as the title proposes, remains to be seen."

Lance Leopold, CMO of **Ascenta Therapeutics Inc.**, was less concerned about the missing puzzle pieces in the mechanism by which Hh inhibits p53. Instead, he wants to see the findings expanded.

"The paper helps connect the Hh and cancer-related pathways by linking relevant genes to molecules in cell proliferation and cell death pathways," he said. "Next we need a broader survey of human tumor types, to see how extensive Hh signaling is in human cancer."

Ascenta's lead cancer compound is AT-219, a small-molecule inhibitor of HDM2 (the human form of MDM2) that is in preclinical development.

Leopold hypothesized that markers along the Hh pathway could be used to predict responses to cancer therapies in preclinical and clinical studies by distinguishing between cancers that express wild-type p53 and mutant p53. In the former case, HDM2 inhibition could be expected to restore p53's tumor-suppressing functions by decreas-

Figure 2. Hedgehog-mediated suppression of p53-induced apoptosis. A mutant form of the hedgehog signal transducer Smoothened (Smo) was found to simultaneously activate two pathways that enable a nascent tumor to survive and grow.

[a] In one pathway, mutant Smo activated a PI3K/Akt pathway that is thought to upregulate glioma-associated oncogene homologs 1 and 2 (Gli1 and Gli2). Gli1 and Gli2 increase the expression of kinases that induce phosphorylation of MDM2, increasing MDM2 binding of p53. Ubiquitin binds the MDM2-p53 complex, labeling it for degradation. This results in partial inhibition of the accumulation of p53 and of p53-induced apoptosis and senescence, allowing the nascent tumor to survive and grow.

[b] Additionally, mutant Smo was found to enhance expression of cyclin D and cyclin E, resulting in cell proliferation, again allowing the nascent tumor to survive and grow.

ing the HDM2-p53 binding that leads to p53 degradation. In the latter case, HDM2 inhibition would likely not have this restorative effect.

Existing methods for distinguishing wild-type and mutant p53 have multiple drawbacks, he said, including significant false-positive and false-negative rates and the inability to identify the mutations that actually alter the function of p53.

The connection of Hh signaling to p53 inhibition could provide better markers than existing methods, Leopold said. "This could be useful for patient selection in clinical trials, where you want to enrich for patients who could benefit from the therapy and avoid treating patients who are unlikely to respond" because their cancer involves nonfunctional p53, he said.

He also said the findings of Tanaka's team validate Ascenta's strategy of targeting molecules that are downstream of epithelial and epidermal growth factors on tumorigenic pathways.

"Targeting the upstream pathway has been effective for VEGF, ErbB1, and Abl, for example, though we know this may select for mutants that are drug resistant—at least in the case of Gleevec and Tarceva," Leopold said. "So, targeting the upstream cascade could be trumped by downstream mutations" in molecules such as Smo, he said. Thus it makes "additional sense" to target the HDM2-p53 interaction at the end of the cascade, possibly in combination with other Hh pathway inhibitors, he said.

Novartis AG markets Gleevec, a Bcr-Abl tyrosine kinase inhibitor, to treat chronic myelogenous leukemia and gastrointestinal stromal tumors. Tarceva, a small-molecule inhibitor of epidermal growth factor receptor 1 (HER1), is approved to treat non-small cell lung cancer and

pancreatic cancer. Tarceva is marketed by **OSI Pharmaceuticals Inc.** and **Genentech Inc.** in the U.S. and by **Roche** elsewhere.

Going forward, Tanaka told *SciBX* his research team is working to identify the kinases that activate MDM2 and plans to study the role of Hh signaling on cancer development *in vivo*.

"I would like to study transgenic mice expressing mutant Smo or Hh itself in stomach or other tissues, and also the role of hedgehog signaling in chronic inflammation-induced cancer in a mouse model," Tanaka said. "If we also identify the kinases or signaling molecules of Gli-mediated MDM2 activation, we could analyze the protective effect of blocking Gli signaling *in vivo*."

Tanaka said the findings reported in *PNAS* are not patented.

REFERENCES

1. Abe, Y. *et al. Proc. Natl. Acad. Sci. USA*; published online March 10, 2008; doi:10.1073/pnas.0712216105
Contact: Nobuyuki Tanaka, Nippon Medical School, Kawasaki, Kanagawa, Japan
e-mail: nobuta@nms.ac.jp
2. Lauth, M. & Toftgård, R. *Cell Cycle* **6**, 2458–2464 (2007)

COMPANIES & INSTITUTIONS MENTIONED

Ascenta Therapeutics Inc., Malvern, Pa.
Genentech Inc. (NYSE:DNA), South San Francisco, Calif.
Infinity Pharmaceuticals Inc. (NASDAQ:INFI), Cambridge, Mass.
Karolinska Institute, Huddinge, Sweden
National Cancer Center Research Institute, Tokyo, Japan
Nippon Medical School, Kanagawa, Japan
Novartis AG (NYSE:NVS; SWX:NOVN), Basel, Switzerland
OSI Pharmaceuticals Inc. (NASDAQ:OSIP), Melville, N.Y.
Roche (SWX:ROG), Basel, Switzerland
Stanford University, Stanford, Calif.
Weizmann Institute of Science, Rehovot, Israel