Armed & dangerous

By Tim Fulmer, Senior Writer

Researchers at the **Burnham Institute for Medical Research** are taking a second shot at exploiting *Salmonella*'s tumor-homing ability to develop cancer therapies. In work reported in the *Journal of the National Cancer Institute*, they armed an attenuated strain of *Salmonella* with a proapoptotic cytokine and showed the strain's therapeutic activity in mouse cancer models.¹ But given the failure of similar strategies in the past, it remains to be seen whether any companies will pick up the baton and in-license the revamped technology.

It has been known for over a century that certain facultative anaerobic bacteria—bacteria capable of growing with or without oxygen—are able to preferentially home in on tumors when injected intravenously, establish themselves, replicate and inhibit tumor growth.²

Although the use of *Salmonella* vectors to treat cancer is enticing, the only biotech to have taken them into the clinic, **Vion Pharmaceuticals Inc.**, stopped those programs about five years ago due to minimal efficacy in Phase I trials.

The Burnham researchers, led by John Reed, president and CEO of the institute, used an attenuated strain of *S. typhimurium*, one of the better-known tumor-targeting bacteria, to deliver the proapoptotic cytokine Fas ligand (TNF superfamily, member 6) (FasL; CD95L) to solid tumors in mice. The goal of the study was to see whether bacteria could deliver toxic payloads to tumors and whether the inherent tumor-inhibiting ability of *Salmonella* could be enhanced this way.

The Burnham group had previously used the same attenuated *S. typhimurium* strain as a tumor-targeting vector to deliver the immunostimulatory cytokine LIGHT to tumors in mice.³ In that instance the goal had been to trigger an immune response specific to the targeted tumor.

In the latest study, i.v. administration of the attenuated *Salmonella* expressing FasL significantly lowered tumor volume compared with what was seen using saline control in murine models of breast and colon cancer ($p=0.02$ and $p<0.001$, respectively). In the breast cancer models, FasL-expressing *Salmonella* also substantially minimized tumor metastases to the lung compared with saline control ($p=0.001$).

"Our strategy has been to take an attenuated *Salmonella* strain that previously showed safety and tumor-targeting ability in humans and to arm that strain with a gene that expresses an immunostimulatory or proapoptotic ligand in order to enhance its antitumor activity," Reed told *SciBX*.

Indeed, the original selling points of using bacterial vectors instead of other delivery vehicles have not changed. "One potential advantage of using a bacterial vector compared to using inert materials like liposomes and micelles is that bacteria will respond to the appropriate

antibiotics and can thus be turned off relatively quickly in the event toxicity appears," Reed noted.

Christopher Black, executive director of preclinical development at **Celsion Corp.**, agreed. Bacterial vectors "give unique control to the treatment since the toxicity associated with conventional anticancer drugs cannot be reversed once the drug is administered and depends on normal drug excretion pathways to lower a toxic dose," he said.

Celsion's ThermoDox—doxorubicin encapsulated in a heat-activated liposome—is in Phase III trials to treat nonresectable hepatocellular carcinoma. ThermoDox liposomes release chemotherapeutics only when mild hyperthermia is applied to the tumor, thus keeping the active drug confined to the malignant cells.

Another potential advantage of using attenuated *Salmonella* is "the bacterium's capacity for sustained proliferation under hypoxic conditions that often characterize larger tumors," said Douglass Thamm, professor of oncology in the Animal Cancer Center at **Colorado State University**.

Going forward, Thamm said he wants to see data in larger animals and whether the approach "is extendable to multiple transgenes. For example, a proapoptotic ligand like FasL could perhaps be expressed in the same *Salmonella* strain with an immunostimulatory ligand such as IL-2 or GM-CSF [granulocyte/macrophage colony-stimulating factor]. In this way, you not only trigger apoptosis in the tumor but also simultaneously stimulate the host immune system."

Thamm and colleagues have shown that the unarmed version of the same attenuated *Salmonella* strain used by the Burnham team was safe in tumor-bearing dogs. In those studies, 15% of the animals showed an antitumor response, with fever and vomiting being the only dose-limiting toxicities observed.⁴

According to Reed, his group is interested in combining LIGHT and FasL in the same *Salmonella* strain and characterizing it in a number of rodent cancer models. "More generally, we predict that the same bacterial strain could be engineered to express multiple ligands that activate parallel mechanisms in the tumor or tumor microenvironment that contribute to tumor regression," he added.

"It will be important to carry out toxicity studies on the FasL-expressing *Salmonella* strain, and we hope to do so in the near term using primate models and thus prepare the way for an IND application a few years down the road," Reed told *SciBX*.

Blast from the past

The attenuated *Salmonella* strain used by the Reed and Thamm labs was developed by researchers at **Yale University** more than 10 years ago and licensed to Vion.

Based on initial reports of the strain's antitumor activity in pre-clinical animal models,⁵ the company decided to run Phase I trials in patients with advanced solid tumors. The unarmed strain (designated as VNP20009) was safe, but showed minimal efficacy.^{6,7} As a result, the company announced in 2003 it was stopping development.

"The *JNCI* paper illustrates a potential way to improve the efficacy of an attenuated *Salmonella* strain whose safety has already been shown in humans," said John Pawelek, one of the scientists who originally

designed the strain at Yale. “Nevertheless, the study is in mice and therefore raises the question of how well the new strategy addresses the basic problem we encountered—namely, a *Salmonella* strain with good tumor-targeting efficacy in mice failed to turn up in great abundance in human tumors.”

Pawelek is a research affiliate in the Department of Dermatology at the Yale School of Medicine.

Celsion's Black said another potential concern is what happens to *Salmonella* that don't accumulate in the tumor. “Are they being destroyed in the liver, spleen or elsewhere?” he asked.

If the bacteria continue to live, addition of a FASL transgene might induce local accumulation of T cells and cytokines in normal tissues, which could promote unwanted inflammation, according to Black.

In one of Vion's Phase I trials in 24 patients with metastatic melanoma and renal cell carcinoma, the unarmed *Salmonella* strain elicited dose-related increases in circulating proinflammatory cytokines such as IL-1 β and tumor necrosis factor- α (TNF- α). At that time, the authors speculated that the systemic release of these cytokines was likely associated with dose-limiting toxicities that included fever, hypotension, anemia, nausea and diarrhea.⁶

In the *JNCI* paper, histological analysis of mouse lungs, kidneys, spleen and heart showed minimal toxicity associated with the FASL-expressing *Salmonella*.

The Burnham Institute told *SciBX* that the technology described in the *JNCI* paper is available for licensing.

REFERENCES

1. Loeffler, M. *et al.* *J. Natl. Canc. Inst.*; published online Aug. 6, 2008; doi:10.1093/jnci/djn205
Contact: John Reed, Burnham Institute for Medical Research, La Jolla, Calif.
e-mail: reedoffice@burnham.org
2. Chakrabarty, A. J. *Bacteriol.* **185**, 2683–2686 (2003)
3. Loeffler, M. *et al.* *Proc. Natl. Acad. Sci. USA* **104**, 12879–12883 (2007)
4. Thamm, D. *et al.* *Clin. Cancer Res.* **11**, 4827–4834 (2005)
5. Pawelek, J. *et al.* *Cancer Res.* **57**, 4537–4544 (1997)
6. Toso, J. *et al.* *J. Clin. Oncol.* **20**, 142–152 (2002)
7. Heimann, D. & Rosenberg, S. J. *Immunother.* **26**, 179–80 (2003)

COMPANIES AND INSTITUTIONS MENTIONED

Burnham Institute for Medical Research, La Jolla, Calif.

Celsion Corp. (NASDAQ:CLSN), Columbia, Md.

Colorado State University, Fort Collins, Colo.

Vion Pharmaceuticals Inc. (NASDAQ:VION), New Haven, Conn.

Yale University, New Haven, Conn.