This week in techniques

Approach	Summary	Licensing status	Publication and contact information
Assays & screens			
Screen of sequentially administered cancer therapeutics to identify new combination therapies	A screen of drug combinations could help identify new therapeutic combinations to help treat cancer. A cellular screen of combinations of sequentially dosed DNA-damaging agents showed that Tarceva erlotinib given four hours before doxorubicin led to a fivefold increase in apoptosis in a triple-negative breast cancer cell line compared with the combination dosed simultaneously. In mice, the timed combination decreased tumor growth compared with coadministration. Next steps include designing a clinical trial to test the sequential combination treatment and using the screen to look for other sequential combination therapies to treat cancer. Tarceva, an epidermal growth factor receptor (EGFR) inhibitor, is marketed by Astellas Pharma Inc., Chugai Pharmaceutical Co. Ltd. and Roche for non-small cell lung cancer (NSCLC) and pancreatic cancer. SciBX 5(22); doi:10.1038/scibx.2012.583 Published online May 31, 2012	Patent application filed; available for licensing	Lee, M.J. <i>et al. Cell</i>; published online May 11, 2012; doi:10.1016/j.cell.2012.03.031 Contact: Michael B. Yaffe, Massachusetts Institute of Technology, Cambridge, Mass. e-mail: myaffe@mit.edu