

ARTICLE

Received 30 Sep 2014 | Accepted 25 Feb 2015 | Published 10 Apr 2015

DOI: 10.1038/ncomms7771

Roles of lymphatic endothelial cells expressing peripheral tissue antigens in CD4 T-cell tolerance induction

Sherin J. Rouhani¹, Jacob D. Eccles¹, Priscila Riccardi¹, J. David Peske¹, Eric F. Tewalt¹, Jarish N. Cohen¹, Roland Liblau^{2,3,4,5}, Taija Mäkinen⁶ & Victor H. Engelhard¹

Lymphatic endothelial cells (LECs) directly express peripheral tissue antigens and induce CD8 T-cell deletional tolerance. LECs express MHC-II molecules, suggesting they might also tolerize CD4 T cells. We demonstrate that when β -galactosidase (β -gal) is expressed in LECs, β -gal-specific CD8 T cells undergo deletion via the PD-1/PD-L1 and LAG-3/MHC-II pathways. In contrast, LECs do not present endogenous β -gal in the context of MHC-II molecules to β -gal-specific CD4 T cells. Lack of presentation is independent of antigen localization, as membrane-bound haemagglutinin and I-E α are also not presented by MHC-II molecules. LECs express invariant chain and cathepsin L, but not H2-M, suggesting that they cannot load endogenous antigenic peptides onto MHC-II molecules. Importantly, LECs transfer β -gal to dendritic cells, which subsequently present it to induce CD4 T-cell anergy. Therefore, LECs serve as an antigen reservoir for CD4 T-cell tolerance, and MHC-II molecules on LECs are used to induce CD8 T-cell tolerance via LAG-3.

¹ Carter Immunology Center, Department of Microbiology, Immunology and Cancer Biology, University of Virginia School of Medicine, Charlottesville, Virginia 22908, USA. ²INSERM, Unité 1043, F-31300 Toulouse, France. ³Centre National de la Recherche Scientifique, Unité 5282, F-31300 Toulouse, France. ⁴Centre de Physiopathologie Toulouse-Purpan, Université de Toulouse, Université Paul Sabatier, F-31300 Toulouse, France. ⁵Département d'Immunologie, Centre Hospitalier Universitaire de Toulouse, Hôpital Purpan, F-31300 Toulouse, France. ⁶Department of Immunology, Genetics and Pathology, Rudbeck Laboratory, Uppsala University, Dag Hammarskjöldsv. 20, 751 85 Uppsala, Sweden. Correspondence and requests for materials should be addressed to V.H.E. (email: vhe@virginia.edu).

Immune tolerance is imposed through multiple processes that begin during thymic T-cell development and continue in the periphery. During negative selection in the thymus, medullary thymic epithelial cells (mTECs) and dendritic cells (DCs) present self-antigens to tolerize auto-reactive T cells. Intrinsic tolerance mechanisms induce deletion or anergy of high-affinity self-reactive T cells, whereas lower affinity CD4 cells are converted into regulatory T cells (T_{reg}) that mediate extrinsic tolerance^{1,2}. DCs can acquire antigen in the periphery and migrate into the thymus³, or thymic resident DCs can capture circulating antigen⁴. In addition to presenting ubiquitous antigens, mTECs also transcribe and present a variety of peripheral tissue antigens (PTAs) under the control of the autoimmune regulatory element (Aire)^{5,6}, increasing the diversity of self-antigens presented in the thymus.

Thymic tolerance does not eliminate all self-reactive T cells, necessitating mechanisms of peripheral tolerance. Immature DCs continually survey peripheral tissues to acquire self-antigens, which are presented in the draining lymph nodes (LNs) to induce T-cell deletion, anergy or T_{reg} formation⁷. In contrast to DCs, which are specialized for acquiring antigens from other tissues, several subsets of LN cells transcribe PTAs, analogous to mTECs. Extrathymic Aire-expressing cells transcribe and present PTAs in an Aire-dependent manner, leading to CD8 T-cell deletion and CD4 T-cell anergy^{8,9}. Extrathymic Aire-expressing cells are developmentally related to DCs⁹. PTAs are transcriptionally expressed independently of Aire by several subsets of radioresistant LN stromal cells (LNSCs), including lymphatic endothelial cells (LECs), fibroblastic reticular cells (FRCs) and blood endothelial cells (BECs)^{10,11}. Although the effects of PTAs expressed in BECs have not been tested, LECs and FRCs both induce deletion of CD8 T cells^{10–13}.

We previously showed that LECs transcribe and present an epitope from the melanocyte differentiation protein tyrosinase, leading to proliferation and deletion of tyrosinase-specific CD8 T cells^{10,14}. Proliferating tyrosinase-specific CD8 T cells activated by LECs in the absence of 4-1BB co-stimulation upregulate PD-1, which binds to PD-L1 on a radioresistant stromal cell, inhibits the upregulation of the IL-2 receptor and leads to death¹². LECs express the highest level of PD-L1 among the LNSC. LECs also express herpesvirus entry mediator (HVEM) and major histocompatibility complex (MHC)-II¹², which are ligands for the BTLA/CD160 and LAG-3 inhibitory pathways, respectively¹⁵. Tyrosinase and PD-L1 are more highly expressed by LECs in the LN (LN-LECs) compared with LECs from tissue lymphatics in the diaphragm or colon¹⁶, suggesting the LN microenvironment endows LN-LECs with tolerogenic properties not found in tissue LECs. In this study, we investigated whether MHC-II expression on LN-LECs is related to their tolerogenic role, and whether MHC-II is used to induce CD4 T-cell tolerance.

The MHC-II antigen presentation pathway has been extensively studied in professional antigen-presenting cells (APC) and in cell lines. MHC-II molecules are synthesized in the ER and associated with invariant chain (Ii), which targets the complex into late endosomal MHC-II-loading compartments (MIICs). Next, Ii is cleaved by cathepsins, leaving the class II Ii-associated peptide (CLIP) in the peptide-binding groove. CLIP is exchanged for antigenic peptides by the non-classical MHC-II molecule H2-M. H2-M can be inhibited by H2-O, altering the representation of peptides presented¹⁷. LECs express MHC-II¹², but the ability of LECs to load and present self-peptides on MHC-II molecules has not been investigated. In addition, it is unknown whether PTA expression in LECs leads to CD4 T-cell tolerance.

To investigate whether LECs present PTAs on MHC-II molecules and induce CD4 T-cell tolerance, we created transgenic systems where the model antigens β -galactosidase (β -gal) or

haemagglutinin (HA) are expressed in LECs under the control of LEC-specific *Lyve-1* or *Prox1* promoters. Using these complementary models, we demonstrate that LECs do not directly present these PTAs on MHC-II molecules, but instead provide antigen to DCs to induce CD4 T-cell anergy.

Results

LN but not diaphragm LECs have intermediate levels of MHC-II. We previously showed that LN-LECs express MHC-II molecules¹². To determine whether this was a specialized property of LN-LECs, we compared the level of MHC-II molecules on LN-LECs with those on tissue lymphatic LECs, other LNSC subsets and haematopoietically derived APC. MHC-II molecules are expressed on LN-LECs, but not on LECs from the diaphragm (D-LECs; Fig. 1a). LN-LECs express similar levels of MHC-II molecules as LN-BECs and LN-FRCs. Although the geometric mean fluorescence intensity (gMFI) of the entire LEC population is approximately 10% of the gMFI of the entire macrophage population, most LECs express levels of MHC-II similar to that of the lower half of the macrophage profile. This suggests that LECs express levels of MHC-II sufficient for antigen presentation.

We detected mRNA for the I-A^b β -chain in C57BL/6 (B6) LECs (Fig. 1b), demonstrating that they directly express MHC-II molecules. However, MHC-II molecules can also be acquired from another cell through a process called ‘cross-dressing’^{18,19}. We used reciprocal MHC-II^{-/-} bone marrow chimeras to determine the contribution of endogenous synthesis and acquisition from haematopoietically derived cells to the MHC-II molecules displayed by LN-LECs. The LECs in CD45.1 \rightarrow MHC-II^{-/-} bone marrow chimeras continued to display MHC-II molecules, demonstrating they can acquire them from haematopoietically derived cells (Fig. 1c–e). However, LECs in the reciprocal MHC-II^{-/-} \rightarrow CD45.1 chimeras express similar levels of MHC-II molecules. Although there was some heterogeneity between mice (Fig. 1c, left versus right), the levels of MHC-II and percentages of MHC-II⁺ LECs were similar overall between the two chimera groups (Fig. 1d,e). This suggests that the absence of one source of MHC-II molecules is compensated for by another source to maintain a relatively constant level on LECs. These results demonstrate LN-LECs both endogenously synthesize MHC-II molecules and acquire them from haematopoietically derived cells.

Complementary models to evaluate antigen expression in LECs.

Because there are no CD4 TCR transgenic models directed against endogenous PTAs expressed by LECs, we transgenically expressed β -gal as a model PTA (Supplementary Table 1), as there are β -gal-specific CD4 and CD8 TCR transgenic mice available (Supplementary Table 2). *Rosa26^{stop-LacZ}* (ref 20) mice express β -gal after cre-mediated excision of a floxed stop cassette. These mice were crossed with mice expressing either *Lyve-1-cre*²¹ or *Prox1-creER^{T2}* (ref. 22; Supplementary Fig. 1). *Lyve-1-cre* is used as a specific marker for LECs²⁵. *Lyve-1-cre* is constitutively active in all LECs, and in limited subsets of BECs, lymphoid and myeloid cells²¹. *Prox1* is the master transcriptional regulator inducing LEC differentiation²³. *Prox1-creER^{T2}* is induced by tamoxifen, and mediates high-efficiency recombination in LECs but not in BECs, FRCs, DCs, macrophages, B or T cells (Fig. 2). Since *Prox1-creER^{T2}* can also lead to recombination in *Prox1*⁺ cells in the liver and heart²², we used skin-draining LNs in our analysis to eliminate potential effects of antigen draining from these sites. Because these two models, referred to hereafter as *Lyve-1x β -gal* and *Prox1x β -gal*, differ in their patterns of ectopic expression, they were used as complementary models to investigate LEC-induced tolerance.




Figure 1 | LN-LECs endogenously express intermediate levels of MHC-II. (a) Pooled LNs and diaphragms from B6 or MHC-II^{-/-} mice were enzymatically digested. LECs (DAPI^{neg}CD45^{neg}CD31⁺gp38⁺), BECs (DAPI^{neg}CD45^{neg}CD31⁺gp38^{neg}), FRCs (DAPI^{neg}CD45^{neg}CD31^{neg}gp38⁺), macrophages (DAPI^{neg}CD11c^{neg/low}CD11b⁺F4/80⁺) and B cells (DAPI^{neg}CD19⁺CD11c^{neg}CD11b^{neg}) were stained extracellularly for MHC-II, and gMFI was calculated. Data representative of at least three experiments with LN pooled from 1 to 2 mice. (b) B cells (DAPI^{neg}CD19⁺), macrophages (DAPI^{neg}CD11c^{neg/low}CD11b⁺), LECs (DAPI^{neg}CD45^{neg}CD31⁺gp38⁺), FRCs (DAPI^{neg}CD45^{neg}CD31^{neg}gp38⁺) and BECs (DAPI^{neg}CD45^{neg}CD31⁺gp38^{neg}) from B6 mice were sorted by flow cytometry, and cultured 3T3 cells were harvested. Quantitative PCR for I-A^b was performed. Data shown from 2 to 3 independent experiments with LN pooled from 4 to 5 mice. (c-e) Pooled LNs from B6, MHC-II^{-/-}, CD45.1 → MHC-II^{-/-} or MHC-II^{-/-} → CD45.1 mice were digested and stained extracellularly for MHC-II expression on LECs. Numbers in c represent gMFI. Representative data (c,d) or cumulative data (e) from 2 to 4 experiments with 1-2 mice each are shown. Groups compared using a one-way analysis of variance with Tukey post-test. *P < 0.05; **P < 0.01; ***P < 0.001, NS = not significant. All data shown as mean ± s.e.m.

LECs present endogenous β-gal epitopes on MHC-I molecules.

To determine whether LECs from Lyve-1xβ-gal or Prox1xβ-gal mice express β-gal antigen, we tested whether they activated Bg1 CD8 T cells, which express a transgenic TCR specific for the β-gal₉₆₋₁₀₃ epitope presented by H-2K^b (ref 24). A substantial fraction of Bg1 cells proliferated by day 3 after transfer into Prox1xβ-gal mice, and the majority of proliferating cells were gone by day 7 (Fig. 3a). As *Lyve-1-cre* can induce recombination in haematopoietic cells²¹, we used MHC-I^{-/-} → Lyve-1xβ-gal bone marrow chimeras to restrict β-gal expression to radioresistant LNSC. Bg1 cells also proliferated and deleted after adoptive transfer into these mice (Supplementary Fig. 2a). *In vitro*, Bg1 cells proliferated when co-cultured with purified LN-LECs from both models, whereas no significant proliferation

was induced by purified FRCs or BECs (Fig. 3b,c). Thus, LECs are the only LNSCs in either Lyve-1xβ-gal or Prox1xβ-gal mice that present β-gal₉₆₋₁₀₃ at immunologically relevant levels.

The loss of proliferating Bg1 cells by day 7 *in vivo* is consistent with our earlier data demonstrating that LECs induce deletional tolerance. In keeping with this, proliferating Bg1 cells upregulated Annexin V and the inhibitory ligands PD-1 and LAG-3 (Supplementary Fig. 2b,c). Although blockade of PD-1/PD-L1 or LAG-3/MHC-II inhibitory pathways individually did not inhibit deletion of Bg1 cells, blockade of both pathways did so (Fig. 3d and Supplementary Fig. 2d). This demonstrates that one role for MHC-II molecules on LECs is to promote CD8 T-cell tolerance by engaging LAG-3.




Figure 2 | *Prox1-creER^{T2}* is active in LECs but not other LNSCs or haematopoietically derived cells. *Prox1-CreER^{T2} x EYFP^{stop-flox}* (*Prox1xEYFP*) mice were maintained on tamoxifen chow for 2 weeks. Pooled inguinal, axillary and brachial LNs from B6 or *Prox1xEYFP* mice were enzymatically digested and EYFP expression in LECs (DAPI^{neg}CD45^{neg}CD31⁺gp38⁺), BECs (DAPI^{neg}CD45^{neg}CD31⁺gp38^{neg}), FRCs (DAPI^{neg}CD45^{neg}CD31^{neg}gp38⁺), DCs (DAPI^{neg}CD11c^{high}), macrophages (DAPI^{neg}CD11c^{neg/low}CD11b⁺), B cells (DAPI^{neg}CD19⁺), CD4 T cells (DAPI^{neg}CD4⁺CD19^{neg}) and CD8 T cells (DAPI^{neg}CD8⁺CD19^{neg}) was analysed by flow cytometry. Numbers refer to the percent of EYFP⁺ cells in the *Prox1xEYFP* mice. Data representative of two independent experiments with 1–2 mice each.




Figure 3 | LECs induce deletion of β -gal-specific CD8 T cells via the PD-1/PD-L1 and LAG-3/MHC-II pathways. (a) CTV-labelled Thy1.1⁺ Bg1 cells were adoptively transferred along with CTV-labelled Thy1.1^{neg} cells as an injection control. Skin-draining LNs were analysed for Bg1 proliferation 3 or 7 days later. Plots are gated on total CD8⁺ T cells. Data representative of 2–3 experiments with 1–3 mice each. LNSCs from *Prox1x β -gal* (b) or *Lyve-1x β -gal* (c) mice were sorted and co-cultured with Thy1.1⁺ Bg1 T cells for 4 days. Plots are gated on CD8⁺Thy1.1⁺ Bg1 cells. Data representative of 2–3 experiments with LNs pooled from 4 to 9 mice. Each group was compared with T cells only using a one-way analysis of variance (ANOVA) with Bonferroni post-test. (d) CTV-labelled Thy1.1⁺ Bg1 cells were adoptively transferred along with CTV-labelled Thy1.1^{neg} cells as an injection control. Mice were treated with blocking antibodies as described in Methods. Skin-draining LNs were analysed for Bg1 proliferation 7 days later. Data representative of 2 experiments with 1–2 mice each. Treatment groups were compared with untreated group using a one-way ANOVA with Bonferroni post-test. * $P < 0.05$; ** $P < 0.01$; NS = not significant. All data shown as mean \pm s.e.m.

LECs do not present endogenous β -gal epitopes on MHC-II.

To test whether LECs from *Prox1x β -gal* and *Lyve-1x β -gal* mice present β -gal epitopes on MHC-II molecules, we used Bg2 CD4 T-cells²⁵, which recognize β -gal_{721–739} presented by I-A^b. Bg2 cells transferred into *Lyve-1x β -gal* or *Prox1x β -gal* mice proliferated by day 3 and continued to proliferate and accumulate by day 7 (Fig. 4a,b). However, they did not proliferate in *MHC-II^{-/-} \rightarrow Lyve-1x β -gal* or *MHC-II^{-/-} \rightarrow*

Prox1x β -gal chimeras, in which radioresistant LECs synthesize MHC-II molecules but bone marrow-derived cells do not. This indicates that Bg2 proliferation in non-chimeric mice is induced by bone marrow-derived cells, but not LECs. We considered that Bg2 T cells might recognize antigen but not proliferate because of the lack of costimulatory molecules on LECs¹². However, Bg2 cells adoptively transferred into *MHC-II^{-/-} \rightarrow Prox1x β -gal* chimeras treated with α CD28 agonistic antibodies also did not




Figure 4 | LECs do not present endogenous β -gal on MHC-II. (a) Representative and (b) cumulative data of CTV-labelled Thy1.1⁺ Bg2 cells adoptively transferred into the indicated recipients. CTV-labelled Thy1.1^{neq} cells were co-transferred as an injection control. Skin-draining LNs were analysed 3 (a) or 7 (a,b) days later, and plots are gated on CD4⁺ T cells. Data representative of 1-3 experiments with 1-4 mice each. Indicated groups were compared using a one-way analysis of variance (ANOVA) with Bonferroni post-test. (c) CTV-labelled Bg2 cells were adoptively transferred into B6 and MHC-II^{-/-} \rightarrow Prox1x β -gal mice treated with PBS, α CD28 or IFN- γ , and proliferation was analysed 3 days later. Plots are gated on Thy1.1⁺CD4⁺ cells. Data representative of 2 experiments with 2 mice each. (d) CTV-labelled Thy1.1⁺ Bg2 cells were transferred into the indicated recipients and activation markers were analysed 16 h later. Plots are gated on Thy1.1⁺CD4⁺ cells. Data from one experiment. (e) LNs from MHC-II^{-/-} and PBS- or IFN- γ -treated B6 mice were enzymatically digested 24 h after treatment, and MHC-II on LECs was analysed by flow cytometry. Data representative of 3 experiments with 1-2 mice each. (f) LNSCs, DCs and macrophages from B6 mice were sorted by flow cytometry, pulsed with 50 μ M Bg2 peptide for 3 h, washed and co-cultured with CPD eF670-labelled Thy1.1⁺ Bg2 T cells for 4 days. (g) LNSCs, DCs and macrophages from Prox1x β -gal mice were sorted by flow cytometry and co-cultured with CPD eF670-labelled Thy1.1⁺ Bg2 T cells for 4 days. (f,g) Plots are gated on DAPI^{neq}CD4⁺ Thy1.1⁺ cells. Data representative of 2 experiments with pooled LNs from 5 to 9 mice. All data shown as mean \pm s.e.m.

proliferate (Fig. 4c). We also considered that LECs might present β -gal, but rapidly induce anergy or suppress T-cell proliferation through nitric oxide^{26,27}. Therefore, we examined Bg2 cells for upregulation of CD69, CD25 and CD44, and downregulation of CD62L, one day after adoptive transfer. After transfer into non-chimeric Prox1x β -gal mice, CD69, CD25 and CD44 were all upregulated and CD62L was downregulated (Fig. 4d). However, these markers remained unchanged and identical on Bg2 cells

transferred into MHC-II^{-/-} \rightarrow Prox1x β -gal chimeras and antigen-free B6 mice (Fig. 4d), indicating the Bg2 cells were not activated. We conclude that LECs do not present MHC-II-restricted β -gal epitopes to Bg2 cells *in vivo*, even though they express the source protein and the restriction element.

We next considered that the level of MHC-II on the surface of LECs might be too low to induce CD4 proliferation. In MHC-II^{-/-} \rightarrow Prox1x β -gal mice treated with interferon (IFN)- γ , the

gMFI of MHC-II on LECs was upregulated more than sixfold (Fig. 4e). However, Bg2 cells transferred into these mice also did not proliferate (Fig. 4c). Finally, we pulsed FACS-sorted LECs from B6 mice with β -gal₇₂₁₋₇₃₉ peptide. Peptide-pulsed LECs induced Bg2 proliferation *in vitro* (Fig. 4f), as did peptide-pulsed FRCs, BECs, DCs and macrophages. The level of proliferation induced by the LNSCs was lower than that induced by DCs, which could reflect lower levels of MHC-II molecules on these cells and/or lack of costimulatory molecules. However, the results demonstrate that the level of MHC-II on LECs is sufficient to present antigen to Bg2 T cells.

As LECs were not responsible for the *in vivo* proliferation of Bg2 cells in non-chimeric Prox1x β -gal mice (Fig. 4a), we used *in vitro* co-cultures to determine which cells were. LECs, FRCs and BECs sorted by flow cytometry from Prox1x β -gal mice did not induce Bg2 proliferation (Fig. 4g), confirming the *in vivo* results. However, DCs from these mice induced strong Bg2 proliferation, and macrophages induced a less robust, but still highly significant, response. The basis for antigen presentation by these cells is explored further below.

Antigen location does not affect MHC-II presentation in LECs.

MHC-II presentation of epitopes from cytoplasmic proteins such as β -gal depends on autophagy²⁸. Therefore, we investigated whether LECs could present an autophagy-independent I-E^d-restricted epitope from the membrane protein influenza HA²⁹. We expressed HA in LECs using *Prox1-creER^{T2} x Rosa26^{stop-HA}* (Prox1xHA) (B6 x Balb/c) F1 mice and assessed antigen presentation using HA-specific CD4 or CD8 TCR transgenic cells (TS1 (ref. 30) or Clone 4 (ref. 31), respectively, also [B6 x Balb/c] F1). LECs from Prox1xHA mice induced proliferation of Clone 4 CD8 T cells *in vitro* (Fig. 5a and Supplementary Fig. 3a), indicating that HA is expressed in LECs and presented on H-2K^d MHC-I molecules. However, Prox1xHA LECs did not induce TS1 CD4 T-cell proliferation *in vitro* (Fig. 5a and Supplementary Fig. 3a). A recent report demonstrated that LECs that had acquired peptide:MHC-II complexes from DCs induced CD4 apoptosis *in vitro* without prior proliferation¹⁹. However, we saw no difference in death between TS1 cells co-cultured with or without Prox1xHA LECs (Fig. 5b). In contrast to LECs, both CD11c⁺CD11b⁺ and CD11c⁺CD11b^{neg} DCs presented HA on both MHC-I and MHC-II molecules (Fig. 5a and Supplementary Fig. 3a). Corresponding results were seen *in vivo*, where TS1 cells upregulated CD69 and proliferated in non-chimeric Prox1xHA mice but not in MHC-II^{-/-} → Prox1xHA chimeras (Supplementary Fig. 3b,c). Therefore, like β -gal, HA is presented by LECs on MHC-I molecules, but is not presented on MHC-II molecules. Because *Prox1-creER^{T2}* does not induce HA expression in haematopoietic cells, the presentation of HA by two DC subsets indicates that it has been transferred from LECs.

We also used a T-cell independent assay to directly test whether LECs can form peptide:MHC-II complexes. The Y-Ae antibody recognizes I-A^b molecules presenting an epitope derived from the I-E α chain³², which is generated in endosomes³³. I-E α , encoded by the *H2-Ea-ps* gene, was expressed in BALB/c LECs but is a pseudogene and was not expressed in B6 LECs (Supplementary Fig. 4). However, although DCs, macrophages and B cells from (B6 x BALB/c) F1 mice expressed Y-Ae epitopes on the cell surface, LECs did not (Fig. 5c). In addition, treating mice with IFN- γ to upregulate I-A^b and I-E^d did not induce Y-Ae staining on LECs. These results demonstrate that LECs fail to present three separate antigens, which included both cytoplasmic and membrane-bound proteins, and which were restricted by two different MHC-II alleles. Both HA and I-E α can be presented in the absence of autophagy^{29,34}, suggesting these results are not due

to any impairment of that process in LECs. Instead, the results suggest that there is a fundamental deficiency in the MHC-II processing pathway in LECs.

LECs express Ii and Cathepsin L, but not H2-M or H2-O. We next investigated whether LECs were deficient in any components of the MHC-II processing pathway. LECs express high levels of Ii (Fig. 6a), suggesting that MHC-II molecules are correctly targeted to the MIIC. Ii is degraded by cathepsins S or L, leaving CLIP, which blocks the MHC-II peptide-binding groove. On average, one-third of LAMP1⁺ endosomes in LECs co-expressed cathepsin L by immunofluorescence microscopy, but there was substantial variation among individual cells and some LN-LECs did not contain detectable Cathepsin L (Fig. 6b and Supplementary Fig. 5a). Cathepsin L in LECs was active and cleaved the substrate (CBZ-Phe-Arg)₂-Rhodamine110 (Supplementary Fig. 5b). In contrast, D-LECs did not generate a signal above the background level in cells pretreated with a cathepsin L inhibitor, suggesting that MHC-II^{neg} D-LECs do not express active cathepsin L. The presence of active cathepsin L in some endosomes suggests that LN-LECs can digest Ii into CLIP, although the efficiency of this process may vary among individual cells and endosomes.

The peptide editor H2-M exchanges CLIP for antigenic peptides, and also optimizes the repertoire of peptides bound to various MHC-II alleles³⁵⁻³⁷. Its activity is antagonized by H2-O (ref. 17). Therefore, low levels of H2-M or high levels of H2-O could explain the inability of LECs to present antigenic peptides. We did not detect either H2-M α β 2 or H2-O expression in LECs by flow cytometry (Fig. 6c and Supplementary Fig. 5c). We also did not detect significant levels of *H-2DM α* or *H-2DM β* mRNA (Fig. 6d). This suggests that LECs might inefficiently exchange CLIP or suboptimal self-peptides for antigenic peptides. We attempted to measure the level of CLIP bound to I-A^b MHC-II molecules using the 15G4 antibody that specifically recognizes I-A^b:CLIP complexes. However, background staining in the LEC MHC-II^{-/-} negative control was extremely high, compromising our ability to detect a positive signal in MHC-II^{+/+} LECs (Fig. 6e). With this limitation, our data nonetheless suggest that the most likely explanation(s) for the lack of antigen presentation by LN-LECs are either that Ii associated with MHC-II molecules is not efficiently cleaved into CLIP or that the lack of H2-M prevents the exchange of CLIP or suboptimal self-peptides for optimal antigenic peptides.

DCs acquire antigen from LECs to induce anergy. Although LECs do not directly present β -gal or HA on MHC-II, these antigens are presented by DCs and macrophages (Figs 4 and 5). As *Prox1-creER^{T2}* does not induce genetic recombination in DCs or macrophages (Fig. 2), this indicates that these cells acquired antigen transcribed in LECs. However, antigen presentation did not occur in MHC-II^{-/-} → Prox1x β -gal (Fig. 4a-d) or MHC-II^{-/-} → Prox1xHA (Supplementary Fig. 3b,c) chimeras, indicating that antigen is loaded onto MHC-II molecules in haematopoietically derived cells, rather than being transferred as preformed peptide:MHC-II complexes. To formally demonstrate this, we used bone marrow chimeras where genetic expression of β -gal is restricted to either the haematopoietically derived or radioresistant cells. Bg2 cells transferred into Prox1x β -gal → CD45.1 chimeras did not proliferate (Fig. 7a), demonstrating that β -gal is not genetically expressed in haematopoietically derived cells. In contrast, Bg2 T cells transferred into CD45.1 → Prox1x β -gal chimeras proliferated, confirming that β -gal from LECs is transferred to haematopoietically derived cells.

The presentation of β -gal or HA by haematopoietically derived cells *in vivo* led to proliferation and accumulation of CD4 T cells




Figure 5 | LECs do not present MHC-II peptides from the membrane proteins HA and I-E α . (a,b) LNSCs and DCs from Prox1xHA mice were sorted by flow cytometry and co-cultured with CFSE-labelled Thy1.1⁺ Clone 4 CD8 cells (left) or TS1 CD4 cells (right) for 4 days, and the percent of T cells proliferating was calculated. Each group was compared with T cells alone using a one-way analysis of variance (ANOVA) with Bonferroni post-test. *** $P < 0.001$. Data representative of 2 experiments with LN pooled from 8 mice. (b) The percent of DAPI⁺ TS1 cells from the co-culture in a was calculated. Plots are gated on CD4⁺Thy1.1⁺ TS1 cells. No significant differences were seen when each group was compared with T cells alone using a one-way ANOVA with Bonferroni post-test. (c) LNs from PBS- or IFN- γ -treated B6 or (B6 x BALB/c) F1 mice were enzymatically digested, and extracellular I-A^b:I-E α peptide complex expression (top) or MHC-II levels (bottom) were analysed on the indicated populations by flow cytometry. Data representative of 2-3 experiments with 1-3 mice. All data shown as mean \pm s.e.m.




Figure 6 | LECs express Ii and cathepsin L, but do not express the peptide editor H2-M. (a) B6 LECs (DAPI^{neg}CD45^{neg}CD31⁺10.1.1⁺) and B cells (DAPI^{neg}CD19⁺) were stained intracellularly for Ii and MHC-II or isotype control. The LEC-specific marker 10.1.1 (refs 10,12) was used instead of gp38 since 10.1.1 is compatible with intracellular staining protocols. (b) LNs were enzymatically digested, depleted of CD45⁺ cells, stained with 10.1.1 (cyan) extracellularly and cathepsin L (white), LAMP1 (magenta) and DAPI (blue) intracellularly before being cytopspun onto slides. Scale bar, 5 μm. (c) LECs and B cells were stained for MHC-II and H2-Mαβ2 intracellularly. (d) LECs, DCs and B cells from B6 mice were sorted by flow cytometry, and 3T3 cells were harvested from culture as in Fig. 1. mRNA was purified and quantitative PCR was performed for the indicated genes. LECs were compared with the other groups using a one-way analysis of variance (ANOVA) with Bonferroni post-test. **P* < 0.05; ***P* < 0.01; ****P* < 0.001. (e) LECs and B cells from B6 or MHC-II^{-/-} (negative control) mice were stained extracellularly for I-A^b:CLIP complexes and MHC-II. Data in a-c, e are representative of at least 2 experiments with 1–2 mice each, and data in d is from 2 experiments with LN pooled from 4 to 5 mice. All data shown as mean ± s.e.m.




Figure 7 | Haematopoietically derived cells acquire β -gal from LECs and induce Bg2 anergy. (a) CTV-labelled Thy1⁺ Bg2 cells were transferred into the indicated recipients, along with CTV-labelled Thy1.1^{neg} cells as an injection control. Proliferation in skin-draining LNs was measured 7 days later. Data representative of 1–2 experiments with 2–5 mice each. (b) CTV-labelled Thy1⁺ Bg2 cells were transferred into the indicated recipients, and T_{reg} development was measured 3 days later. Data representative of 2 experiments with 2–3 mice each. (c) Experimental design to measure tolerance induction. CTV-labelled CD25^{neg}Thy1.1⁺ Bg2 CD4 T cells were adoptively transferred, and recipient mice were challenged 28 days later with peptide-pulsed BMDCs. Proliferation was measured 5 days later by Ki67 upregulation, and FoxP3 and CD25 were measured to assess T_{reg} formation. (d) Representative and (e) cumulative data gated on CD4⁺Thy1.1⁺ Bg2 cells from skin-draining LN from 2 experiments with 1–7 mice each. **P* < 0.05; ***P* < 0.01; ****P* < 0.001, NS = not significant. Indicated comparisons were made using a one-way analysis of variance with Bonferroni post-test. All data shown as mean \pm s.e.m.

over 7 days (Figs 4a and 7a, and Supplementary Fig. 3c). To determine the fate of these cells, we tested whether they developed into either T_{reg} or anergic cells. After 3 days, Bg2 cells transferred into Prox1x β -gal, MHC-II^{-/-} → Prox1x β -gal and B6 mice all had equivalently low levels of FoxP3 staining (Fig. 7b), indicating they were not differentiating into T_{reg}. To test for anergy, we adoptively transferred CD25^{neg} CTV-labelled Bg2 cells into recipient mice and re-challenged them 28 days later with β -gal₇₂₁₋₇₃₉-pulsed bone marrow-derived DCs (BMDCs). Proliferation was measured 5 days later by Ki67 upregulation (Fig. 7c). In this experimental setup, CTV^{dilute}Ki67^{pos} cells are actively proliferating in response to BMDCs, whereas CTV^{dilute}Ki67^{neg} cells proliferated initially after adoptive transfer but did not respond subsequently to the β -gal-pulsed BMDCs, and are thus anergic. As expected, Bg2 cells transferred into B6 mice remained naïve and proliferated strongly upon subsequent BMDC stimulation, as evidenced by their dilution of CTV and >85% expression of Ki67 (Fig. 7d,e). Consistent with previous work³⁸, Bg2 cells transferred into B6 mice treated with 300 μ g of β -gal₇₂₁ peptide 0 and 3 days after adoptive transfer were CTV^{dilute} and Ki67^{neg}, and thus anergic. Bg2 cells transferred into Prox1-x β -gal

mice were also CTV^{dilute}Ki67^{neg}, and therefore were also anergic. Bg2 cells transferred into either MHC-II^{-/-} → Prox1-x β -gal or control antigen-free MHC-II^{-/-} → CD45.1 bone marrow chimeras were strongly and equivalently CTV^{dilute}Ki67^{pos}, indicating that anergy does not occur without antigen presentation by haematopoietically derived cells. We also tested whether Bg2 cells differentiate into T_{reg} cells at this later timepoint. We found no increase of FoxP3⁺CD25⁺ Bg2 cells in Prox1-x β -gal mice compared with B6 mice (Fig. 7d,e), further demonstrating that T_{reg} formation is not a form of tolerance induced in this model. These data demonstrate that the antigen transcribed in LECs that is provided to DCs is responsible for anergy induction in CD4 T cells.

Discussion

LECs express a variety of PTAs and directly induce CD8 T-cell tolerance, but their ability to induce CD4 T-cell tolerance to PTAs has not previously been examined. In this study, we demonstrate that MHC-II expression on LECs is sufficient for presentation of exogenously pulsed peptides, but LECs do not present peptides

from three endogenous proteins. However, LECs provide antigens to DCs, which then induce anergy. Therefore, LECs serve as a reservoir of PTAs in the LN, which are acquired by DCs for the induction of CD4 T-cell tolerance. In addition, MHC-II plays a role in the deletional tolerance of Bg1 CD8 T-cells, which delete through either the PD-1/PD-L1 or LAG-3/MHC-II pathways after recognizing antigen presented by LECs on MHC-II molecules.

LECs do not present epitopes derived from β -gal, HA or I-E α on their MHC-II molecules. This is independent of antigen source protein localization to the cytoplasm, plasma membrane or endosome, respectively. Because two of these antigens are presented by an autophagy-independent mechanism^{29,34}, lack of presentation is not due to a defect in autophagy. In addition, the absence of the Y-Ae epitope on LECs suggests that the failure to present is not due to a deficiency in loading source proteins into the MHC-II processing pathway, as I-E α will be endogenously targeted to the MIIC by Ii as part of the I-E^d molecule. LECs express high levels of Ii, suggesting that MHC-II molecules are appropriately chaperoned into the MIIC. In addition, cathepsin L is active and expressed in a subset of LEC LAMP-1⁺ endosomes. This suggests that Ii can be cleaved into CLIP in many cells, but it is also possible that MHC-II molecules maturing in cathepsin L^{neg} endosomes contain Ii instead of CLIP. However, LECs also do not express H2-M, suggesting that even if CLIP is generated it may be inefficiently exchanged for antigenic peptides. Unfortunately, a high level of nonspecific background CLIP staining in LECs prevented direct evaluation of this latter hypothesis. It has been shown that for MHC-II alleles with a relatively low affinity for CLIP, competition from non-optimal endogenous peptides in the absence of H2-M can limit effective presentation of exogenous epitopes^{35–37}. Consistent with this, Y-Ae expression and presentation of HA_{109–119} on I-E^d are both dependent on H2-M^{39,40}. Thus, lack of MHC-II-restricted epitope presentation by LECs is likely due to deficiencies in expression of H2-M, and potentially that of cathepsin L as well.

Although class II transactivator (CIITA) usually drives co-expression of MHC-II, Ii and H2-M, several papers have shown that MHC-II^{41–47} and invariant chain⁴⁸ can be expressed independently of CIITA, and invariant chain can be expressed independently of MHC-II⁴⁹. CIITA-independent transcription of MHC-II molecules in endothelial cells has also been described⁴⁴. Most importantly, kidney glomerular endothelial cells have been shown to express HLA-DR MHC-II molecules in the absence of the human homologue of H2-M, HLA-DM⁵⁰. However, the mechanism(s) of this differential regulation have not been defined. Alternatively, H2-M mRNA could be degraded in LECs by microRNA, preventing translation. It is interesting to speculate that MHC-II molecules may be expressed independently of H2-M as a means to separate the generation of a LAG-3 ligand from generation of antigens. Although the transcriptional mechanisms used in LECs to induce MHC-II molecules and Ii in the absence of H2-M remain to be determined, our data suggest that the most likely model is that MHC-II:Ii complexes are correctly targeted into the MIIC, but antigenic peptides are inefficiently loaded for presentation due to a lack of H2-M and possibly inefficient Ii cleavage by cathepsin L.

Our work demonstrating a lack of endogenous antigen presentation by LECs is relevant to three other recently published studies. Onder *et al.*⁵¹ used *podoplanin-cre* transgenic mice to induce β -gal expression in FRCs and LECs, analogous to our models, and saw *in vivo* proliferation of Bg2 cells transferred into these mice. However, they did not evaluate the cell populations that presented antigen. Our results suggest that their observations were due to antigen transfer from LECs to DCs and macrophages, rather than direct presentation by either FRCs or LECs. In a

second study, Dubrot *et al.*¹⁹ showed that LECs acquired preloaded peptide:MHC-II complexes from DCs and directly induced apoptosis of antigen-specific T cells *in vitro* in the absence of proliferation¹⁹. Our data support their conclusion that LECs can acquire MHC-II molecules from haematopoietic cells through cross-dressing. However, we also found that LECs synthesize a large fraction of the MHC-II molecules that they express. Importantly, our results establish that LECs provide source proteins to DCs for efficient presentation by both MHC-I and MHC-II molecules. However, in contrast to Dubrot *et al.*, we saw no evidence that these particular MHC-II:peptide complexes were transferred back to the LECs *in vivo* in sufficient quantities to induce CD4 T-cell recognition, either *in vitro* or *in vivo*. Thus, our results argue against a general role for LECs in tolerance induction through presentation of cross-dressed MHC-II molecules acquired from DCs. One possible explanation for the difference between our study and theirs is that Dubrot *et al.* used an antigen genetically expressed in DCs, potentially expanding the amount of antigen available as well as the number of DCs expressing it, which may have increased the level of cross-dressed antigen:MHC-II complexes to unusually high levels. However, they did not specifically demonstrate CD4 T-cell recognition of antigens displayed on MHC-II by LECs *in vivo*, and the form of apoptotic death demonstrated *in vitro* was unusual. It is clear that much remains to be done to establish the relevance of LEC cross-dressing for *in vivo* tolerance induction.

A third study recently published by Baptista *et al.*⁵² showed that the transplantation of MHC-II^{neg} LN (and thus MHC-II^{neg} LNSCs) into wild-type mice reduces the frequency of T_{reg} cells and increases CD4 and CD8 T-cell activation, suggesting that one or more LNSC populations present antigens on MHC-II. Although LECs were not specifically evaluated, an alternative explanation not explored in this paper is that the effect could be due to loss of the LAG-3/MHC-II pathway. This pathway has been shown to enhance T_{reg} function and inhibit T-cell effector activity^{15,53}, and our work demonstrates that LECs use it for the induction of CD8 T-cell tolerance. In addition, Baptista *et al.*⁵² elegantly demonstrate that transplantation of OVA⁺ LNs induces tolerance to delayed type hypersensitivity and skin transplantation. However, they do not investigate whether this was due to direct antigen presentation by LNSCs, or to antigen acquisition and presentation by CD45⁺ haematopoietic cells. As our work shows that CD45⁺ cells can induce tolerance using antigen acquired from LECs, this mechanism also needs to be considered.

Our results provide compelling evidence that LECs indirectly function in peripheral CD4 tolerance by serving as a reservoir of PTAs in the LN, which are acquired and presented by DCs to induce anergy. Similarly, mTECs and DCs share responsibility for CD4 tolerance induction in the thymus. mTECs express high levels of MHC-II, and directly present some antigens to CD4 T cells, whereas others are transferred to DCs^{1,54–59}. It is unclear what determines whether an antigen is presented by mTECs, DCs or both. mTECs are not highly phagocytic⁶⁰, but do constitutively undergo autophagy³⁴, and disrupting autophagy shifts presentation of a model antigen from mTECs to DCs⁶¹. Thus, one model is that PTAs not efficiently incorporated into autosomes are presented by DCs instead of mTECs. Although it is unknown whether LECs constitutively undergo autophagy, presentation of HA and I-E α on MHC-II molecules does not require autophagy^{29,34}, suggesting this alone would not preclude presentation. Our work demonstrates LECs transfer PTAs to DCs for the induction of tolerance, a process analogous to that occurring in the thymus. Interestingly, a recent report demonstrated that viral antigens are archived in LECs and subsequently transferred to DCs for the maintenance of memory

T-cells and enhanced protective immunity⁶². This indicates extensive cross-talk between LECs and DCs in the LN, and future studies may determine how tolerogenic versus immunogenic antigens are separated in LECs and transferred to DCs.

A variety of different mechanisms are used to transfer antigens among different cell types. Gut macrophages transfer antigens to CD103⁺ DCs through gap junctions during oral tolerance⁶³. Antigen transfer can also occur through exosomes⁶⁴ or phagocytosis of apoptotic cells^{65,66}. Transfer of peptide:MHC-II complexes from DCs to LECs occurs through cell contact and/or exosomes¹⁹. LECs could potentially transfer antigens to DCs through exosomes, gap junctions or DC phagocytosis of apoptotic LECs. We found that transfer of β -gal from LECs to DCs does not solely rely on phagocytosis, as blocking phosphatidylserine recognition or inhibiting DC phagocytosis does not block antigen transfer *in vivo* (unpublished data). Although the CD11b⁺CD11c⁺ DC subset is generally considered to be specialized for antigen presentation to CD4 T cells and CD11b^{neg}CD11c⁺ DCs are specialized for antigen presentation to CD8 T cells⁷, we found that HA is presented to CD4 and CD8 T cells by both CD11b^{neg}CD11c⁺ and CD11b⁺CD11c⁺ DC subsets. This suggests that HA is non-specifically acquired and presented by multiple subsets of DCs. Further work is needed to elucidate the exact mechanism by which PTAs are transferred from LECs to DCs.

Although MHC-II molecules on LECs do not directly present peptides for CD4 T-cell tolerance, they play an important role in CD8 T-cell tolerance. We previously demonstrated LECs induce deletion of tyrosinase-specific CD8 T cells through PD-1/PD-L1 (ref. 12). Here, we show that both the PD-1/PD-L1 and LAG-3/MHC-II pathways are used in combination to tolerize β -gal-specific CD8 T cells. LAG-3 and PD-1 act synergistically in other tolerance models¹⁵. Although we have not specifically shown that MHC-II on LECs is used as the ligand for LAG-3, this is likely as LECs are the only cells presenting β -gal₉₆₋₁₀₃ in MHC-I^{-/-} \rightarrow Lyve-1 β -gal bone marrow chimeras. Our results demonstrate that LECs use multiple inhibitory pathways to enforce CD8 T-cell tolerance.

In summary, we have demonstrated that although LN-LECs express intermediate levels of MHC-II molecules, they do not present MHC-II β -gal, HA or I-E α peptides. Instead, MHC-II is important to enforce CD8 T-cell tolerance through the LAG-3 pathway. LECs indirectly induce CD4 anergy by transcribing PTAs and transferring them to DCs for MHC-II presentation, a process analogous to what occurs in the thymus. As thymic tolerance is incomplete, peripheral transcription of PTAs by LECs ensures that self-reactive T cells have multiple opportunities to be tolerized. By directly inducing CD8 T-cell deletion as well as transferring antigens to DCs for CD4 anergy, LECs play multiple tolerogenic roles.

Methods

Mice. C57Bl/6 (B6) and CD45.1 (B6-LY5.2/Cr) were from NCI. MHC-II^{-/-} (B6.129-H2-Ab1^{tm1Gru}N12) and MHC-I^{-/-} (B6.129P2-H2-Kb^{tm1}H2-Db^{tm1}N12)⁶⁷ mice were from Taconic (NIAID Exchange Program). *Lyve-1-cre* (B6.129P2-Lyve1^{tm1.1}(EGFP/cre)^{Cys}/J)²¹, β -gal (B6.129S4-Gt(ROSA)26Sor^{tm1}Sor/J)²⁰, EYFP (B6.129X1-Gt(ROSA)26Sor^{tm1}(EYFP)^{Cos}/J) and Clone 4 (CBy.Cg-Thy1a Tg(Tcr α Cl4,Tcr β Cl4)1Shrm/Shrm, Jax)³¹ mice were from Jax. Bg1 (ref. 24) and Bg2 (ref. 25) mice from Christopher Norbury (Pennsylvania State University) were crossed to Thy1.1 mice (Jax). *Prox1-creERT2* (ref. 22) and *Rosa26tm(HA)1Lilb1* mice⁶⁸ have been described. TS1 (ref. 30) mice were from Andrew Caton (University of Pennsylvania). All mice other than Clone 4, Clone 6.5 and *Rosa26tm(HA)1Lilb1* are on a B6 background. The BALB/c Clone 4 and Clone 6.5 mice were crossed to B6 Thy1.1 mice, and the resulting (BALB/c x B6) background mice were used for adoptive transfers into *Prox1-creERT2* x *Rosa26tm(HA)1Lilb1* (BALB/c x B6) recipients. *Prox1-creERT2* was induced with tamoxifen chow (TAM400, Harlan) for 2 weeks before experiments, and mice were maintained on TAM400 throughout the course

of the experiment. Male and female mice were used between the ages of 6 and 32 weeks. Animals were maintained in pathogen-free facilities. This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Procedures were approved by the University of Virginia Animal Care and Use Committee.

LNCS and DC isolation. LNs (pooled inguinal, axillary, brachial, cervical and mesenteric) were digested for 45–60 min at 37 °C using 0.42 U ml⁻¹ Liberase TM (Roche) and 60 U ml⁻¹ DNase I (Sigma) in DMEM (Cellgro) with 2% FCS, essential and non-essential amino acids, sodium pyruvate and HEPES. Red blood cells were lysed (Sigma). CD45⁺ cells were labelled with CD45 magnetic beads in AwesomeMACS (PBS with 0.5% BSA, 2 mM EDTA, 2 mM L-glutamate, 10 mM sodium pyruvate, 1 \times essential and non-essential amino acids, and 4.5 g l⁻¹ dextrose) and separated using the Deplete AutoMACS protocol (Miltenyi Biotec). For *in vitro* co-cultures, DCs and macrophages were enriched using CD11c⁺ and CD11b⁺ beads (Miltenyi Biotec) before CD45 depletion. Cells were stained and analysed by flow cytometry (see below), or sorted by flow cytometry using Influx (BD), FACSVantage (BD) or iCyt Reflection (Sony) cell sorters for *in vitro* co-cultures and quantitative PCR.

In vitro co-culture. Antigen-specific T cells were enriched using CD4 or CD8 magnetic beads (Miltenyi Biotec) and labelled with 1 μ M carboxyfluorescein succinimidyl ester (CFSE) or cell proliferation dye eF670 (CPD eF670, eBioscience). T cells were co-cultured with FACS-sorted LNCS, DCs or macrophages at a 1:2 (T-cell/APC) ratio for 4 days in RPMI-1640 with 10% FCS, 2 mM L-glutamine, 30 mM HEPES, 10 mM sodium pyruvate, 0.05 mM β -mercaptoethanol, 1 \times essential and non-essential amino acids, and 0.05 mg ml⁻¹ gentamycin containing IL-2 (10 U ml⁻¹) and IL-7 (1 pg ml⁻¹) for CD8 T cells and 1 ng ml⁻¹ for CD4 T cells. CD4 co-cultures also received 0.25 μ g ml⁻¹ α CD28 (clone 37.51).

Adoptive transfer. Thy1.1⁺ antigen-specific T cells were positively selected with CD4 or CD8 beads or CD25^{neg} CD4⁺ cells were negatively selected using a regulatory T-cell isolation kit (Miltenyi Biotec). Cells were labelled with Cell Trace Violet (CTV, Invitrogen), and 1 \times 10⁶ antigen-specific T cells were injected intravenously. Thy1.2⁺ CTV-labelled congenic cells were used as an injection control. Skin-draining LN (inguinal, axillary, brachial) were harvested from recipient mice 1, 3, 7 or 33 days after adoptive transfer.

In vivo antibody and IFN- γ treatment. Mice were injected with 100 μ g anti-CD28 (clone 37.51, BioXCell) IP at days 0 and 2. 100 μ g anti-LAG-3 (C9B7W, BioXCell) and/or anti-PD-L1 (10B5, University of Virginia Lymphocyte Culture Center or BioXCell) were injected intraperitoneally (IP) at days -1, 1, 3 and 5. IFN- γ (5 \times 10⁴ U, Peprotech) was injected intravenously (IV) at days -1 and 1 for adoptive transfer experiments, and 24 h before harvest for Y-Ae staining experiments. Mice were randomly assigned to treatment groups.

Bone marrow chimeras. Mice were irradiated (6.5 Gy \times 2) and reconstituted with a minimum of 2 \times 10⁶ bone marrow cells depleted of CD4⁺ and CD8⁺ T cells (Miltenyi Biotec) to prevent graft-versus-host disease. Chimeras were maintained on sulfa water for 3 weeks, and allowed to reconstitute for at least 8 weeks before use.

Flow cytometry. Antibodies used include: Ii (In-1, 1:100), H2-M (2E5A, 1:50), Rat IgG1 (R3-34, 1:50; all from BD Biosciences); podoplanin (8.1.1, BioLegend, 1:500); CLIP (15G4, Santa Cruz, 1:5); H2-O (ref. 69); Mags.Ob3, Lisa Denzin, 1:400; 10.1.1 (UVA lymphocyte culture centre, 1:1,000); CD45 (30-F11, 1:1,000), CD31 (390, 1:1,000), MHC-II (M5/114.15.2, 1:50 (Fig. 1) to 1:500 (Fig. 6)), CD8 (53-6.7, 1:1,000), CD4 (GK1.5, 1:1,000), Thy1.1 (HIS51, 1:1,000), Thy1.2 (53-2.1, 1:1,000), CD45.1 (A20, 1:500), CD11c (N418, 1:500), CD11b (M1/70, 1:500), CD69 (H1.2F3, 1:500), CD62L (MEL-14, 1:1,000), CD44 (IM7, 1:1,000), CD25 (PC61.5, 1:500), Y-AE (eBioY-Ae, 1:200), PD-1 (RMP1-30, 1:100), LAG-3 (eBioC9B7W, 1:100), Rat IgG2b (eB149/10H5; all from eBioscience). Intracellular staining for PD-1, LAG-3, Ii, H2-M and H2-O was done using the Cytofix/Cytoperm kit (BD Bioscience), and Ki67 (Sola15, 1:500) and FoxP3 (FJK-16s, 1:100) were stained using Treg permeabilization buffers (eBioscience). Annexin V (1:20) was stained using the eBioscience kit. 4,6-Diamidino-2-phenylindole (DAPI; Sigma) or live/dead aqua (Invitrogen, 1:200) were used to distinguish live cells. Cells were acquired on a FACSCanto II (BD Biosciences) and data analysed using FlowJo (Tree Star).

Cathepsin L assay. LN-LECs or D-LECs were preincubated with 1 mM of the cathepsin L inhibitor 1-naphthalenesulfonyl-IW-CHO (Calbiochem) or dimethylsulphoxide vehicle control for 20 min at 37 °C, followed by a 20-min incubation with 50 μ M of the cathepsin L substrate (CBZ-Phe-Arg)₂-rhodamine 110 (Invitrogen) at 37 °C before being acquired on a FACSCanto II.

Immunofluorescence microscopy. LNs were enzymatically digested, depleted of CD45⁺ cells, stained with 10.1.1 extracellularly and CD107a (1D48, BioLegend), cathepsin L (EPR8011, Abcam), biotin-anti-rabbit (Vector labs) and streptavidin Dylite 550 (Thermo Scientific) intracellularly before being cytospun onto slides and mounted with Slow Fade Gold with DAPI. Images were taken using an Axio Imager 2 with Apotome (Carl Zeiss), and brightness and contrast were adjusted using Zen software (Carl Zeiss).

LAMP1 and cathepsin L co-localization. Single colour images of LAMP1 and cathepsin L staining were imported into ImageJ and thresholded to identify positive vesicles as greater than 10 pixels in area. The percent of these vesicles co-expressing cathepsin L per cell was measured. Histograms were computed using GraphPad Prism.

Quantitative PCR. Cells were sorted by flow cytometry into RNA Protect (Qiagen). mRNA was purified using either the RNeasy Mini or Plus Micro kits (Qiagen), and cDNA was synthesized using the iScript cDNA synthesis kit (Bio-Rad). Amplification of I-Ab (*H2-Ab1*), *H2-Mx*, *H2-DMβ2* and *HPRT* was performed using iQ SYBR Green Supermix (Bio-Rad) and amplification of *I-Ez* and *HPRT* was performed using TaqMan Master Mix (Life Technologies) on a MyiQ qPCR Detection System (Bio-Rad). SYBR Green reactions were run at 95 °C for 10 min followed by 40 cycles of 95 °C for 30 s and 60 °C for 1 min. TaqMan reactions were run at 50 °C for 2 min, 95 °C for 10 min and 40 cycles of 95 °C for 15 s and 60 °C for 1 min. Primers used were the following: *Hprt* forward, 5'-AGG TTGCAAGCTTGCTGGT-3', and reverse, 5'-TGAAGTACTCATTATAGTCAA GGGCA-3'; *H2-Ma* forward, 5'-CTCGAAGCATCTACACCAGTG-3', and reverse, 5'-TCCGAGAGCCCTATGTTGGG-3'; *H2-DMβ2* forward, 5'-GTTGGCTTCTT CAGATGGCG-3', and reverse, 5'-TGCCGTCCTTCTGGGTAGG-3'; *H2-Ab1* forward, 5'-GGTGTGCAGACACAACACTACG-3', and reverse, 5'-CGACATTGGG CTGTTCAAGC. Taqman probes (Life Technologies) used were Mm00446968_m1 for *HPRT* and Mm00772352_m1 for *H2-Ea-ps* (*I-Ez*). Ct values were normalized to *HPRT* and relative expression compared with B cells or dendritic cells was reported as $2^{-\Delta\Delta Ct}$.

BMDCs. Bone marrow cells were isolated from the tibias and femurs of B6 mice, cultured *in vitro* in RPMI-1640 with 10% FCS, 2 mM L-glutamine, 30 mM HEPES, 0.05 mM β-mercaptoethanol and 0.05 mg ml⁻¹ gentamycin, supplemented with 1,000 U ml⁻¹ GM-CSF (BD Biosciences) and 100 U ml⁻¹ IL-4 (eBioscience). Medium was changed on day 2 and the top half of the medium was changed on day 5 of culture. On day 7, BMDCs were harvested, enriched for CD11c⁺ cells (Miltenyi Biotec) and activated overnight with NIH 3T3 cells expressing CD40L⁷⁰ (provided by R. Lapoint, University of Montreal). Activated BMDCs were pulsed with 50 μg ml⁻¹ of β-gal₇₂₁₋₇₃₉ (AENLSVTLPAASHAIPHLT, GenScript) for 3 h, washed and 1 × 10⁵ BMDCs were injected IV.

Statistical analysis. Statistical analysis of data was performed using GraphPad Prism 5. *P* values <0.05 were considered statistically significant; **P*<0.05; ***P*<0.01; ****P*<0.001, NS = not significant. Two-tailed unpaired *t*-tests were used for comparisons between two groups, and a one-way analysis of variance with Tukey or Bonferroni post-test was used for three or more groups. All data shown as mean ± s.e.m.

References

- Hinterberger, M. *et al.* Autonomous role of medullary thymic epithelial cells in central CD4(+) T cell tolerance. *Nat. Immunol.* **11**, 512–519 (2010).
- Ramsdell, F. & Fowlkes, B. J. Clonal deletion versus clonal anergy: the role of the thymus in inducing self tolerance. *Science* **248**, 1342–1348 (1990).
- Bonasio, R. *et al.* Clonal deletion of thymocytes by circulating dendritic cells homing to the thymus. *Nat. Immunol.* **7**, 1092–1100 (2006).
- Baba, T., Nakamoto, Y. & Mukaida, N. Crucial contribution of thymic Sirp alpha+ conventional dendritic cells to central tolerance against blood-borne antigens in a CCR2-dependent manner. *J. Immunol.* **183**, 3053–3063 (2009).
- Derbinski, J., Schulte, A., Kyewski, B. & Klein, L. Promiscuous gene expression in medullary thymic epithelial cells mirrors the peripheral self. *Nat. Immunol.* **2**, 1032–1039 (2001).
- Anderson, M. S. *et al.* Projection of an immunological self shadow within the thymus by the aire protein. *Science* **298**, 1395–1401 (2002).
- Belz, G. T. *et al.* The CD8α+ dendritic cell is responsible for inducing peripheral self-tolerance to tissue-associated antigens. *J. Exp. Med.* **196**, 1099–1104 (2002).
- Gardner, J. M. *et al.* Deletional tolerance mediated by extrathymic Aire-expressing cells. *Science* **321**, 843–847 (2008).
- Gardner, J. M. *et al.* Extrathymic Aire-expressing cells are a distinct bone marrow-derived population that induce functional inactivation of CD4(+) T cells. *Immunity* **39**, 560–572 (2013).
- Cohen, J. N. *et al.* Lymph node-resident lymphatic endothelial cells mediate peripheral tolerance via Aire-independent direct antigen presentation. *J. Exp. Med.* **207**, 681–688 (2010).
- Fletcher, A. L. *et al.* Lymph node fibroblastic reticular cells directly present peripheral tissue antigen under steady-state and inflammatory conditions. *J. Exp. Med.* **207**, 689–697 (2010).
- Tewalt, E. F. *et al.* Lymphatic endothelial cells induce tolerance via PD-L1 and lack of costimulation leading to high-level PD-1 expression on CD8 T cells. *Blood* **120**, 4772–4782 (2012).
- Lee, J. W. *et al.* Peripheral antigen display by lymph node stroma promotes T cell tolerance to intestinal self. *Nat. Immunol.* **8**, 181–190 (2007).
- Nichols, L. A. *et al.* Deletional self-tolerance to a melanocyte/melanoma antigen derived from tyrosinase is mediated by a radio-resistant cell in peripheral and mesenteric lymph nodes. *J. Immunol.* **179**, 993–1003 (2007).
- Pardoll, D. M. The blockade of immune checkpoints in cancer immunotherapy. *Nat. Rev. Cancer* **12**, 252–264 (2012).
- Cohen, J. N. *et al.* Tolerogenic properties of lymphatic endothelial cells are controlled by the lymph node microenvironment. *PLoS One* **9**, e87740 (2014).
- Denzin, L. K., Sant'Angelo, D. B., Hammond, C., Surman, M. J. & Cresswell, P. Negative regulation by HLA-DO of MHC class II-restricted antigen processing. *Science* **278**, 106–109 (1997).
- Dolan, B. P., Gibbs, K. D. & Ostrand-Rosenberg, S. Tumor-specific CD4+ T cells are activated by 'cross-dressed' dendritic cells presenting peptide-MHC class II complexes acquired from cell-based cancer vaccines. *J. Immunol.* **176**, 1447–1455 (2006).
- Dubrot, J. *et al.* Lymph node stromal cells acquire peptide-MHCII complexes from dendritic cells and induce antigen-specific CD4+ T cell tolerance. *J. Exp. Med.* **211**, 1153–1166 (2014).
- Soriano, P. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nature Genet.* **21**, 70–71 (1999).
- Pham, T. H. *et al.* Lymphatic endothelial cell sphingosine kinase activity is required for lymphocyte egress and lymphatic patterning. *J. Exp. Med.* **207**, 17–27 (2010).
- Bazigou, E. *et al.* Genes regulating lymphangiogenesis control venous valve formation and maintenance in mice. *J. Clin. Invest.* **121**, 2984–2992 (2011).
- Karpanen, T. & Alitalo, K. Molecular biology and pathology of lymphangiogenesis. *Annu. Rev. Pathol. Mech. Dis* **3**, 367–397 (2008).
- Donohue, K. B. *et al.* Cross-priming utilizes antigen not available to the direct presentation pathway. *Immunology* **119**, 63–73 (2006).
- Tewalt, E. F. *et al.* Viral sequestration of antigen subverts cross presentation to CD8(+) T cells. *PLoS Pathog.* **5**, e1000457 (2009).
- Lukacs-Kornek, V. *et al.* Regulated release of nitric oxide by nonhematopoietic stroma controls expansion of the activated T cell pool in lymph nodes. *Nat. Immunol.* **12**, 1096–1104 (2011).
- Khan, O. *et al.* Regulation of T cell priming by lymphoid stroma. *PLoS One* **6**, e26138 (2011).
- Strawbridge, A. B. & Blum, J. S. Autophagy in MHC class II antigen processing. *Curr. Opin. Immunol.* **19**, 87–92 (2007).
- Comber, J. D., Robinson, T. M., Siciliano, N. A., Snook, A. E. & Eisenlohr, L. C. Functional macroautophagy induction by influenza A virus without a contribution to major histocompatibility complex class II-restricted presentation. *J. Virol.* **85**, 6453–6463 (2011).
- Kirberg, J. *et al.* Thymic selection of CD8+ single positive cells with a class II major histocompatibility complex-restricted receptor. *J. Exp. Med.* **180**, 25–34 (1994).
- Morgan, D. J. *et al.* CD8(+) T cell-mediated spontaneous diabetes in neonatal mice. *J. Immunol.* **157**, 978–983 (1996).
- Rudensky, A. Y., Rath, S., Preston-Hurlburt, P., Murphy, D. B. & Janeway, C. A. On the complexity of self. *Nature* **353**, 660–662 (1991).
- Dani, A. *et al.* The pathway for MHCII-mediated presentation of endogenous proteins involves peptide transport to the endo-lysosomal compartment. *J. Cell Sci.* **117**, 4219–4230 (2004).
- Nedjic, J., Aichinger, M., Emmerich, J., Mizushima, N. & Klein, L. Autophagy in thymic epithelium shapes the T-cell repertoire and is essential for tolerance. *Nature* **455**, 396–400 (2008).
- Koonce, C. H. *et al.* DM loss in k haplotype mice reveals isotype-specific chaperone requirements. *J. Immunol.* **170**, 3751–3761 (2003).
- Lazaraki, C. A., Chaves, F. A. & Sant, A. J. The impact of DM on MHC class II-restricted antigen presentation can be altered by manipulation of MHC-peptide kinetic stability. *J. Exp. Med.* **203**, 1319–1328 (2006).
- Menges, P. R. *et al.* An MHC class II restriction bias in CD4 T cell responses toward I-A is altered to I-E in DM-deficient mice. *J. Immunol.* **180**, 1619–1633 (2008).
- Pape, K. A., Merica, R., Mondino, A., Khoruts, A. & Jenkins, M. K. Direct evidence that functionally impaired CD4+ T cells persist *in vivo* following induction of peripheral tolerance. *J. Immunol.* **160**, 4719–4729 (1998).
- Grubin, C. E., Kovats, S., deRoos, P. & Rudensky, A. Y. Deficient positive selection of CD4 T cells in mice displaying altered repertoires of MHC class II-bound self-peptides. *Immunity* **7**, 197–208 (1997).

40. Chianese-Bullock, K. A. *et al.* Antigen processing of two H2-IEd-restricted epitopes is differentially influenced by the structural changes in a viral glycoprotein. *J. Immunol.* **161**, 1599–1607 (1998).
41. Zhou, H., Su, H. S., Zhang, X., Douhan, J. & Glimcher, L. H. CIITA-dependent and -independent class II MHC expression revealed by a dominant negative mutant. *J. Immunol.* **158**, 4741–4749 (1997).
42. Buch, T. *et al.* MHC class II expression through a hitherto unknown pathway supports T helper cell-dependent immune responses: implications for MHC class II deficiency. *Blood* **107**, 1434–1444 (2006).
43. Hake, S. B., Tobin, H. M., Steimle, V. & Denzin, L. K. Comparison of the transcriptional regulation of classical and non-classical MHC class II genes. *Eur. J. Immunol.* **33**, 2361–2371 (2003).
44. Arancibia-Carcamo, C. V. *et al.* A CIITA-independent pathway that promotes expression of endogenous rather than exogenous peptides in immune-privileged sites. *Eur. J. Immunol.* **34**, 471–480 (2004).
45. Collinge, M., Pardi, R. & Bender, J. R. Class II transactivator-independent endothelial cell MHC class II gene activation induced by lymphocyte adhesion. *J. Immunol.* **161**, 1589–1593 (1998).
46. Magner, W. J. *et al.* Activation of MHC class I, II, and CD40 gene expression by histone deacetylase inhibitors. *J. Immunol.* **165**, 7017–7024 (2000).
47. Landmann, S. *et al.* Maturation of dendritic cells is accompanied by rapid transcriptional silencing of class II transactivator (CIITA) expression. *J. Exp. Med.* **194**, 379–391 (2001).
48. Tai, A. K.-F., Zhou, G., Chau, K. & Ono, S. J. Cis-element dependence and occupancy of the human invariant chain promoter in CIITA-dependent and -independent transcription. *Mol. Immunol.* **36**, 447–460 (1999).
49. Momburg, F. *et al.* Differential expression of Ia and Ia-associated invariant chain in mouse tissues after *in vivo* treatment with IFN- γ . *J. Immunol.* **136**, 940–948 (1986).
50. Muczynski, K. A., Cotner, T. & Anderson, S. K. Unusual expression of human lymphocyte antigen class II in normal renal microvascular endothelium. *Kidney Int.* **59**, 488–497 (2001).
51. Onder, L. *et al.* A novel bacterial artificial chromosome-transgenic podoplanin-cre mouse targets lymphoid organ stromal cells *in vivo*. *Front. Immunol.* **2**, 50 (2011).
52. Baptista, A. P. *et al.* Lymph node stromal cells constrain immunity via MHC class II self-antigen presentation. *eLife* **3**, e04433 doi:10.7554/eLife.04433 (2014).
53. Segal, E. I. *et al.* Role of lymphocyte activation gene-3 (Lag-3) in conventional and regulatory T cell function in allogeneic transplantation. *PLoS One* **9**, e86551 (2014).
54. Taniguchi, R. T. *et al.* Detection of an autoreactive T-cell population within the polyclonal repertoire that undergoes distinct autoimmune regulator (Aire)-mediated selection. *Proc. Natl Acad. Sci. USA* **109**, 7847–7852 (2012).
55. Gallegos, A. M. & Bevan, M. J. Central tolerance to tissue-specific antigens mediated by direct and indirect antigen presentation. *J. Exp. Med.* **200**, 1039–1049 (2004).
56. Hubert, F. X. *et al.* Aire regulates the transfer of antigen from mTECs to dendritic cells for induction of thymic tolerance. *Blood* **118**, 2462–2472 (2011).
57. Koble, C. & Kyewski, B. The thymic medulla: a unique microenvironment for intercellular self-antigen transfer. *J. Exp. Med.* **206**, 1505–1513 (2009).
58. Aschenbrenner, K. *et al.* Selection of Foxp3⁺ regulatory T cells specific for self antigen expressed and presented by Aire⁺ medullary thymic epithelial cells. *Nat. Immunol.* **8**, 351–358 (2007).
59. Perry, J. S. A. *et al.* Distinct contributions of Aire and antigen-presenting-cell subsets to the generation of self-tolerance in the thymus. *Immunity* **41**, 414–426 (2014).
60. Klein, L., Roettinger, B. & Kyewski, B. Sampling of complementing self-antigen pools by thymic stromal cells maximizes the scope of central T cell tolerance. *Eur. J. Immunol.* **31**, 2476–2486 (2001).
61. Aichinger, M., Wu, C., Nedjic, J. & Klein, L. Macroautophagy substrates are loaded onto MHC class II of medullary thymic epithelial cells for central tolerance. *J. Exp. Med.* **210**, 287–300 (2013).
62. Tamburini, B. A., Burchill, M. A. & Kedl, R. M. Antigen capture and archiving by lymphatic endothelial cells following vaccination or viral infection. *Nat. Commun.* **5**, 3989 (2014).
63. Mazzini, E., Massimiliano, L., Penna, G. & Rescigno, M. Oral tolerance can be established via gap junction transfer of fed antigens from CX3CR1⁺ macrophages to CD103⁺ dendritic cells. *Immunity* **40**, 248–261 (2014).
64. Testa, J. S., Apcher, G. S., Comber, J. D. & Eisenlohr, L. C. Exosome-driven antigen transfer for MHC class II presentation facilitated by the receptor binding activity of influenza hemagglutinin. *J. Immunol.* **185**, 6608–6616 (2010).
65. Steinman, R. M., Turley, S., Mellman, I. & Inaba, K. The induction of tolerance by dendritic cells that have captured apoptotic cells. *J. Exp. Med.* **191**, 411–416 (2000).
66. Albert, M. L., Sauter, B. & Bhardwaj, N. Dendritic cells acquire antigen from apoptotic cells and induce class I-restricted CTLs. *Nature* **392**, 86–89 (1998).
67. Pérarnau, B. *et al.* Single H2Kb, H2Db and double H2KbDb knockout mice: peripheral CD8⁺ T cell repertoire and antilymphocytic choriomeningitis virus cytolytic responses. *Eur. J. Immunol.* **29**, 1243–1252 (1999).
68. Saxena, A. *et al.* Cutting edge: multiple sclerosis-like lesions induced by effector CD8^T cells recognizing a sequestered antigen on oligodendrocytes. *J. Immunol.* **181**, 1617–1621 (2008).
69. Fallas, J. L., Yi, W., Draghi, N. A., O'Rourke, H. M. & Denzin, L. K. Expression patterns of H2-O in mouse B cells and dendritic cells correlate with cell function. *J. Immunol.* **178**, 1488–1497 (2007).
70. Kershaw, M. H. *et al.* Immunization against endogenous retroviral tumor-associated antigens. *Cancer Res.* **61**, 7920–7924 (2001).

Acknowledgements

We thank Holly Davis and Kara Cummings for excellent technical support, and the UVA Flow Cytometry and Lymphocyte Culture Cores, which were supported by NCI Cancer Center Support Grant 5 P30 CA044579. This work was supported in part by USPHS grant AI068836, AI099033 and AI109250 to V.H.E., USPHS training grants AI07496 to S.J.R., E.F.T. and J.N.C.; GM007267 to S.J.R., J.D.P. and J.N.C.; CA009109 to J.D.P., and PF-10-156-01-LIB from the American Cancer Society to E.F.T. P.R. was supported by the CAPES Foundation, Ministry of Education of Brazil, Brasília—DF, 70.040-020. T.M. was supported by Cancer Research UK.

Author contributions

S.J.R., J.D.E., J.D.P., E.F.T., J.N.C. and V.H.E. conceived and designed the experiments. S.J.R., J.D.E., P.R., J.D.P., E.F.T. and J.N.C. performed and analysed the experiments. R.L. and T.M. contributed reagents. S.J.R. and V.H.E. wrote the paper.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no conflict of interest.

Reprints and permission information is available online at <http://npng.nature.com/reprintsandpermissions/>

How to cite this article: Rouhani, S. J. *et al.* Roles of lymphatic endothelial cells expressing peripheral tissue antigens in CD4 T-cell tolerance induction. *Nat. Commun.* **6**:6771 doi: 10.1038/ncomms7771 (2015).