

SCIENTIFIC REPORTS



OPEN

Association of a rare variant of the *TNFSF13B* gene with susceptibility to Rheumatoid Arthritis and Systemic Lupus Erythematosus

David González-Serna¹, Lourdes Ortiz-Fernández¹, Sofía Vargas¹, Antonio García², Enrique Raya³, Benjamín Fernández-Gutierrez⁴, Francisco Javier López-Longo⁵, Alejandro Balsa⁶, Isidoro González-Álvaro⁷, Javier Narvaez⁸, Carmen Gómez-Vaquero⁸, José Mario Sabio⁹, Rosa García-Portales¹⁰, María Francisca González-Escribano¹¹, Carles Tolosa¹², Patricia Carreira¹³, Lambertus Kiemeny¹⁴, Marieke J. H. Coenen¹⁵, Torsten Witte¹⁶, Matthias Schneider¹⁷, Miguel Ángel González-Gay¹⁸ & Javier Martín¹

A rare variant (BAFF-var) of the tumor necrosis factor superfamily 13b (*TNFSF13B*) gene has been recently associated with multiple sclerosis (MS) and systemic lupus erythematosus (SLE). The aim of this study was to investigate the association between *TNFSF13B* BAFF-var and susceptibility to rheumatoid arthritis (RA) and replicate that association in SLE. 6,218 RA patients, 2,575 SLE patients and 4,403 healthy controls from three different countries were included in the study. *TNFSF13B* BAFF-var was genotyped using TaqMan allelic discrimination assay. PLINK software was used for statistical analyses. *TNFSF13B* BAFF-var was significantly associated with RA ($p = 0.015$, OR = 1.21, 95% CI = 1.03–1.41) in the Spanish cohort. A trend of association was observed in the Dutch ($p = 0.115$) and German ($p = 0.228$) RA cohorts. A meta-analysis of the three RA cohorts included in this study revealed a statistically significant association ($p = 0.002$, OR = 1.24, 95% CI = 1.10–1.38). In addition, *TNFSF13B* BAFF-var was significantly associated with SLE in the Spanish ($p = 0.001$, OR = 1.41, 95% CI = 1.14–1.74) and the German cohorts ($p = 0.030$, OR = 1.86, 95% CI = 1.05–3.28), with a statistically significant p-value obtained in the meta-analysis ($p = 0.0002$, OR = 1.46, 95% CI = 1.09–2.32). The results obtained confirm the known association of *TNFSF13B* BAFF-var with SLE and, for the first time, demonstrate that this variant contributes to susceptibility to RA.

¹Institute of Parasitology and Biomedicine López-Neyra, CSIC, Granada, Spain. ²Rheumatology Department, Hospital Virgen de las Nieves, Granada, Spain. ³Systemic Autoimmune Diseases Unit, Hospital Campus de la Salud, Granada, Spain. ⁴Rheumatology Service, Hospital Clínico San Carlos, IDISSC, Madrid, Spain. ⁵Rheumatology Service, Hospital General Universitario Gregorio Marañón, Madrid, Spain. ⁶Rheumatology Department, Instituto de Investigación Hospital Universitario La Paz (IDIPAZ), Madrid, Spain. ⁷Rheumatology Department, Instituto de Investigación del Hospital de La Princesa (IIS-IP), Madrid, Spain. ⁸Rheumatology Service, Hospital Universitario de Bellvitge, Barcelona, Spain. ⁹Systemic Autoimmune Diseases Unit, Department of Internal Medicine, Hospital Virgen de las Nieves, Granada, Spain. ¹⁰Rheumatology Department, Hospital Virgen de la Victoria, Málaga, Spain. ¹¹Department of Immunology, Hospital Universitario Virgen del Rocío, IBIS, Sevilla, Spain. ¹²Department of Internal Medicine, Hospital Parc Taulí, Sabadell, Spain. ¹³Rheumatology Service, Hospital Universitario 12 de Octubre, Madrid, Spain. ¹⁴Department for Health Evidence, Radboud Institute for Health Sciences, Radboud university medical center, Nijmegen, The Netherlands. ¹⁵Department of Human Genetics, Radboud Institute for Health Sciences, Radboud university medical center, Nijmegen, The Netherlands. ¹⁶Department for Clinical Immunology and Rheumatology, Hannover Medical School, Hannover, Germany. ¹⁷Policlinic and Hiller Research Unit for Rheumatology, UKD, Heinrich-Heine-University Dusseldorf, Dusseldorf, Germany. ¹⁸Epidemiology, Genetics and Atherosclerosis Research Group on Systemic Inflammatory Diseases IDIVAL, Hospital Universitario Marqués de Valdecilla, Santander, Spain. Correspondence and requests for materials should be addressed to J.M. (email: javiermartin@ipb.csic.es)

Rheumatoid arthritis (RA) and systemic lupus erythematosus (SLE) are complex autoimmune diseases influenced by both genetic and environmental factors^{1,2}. Genome-wide association studies (GWAS) have revealed more than 100 independent signals for RA and 50 for SLE associated with these diseases^{3,4}. In a recent study Steri *et al.*⁵ identified an association signal in the *TNFSF13B* gene which was associated with multiple sclerosis (MS) and SLE. *TNFSF13B* encodes tumor necrosis factor superfamily member 13b (also known as B-cell activating factor (BAFF)). BAFF is a cytokine that is primarily produced by monocytes and neutrophils and plays a crucial role in B-cell homeostasis and the regulation of B-cell maturation, differentiation and survival⁶. This cytokine plays a particularly important role in the pathogenesis of SLE, to such an extent that the first targeted therapy approved for SLE was Belimumab, a monoclonal antibody targeting human BAFF^{7–9}. The causal variant identified by Steri *et al.*⁵ for this association, “BAFF-var”, is an insertion-deletion in which five nucleotides are replaced by one (GCTGT > A), being this A the risk allele. Persons carrying the A allele have an increased risk for the disease. This risk allele results in a shorter transcript that escapes microRNA inhibition thus leading to an increase in the production of soluble BAFF. Besides, it has been observed that *TNFSF13B* BAFF-var is strongly associated with augmented levels of total IgG and IgM and with reduced monocyte counts⁵. Thus, *TNFSF13B* BAFF-var is considered a shared genetic risk variant for autoimmune diseases based on its association with MS and SLE, and it plays an important role in autoimmune processes⁵. Considering that replication of previous results is of vital importance for the correct development of scientific knowledge, we performed a replication study to corroborate the association of *TNFSF13B* BAFF-var with susceptibility to SLE.

Otherwise, BAFF system dysregulation is involved in the pathogenesis of RA^{1,10} and abnormal levels of BAFF have been detected in serum, synovial fluid and saliva from RA patients¹¹. It is noteworthy that more than 10 shared risk loci have been identified for SLE and RA^{12,13}. Taking into considerations above, we aimed to assess for the first time the potential association of *TNFSF13B* BAFF-var with RA in patients from Spain, the Netherlands and Germany.

Materials and Methods

Study subjects. Overall, 6,218 RA patients and 4,403 healthy controls from Spain, Germany and the Netherlands were enrolled in our study. Regarding the Spanish cohort, DNA was obtained from 4,429 RA patients from the Xeral-Calde University Hospital, Lugo; Marqués de Valdecilla University Hospital, Santander; Bellvitge University Hospital, Barcelona; Virgen del Rocío University Hospital, Seville; Campus de la Salud University Hospital, Granada; and Hospital Clínico San Carlos, La Princesa University Hospital, La Paz University Hospital, 12 de Octubre University Hospital and Gregorio Marañón University Hospital, all located in Madrid. DNA from 3,200 healthy controls was obtained from Banco Nacional de ADN. DNA was obtained from 890 RA patients and 733 controls from the Netherlands at the Radboud University Medical Center (Nijmegen) and the Nijmegen Biomedical Study (NBS)¹⁴ respectively. DNA was obtained from 890 RA patients and 470 controls of German descent at the Hannover Medical School (Hannover) and Düsseldorf University Hospital. All the patients included in this study were of European descent and had been diagnosed with RA according to the 1987 classification criteria of the American College of Rheumatology¹⁵.

A total of 2,575 SLE patients from Spain and Germany were included in this study. DNA from 1,160 patients from Spain was obtained at Xeral-Calde University Hospital, Lugo; Virgen del Rocío University Hospital, Sevilla; Virgen de las Nieves University Hospital, Granada; Virgen de la Victoria University Hospital, Málaga, and Parc Taulí University Hospital, Sabadell. DNA from 460 German patients was obtained at the Hannover Medical School, Hannover. All SLE patients were of European descent and fulfilled the American College of Rheumatology criteria for the classification of SLE¹⁶.

The study was approved by the local ethical committees of the different participating centers (Ethics Committee of the Spanish National Research Council, Xeral-Calde University Hospital, Marqués de Valdecilla University Hospital, Bellvitge University Hospital, Virgen del Rocío University Hospital, Campus de la Salud University Hospital, Hospital Clínico San Carlos, La Princesa University Hospital, La Paz University Hospital, 12 de Octubre University Hospital and Gregorio Marañón University Hospital, Virgen de las Nieves University Hospital, Virgen de la Victoria University Hospital, Parc Taulí University Hospital, Radboud University Medical Center, Hannover Medical School and Düsseldorf University Hospital) in accordance with the tenets of the Declaration of Helsinki. Written informed consent was obtained from all participants prior to enrolment in the study.

Genotyping. Genomic DNA was isolated from peripheral blood samples using standard salting-out techniques. *TNFSF13B* BAFF-var was genotyped using a TaqMan allelic discrimination custom assay (assay ID: AH0JGPG) (Applied Biosystems, Foster City, California, USA) on an ABI 7900HT Fast Real-Time PCR System (Thermo Fisher). Genotyping call rate was >95% in all sample sets.

Statistical analysis. All the statistical analyses were carried out with PLINK v1.9 (<http://pngu.mgh.harvard.edu/purcell/plink/>)¹⁷. For all groups of individuals, Hardy-Weinberg equilibrium was determined at a significance level of 0.01. Differences between allele and genotype frequencies in individuals were determined by χ^2 test. Odds ratios (ORs) and 95% confidence intervals were calculated according to Woolf’s method. Meta-analyses were performed with inverse-variance method under a fixed-effects model. Heterogeneity of the ORs across cohorts was assessed using both Cochran’s Q test and inconsistency index (I^2). The statistical power of our study was calculated by using Power Calculator for Genetic Studies 2006 (CaTS) software (<http://www.sph.umich.edu/csg/abecasis/CaTS/>)¹⁸, under an additive model. The statistical power of the study for determining significance at a p value of 0.05 is shown in Supplementary Table S1.

Subgroup (N)	Genotype (N)			MAF (%)	Allele test	
	A A	A GCTGT	GCTGT GCTGT		P-value	OR [95% CI] ^a
Spain						
Controls (3,200) ^b	10	249	2941	4.20		
RA (4,429)	11	424	3994	5.03	0.015	1.21 [1.03–1.41]
Netherlands						
Controls (733)	0	28	705	1.91		
RA (890)	0	49	841	2.75	0.115	1.47 [0.91–2.36]
Germany						
Controls (470) ^b	0	19	451	2.02		
RA (899)	3	44	852	2.78	0.228	1.39 [0.81–2.36]
Meta-analysis^c						
Controls (4,403)						
RA (6,218)					0.002	1.24 [1.10–1.38]

Table 1. Association analysis of the *TNFSF13B* BAFF-var in three independent RA cohorts and meta-analysis. Abbreviations: N: Number of individuals; MAF: Minor Allele Frequency; OR: Odds Ratio; CI: Confidence Interval. ^aOR for the minor allele. ^bHealthy controls were used as reference for RA and SLE patients in Spanish and German subgroups. ^cHeterogeneity q value = 0.64, I² = 0.

Data availability. The complete datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Ethical use of human participants statement. All participants signed a written informed consent prior to their enrolment in the study for identified data to be published. All procedures were performed after signed informed consent was obtained from patients and in accordance with the local (Spanish National Research Council) ethics committee and the Declaration of Helsinki of 1975, as revised in 1983.

Results

First, in order to evaluate the association between the *TNFSF13B* gene variant (BAFF-var) and susceptibility to RA, a comparative analysis of allelic and genotype frequencies in RA patients vs. healthy individuals was performed in the Spanish, Dutch and German populations. Results are shown in Table 1. The minor allele frequency (A) of *TNFSF13B* BAFF-var was higher in RA patients in the three populations included in this study. This difference was statistically significant in the Spanish subgroup ($p = 0.015$, OR = 1.21, 95% CI = 1.03–1.41), and suggestive associations were observed in the Dutch ($p = 0.115$, OR = 1.47, 95% CI = 0.91–2.36) and German ($p = 0.228$, OR = 1.39, 95% CI = 0.81–2.36) subgroups. In addition, a statistically significant p value was obtained in the meta-analysis of the three RA populations included in this study ($p = 0.002$, OR = 1.24, 95% CI = 1.10–1.38) with low heterogeneity levels (heterogeneity q value = 0.64, I² = 0) (Table 1).

Secondly, to try to replicate the results previously published by Steri *et al.*⁵, a comparative analysis of allelic and genotype frequencies in SLE patients vs. healthy individuals from the Spanish and German populations was also performed (Table 2). The minor allele frequency (A) in *TNFSF13B* BAFF-var was statistically significantly higher in SLE patients than in controls in both, the Spanish ($p = 0.0001$, OR = 1.41, 95% CI = 1.14–1.74) and German ($p = 0.030$, OR = 1.86, 95% CI = 1.05–3.28) subgroups. Meta-analysis of the two SLE populations included in this study revealed a statistically significant p value ($p = 0.0002$, OR = 1.46, 95% CI = 1.09–2.32) with low heterogeneity levels (heterogeneity q value = 0.37, I² = 0) (Table 2).

Discussion

We confirm the recently reported association of the *TNFSF13B* BAFF-var with SLE in two independent populations of Spanish and German origin. In addition, this is the first time that evidence is provided of an association between *TNFSF13B* BAFF-var and RA. These findings, together with the data reported by Steri *et al.*⁵, suggest that *TNFSF13B* BAFF-var is a common genetic risk factor for autoimmunity.

SLE and RA mice models showed increased serum BAFF levels, and BAFF blockade reduced diseases manifestations^{19,20}. Nevertheless, although clinical trials are being performed on BAFF in RA and SLE patients, only moderate results are emerging. An example of these modest results is the BAFF-inhibitor drug belimumab, which has been found to be only moderately effective in a small number of patients with RA and SLE^{8,21}. Because RA and SLE are heterogeneous diseases with different biological subsets, the efficacy of different drugs could be divergent in specific patient subgroups. Therefore, our new findings could help in the advance of RA therapies, as patients stratified by *TNFSF13B* BAFF-var status may show a differential benefit from anti-BAFF therapies. In this sense, a recent study showed that a *TNFSF13B* genetic variant influenced response to B-cell targeted therapy (rituximab) in patients with RA²². Furthermore, B-cell-depleting therapies in patients carrying *TNFSF13B* BAFF-var would have a weaker effect due to a rapid resurgence of memory B cells induced by high soluble BAFF levels, which could increase the risk of inadequate response and/or relapse²³.

Genome-wide association studies (GWAS) have identified multiple risk genetic factors associated with human autoimmune disease. Of note, GWAS are designed to detect only common genetic variants (>5%); however, low-frequency and rare variants may represent an important component of autoimmune risk and provide a key

Subgroup (N)	Genotype (N)			MAF (%)	Allele test	
	A A	A GCTGT	GCTGT GCTGT		P-value	OR [95% CI] ^a
Spain						
Controls (3,200) ^b	10	249	2941	4.20		
SLE (1,160)	9	117	1034	5.81	0.001	1.41 [1.14–1.74]
Germany						
Controls (470) ^b	0	19	451	2.02		
SLE (460)	1	32	427	3.69	0.030	1.86 [1.05–3.28]
Meta-analysis^c						
Controls (4,403)						
SLE (6,218)					0.0002	1.46 [1.09–2.32]

Table 2. Association analysis of the *TNFSF13B* BAFF-var in two independent SLE cohorts and meta-analysis. Abbreviations: N: Number of individuals; MAF: Minor Allele Frequency; OR: Odds Ratio; CI: Confidence Interval. ^aOR for the minor allele. ^bHealthy controls were used as reference for RA and SLE patients in Spanish and German subgroups. ^cHeterogeneity q value = 0.37, I² = 0.

insight into both novel and previously implicated immunological pathways that are disrupted in autoimmune diseases²⁴. Thus, large genetic studies have identified low frequency variants with relevant functional significance associated with autoimmune diseases (e.g. *TREX1* in SLE²⁵ and *TYK2* in several autoimmune diseases^{26,27}). Our findings support the important role of rare variants, such as *TNFSF13B* BAFF-var, in understanding the unknown mechanisms of autoimmunity.

For the first time, our results showed that allele A of *TNFSF13B* BAFF-var was associated with the risk of RA in the Spanish population. Trends of association were observed in the Dutch and German RA cohorts. This could be due to a low statistical power as the Dutch and the German cohorts sample sizes are relatively small as compared to the Spanish cohort, as well as the minor allele frequency is less common in northern Europe (2.02% in Germany and 1.91% in the Netherlands) than in southern Europe (4.2% in Spain). This north-south gradient in the minor allele frequency of *TNFSF13B* BAFF-var has been previously reported, with higher frequencies observed in southern Europe (5.7% in Italy and 4.9% in Spain) than in northern Europe (1.8% in United Kingdom and Sweden)⁵. However, the three RA cohorts showed higher levels of allele A in cases than in controls and statistically significant association was observed in the meta-analysis. Thus, further studies involving larger cohorts would be necessary to elucidate if this variant is associated with autoimmune risk in northern populations.

Regarding the SLE replication study, our results confirm the association of allele A of *TNFSF13B* BAFF-var with the risk of SLE in the Spanish and German populations. The association of the variant was found to be stronger in SLE than in RA and, in both cases, the allele A acts as a risk factor.

GWAS have revealed a significant shared genetic component in autoimmunity^{28,29}. The association of *TNFSF13B* BAFF-var with RA and SLE observed in our study extends the number of overlapping genetic risk factors across different autoimmune diseases. In this regard, performing further studies to elucidate the specific association of *TNFSF13B* BAFF-var with other autoimmune diseases would be of great interest as new therapy approaches involving BAFF could emerge.

In conclusion, our results suggest that *TNFSF13B* BAFF-var plays an important role in susceptibility to RA, and confirm the association of this variant with SLE.

References

- McInnes, I. B. & Schett, G. The Pathogenesis of Rheumatoid Arthritis. *N. Engl. J. Med.* **365**, 2205–2219 (2011).
- Tsokos, G. C. Systemic Lupus Erythematosus. *N. Engl. J. Med.* **365**, 2110–2121 (2011).
- Okada, Y. *et al.* Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature*. **506**, 376–381 (2013).
- Morris, D. L. *et al.* Genome-wide association meta-analysis in Chinese and European individuals identifies ten new loci associated with systemic lupus erythematosus. *Nat. Genet.* **48**, 940–946 (2016).
- Steri, M. *et al.* Overexpression of the Cytokine BAFF and Autoimmunity Risk. *N. Engl. J. Med.* **376**, 1615–1626 (2017).
- Woodland, R. T., Schmidt, M. R. & Thompson, C. B. BlyS and B cell homeostasis. *Semin. Immunol.* **18**, 318–326 (2006).
- Vincent, F. B., Morand, E. F., Schneider, P. & Mackay, F. The BAFF/APRIL system in SLE pathogenesis. *Nat. Rev. Rheumatol.* **10**, 365–373 (2014).
- Furie, R. *et al.* A phase III, randomized, placebo-controlled study of belimumab, a monoclonal antibody that inhibits B lymphocyte stimulator, in patients with systemic lupus erythematosus. *Arthritis Rheum.* **63**, 3918–3930 (2011).
- Navarra, S. V., Guzman, R. M. & Gallacher, A. E. Efficacy and safety of belimumab in patients with active systemic lupus erythematosus: a randomised, placebo-controlled, phase 3 trial. *Lancet*. **377**, 721–731 (2011).
- Wei, F., Chang, Y. & Wei, W. The role of BAFF in the progression of rheumatoid arthritis. *Cytokine*. **76**, 537–544 (2015).
- Bosello, S. *et al.* Concentrations of BAFF Correlate with Autoantibody Levels, Clinical Disease Activity, and Response to Treatment in Early Rheumatoid Arthritis. *J. Rheumatol.* **35**, 1256–1264 (2008).
- Márquez, A. *et al.* A combined large-scale meta-analysis identifies COG6 as a novel shared risk locus for rheumatoid arthritis and systemic lupus erythematosus. *Ann. Rheum. Dis.* **76**, 286–294 (2017).
- Orozco, G. *et al.* Study of the common genetic background for rheumatoid arthritis and systemic lupus erythematosus. *Ann. Rheum. Dis.* **70**, 463–8 (2011).
- Wetzels, J. F., Kiemeny, L. A., Swinkels, D. W., Willems, H. L. & den Heijer, M. Age- and gender-specific reference values of estimated GFR in Caucasians: the Nijmegen Biomedical Study. *Kidney Int.* **72**, 632–637 (2007).
- Arneti, F. C. *et al.* The American Rheumatism Association 1987 Revised Criteria for the Classification of Rheumatoid Arthritis. *Arthritis Rheum.* **31**, 316–324 (1988).

16. Tan, E. M. *et al.* The 1982 Revised Criteria for the Classification of Systemic Lupus Erythematosus. *Arthritis Rheum.* **25**, 1271–1277 (1982).
17. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).
18. Skol, A. D., Scott, L. J., Abecasis, G. R. & Boehnke, M. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat. Genet.* **38**, 209–213 (2006).
19. Ding, H. *et al.* Blockade of B-cell-activating factor suppresses lupus-like syndrome in autoimmune BXSB mice. *J. Cell Mol. Med.* **14**, 1717–1725 (2009).
20. Lai Kwan Lam, Q., King Hung Ko, O., Zheng, B. J. & Lu, L. Local BAFF gene silencing suppresses Th17-cell generation and ameliorates autoimmune arthritis. *Proc. Natl. Acad. Sci. USA.* **105**, 14993–14998 (2008).
21. Stohl, W. *et al.* Efficacy and Safety of Belimumab in Patients with Rheumatoid Arthritis: A Phase II, Randomized, Double-blind, Placebo-controlled, Dose-ranging Study. *J. Rheumatol.* **40**, 579–589 (2013).
22. Juge, P. A. *et al.* Variants of genes implicated in type 1 interferon pathway and B-cell activation modulate the EULAR response to rituximab at 24 weeks in rheumatoid arthritis. *RMD Open.* **3**, e000448, <https://doi.org/10.1136/rmdopen-2017-000448> (2017).
23. Dass, S. *et al.* Highly sensitive B cell analysis predicts response to rituximab therapy in rheumatoid arthritis. *Arthritis Rheum.* **58**, 2993–2999 (2008).
24. Massey, J. & Eyre, S. Rare variants and autoimmune disease. *Brief. Funct. Genomics.* **13**, 392–397 (2014).
25. Lee-Kirsch, M. A. *et al.* Mutations in the gene encoding the 3'-5' DNA exonuclease TREX1 are associated with systemic lupus erythematosus. *Nat. Genet.* **39**, 1065–1067 (2007).
26. Dendrou, C. A. *et al.* Resolving TYK2 locus genotype-to-phenotype differences in autoimmunity. *Sci. Transl. Med.* **8**, 363–149 (2016).
27. Diogo, D. *et al.* TYK2 protein-coding variants protect against rheumatoid arthritis and autoimmunity, with no evidence of major pleiotropic effects on non-autoimmune complex traits. *PLoS One.* **10**, e0122271, <https://doi.org/10.1371/journal.pone.0122271> (2015).
28. Cho, J. H. & Gregersen, P. K. Genomics and the multifactorial nature of human autoimmune disease. *N. Engl. J. Med.* **365**, 1612–1623 (2011).
29. Cho, J. H. & Feldman, M. Heterogeneity of autoimmune diseases: pathophysiologic insights from genetics and implications for new therapies. *Nat. Med.* **21**, 730–738 (2015).

Acknowledgements

We thank Sonia Garcia, and Gema Robledo for their excellent technical assistance, and all the patients and healthy controls for kindly accepting participating in this study. This work was supported by the following grants: P12-BIO-1395 from Consejería de Innovación, Ciencia y Tecnología, Junta de Andalucía (Spain), and the Cooperative Research Thematic Network (RETICS) program, RD16/0012/0004 (RIER), from Instituto de Salud Carlos III (ISCIII, Health Ministry, Madrid, Spain). DGS was supported by the Spanish Ministry of Economy and Competitiveness through the program FPI (SAF2015-66761-P).

Author Contributions

D.G.S., L.O.F. and J.M. were involved in the conception and design of the study as well as in the interpretation of data. D.G.S. performed the statistical analyses and drafted the manuscript. J.M. critically revised the manuscript for important intellectual content. S.V., A.G., E.R., B.F.G., F.J.L.L., A.B., I.G.A., J.N., C.G.V., J.M.S., R.G.P., M.F.G.E., C.T., P.C., B.K., M.J.H.C., T.W., M.S. and M.A.G.G. contributed samples and/or data acquisition and all authors revised and approved the final version of the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-26573-4>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018