

SCIENTIFIC REPORTS



OPEN

Hagfish: Champions of CO₂ tolerance question the origins of vertebrate gill function

Daniel W. Baker¹, Brian Sardella², Jodie L. Rummer³, Michael Sackville⁴ & Colin J. Brauner⁴

Received: 02 December 2014

Accepted: 07 April 2015

Published: 09 June 2015

The gill is widely accepted to have played a key role in the adaptive radiation of early vertebrates by supplanting the skin as the dominant site of gas exchange. However, in the most basal extant craniates, the hagfishes, gills play only a minor role in gas exchange. In contrast, we found hagfish gills to be associated with a tremendous capacity for acid-base regulation. Indeed, Pacific hagfish exposed acutely to severe sustained hypercarbia tolerated among the most severe blood acidoses ever reported (1.2 pH unit reduction) and subsequently exhibited the greatest degree of acid-base compensation ever observed in an aquatic chordate. This was accomplished through an unprecedented increase in plasma [HCO₃⁻] (>75 mM) in exchange for [Cl⁻]. We thus propose that the first physiological function of the ancestral gill was acid-base regulation, and that the gill was later co-opted for its central role in gas exchange in more derived aquatic vertebrates.

In terrestrial vertebrates the division of gas exchange and ion/acid-base homeostasis between the lungs and kidney, respectively, is well established. In the marine environment, where vertebrate life first evolved, the vertebrate gill is used for all of these processes¹. The gill was crucial to the adaptive radiation of early vertebrates, allowing for increased body size, activity level and skin mineralization². However, the gill could not have initially evolved to satisfy all of these processes simultaneously. Indeed, it has long been held that the primary selective pressure driving early gill function was for increased O₂ uptake, and thus gas exchange (termed the O₂ hypothesis²⁻⁴). However, as protovertebrates increased in size and activity, the inability of the skin to maintain acid-base and ionoregulatory homeostasis may have been more limiting than that for gas exchange⁴. This appears to be the case during development in teleosts, where these processes shift from the skin to the gills long before gas exchange^{5,6}. We thus propose that the primary selective pressure shaping early vertebrate gill evolution was for an increased acid-base relevant ionoregulatory capacity, which is supported by our findings presented here on hagfish, the most basal extant craniate (Fig. 1).

While hagfish have many derived traits, they also retain several key ancestral features deemed representative of protovertebrates⁷. Three that are particularly relevant to our hypothesis are the following: 1) they are the most basal extant deuterostome to have filamentous gills that possess a semi-permeable barrier separating blood from the external environment, 2) they have a high epidermal surface area to volume ratio (SA:V) due to a long, thin body plan (Figs. 1 and 3) they are osmo-ionoconformers. In fact, hagfish are the only extant craniates that are both osmo- and ionoconformers, with their plasma composition closely resembling seawater^{8,9}. This is thought to represent the ancestral state because hagfish probably never invaded freshwater, and all of the more basal, extant deuterostomes (e.g., hemichordata, echinodermata) are also osmo-ionoconformers, in contrast to all of the more derived craniates^{1,7}.

¹Department of Fisheries and Oceans, Vancouver Island University, Nanaimo, BC, Canada, V9R 5S5. ²Department of Biological Sciences, California State University, Stanislaus, California, One University Circle, Turlock, CA, USA.

³ARC Centre of Excellence for Coral Reef Studies, James Cook University, 1 James Cook Drive, Townsville, QLD 4811 Australia. ⁴Department of Zoology, University of British Columbia, 6270 University Blvd., Vancouver, B.C., Canada, V6T 1Z4. Correspondence and requests for materials should be addressed to D.W.B. (email: dan.baker@viu.va)




Figure 1. Chordate phylogeny of major extinct (boxed) and extant (unboxed) taxa (modified^{34,35}) with maximum estimated total lengths (length data³⁶). Divergence times are based on best estimates of combined molecular and fossil data^{35,37}. The star represents the approximate appearance of gill filaments in the fossil record, which we propose were predominantly associated with acid-base regulation. Major increases in body size and/or reductions in body surface area to volume ratio occurred after this event and likely signify the transition of acid/base ion regulation from the skin to the gills.

As osmo-ionoconformers, hagfish gills are not involved in ionoregulation for the purposes of ion (i.e., Na^+ and Cl^-) balance as in other fishes and interestingly, their gills do not appear to play much of a role in gas exchange. Instead, 80–90% of oxygen uptake in hagfishes occurs across the skin at rest^{10,11} due to the high epidermal SA:V characteristic of all extant or extinct early fishes (Fig. 1). Given the high dependence upon cutaneous respiration, perhaps it is not surprising that hagfish have the lowest oxygen consumption rate (MO_2) of any craniate at rest or following severe stress^{12,13}. A few studies have provided evidence for acid-base relevant ionoregulation in the hagfish gill^{14–16} by identifying associated gill transporters^{17–19} and cell types with a recent study providing the first direct support for acid extrusion *via* the gills through use of a divided chamber system¹⁶. However, acid-base regulatory capacity (as informed by the rate and degree of pH compensation following a disturbance) in response to an environmentally induced acidosis (as opposed to acid injection) has not been specifically investigated for comparison with other aquatic craniates.

Acid-base regulation is one of the most tightly regulated physiological processes in animals²⁰. Changes in blood and cellular pH impact protein charge, and the consequences to protein function can impair everything from enzyme function, cellular ion transport, muscle contractility and metabolism through to survival²¹. While the mechanisms and capacity for acid-base regulation *via* the teleost gill have been reasonably well studied, much less is known about the hagfish gill²². Given the intimate association between ion and acid equivalent exchange during acid-base compensation at the teleost gill¹ and that hagfish are ionoconformers⁹, one might expect limited acid-base compensatory capacity. Contrastingly, aspects of hagfish life history suggest otherwise: hagfish burrow in soft sediments and use their toothed tongue to enter dead animal carcasses while feeding where they may remain for extended periods²³. These conditions promote aquatic hypoxia (low environmental O_2) and hypercarbia (elevated environmental CO_2), both of which can severely disrupt acid-base status. Indeed, Pacific hagfish, *Eptatretus stoutii*, are supremely tolerant of hypoxia and anoxia¹³. However, hypercarbia tolerance in hagfishes is unknown¹⁴, and while elevated CO_2 is an environmentally relevant challenge, hypercarbia exposure can also be used as a tool to quantify acid-base regulatory capacity. Here we exposed hagfish to sustained hypercarbia to induce a rapid acidosis. Subsequent recovery was used to quantify the rate and degree of blood acid-base compensation for direct comparison with other aquatic craniates, which has not been previously investigated.




Figure 2. The effect of elevated water pCO₂ (10 (●), 30 (▼) and 50 (■) mm Hg pCO₂) for up to 96 h on A) blood pH (pHe) and intracellular pH of B) red blood cell, C) liver, D) heart and E) parietal muscle homogenates of the Pacific hagfish. All fish survived exposure until the time of sampling. Values are means ± SEM. “*” indicates a statistically significant difference from time 0 values (p < 0.05). In panel A, “+” indicates a statistically significant pH recovery from lowest pH point within a CO₂ exposure.

Results and Discussion:

Pacific hagfish were acutely exposed to a water pCO₂ of 10, 30 or 50 mm Hg (1.3, 4 and 6 kPa CO₂) for up to 96 h, to simulate the most extreme conditions that hagfish could possibly experience in a benthic burrow, within a decaying carcass, in proximity to deep sea hydrothermal vents²⁴, or near point sources such as CO₂ injection sites²⁵. As expected, hypercarbia triggered an immediate acidosis in Pacific hagfish that varied with the severity of hypercarbia. Within 3 h, blood pH (pHe) decreased from 7.99 (±0.02) to 7.62 (±0.01), 7.04 (±0.03) and 6.81 (±0.03) with exposure to 10, 30 and 50 mm Hg pCO₂, respectively (Fig. 2). Tissue (muscle, heart and liver) intracellular pH (pHi) decreased in parallel with pHe (Fig. 2), but to a lesser degree, in accordance with their higher buffering capacity (Supplementary Fig. 1) relative to blood (Fig. 3). Despite the severe acidosis, among the greatest ever reported for surviving water breathers, hagfish compensated by elevating pHe as early as 6 h but always well before the end of the 96-h exposure. Compensation of pHe was both rapid and extensive, resulting in 95% (at 24 h), 70% (at 48 h),




Figure 3. A pH/HCO₃⁻/CO₂ diagram of pHe compensation in Pacific hagfish during exposure to 10 (●), 30 (▼) and 50 (■) mm Hg pCO₂. Values are means ± SEM. Curved long-dashed lines represent pCO₂ isopleths and the short-dashed line represents the blood buffer line (5 mM [HCO₃⁻] pH-1 blood). Data labels represent duration of CO₂ exposure (h). Shaded area represents the “bicarbonate concentration threshold” associated with pHe compensatory limits in other fish as described previously²⁰.

and 65% (at 96 h) recovery at 10, 30, and 50 mm Hg pCO₂, respectively [calculated as % return of pH to pre-exposure (i.e., 0 hr) values from the 3 h value at the respective CO₂ tension].

Blood pH recovery was associated with elevated plasma [HCO₃⁻] (Figs. 3,4) and an equimolar reduction in plasma [Cl⁻] (Fig. 4). Net plasma HCO₃⁻/Cl⁻ exchange during exposure to hypercarbia is the typical pattern observed in teleosts²⁰, and these data imply it likely represents the basal condition. Other plasma ions were unchanged ([Na⁺], [Mg²⁺], and [Ca²⁺]; Fig. 3). The calculated net acid excretion rate for hagfish was similar to that of other fish species investigated (Supplementary Table 1), but what stands out in the physiological response to hypercarbia is the degree of pHe compensation, as well as the associated quantitative changes in plasma [HCO₃⁻] and [Cl⁻]. No other water-breathing craniate has been reported to either tolerate ~1.2 pH blood acidosis or to recover pHe to this degree. This impressive pHe compensation during acute hypercarbia was driven entirely by an unprecedented increase in plasma [HCO₃⁻], in exchange for [Cl⁻], which reached 78.2 (±4.5) and 75.4 (±8.2) mM during exposure to 30 and 50 mm Hg pCO₂ (Figs. 3,4), respectively. These values are over twice the next highest plasma [HCO₃⁻] ever reported for a water-breathing vertebrate during acute exposure to hypercarbia²⁰. Typically, water-breathing fish exposed to acute (<96 h) hypercarbia are unable to elevate blood HCO₃⁻ beyond 25–33 mM, termed the “bicarbonate concentration threshold”²⁰ (Fig. 3). In any case, the gills of hagfish appear to be an efficacious structure for acid-base regulation with a compensatory capacity that far exceeds that of any other aquatic craniate investigated to date.

We believe that the hagfish’s tremendous upper limit for blood acid-base compensation may be associated with its’ osmo-ionoconforming strategy and consequent high plasma [Cl⁻] (458 mM; Fig. 3B) providing more anions available in the blood for HCO₃⁻ exchange. Teleosts typically have plasma [Cl⁻] of 130–150 mM¹, and during acute hypercarbia, about 17–20% of the plasma [Cl⁻] can be exchanged with HCO₃⁻ before the bicarbonate concentration threshold is reached. Thus, complete pHe compensation during acute hypercarbia is limited to 10–16 mm Hg pCO₂. In hagfish, the increase in plasma [HCO₃⁻] during hypercarbia reached a value of 78.2 ± 4.5 mM (Fig. 4A), which corresponded to 17% of control plasma [Cl⁻] (Fig. 4B), a similar proportion to that observed in teleosts that are able to compensate for acute hypercarbia. We suggest therefore that the degree of pH compensation attainable in fishes during acute hypercarbia may be limited by the relative decrease in plasma Cl⁻ levels and so linked to the importance of avoiding hypochloremia. While there are few opportunities to test this hypothesis, the greater CO₂ tolerance of elasmobranchs compared to teleosts²⁶ may also be a result of higher plasma [Cl⁻], which is intermediate between that of teleosts and hagfish¹. Lamprey, another agnathan, have




Figure 4. The effect of elevated water $p\text{CO}_2$ (10 (●), 30 (▼) and 50 (■) mm Hg $p\text{CO}_2$) on plasma A) $[\text{HCO}_3^-]$, B) $[\text{Cl}^-]$, D) $[\text{Na}^+]$, E) $[\text{Ca}^{2+}]$, and F) $[\text{Mg}^{2+}]$ in Pacific hagfish (mean \pm SEM). In C), bars represent the change (relative to controls) in $[\text{HCO}_3^-]$ (hatched), $[\text{Cl}^-]$ (cross hatched), and net difference between $\text{HCO}_3^- - \text{Cl}^-$ (open) during exposure to a water $p\text{CO}_2$ of 10 (blue), 30 (yellow) and 50 (red) mm Hg (approximately 1.5, 4 and 6 kPa). “*” indicates a statistically significant difference from time 0 values ($p < 0.05$).

similar plasma $[\text{Cl}^-]$ to teleosts and have a very limited ability to compensate for acute hypercarbia (R. Shartau, personal communication), indicating that the exceptional hypercarbia tolerance of hagfish is not necessarily an agnathan trait.

In contrast, the ancestral vertebrate was likely an osmo-ionoconformer, as discussed above, where plasma $[\text{Cl}^-]$ may have limited the extent of acid-base compensation, and the gill dictated the rate of acid extrusion. If so, the gradual lowering of plasma $[\text{Cl}^-]$ associated with the evolution of ionoregulation, although leading to niche expansion and radiation into environments of varying salinity^{27,28}, may also have reduced the ceiling on acid-base compensation. Indeed, other fishes with lower plasma $[\text{Cl}^-]$ that can tolerate extreme hypercarbia to levels similar to those tolerated by hagfish in this study do so using a very different strategy. The most basal actinopterygian (white sturgeon; *Acipenser transmontanus*²⁹) and a few species of air breathing teleosts (marbled swamp eel; *Synbranchus marmoratus*³⁰; armoured catfish; *Pterygoplichthys pardalis*³¹) can tolerate hypercarbia (~ 40 mmHg; 5–6 kPa $p\text{CO}_2$) in the absence of pHe compensation (pHe is depressed and remains low) or active accumulation of plasma $[\text{HCO}_3^-]$ and instead, preferentially and completely regulate tissue pHi ^{29,31}. Thus, preferential pHi regulation may have

evolved to tolerate acid-base disturbances in the face of reduced plasma $[Cl^-]$ within the actinopterygians (Fig. 1) and may be a trait associated with the evolution of air breathing in fish³¹.

Recently, there has been considerable interest in CO_2 tolerance in marine animals, and an increased effort in estimating historical atmospheric CO_2 levels to help predict the effects of climatic trends²⁵. A few strategies have been proposed to sequester CO_2 into deep ocean sites to reduce atmospheric “greenhouse” gases, but these methods could generate historically unprecedented local CO_2 tensions to which deep-sea animals may be exposed²⁵. In particular, there is concern that deep-sea animals may be especially sensitive to acid-base disturbances because they often have relatively low blood buffer capacity, low metabolic rates, and limited ion exchange capacity^{32,33}. Accordingly, hagfish embody one of the most “at-risk” deep-sea animals, having among the lowest metabolic rate of any fish investigated to date¹², relatively poorly buffered blood, and are thought to have extremely limited ion exchange capacity⁹. In contrast, this study indicates that hagfish may be among the most capable of aquatic vertebrates to cope with acid-base disturbances and tolerate high CO_2 levels. Given that hagfishes are abundant, demersal fishes and play an important role in nutrient cycling, their exceptional CO_2 tolerance may prove significant given some of the proposed CO_2 disposal scenarios. It is intriguing that the most basal extant craniate, which may have remained relatively unchanged for hundreds of millions of years⁷, may turn out to be the most suited aquatic animal to survive in a high CO_2 world.

In summary, the findings of this research indicate that the ancestral function of the vertebrate gill may have been predominantly acid-base regulation with a small role in gas exchange. From this we propose that increased capacity for acid-base regulation, rather than gas exchange, may have been the primary selective pressure shaping early evolution of the vertebrate gill. Clearly, more research on hagfish and other phylogenetically relevant animals is warranted to further test the hypothesis that the first physiological function of the ancestral vertebrate gill was acid-base relevant ionoregulation, and the gill was later co-opted for its central role in gas exchange in more derived vertebrate species.

Methods:

Pacific hagfish (*Eptatretus stoutii*; 100–400 g) were exposed to seawater equilibrated with approximately 10, 30 and 50 mm Hg pCO_2 and then sampled either a) immediately after transfer, time 0, or b) after 3, 6, 12, 24, 48 (only 30 and 50 mm Hg pCO_2) or 96 (only 50 mm Hg pCO_2) h of exposure to elevated pCO_2 . Blood was obtained from anaesthetized animals for pH, hematocrit, haemoglobin, and mean cell haemoglobin concentration (MCHC) as previously described²⁹. Plasma total CO_2 (TCO_2), plasma ion composition, and RBC pH_i were also measured²⁹. Tissue pH_i was measured from frozen tissues using the metabolic inhibitor method²⁹. Non-bicarbonate whole blood buffer capacity and tissue non-bicarbonate buffer capacity was determined as described previously²⁹, calculated from the slope of $\Delta[HCO_3^-] \Delta pH^{-1}$, and then expressed in $mmol HCO_3^- pH^{-1} l^{-1}$ of blood or kg^{-1} of intracellular tissue water, over an *in vivo* relevant pH range. Data are presented as mean \pm SEM ($n = 8$ in all cases except one, where $n = 7$). All data was analyzed for normality and equal variance before statistical analysis. Statistical differences were detected using a one-way ANOVA and, when necessary, a post-hoc Dunnett’s test. All statistical analyses were conducted using SigmaStat for Windows 3.5.0.54 (Systat Software, Inc., 2006), and all analyses were 2-tailed and interpreted using $\alpha = 0.05$ to determine statistical significance.

For comparison with other species, an estimate of the increase in whole animal net acid excretion rates in hagfish exposed to hypercarbia was calculated (Supplementary Table 1) as the inverse of the net increase in whole body $[HCO_3^-]$ following CO_2 exposure relative to pre-exposure (i.e., time 0) values as has been done previously for other aquatic species^{29,31} (see supplemental information for details^{38–48}). Overall, net acid excretion rates in hagfish were similar to those determined in other fish as has been observed previously¹⁵ (Supplementary Table 1).

References

- Evans, D. H., Piermarini, P. M. & Choe, K. P. The multifunctional fish gill: Dominant site of gas exchange, osmoregulation, acid-base regulation and excretion of nitrogenous waste. *Physiol. Rev.* **85**, 97–177 (2005).
- Gans, C. Stages in the origin of vertebrates: analysis by means of scenarios. *Biol. Rev.*, **64**, 221–268 (1989).
- Krogh, A. Ed. *The comparative physiology of respiratory mechanisms* (Univ. Pennsylvania Press, PA, 1941).
- Rombough, P. The functional ontogeny of the teleost gill: which comes first, gas or ion exchange? *Comp. Biochem. Physiol. A*, **148**, 732–742 (2007).
- Fu, C., Wilson, J. M., Rombough, P. J. & Brauner, C. J. Ions first: Na^+ uptake shifts from the skin to the gills before O_2 uptake in developing rainbow trout, *Oncorhynchus mykiss*. *Proc. Roy. Soc. B. Biol. Sci.* **277**, 1553–1560 (2010).
- Brauner, C. J. & Rombough, P. J. Ontogeny and paleophysiology of the gill: new insights from larval and air-breathing fish. *Respir. Physiol. Neurobiol.* **184**, 293–300 (2012).
- Janvier, P. Living primitive fishes and fishes from deep time, in *Primitive Fishes* [McKenzie, D. J., Farrell, A. P. & Brauner, C. J. (eds.)], Fish Physiology Series, [vol. 26, 1–51] (Elsevier, New York, 2007).
- Robertson, J. D. Chemical composition of the body fluids and muscle of the hagfish *Myxine glutinosa* and the rabbit-fish *Chimaera monstrosa*. *J. Zool. Lond.* **178**, 261–277 (1976).
- Sardella, B. A., Baker, D. W. & Brauner, C. J. The effects of variable water salinity and ionic composition on the plasma status of the Pacific hagfish (*Eptatretus stoutii*). *J. Comp. Physiol. B*, **179**, 721–728 (2009).
- Steffensen, J. F., Johansen, K., Sindberg, C. D., Sorensen, J. H. & Moller, J. L. Ventilation and oxygen consumption in the hagfish, *Myxine glutinosa* L. *J. Expt. Mar. Biol. Ecol.* **84**, 173–178 (1984).
- Lesser, M. P., Martini, F. H. & Heiser, J. B. Ecology of the hagfish, *Myxine glutinosa* L. in the Gulf of Maine I. Metabolic rates and energetics. *J. Expt. Mar. Biol. Ecol.* **208**, 215–225 (1996).

12. Forster, M. E. Confirmation of the low metabolic rate of hagfish. *Comp. Biochem. Physiol. A*, **96**, 113–116 (1990).
13. Cox, G. K., Sandblom, E., Richards, J. G. & Farrell, A. P. Anoxic survival of the Pacific hagfish (*Eptatretus stoutii*). *J. Comp. Physiol. B* **181**, 361–371 (2011).
14. Evans, D. H. Gill Na^+/H^+ and $\text{Cl}^-/\text{HCO}_3^-$ exchange systems evolved before the vertebrates entered fresh water. *J. Exp. Biol.* **113**, 465–469 (1984).
15. McDonald, D. G., Cavdek, V., Calvert, L. & Milligan, C. L. Acid-base regulation in the Atlantic hagfish *Myxine glutinosa*. *J. Exp. Biol.* **161**, 201–215 (1991).
16. Clifford, A. M., Guffey, S. C. & Goss, G. G. Extrabranchial mechanisms of systemic pH recovery in hagfish (*Eptatretus stoutii*). *Comp. Biochem. Physiol. A* **168**, 82–89 (2014).
17. Mallat, J., Conley, D. M. & Ridgway, R. L. Why do hagfish have gill “chloride cells” when they need not regulate plasma NaCl concentration? *Can. J. Zool.* **65**, 1956–1965 (1987).
18. Edwards, S. L., Claiborne, J. B., Morrison-Shetlar, A. I. & Toop, T. Expression of Na^+/H^+ exchanger mRNA in the gills of the Atlantic hagfish (*Myxine glutinosa*) in response to metabolic acidosis. *Comp. Biochem. Physiol. A* **130**, 81–91 (2001).
19. Tresguerres, M., Parks, S. K. & Goss, G. G. V-H^+ -ATPase, Na^+ , K^+ -ATPase and NHE_2 immunoreactivity in the gill epithelium of the Pacific hagfish (*Eptatretus stoutii*). *Comp. Biochem. Physiol. A* **145**, 312–321 (2006).
20. Heisler, N. Acid-base regulation in fishes in *Acid-Base Regulation in Animals* [Heisler, N. (ed.)] 1st Edition [309–356] (Elsevier, Amsterdam, 1986).
21. Putnam, R. & Roos, A. Intracellular pH in *Handbook of Physiology* [Hoffman, J., Jamieson, J. (eds.)] [vol. **14**, 389–440] (Oxford: Oxford Univ. Press, 1997).
22. Wright, P. A. Ionic, osmotic and nitrogenous waste excretion in *Fish Physiology* [McKenzie, D. J., Farrell, A. P. & Brauner, C. J. (eds.)], [vol. **26**, 283–319] (Elsevier, New York, 2007).
23. Strahan, R. The behaviour of myxinoids. *Acta zoologica*, **44**, 73–102 (1963).
24. Møller, P. R. & Jones, J. J. *Eptatretus strickrotti* (sp. Myxinidae): First hagfish captured from a hydrothermal vent. *Biological Bulletin*, **212**, 55–66 (2007).
25. Seibel, B. A. & Walsh, P. J. Potential impacts of CO_2 injection on deep-sea biota. *Science*, **294**, 319–320 (2001).
26. Ishimatsu, A., Kikkawa, T., Hayashi, M., Lee, S. K. & Kita, J. Effects of CO_2 on marine fish: Larvae and adults. *J. Oceanogr.* **60**, 731–741 (2004).
27. Piermarini, P. M. & Evans, D. H. Immunochemical analysis of the vacuolar proton-ATPase B-subunit in the gills of a euryhaline stingray (*Dasyatis sabina*): effects of salinity and relation to Na^+/K^+ -ATPase. *J. Exp. Biol.* **204**, 3251–3259 (2001).
28. Bray, A. A. The evolution of terrestrial vertebrates: environmental and physiological conditions. *Phil. Trans. Roy. Soc. Lond. B* **309**, 289–322. (1985).
29. Baker, D. W. *et al.* Complete intracellular pH protection during extracellular pH depression is associated with hypercapnia tolerance in white sturgeon, *Acipenser transmontanus*. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **296**, R1868–R1880 (2009).
30. Heisler, N. Intracellular and extracellular acid-base regulation in the tropical fresh-water teleost fish *Synbranchus marmoratus* in response to the transition from water breathing to air breathing. *J. Exp. Biol.*, **99**, 9–28 (1982).
31. Brauner, C. J. *et al.* Limited extracellular but complete intracellular acid-base regulation during short-term environmental hypercapnia in the armoured catfish, *Liposarcus pardalis*. *J. Exp. Biol.* **207**, 3381–3390 (2004).
32. Seibel, B. A. & Walsh, P. J. Biological impacts of deep-sea carbon dioxide injection inferred from indices of physiological performance. *J. Exp. Biol.* **206**, 641–650 (2003).
33. Pane, E. F. & Barry, J. P. Extracellular acid-base regulation during short-term hypercapnia is effective in a shallow-water crab, but ineffective in a deep-sea crab. *Mar. Ecol. Prog. Ser.* **334**, 1–9 (2007).
34. Donoghue, P. C., Forey, P. L. & Aldridge, R. J. Conodont affinity and chordate phylogeny. *Biol. Rev.* **75**, 191–251 (2000).
35. Kardong, K. V. *Vertebrates: Comparative Anatomy, Function, Evolution*. Sixth Edition, (McGraw-Hill, New York, 2012).
36. Albert, J. S., Johnson, D. M. & Knouft, J. H. Fossils provide better estimates of ancestral body size than do extant taxa in fishes. *Acta Zool.*, **90**, 357–384 (2009).
37. Erwin, D. H. *et al.* The Cambrian Conundrum: Early Divergence and Later Ecological Success in the Early History of Animals. *Science*, **334**, 1091–1097 (2011).
38. Zeidler, R. & Kim, H. D. Preferential hemolysis of postnatal calf red cells induced by internal alkalization. *J. Gen. Physiol.* **70**, 385 (1977).
39. Pörtner, H. O., Boutilier, R. G., Tang, Y. & Toews, D. P. Determination of intracellular pH and PCO_2 after metabolic inhibition by fluoride and nitrilotriacetic acid. *Respir. Physiol.* **81**, 255–273 (1990).
40. Baker, D. W., May, C. & Brauner, C. J. A validation of intracellular pH measurements in fish exposed to hypercapnia: The effect of duration of tissue storage and efficacy of the metabolic inhibitor tissue homogenate method. *J. Fish. Biol.* **75**, 268–275 (2009).
41. Boutilier, R. G., Heming, T. A., & Iwama, G. K. Appendix: Physicochemical parameters for use in fish respiratory physiology in *Fish Physiology* [Hoar, W. S. & Randall, D. J. (eds.)] [vol. **10A**, 403–430] (Academic Press Inc., New York, 1984).
42. Forster, M. E., Russell, M. J., Hambleton, D. C. & Olson, K. R. Blood and extracellular fluid volume in whole body and tissues of the Pacific hagfish (*Eptatretus stoutii*). *Physiol. Biochem. Zool.* **74**, 750–756 (2001).
43. Wilkie, M., Couturier, J. & Tufts, B. Mechanisms of acid-base regulation in migrant sea lampreys (*Petromyzon marinus*) following exhaustive exercise. *J. Exp. Biol.* **201**, 1473–1482 (1998).
44. Heisler, N., Weitz, H. & Weitz, A. M. Extracellular and intracellular pH with changes of temperature in the Dogfish *Scyliorhinus stellaris*. *Respir. Physiol.* **26**, 249–263 (1976).
45. Toews, D. P., Holeton, G. F. & Heisler, N. Regulation of the acid-base status during environmental hypercapnia in the marine teleosts *Conger conger*. *J. Exp. Biol.* **107**, 9–20 (1983).
46. Evans, D. H. Mechanisms of acid extrusion by two marine fishes: the teleost, *Opsanus beta*, and the elasmobranch, *Squalus acanthias*. *J. Exp. Biol.* **97**, 289–299 (1982).
47. Claiborne, J. B., Perry, E., Bellows, S. & Campbell, J. Mechanisms of acid-base excretion across the gills of a marine fish. *J. Exp. Zool.* **279**, 509–520 (1997).
48. Claiborne, J. B. & Heisler, N. Acid-base regulation and ion transfers in the carp (*Cyprinus carpio*): pH compensation during graded long- and short-term environmental hypercapnia and the effect of bicarbonate infusion. *J. Exp. Biol.* **126**, 41–61 (1986).

Acknowledgments

This project was funded by a Natural Sciences and Engineering Research Council Discovery grant to CJB. The authors thank Bruce Cameron and the Bamfield Marine Sciences Centre for their technical expertise and use of facilities respectively. Thanks are due to Erin Walsh for artistic enhancement of Fig. 1. We also acknowledge K.T. Huynh for technical support and P.J. Rombough for many stimulating discussions.

Author Contributions

D.W.B., B.A.S., J.L.R. and C.J.B. designed and performed the experiments. D.W.B. measured and analyzed blood and tissue parameters. J.L.R. performed blood buffer experiments and developed figures. M.A.S. provided Fig. 1 and helped expand evolutionary concepts. All authors discussed the results and commented on the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Baker, D. W. *et al.* Hagfish: Champions of CO₂ tolerance question the origins of vertebrate gill function. *Sci. Rep.* 5, 11182; doi: 10.1038/srep11182 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>