

SCIENTIFIC REPORTS



OPEN

Two types of soybean diacylglycerol acyltransferases are differentially involved in triacylglycerol biosynthesis and response to environmental stresses and hormones

Received: 09 December 2015

Accepted: 06 June 2016

Published: 27 June 2016

BeiBei Chen¹, Junejie Wang¹, Gaoyang Zhang¹, Jiaqi Liu², Sehrish Manan¹, Honghong Hu³ & Jian Zhao¹

Diacylglycerol acyltransferases (DGATs) play a key role in plant triacylglycerol (TAG) biosynthesis. Two type 1 and 2 DGATs from soybean were characterized for their functions in TAG biosynthesis and physiological roles. *GmDGAT1A* is highly expressed in seeds while *GmDGAT2D* is mainly expressed in flower tissues. They showed different expression patterns in response to biotic and abiotic stresses. *GmDGAT2D* was up-regulated by cold and heat stress and ABA signaling, and repressed by insect biting and jasmonate, whereas *GmDGAT1A* show fewer responses. Both *GmDGAT1A* and *GmDGAT2D* were localized to the endoplasmic reticulum and complemented the TAG deficiency of a yeast mutant H1246. *GmDGAT2D*-transgenic hairy roots synthesized more 18:2- or 18:1-TAG, whereas *GmDGAT1A* prefers to use 18:3-acyl CoA for TAG synthesis. Overexpression of both *GmDGATs* in Arabidopsis seeds enhanced the TAG production; *GmDGAT2D* promoted 18:2-TAG in wild-type but enhanced 18:1-TAG production in *rod1* mutant seeds, with a decreased 18:3-TAG. However, *GmDGAT1A* enhanced 18:3-TAG and reduced 20:1-TAG contents. The different substrate preferences of two DGATs may confer diverse fatty acid profiles in soybean oils. While *GmDGAT1A* may play a role in usual seed TAG production and *GmDGAT2D* is also involved in usual TAG biosynthesis in other tissues in responses to environmental and hormonal cues.

Soybean [*Glycine max* (L.) Merr.] is the largest oilseed crop in the world, providing almost 28% edible vegetable oils for human^{1,2}. However, compared to rapeseeds and sunflower seeds that usually contain around 40% oils, there might be a large space to improve the oil content in soybean seeds^{1,2}. The limited understanding of soybean oil biosynthesis has become a major obstacle for the improvement of soybean oil production³. In higher plants triacylglycerols (TAGs) are the most common storage lipids in seed as the form of lipid droplets. The biosynthesis of TAGs is not restricted to seeds, but widely occurs in most vegetative tissues, although their exact function remains to be determined^{2,4,5}. However, an increasing body of evidence suggests TAG and biosynthetic genes participate in plant stress or defense response^{2,4-6}. The plastid is the major place where the *de novo* biosynthesis of fatty acids (FAs) occurs; then these FAs are transported out of plastids in active forms into the cytosol, where fatty acyl-CoAs form precursor pools. Fatty acyl-CoAs and glycerol 3-phosphate are transported into the endoplasmic reticulum (ER) to synthesize TAGs and then assembled into oil bodies^{2,5,7}. The synthesis of phospholipids and TAGs in ER contains various acyltransferases, which have substrate specificities towards fatty acyl-CoAs as

¹National Key Laboratory of Crop Genetic Improvement, College of Plant Science & Technology, Huazhong Agricultural University, Wuhan, 430070, China. ²College of Agronomy, Jilin Agricultural University, Changchun, 130047, China. ³National Key Laboratory of Crop Genetic Improvement, College of Life Science & Technology, Huazhong Agricultural University, Wuhan, 430070, China. Correspondence and requests for materials should be addressed to J.Z. (email: 2948308723@qq.com)

donors and different glycerol backbones as acceptors. Thus acyltransferases not only conduct the acyl editing on DAG and eventual product TAGs in the ER, but also manipulation of these acyltransferases leads to increased TAG production in plants⁷.

Diacylglycerol acyltransferases (DGATs) catalyze the esterification of the acyl group of acyl-CoAs at the sn-3 position of sn-1, 2-diacylglycerol (DAG) molecules, the final step in the Kennedy pathway leading to TAG production. DGATs play essential roles in determining the quality and quantity of acyl-CoA flux into TAG synthesis through Kennedy pathway. At least four classes of DGATs have been identified based on structure and activity⁷. The most common is type 1 (DGAT1), which contributes to the synthesis of the majority of TAGs⁷. DGAT1 is usually predicted to possess six or more transmembrane domains and belongs to the membrane-bound O-acyltransferase (MBOAT) family⁸. Studies have confirmed that DGAT1 affects the TAG levels in seeds of many species^{9–13}. DGAT2 proteins possess only two to three predicted transmembrane domains and belongs to the monoacylglycerol acyltransferase (MGAT) family. A third type of DGATs is similar to DGAT2 with both wax ester synthase (WS) and DGAT activity, including ADP1 from *Acinetobacter calcoaceticus*¹⁴ and ADP1 homologs from Arabidopsis and petunia (*Petunia hybrida*)^{15,16}. A fourth type, a soluble cytosolic DGAT (DGAT3), was identified in peanut (*Arachis hypogaea*) and Arabidopsis^{17,18}.

DGAT1s from different plants, such as Arabidopsis, have been shown to have major contribution to TAG synthesis, whereas the role of DGAT2s from these plants is not fully understood^{19,20}. Arabidopsis DGAT2 has recently been reported for a much weaker function in TAG synthesis compared with DGAT1^{16,21–23}. However, the mostly characterized plant DGAT2s are active for synthesis of unusual TAG using hydroxyl, conjugated or epoxy FAs. For examples, DGAT2 in tung tree (*Vernicia fordii*) uses eleostearic acid, CpDGAT2 in *Claviceps purpurea* and RcDGAT2 in castor bean (*Ricinus communis*) use ricinoleate, and VgDGAT in *Vernonia galamensis* uses vernolic acid for production of unusual TAGs, including trioleostearin, triricinolein, or epoxy-TAG^{10,24–26}. DAGT2s from fungi and animals were shown active for TAG synthesis^{25,27–30}. Thus, knowledge about the redundant functions of DGAT1 and 2s from plants that do not synthesize hydroxy TAGs, such as Arabidopsis and soybean is still very limited²⁰. In particular, tetraploid Soybean genome contains at least 10 putative DGAT genes: 3 *DGAT1s*, 5 *DGAT2s*, and 2 *DGAT3s*^{16,31,32} (Supplementary Fig. S1). GmDGAT1A and GmDGAT1B are active in TAG synthesis in yeast and their transcripts are abundant in developing seeds and associated with maximal TAG accumulation¹⁶. GmDGAT2C played role in seed yield, whereas GmDGAT1B was associated with seed yield and protein concentration³². Since soybean seeds do not produce unusual oil but with such abundant copies of DGATs, it is of great interest and needs to be understood their functional redundancy, conservation, and divergence. Here, we report the functional comparison study on type 1 and 2 soybean diacylglycerol acyltransferases, GmDGAT1A and GmDGAT2D. The expression of both *GmDGAT1A* and *GmDGAT2D* restored the *in vivo* synthesis of TAG in yeast strain *Saccharomyces cerevisiae* H1246. *GmDGAT2D* expression in soybean hairy roots and Arabidopsis lead to similar function in regular TAG production like fungal and animal DGAT2, but differ from other known plant DGAT2s, which synthesize unusual TAGs with hydroxy FAs. Our study also suggests that *GmDGAT1A* and *2D* are differentially regulated by jasmonate during insect and wounding responses and ABA for cold and heat stress response, indicating their different functions in soybean stress responses.

Results

Identification of diacylglycerol acyltransferase 1A and 2D from *Glycine max*. To understand the functions of DGATs in biosynthesis of soybean oils that do not contain unusual TAGs, such as hydroxyl TAGs, we cloned *GmDGAT1A* and *GmDGAT2D*. Homology searches revealed GmDGAT2D shares highest identity with other plant DGAT2 genes from *Glycine soja* and *Phaseolus vulgaris*. GmDGAT2D shares 68% identity and 85% similarity with OeDGAT2 from olive (*Olea europaea*), 68% identity and 83% similarity with AtDGAT2, 67% identity and 81% similarity with HaDGAT2 from sunflower (*Helianthus annuus*), and 64% identity and 80% similarity with *Ricinus communis* RcDGAT2 (Supplementary Fig. S1). GmDGAT1A has nine predicted transmembrane (TM) domains whereas GmDGAT2D has only two TM domains, with a short head and a long tail inside the ER, and a cytosolic motif between TMs (Supplementary Fig. S1). These features are similar to fungal DGATs^{10,20,33}.

Quantitative RT-PCR (qRT-PCR) data indicated that *GmDGAT1A* is mainly expressed in seeds, but also in tissues like roots, leaves, flowers, at relatively lower levels. *GmDGAT2D* transcripts were high in flowers and seeds, but also present in other tissues at lower levels, such as roots, leaves and pods (Fig. 1a, Supplementary Fig. S2). The transcripts of *GmDGAT1A* and *2D* at soybean developing seeds were also examined (Fig. 1c). Both expression patterns of *GmDGAT1A* and *2D* are consistent with oil accumulation (Fig. 1d, Supplementary Fig. S2). The patterns seem also consistent with our transcriptomic data from soybean seeds at two developmental stages. The transcriptomic data showed that both *GmDGAT1A* and *2D* transcripts increased with development, and were higher in yellow seeds than in black seeds, which is in line with higher oil contents in yellow seeds than in black seeds (Fig. 1h). These results suggest that GmDGAT1A and 2D may play role in seed oil biosynthesis.

Analysis of *GUS* expression driven by *GmDGAT2D* promoter (pro*GmDGAT2D*::*GUS*) in transformed Arabidopsis suggested that the pro*GmDGAT2D*::*GUS* is mainly expressed in flower tissues, such as anther filament, stigma and petals. The *GUS* activity in leaf vascular tissues was also observed (Fig. 1e,g). Its expression in immature seeds was not observed. These results indicate that *GmDGAT2D* expression in non-seed tissues is relatively higher.

To further understand the functions of GmDGAT1A and 2D, we examined their subcellular localization. GmDGAT2D has two TM domains and GmDGAT1A with nine TM domains (Supplementary Fig. S1). GmDGAT1A - and 2D-green fluorescent protein (GFP) was co-localized with the CD3-959-mCherry, a well-known endoplasmic reticulum (ER) marker (Fig. 2a–h)³³. The distribution of GmDGAT2D-GFP fusion showed overlapping patterns with the ER. Thus two GmDGATs were primarily associated with the ER, as further referenced by chloroplast autofluorescence (Fig. 2i–l).




Figure 1. *GmDGAT1A* and *GmDGAT2D* expression pattern. The tissue expression patterns of *GmDGAT1A* and *GmDGAT2D* in nodule, root, stem, leaf, flower, pod, seed was tested by qRT-PCR with actin as internal standard. Seeds at different developmental stages were harvested. Six developmental stages are: stage 1. 40–70 mg; stage 2. 80–100 mg; stage 3. 150–200 mg; stage 4. 250–300 mg; stage 5. 350–430 mg; stage 6. 320–350 mg. (a) Tissue specific expression of *GmDGAT2D*. (b) Tissue specific expression of *GmDGAT1A*. (c) *GmDGAT1A* and *GmDGAT2D* expression at various seed developmental stages. (d) TAG accumulation and fatty acid composition changes in various seed developmental stages (e–g) *GmDGAT2D* promoter-driven *GUS* expression in Arabidopsis flower (e), *GmDGAT2D* promoter-driven *GUS* expression in Arabidopsis silique (f) and leaf (g). (h) *GmDGAT2D* and *GmDGAT1A* expression patterns in comparison of other DAGT genes. Soybean Illumina expression data obtained from developing seeds of two varieties with PI 84970 (Hokkaido, seed with a black seed coat) and PI 518671 (Williams 82, seed with yellow seed coat) at the developmental stages of 2 and 5. Data are presented as averages of three biological replicates \pm SD.




Figure 2. Subcellular localization of GmDGAT-1A and -2D. *GmDGAT1A-GFP*, *GmDGAT2D-GFP*, and an ER marker *CD3-959-mCherry*, driven by the cauliflower mosaic virus 35S promoter, were transiently expressed in tobacco leaf epidermal cells. Materials were viewed by confocal microscopy. Representative photos were shown. (a–d) Localization of GmDGAT1A-GFP. Image of GmDGAT1A-GFP (a), the blue fluorescence image of ER marker CD3-959-mCherry (b), and the merged image of GmDGAT1A-GFP and CD3-959-mCherry (c), and the bright field image (d). Bars = 20 μm . (e–h) Localization of GmDGAT2D-GFP. Image of GmDGAT2D-GFP (e), image of ER marker CD3-959-mCherry (f), the merged image of GmDGAT2D-GFP and ER marker CD3-959-mCherry (g), and the bright field image of cells (h). Bars = 30 μm . (i–l) Enlarged image of GmDGAT2D-GFP (i), image of ER marker (j), the chloroplast autofluorescence images (k), and merged image of GmDGAT2D-GFP, ER marker, and chloroplast autofluorescence in a bright field (l). Bars = 30 μm .

GmDGAT1A and GmDGAT2D are functional DGATs when expressed in yeast. The ORFs of *GmDGAT1A* and *GmDGAT2D* were expressed in the TAG-deficient yeast (*Saccharomyces cerevisiae*) strain H1246, in which all four acyltransferase genes contributing to TAG synthesis were deleted³⁴. When the *GmDGAT1A* and *GmDGAT2D* driven by a galactose-inducible GAL10 promoter in the vector pYESDEST52 were expressed in H1246 cells, many large and distinct oil bodies were observed in the Nile red-stained H1246 mutant cells after 24 h of galactose induction (Fig. 3a). Nile red has been broadly used for detecting TAG levels in numerous organisms, and used as a proper indicator of DGAT activity in H1246 strain³⁵. As expected, no fluorescence was observed in H1246 yeast cells transformed with empty vector. Furthermore, overexpression of *GmDGAT2D* in a wild-type yeast strain also showed enhanced density and size of intracellular oil bodies (Fig. 3a). Quantification of Nile red fluorescence intensity of these cells suggested that, the fluorescence intensity was very low in the control H1246 cells, but double in *GmDGATs*-expressing H1246 cells (Fig. 3b). Furthermore, *GmDGAT1A*- and *2D*-expressing wild-type cells also showed a stronger intensity than the empty vector-expressing wild-type cells. Nile red has been broadly used for detecting TAG levels in numerous organisms, and also used as a proper indicator of DGAT activity in H1246 strain³⁵. TLC analysis of extracted lipids from these *GmDGAT*-expressing cells revealed a strong accumulation of TAG that was almost absent from H1246 cells only expressing empty vector (Fig. 3c). The strong FA bands shown on preparative TLC plates may come from hydrolysis of lipids during extraction process, rather than endogenous fatty acids³⁴. Further analysis of the contents and compositions of TAGs separated by TLC confirmed that the total TAGs increased in *GmDGAT1A*- and *2D*-expressing yeast cells (Fig. 3d). Both *GmDGAT1A* and *2D* showed DGAT activity sufficient to restore TAG synthesis in H1246 strain. The endogenous FA profile of H1246 consists mainly of palmitic acid (16:0), palmitoleic acid (16:1), stearic acid (18:0), and oleic acid (18:1). *GmDGAT1A* and *2D* expression triggered an increase in TAGs with 16:1 and 18:1 fatty acyl chains, but decreases in 16:0 and 18:0 fatty acyl chains (Fig. 3e), suggesting that both *GmDGAT1A* and *2D* preferred to utilize 16:1 and 18:1 acyl-CoAs to synthesize TAGs in yeast. Overexpression of *GmDGAT2D* in wild-type yeast strain slightly increased 16:1/16:0 ratios.

Ectopic expression of *GmDGATs* in soybean hairy roots promotes TAG accumulation. To further understand the roles of *GmDGAT1A* and *2D* in oil production, we expressed *GmDGAT1A* and *2D*, driven by a CaMV 35S promoter, in hairy roots derived from soybean cotyledons, which has been used as an efficient approach to verify functions of soybean genes³⁶, giving that the lower transformation rate and time-consuming of soybean plant transformation. Endogenous expression of *GmDGAT1A* and *2D* in hairy roots is low, and TAG contents in these hairy roots were also much lower than in soybean seeds. The ectopic expression of *GmDGAT1A* and *2D* was confirmed with semi-quantitative RT-PCR and qRT-PCR; and TAG contents were confirmed with TLC analysis in combination with gas chromatography (GC) measurement (Fig. 4). We analyzed more than 10 independent hairy root lines for the TAG accumulation as a function of *GmDGAT1A* and *2D* over-expression. The ectopic expression of *GmDGAT1A* and *2D* increased TAG contents in these transgenic hairy roots by about 2 to 4 folds as compared with GFP vector control, indicating that overexpression of *GmDGAT1A* and *2D* triggered a *de novo* TAG biosynthesis (Fig. 4). Analysis of FA compositions in these TAGs suggested that *GmDGAT2D* expressed in soybean hairy roots preferred to synthesize TAGs with linoleic acid (18:2) acyl chains; meanwhile slight decreases in 18:0- and 18:1-TAGs were observed. The 16:0 content decreased averagely by 11%, 18:0 decreased averagely by 21%, 18:1 decreased by 28%, but 18:2 averagely increased by 55% (Fig. 4d). *GmDGAT1A*-hairy roots preferred to synthesize TAGs with linolenic acid (18:3) acyl chains and to less extent, 18:2 acyl chains, at the expense of 18:1 (Fig. 4f,g). The portion of 18:2 in *GmDGAT1A*-transgenic hairy roots increased by 13–40%, whereas 18:3 increased by 46–120%, and meanwhile, 18:1 decreased averagely by about 47% (Fig. 4g).

***GmDGAT* expression in *Arabidopsis thaliana* alters seed oil production and TAG composition.**

In order to verify the function of *GmDGAT1A* and *2D* in seed oil production, *GmDGAT1A* and *2D* was also expressed in *Arabidopsis Col-0*. Analysis of seeds from T3 plants of more than 10 independent lines expressing *GmDGAT1A* and *2D* as confirmed by qRT-PCR (Supplementary Fig. S3). From more than 10 independent lines, the markedly increased amounts of 18:2 were detected in *GmDGAT2D*-expressing *Arabidopsis* seeds relative to the seeds from control plants. The 18:1 content also increased in all lines, but 18:3 drastically reduced in transgenic lines, averagely by 18%. The highest total contents of 18:1 and 18:2 detected in seeds of transgenic *Arabidopsis* lines were approximately 20% more than these in seeds of control plants (Fig. 5).

For *GmDGAT1A*, the markedly increased amounts of 18:3 (averagely by 15%) were detected in *GmDGAT1A*-expressing *Arabidopsis* wild-type *Col-0* seeds relative to the seeds from control plants (Fig. 5b). 18:2 content also increased by 4%, whereas 20:1 decreased markedly by 17%. The total TAG contents in these *GmDGAT1A*- or *GmDGAT2D*-transgenic lines are also enhanced to different degrees as compared with the wild-type *Arabidopsis* seeds (Fig. 5a,c).

To gain more insights into the properties of *GmDGAT2D*, *GmDGAT2D* was overexpressed in *Arabidopsis* mutant *rod1* (*reduced oleate desaturation 1*) plants. *ROD1* encodes a phosphatidylcholine:diacylglycerol cholinephosphotransferase, which catalyzes a major reaction transferring 18:1 into PC for desaturation and also for reversely transferring 18:2 and 18:3 into DAG for the TAG synthesis pathway³⁷. Therefore, the seeds of *rod1* mutant have an averagely 60% reduction in 18:2 plus 18:3 but a similar portion of increase in 18:1TAG³⁷ (Fig. 6c). When *GmDGAT2D* was over-expressed in *rod1* mutant, although total oil production averagely increased by only 5.7% in these transgenic seeds of T3 generation, the 18:1 increased by about 14% averagely than these in *rod1* seeds, or increased by about 2.4-fold compared with the wild-type control. Meanwhile, the contents of 16:0 were reduced by 14%, compared with these in *rod1* mutant seeds (Fig. 6b,c).




Figure 3. Functional expression of *GmDGATs* in yeast cells. The quadruple mutant *Saccharomyces cerevisiae* strain H1246 and the wild-type strain YPH499a were transformed with *GmDGAT* genes and vector control. **(a)** Nile red staining of oil drops in *GmDGAT1A* (i, vi)- and *2D* (iii, viii)- or *pYESDEST52* vector (ii, vii)-expressing yeast mutant strain H1246, and vector (iv, ix) and *GmDGAT2D* (v, x) expressed in wild-type strain YPH499a cells. TAG was extracted and separated by thin-layer chromatography and TAG spots were scraped for analyzing compositions of fatty acids by GC. **(b)** Nile Red Assay on oil drops and fluorescence-based TAG content. **(c)** Thin-layer chromatography of lipids extracted from yeast transformants. DAG, fatty acids (FAs), and TAG spots were labeled out. TAG spots were scraped off from the thin-layer chromatography (TLC) plates and were extracted for analysis of TAG contents and compositions with GC. **(d)** Total TAG production in yeast mutant strain H1246 and wild-type strain YPH499a, Total TAG was measured by GC. **(e)** Fatty acid compositions of TAG produced in *GmDGAT2D*- or *pYESDEST52* empty vector-expressing yeast mutant strain H1246 and wild-type strain YPH499a. All data are presented by mean of at least three biological replicates \pm SD. * $P < 0.05$ and ** $P < 0.01$ by Student's *t* test for significant difference.




Figure 4. Ectopic expression of *GmDGATs* in soybean hairy roots. The cotyledons of germinating soybean seeds were used for infection by *Agrobacteria* K599 harboring *pB2GW7-GmDGAT2D* or *-GmDGAT1A*. Generated transgenic hairy roots selected on ppt-medium were confirmed by qRT-PCR for target gene expression, and analyzed for TAG production, as compared with GFP-expressing hairy roots as control. Total neutral lipids were extracted and separated by TLC. TAG spots were scraped off from TLC plates and extracted for determination of total contents and compositions by using GC. (a) qRT-PCR confirmation of *GmDGAT2D* expression in soybean transgenic hairy roots. (b) TLC analysis of neutral lipids extracted from hairy root transformants. DAG, fatty acids (FAs), and TAG spots were labeled out. Bottom panel shows the photos of soybean transgenic hairy roots. Hairy roots expressing free GFP vector were used as a control. (c) Quantification of total TAGs from hairy roots ectopically expressing *GmDGAT2D* compared with GFP control. (d) Fatty acid composition of TAGs from hairy roots expressing *GmDGAT2D* and GFP control. (e) qRT-PCR confirmation of *GmDGAT1A* expression in soybean transgenic hairy roots. (f) Total TAGs in *GmDGAT1A*-expressing hairy roots, GFP-hairy roots are used as a control. (g) Fatty acid composition of TAGs from hairy roots expressing *GmDGAT1A* and GFP control. All data are expressed as mean \pm SD ($n > 4$); three lines are representative of more than 10 transgenic hairy root lines were analyzed. * $P < 0.05$ and ** $P < 0.01$ by Student's *t* test for significant difference.




Figure 5. Functional expression of *GmDGAT-1A* and *-2D* in *Arabidopsis thaliana* seeds. (a) Total TAGs in *GmDGAT1A*-expressing *Arabidopsis* seeds. (b) Fatty acid composition of TAGs from *Arabidopsis* seeds expressing *GmDGAT1A*. (c) Total TAGs in *GmDGAT2D*-expressing *Arabidopsis* seeds. (d) Fatty acid composition of TAGs from *Arabidopsis* seeds expressing *GmDGAT2D*. All data are expressed as mean \pm SD from at least three biological duplicates. * $P < 0.05$ and ** $P < 0.01$ by Student's *t* test for significant difference. T2 transgenic *Arabidopsis* seeds were used for analysis.

Expression of *GmDGAT1A* and *2D* in response to environmental and hormonal stresses.

GmDGAT2D was up-regulated by abiotic stresses, such as heat, and cold stress and biotic stresses (Fig. 7, Supplementary Figs S4 and 5). Upon the cold treatment (4 °C), *GmDGAT2D* transcript levels in developing seeds initially increased at 12 h after cold treatment, and then dropped markedly, whereas it was not the case for *GmDGAT1A*, which was down-regulated upon cold treatment (Fig. 7a,b). Also, under heat stress at 42 °C, *GmDGAT1A* and *2D* expression in seed also showed different patterns: *GmDGAT2D* transcripts increased firstly and then fell back to the basal level, whereas *GmDGAT1A* transcript level increased and then dropped to a level




Figure 7. Expression of *GmDGATs* in response to hormonal and environmental stresses. Soybean seedlings with 9 trifoliolate (4 weeks old) were subject to insect biting, spraying with hormones (100 μ M ABA), or their detached leaves were floating on 50 μ M MeJA solution and water (control). Eight-week old soybean plant bearing pods were moved into a cold (4 $^{\circ}$ C) or heat (42 $^{\circ}$ C) stress in incubator for indicated time. Leaf samples or pods were harvested at the indicated time. qRT-PCR analysis of *GmDGAT1A* (a,c,e,g,i,k) and *GmDGAT2D* (b,d,f,h,j,l) expression under abiotic and biotic stresses. Soybean developing seeds treated under 4 $^{\circ}$ C (a,b) or 42 $^{\circ}$ C (c,d) stresses. Total TAG contents and fatty acid composition of soybean seeds exposed to 4 $^{\circ}$ C (e) or 42 $^{\circ}$ C (f) for the indicated time. Soybean seedlings were subject to insect biting (g,h), JA treatment (i,j), ABA treatment (k,l). Data are expressed as mean \pm SD (n > 3). *P < 0.05 and **P < 0.01 by Student's *t* test for significant difference.

distinguishable preferences to utilize 18:1 and 18:2 acyl CoA (GmDGAT2D) or 18:3 acyl CoA (GmDGAT1A) as acyl donors. Since most reported plant DGAT2s are involved in biosynthesis of unusual TAG with hydroxy FAs, this soybean GmDGAT2D is thus different from these plant DGAT2s, but more close to animal, fungal or algae DGAT2s, which have potential for enhancing oil production and improve oil compositions in crops^{7,28,38}.

GmDGAT1A and 2D are ER-localized functionally active short polypeptides. Both GmDGAT1A and 2D can complement TAG-deficiency mutant H1246 phenotype, suggesting they are functional DGATs. GmDGAT2D had much stronger activity in yeast TAG biosynthesis than GmDGAT1A did. Both GmDGAT1A and GmDGAT2D do not possess the known ER retrieval motifs –K(X) KXX–COOH, nor the motif YKSKW of the C-terminus²⁰. GmDGAT2D is a shorter polypeptide of 329 amino acids, as compared with GmDGAT1A of 500 amino acids. GmDGAT2D only has two TM domains whereas GmDGAT1A has eight TMs. We showed that GmDGAT1A- and GmDGAT2D-GFP similarly locate to the ER when expressed in tobacco leaf epidermal cells (Fig. 2). Unlike fungal DGAT2s, such as NcDGAT2 and TaDGAT2, that usually have longer and unique N-terminal extension (around 200 aa), GmDGAT2D does not have it^{20,38}. The histidines in active sites and N-terminus being essential for formation of DGAT dimers and tetramers, and a phenylalanine essential for a maize DGAT activity are present in GmDGAT1A and 2D^{20,39}. From results in yeast, soybean, and Arabidopsis expression of *GmDGAT2D*, we conclude that GmDGAT2D is different from other known plant DGAT2 for unusual oil synthesis: GmDGAT2D is more similar to DGATs from fungi, animal, and some plants such as Arabidopsis and flax^{10,11,24,26}, for regular TAG biosynthesis^{28,29}. Also, GmDGAT2D appears to have subdomain localizations in the ER similar to these of GmDGAT1A, which may suggest similar functions of Type 1 and 2 DGATs in TAG biosynthesis. This is different from a previous report on Tung tree DGAT1 and DGAT2¹⁰.

GmDGAT1A and 2D have different substrate preferences for regular TAG biosynthesis.

GmDGAT1A and 2D restored TAG biosynthesis in *S. cerevisiae* strain H1246, they are functional DGAT enzymes. We also found that expression of *GmDGAT2D* in both soybean hairy roots and Arabidopsis wild-type plants increased TAG contents and changed TAG compositions. An increased 18:2 and 18:3 acyl chains in TAGs was observed in *GmDGAT2D* and *GmDGAT1A* transgenic hairy roots, respectively. Overexpression of *GmDGAT1A* and 2D in Arabidopsis seeds results in substantial changes in TAG contents. A significant increase in 18:2 and slight increase in 18:1 acyl chains in TAGs, accompanied by a pronounced decrease in 18:3, were observed with *GmDGAT2D* transgenic lines. In contrast, 18:3 was significantly increased, accompanied with decrease in 20:1 or 18:1 in *GmDGAT1A*-transgenic Arabidopsis seeds or soybean hairy roots, respectively. These data suggest that GmDGAT1A has a strong preference to 18:3 acyl-CoA and GmDGAT2D prefers to use 18:2 and then 18:1 acyl-CoAs. However, when *GmDGAT2D* was over-expressed in *rod1* mutant, whose seeds accumulate much more 18:1 than 18:2, *GmDGAT2D/rod1* seeds produced even more 18:1 than *rod1* mutant seeds, but 18:2 contents were unchanged. The data suggest that substrate preference for DGATs, at least for GmDGAT2D, is dependent of availability of preferred substrates. GmDGAT2D is more similar to DGAT2s from fungi and animals in regular TAG synthesis in plants^{20,38,40}, but unlike other DGAT2s in plants that are involved in synthesis of unusual TAGs containing hydroxy FAs^{7,24,26}.

The substrate preferences of DGATs can significantly edit the acyl chains in TAGs, which may mark plant species with specific features in storage TAGs. The analysis of the neutral lipids from the *GmDGAT1A* and 2D-transformed yeast strain revealed that GmDGAT1A and 2D-catalized TAG production with more 16:1 and 18:1 unsaturated FA chains, as compared with native TAGs with 16:0 and 18:0 saturated and 18:1 unsaturated FAs in yeast. When expressed in Arabidopsis seeds, *GmDGAT2D* render the transgenic Arabidopsis to accumulate significantly more 18:2 and slightly increased 18:1 TAGs in seeds, in expense of a reduced 18:3 TAGs. However, *GmDGAT1A* renders transgenic Arabidopsis to accumulate significantly more 18:3 in seeds, but a reduced 20:1 TAGs. Thus two DGATs showed very different substrate preference and therefore transgenic consequences. Results from transgenic soybean hairy roots also provided unambiguous evidence that GmDGAT2D preferentially utilizes 18:2 acyl-CoA as acyl donors for TAG biosynthesis, whileas GmDGAT1A prefers to utilize 18:3 acyl-CoA as acyl donors for TAG biosynthesis. This substrate preference may not be unexpected, given that no 18:2 available in yeast cells, but 18:2 accounts for around 25–30%, and 18:3 (~20%), of the total FAs in soybean and Arabidopsis seeds, whereas 18:1 are relatively minor unsaturated FAs (~17%)^{2,20}. However, in soybean hairy roots, 16:0 and 18:2 FAs account for about 30% and 25%, respectively, of total FAs, whereas 18:0, 18:1, and 18:3 each accounts for only around 15–18%. Therefore, GmDGAT2D selects 18:2 for TAG synthesis in soybean hairy roots, but selects 18:1 in *rod1* mutant seeds (accounting for more than 40% of total FAs). GmDGAT1A prefers to use 18:3 for TAG biosynthesis, although 18:3 FAs are small portion. Therefore, both GmDGATs prefer to use unsaturated FAs as substrates in yeast and plant cells. The different substrate preferences of GmDGAT2D and GmDGAT1A may confer the diverse FA profiles in soybean oils. Meanwhile, their differential expression patterns also indicate that they may have other additional functions in non-seed tissues upon stresses.

Role of GmDGAT2D in plant response to stresses and GmDGAT1A in seed TAG biosynthesis.

It is well-known that oil production in oilseed is regulated by ABA signaling, which regulates seed maturation at least through abscisic acid insensitive 4 (ABI4) and downstream transcriptional machineries^{5,41}. *GmDGAT2D* transcripts are also found in non-seed tissues, such as flower, root and leaf, where it may play a role in glycerolipid biosynthesis and physiological response to adverse environments, although functions of TAG and DGATs in these tissues remain ambiguous^{42,43}. In flower, coat materials in the pollen grain surface include TAGs, other lipids, and flavonoids synthesized by the tapetum cells for keeping pollen viability under stress conditions^{6,19}. Anther filament elongation plays an essential role in plant pollination. Failure in anther filament growth, one of the reasons for plant male sterility, may be attributable to JA signaling pathway⁴⁴. JA is required for anther filament development, and blocking JA synthesis or signaling (e.g. *coi1* mutant) confers male sterile as a result of

improper anther filament elongation^{44,45}. COI1 is specifically expressed in the anther filament where *GmDGAT2D* is also highly expressed, indicating a possible connection between JA signaling and *GmDGAT2D* and TAG biosynthesis. *GmDGAT2D* transcript levels coincidentally decreased followed by insect biting, wounding, and MeJA treatments. The insect biting caused-wounding effect usually induces an endogenous JA increase in plant tissues, as one of mechanisms to initiate plant responses^{6,44–46}. This may explain that both insect biting and MeJA treatment similarly repressed *GmDGAT2D* expression.

Since plants normally accumulate more polyunsaturated FAs for glycerolipid biosynthesis in response to cold stress⁴⁷, and *GmDGAT1A* and *2D* prefer to utilize 18:3- and 18:2-CoA for TAG biosynthesis, it is plausible that *GmDGAT2D* was up regulated in cold and heat stresses and contributed to adaptation of plants to these stresses by modifying the polyunsaturated TAG synthesis. *GmDGAT2D* promoter contains several ABA and JA responsive *cis*-elements, such as DPBFCOREDCDC3, MYCCONSENSUSAT, MYCATRD22, and PYRIMIDINEBOX that are responsible to cold stress and ABA signaling (Supplementary Table S2). However, the promoter of *GmDGAT1A* has fewer ABA and JA responsive *cis*-elements, consistently; *GmDGAT1A* transcripts were less changed upon JA treatment.

TAG usually represents <1% of leaf glycerolipids but can accumulate more under stresses, such as heat and cold treatments, or other conditions such as expression of structural or regulatory genes^{6,43,47}. Recently, it was found that DAG and TAG are also identified from phloem exudates under normal and stressed conditions, suggesting that they might act as long distance signals⁴⁶. However, what is TAG biological functions under these conditions and which DGAT is responsible for spatial and temporal TAG biosynthesis under normal or stressed conditions is not understood^{43,47}. Given stress-responsive *GmDGAT2D* or *GmDGAT1A* with such dramatically changed expression patterns under various conditions in this study, and recent reports showing lipid metabolism is involved in multiple biotic and abiotic stress responses in plants⁴⁷, DGATs and TAG metabolism may have intriguing connections with plant-environment interactions. However, whether *GmDGAT2D* expression in anther filament is also involved in production of these vascular long-distance traveling TAGs remain to be determined.

The similarly steady increases of *GmDGAT1A* and *2D* expression levels at the stages of active oil accumulation, followed by the decreases when TAG reaches the plateau, suggest that *GmDGAT1A* and *2D* play essential roles in soybean seed oil production. This is in contrast to oleogenic seed crops that contain unusual FAs, where DGAT2 plays a more central role than DGAT1 in unusual oil production^{10,24,26}. Since soybean oil does not contain unusual FAs, as the major contributors of TAG production, *GmDGAT1A* and *2D*, together with other DGATs, function redundantly in TAG biosynthesis in both seeds, flowers, leaves, and other tissues under specific conditions⁴³.

Plant employs different strategies to respond to environmental stresses, such as cold and heat, during the seed development and seed filling, when metabolism of polyunsaturated FAs is believed to play an essential role in adaptation to these stresses^{4,47}. The differential responses of various DGATs to heat and cold stresses suggest that they may function in fine tune to change lipid metabolic pathways correspondingly to ensure the proper TAG biosynthesis for adapting them to temperature changes^{4,6,47}. Given the increasing concerns on effects of unusual global climate changes, such as frequently high temperature and cold stresses, on soybean production, further investigation on regulation of DGATs and TAG metabolism may provide insights into how plant glycerolipid production is affected by environmental stresses^{6,47}.

Potential application of two *GmDGATs* for metabolic engineering of soybean oil composition. The considerable attention has been paid to DGATs as biotechnological targets for improvement of the TAG production in oilseed crops such as soybean, maize, and rapeseed to meet the growing demand for vegetable oils or biofuels^{13,27,38}. Seed-specific overexpression of Arabidopsis DGAT1 increased oil content by up to 15 and 46% in seeds of transgenic *Brassica napus* and Arabidopsis, respectively^{13,48}. The increased oil contents in maize and soybean seeds were also achieved by manipulating the expression of the fungus *Umbelopsis ramanniana* UrDGAT2 or algae DGAT2^{7,27,38}. Most DGAT2s from plants such as *R. communis* and *Vernonia galamensis* are capable of using hydroxyl or epoxy fatty acyl-CoAs (e.g. ricinoleic acid, vernolic acids) into TAGs to produce unusual TAGs for industrial purposes^{24,26}, while we show that *GmDGAT2D* is different from them.

Because higher levels of 18:3 in soybean seeds is a major cause for oxidation instability of vegetable oils, and *trans*-fats generated primarily in hydrogenation of vegetable oils during food processing have negative impact on human health, enriching 18:1 and 18:2 and meanwhile minimizing 18:3 in vegetable oils has become one of targets for crop biotechnology². Our study here revealed two soybean DGATs, *GmDGAT2D* are responsible for 18:1 and 18:2 production and *GmDGAT1A* for 18:3 in soybean seeds. The demonstration provides possible strategies for either increasing 18:1-TAG and repressing 18:3-TAG production in a more specific way than the reported technology for high-oleic acid soybean variety^{37,38,40}.

The study sheds light on our understanding of TAG biosynthesis and metabolic engineering of soybean oil with appropriate DGATs. Using a stronger seed-specific promoter may further improve the performance of *GmDGATs* for creating soybean oils with desired composition. Moreover, our study suggests roles of TAG biosynthesis and DGATs in other non-seed tissues during plant responses to hormones and various stresses, which may open a door for exploring and understanding more fundamental lipid biology in crops.

Methods

Plant growth conditions and treatments. Soybean (*Glycine max* L.) seeds were germinated in three-gallon pots containing soils. The seedlings were grown in growth chamber (26/20 °C day/night temperature, photoperiod of 14/10 h, 800 μmol m⁻² s⁻¹ light intensity and 60% humidity). Roots, leaves, stems, flowers, seeds, and drupes at different developmental stages were harvested from “Tianlong No.1”, a soybean (*Glycine max* L.) cultivar, grown in a growth chamber under the same conditions as above or a natural environment at the fields of Huazhong Agricultural University at Wuhan, China. For methyl jasmonate (MeJA) treatment, soybean leaves

was detached and floated on the water containing or not containing 100 μ M MeJA. For insect biting experiments, the chewing insect measuring worms (*Geometridae*) were put on four-week old soybean plants with 3- trifoliolate leaves for insect biting. For cold and heat treatments, eight-week old soybean plants bearing pods were moved into 4 °C refrigerator with regulator lights or into oven set at 42 °C for treatments. Leaves or seeds (at stage 4) were collected at different time and immediately frozen in liquid nitrogen and stored at -80 °C after treatments for RNA extraction and TAG analysis.

Vector construction. The open reading frames (ORFs) of *GmDGAT1A* (Glyma13G106100) and *GmDGAT2D* (Glyma.01G156000) were amplified with the cDNA made from soybean developing seeds using pairs of primers for *GmDGATs* (Supplementary Table S1). Total RNA was extracted from *G. max* developing seeds, and 10 μ g of total RNA was used to synthesize first-strand cDNA using the first-strand synthesis system (Invitrogen). The cDNA was synthesized as a template for *GmDGAT2D* cDNA amplification. After *GmDGAT1A* and *GmDGAT2D* ORFs was cloned into T-easy vector and sequenced, the cDNAs in pDONOR221, pDONOR221-*GmDGAT1A* or -*GmDGAT2D*, was recombined into different destination vectors such as pYEST-DEST52 and pB2GW7 by using LR recombinase (Invitrogen), and were sequenced. For *GmDGAT2D* promoter analysis, a 1.5 K bp promoter region was amplified with primers proF and R (Supplementary Table S1). The PCR products were cloned into T-easy vector for sequencing; then subcloned into pDONOR221 and subsequently pHGWFS7 by using BP and LR recombinases (Invitrogen).

Soybean hairy root transformation and analysis of TAG in hairy roots. pB2GW7-*GmDGAT1A* or -*GmDGAT2D* was transformed by electroporation into *Agrobacterium rhizogenes* strain K599, which was used to transform soybean cotyledons. Seeds of soybean cultivar “Tianlong” was surface sterilized and germinated in sterilized filter papers in petridishes. The green cotyledons from about 7 days-old germinating soybean seeds were wounded on the surfaces, followed by the infections with *Agrobacterium rhizogenes* K599 bacteria harboring pB2GW7-*GmDGAT1A*, -*GmDGAT2D*, or -*GFP* as a control. Generated hairy roots were selected on MS medium containing antibiotics for controlling *Agrobacterium* and 7 mg/l phosphinothricin (ppt) for transformed hairy roots. The transformed hairy roots expressing pB2GW7-*GmDGAT1A* or -*GmDGAT2D* were confirmed with PCR. Then roots were used for extraction of total lipids, from which TAG was separated on TLC. TAG spots were scraped off the plate and extracted for measurement by using GC.

Expression of GmDGATs in Arabidopsis. The binary vector containing the cassette for 35S::*GmDGAT1A* or 35S::*GmDGAT2D* was transformed into *Agrobacterium tumefaciens* GV3101 by electroporation. Transgenic Arabidopsis Col-0 plants were transformed by using floral dip method. BASTA was used for selection of transformant seedlings which were also PCR confirmed using 18s univ F and 18s univ R, as well as together with gene specific 3' primers (Supplementary Table S1). Expression of the transgene in developing seeds was confirmed by RT-PCR. The dry seeds of T3 Arabidopsis transformants were analyzed for oil contents and FA composition.

Yeast strains and transformation. For the functional expression of *GmDGAT1A* or 2D in yeast, the quadruple mutant *Saccharomyces cerevisiae* strain H1246 (W303; MAT α are1- Δ ::HIS3 are2- Δ ::LEU2 dga1- Δ ::KanMX4 lro1- Δ ::TRP1 ADE2 ura3)³⁴ and the wild-type *S. cerevisiae* strain YPH499a (ura3-52, lys2-801, ade2-101, trp1-63, his3-200, leu2-1) were used as heterologous hosts. The plasmid DNA pYESDEST52-*GmDGAT1A* or 2D was transformed into stationary H1246 cells using the PEG/lithium acetate method⁴⁹. Yeast cells harboring the empty pYESDEST52 vector were used as negative control. Transformants were selected on YNB lacking uracil and functional expression and induction of *GmDGAT1A* or *GmDGAT2D* was performed as described previously⁵⁰.

Yeast expression and Nile red assay. Intracellular lipid bodies were visualized by fluorescent microscopy using Nile Red (9-diethylamino-5H-benzo α -phenoxazine-5-one, 5 μ l Nile Red in 0.8 mg/mL acetone stock) staining method^{51,52}. Fluorescence microscopic images of the stained cells were recorded using excitation wavelength at 485 nm and emission wavelength at 538 nm under Nikon Eclipse 80I microscope.

Nile red assay and optical density determination were performed in 96-well polystyrene flat-bottom microplates (UNIPLATE, Whatman, UK), according to previously described⁵². The Nile red fluorescence assay was performed at excitation and emission filters of 485 and 538 nm, respectively. The assays were performed with ten technical replicates. The measurement of the fluorescence intensity of various yeast cultures were done after adjusting these yeast resuspension cells to OD_{600 nm} to 1 to make sure that all comparisons are in the same cell density.

Quantification of TAG and analysis of FA composition. Total lipid extraction and TAG content and composition determination were done according to previously described methods with slight modifications^{35,52-54}. Briefly, total lipids in yeast pellets (~0.014 g dry yeast) or soybean hairy roots (~0.2 g fresh tissues) were extracted with 4 ml of 4M HCl in glass tubes tightly with Teflon-lined caps at room temperature for 30 min, then in a 100 °C water bath for 10 minute. After cooling tubes were centrifuged, yeast pellets or hairy root powder were extracted with 4 ml of hexane: isopropanol (3:2, v/v). The upper hexane layer of the extractions was removed into a new glass tube and evaporated under a slow stream of N₂. The residues were dissolved in 50 μ l hexane for TLC analysis.

The TAG from non-seed tissues including yeast and soybean hairy roots was resolved by TLC on a silica plate (SIL GF254, 0.25 mm). The plate was developed with hexane/diethyl ether/acetic acid (80:20:1, v/v/v), essentially according to the method as previously described³⁵. Fatty acid methyl esters (FAMES) were prepared by heating the dry TAG materials at 85 °C for 30 min in 1 M HCl in dry methanol, as described by Wang and Benning⁵². FAMES then were dried under nitrogen gas and resuspended in 200 μ l of hexane for GC analysis.

The TAG content and composition from soybean and Arabidopsis seeds were measured according to methods described by Christie⁵⁴. Briefly, ~10 mg of seeds were weighed in a 13 × 100 mm glass screw-cap test tube. To each tube, 1.5 ml of 2.5% sulphuric acid in methanol, 400 µl toluene, and 100 µl of 1 mg/ml triheptadecanoin in toluene (Nu-Chek Prep, Elysian, USA) as an internal standard were added. The tubes were heated at 90 °C for 1 h. FAMES generated by above transesterification reaction were extracted by addition of 1.8 ml H₂O and 1 ml heptanes, and the heptane layer was recovered and analyzed with GC⁵³.

FA content and composition on TAGs from seeds or purified with TLC from total lipids of yeast cells and hairy roots were analyzed with an Agilent 7890A GC system with flame ionization detector (FID), according to the conditions described previously⁵³. Oil content was calculated by FID response of sample components relative to 17:0 methyl ester from the internal standard triheptadecanoin.

Soybean Illumina expression data analysis. Solexa sequencing libraries for developing seed samples covering two genotypes of soybean varieties at two developmental stages (at around 20 and 48 days after pollination) were generated and analyzed. Data were utilized to quantify the transcriptomic expression of soybean genes (i.e. the number of sequence reads/million reads aligned). Read counts used in expression analyses were based on the subset of uniquely aligned reads that also overlapped the genomic spans of the soybean gene predictions. Read counts for a given sample were normalized by using values for a gene's uniquely aligned read counts per million reads uniquely aligning within that sample. A total of 51,529 annotated soybean genes (74.5% of the 69,145 putative, annotated soybean genes) were found to be expressed in at least one condition.

Quantitative RT-PCR (qRT-PCR) analysis of gene expression. Total RNAs from tissues of soybean plants were isolated using TRIzol reagent (Invitrogen, Carlsbad, CA) or RNA isolation kit (Biotech, Beijing) according to the manufacturer's instructions. For each sample, 10 µg of total RNA were digested with RNase-free DNaseI (Promega, Madison, WI) to remove any genomic DNA contamination. After DNaseI treatment, RNA concentration was determined again using a NanoDrop ND-2000 UV spectrophotometer (Thermo Scientific, USA). First-strand cDNA was synthesized from 2 µg total RNA using the Superscript III first strand synthesis system (Invitrogen). All cDNA samples were diluted 50-fold in sterile water for qRT-PCR reaction. Gene specific primers were listed in Supplementary Table. S1. Soybean *ACTIN* was used as internal control. qRT-PCR reactions were performed in 96-well plates (iQ5 Real Time PCR System; Bio-Rad) for all tissues tested, and data were analyzed according to methods described previously⁴⁹.

Histochemical analysis. *pGmDGAT2D::GUS* reporter Arabidopsis lines were generated as described previously⁴⁹. T1 seeds were screened with BASTA for selection of transformants. T2 plants were used for GUS activity. GUS staining of transgenic seedlings was conducted as previously described⁴⁹. Samples were photographed using an Olympus sxx16 microscope.

Subcellular localization of GmDGAT1A- and 2D-GFP in planta. Construction of GmDGAT1A- and 2D-GFP fusion was done by using gateway recombination into pK7WGF2 in fusion with N-terminus of GFP. Determination of the subcellular localization of GmDGAT-GFP was performed by using tobacco leaf infiltration following the method described previously⁵⁰. GmDGAT-GFP and ER marker CD3-959: Cherry DNAs were transformed into *Agrobacterium* strain K599, which were used for plant transformation. The acetosyringone-activated bacteria were co-infiltrated into the abaxial epidermal surface of a tobacco leaf with a syringe, and the plant was grown at room temperature for 2 to 3 d before imaging. Imaging of GmDGAT-GFP fusion proteins was performed using a Leica TCS SP2 inverted confocal microscope using a 63 × water-immersion objective and Leica Confocal software with an excitation wavelength of 488 nm and emissions collected at 500 nm. CD3-959-mCherry-labeled ER membrane was excited at 543 nm with the argon laser, and emission was detected from 620 to 680 nm. Chloroplasts were visualized at 560 to 610 nm under excitation at 543 nm.

Bioinformatics analysis. Protein sequences, GmDGAT2D (GenBank accession number: KP752053) and GmDGAT1A (GenBank accession number: KT878751), were obtained from NCBI and phytozome. Transmembrane regions were predicted by the TMHMM Server ver. 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>). Amino acid multiple alignments were made with the ClustalW program (<http://www.ebi.ac.uk/clustalw/>) under default parameters. A phylogenetic tree was constructed using MEGA4. The significance level of the neighbor-joining analysis was examined by bootstrap testing with 1000 repeats.

Statistic analysis. Most experimental data were obtained from at least three independent experiments and were analyzed using Student's *t*-test. The significant differences between two tails of data represent 95% confidence limits. Histochemical staining and fluorescence imaging experiments, at least two repeat experiments were done, and representatives of photos or images were shown.

References

1. Wilson, R. F. The role of genomics and biotechnology in achieving global food security for high-oleic vegetable oil. *Journal of oleo science* **61**, 357–367 (2012).
2. Singer, S. D., Zou, J. & Weselake, R. J. Abiotic factors influence plant storage lipid accumulation and composition. *Plant Sci* **243**, 1–9 (2016).
3. Clemente, T. E. & Cahoon, E. B. Soybean oil: genetic approaches for modification of functionality and total content. *Plant Physiol* **151**, 1030–1040 (2009).
4. Moellering, E. R., Muthan, B. & Benning, C. Freezing tolerance in plants requires lipid remodeling at the outer chloroplast membrane. *Science* **330**, 226–228 (2000).
5. Yang, Y., Yu, X., Song, L. & An, C. ABI4 activates DGAT1 expression in Arabidopsis seedlings during nitrogen deficiency. *Plant Physiol* **156**, 873–883 (2011).

6. Zhao, J. Phospholipase D and phosphatidic acid in plant defense response: from protein-protein and lipid-protein interactions to hormone signaling. *J Exp Bot* **66**, 1721–1736 (2015).
7. Sanjaya *et al.* Altered lipid composition and enhanced nutritional value of Arabidopsis leaves following introduction of an algal diacylglycerol acyltransferase 2. *Plant Cell* **25**, 677–693 (2013).
8. Cases, S. *et al.* Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis. *Proc Natl Acad Sci USA* **95**, 13018–13023 (1998).
9. Liu, Q., Siloto, R. M., Lehner, R., Stone, S. J. & Weselake, R. J. Acyl-CoA:diacylglycerol acyltransferase: molecular biology, biochemistry and biotechnology. *Prog Lipid Res* **51**, 350–377 (2012).
10. Shockey, J. M. *et al.* Tung tree DGAT1 and DGAT2 have nonredundant functions in triacylglycerol biosynthesis and are localized to different subdomains of the endoplasmic reticulum. *Plant Cell* **18**, 2294–313 (2006).
11. Banilas, G., Karampelias, M., Makariti, I., Kourti, A. & Hatzopoulos, P. The olive DGAT2 gene is developmentally regulated and shares overlapping but distinct expression patterns with DGAT1. *J Exp Bot* **62**, 521–532 (2011).
12. Zou, J. *et al.* The Arabidopsis thaliana TAG1 mutant has a mutation in a diacylglycerol acyltransferase gene. *Plant J* **19**, 645–653 (1999).
13. Jako, C. *et al.* Seed-specific over-expression of an Arabidopsis cDNA encoding a diacylglycerol acyltransferase enhances seed oil content and seed weight. *Plant Physiol* **126**, 861–874 (2001).
14. Kalscheuer, R. & Steinbuechel, A. A novel bifunctional wax ester synthase/acyl-CoA:diacylglycerol acyltransferase mediates wax ester and triacylglycerol biosynthesis in Acinetobacter calcoaceticus ADP1. *J Biol Chem* **278**, 8075–8082 (2003).
15. King, A., Nam, J. W., Han, J. X., Hilliard, J. & Jaworski, J. G. Cuticular wax biosynthesis in petunia petals: cloning and characterization of an alcohol-acyltransferase that synthesizes wax-esters. *Planta* **226**, 381–394 (2007).
16. Li, R. *et al.* Soybean oil biosynthesis: role of diacylglycerol acyltransferases. *Functl Integral Genomics* **13**, 99–113 (2013).
17. Saha, S., Enugutti, B., Rajakumari, S. & Rajasekharan, R. Cytosolic triacylglycerol biosynthetic pathway in oilseeds. Molecular cloning and expression of peanut cytosolic diacylglycerol acyltransferase. *Plant Physiol* **141**, 1533–1543 (2006).
18. Hernandez, M. L. *et al.* A cytosolic acyltransferase contributes to triacylglycerol synthesis in sucrose- rescued Arabidopsis seed oil catabolism mutants. *Plant Physiol* **160**, 215–225 (2012).
19. Zhang, M., Fan, J., Taylor, D. C. & Ohlrogge, J. B. DGAT1 and PDAT1 Acyltransferases have overlapping functions in Arabidopsis triacylglycerol biosynthesis and are essential for normal pollen and seed development. *Plant Cell* **21**, 3885–3901 (2009).
20. Zhang, C. *et al.* A thraustochytrid diacylglycerol acyltransferase 2 with broad substrate specificity strongly increases oleic acid content in engineered Arabidopsis thaliana seeds. *J Exp Bot* **64**, 3189–3200 (2013).
21. Cao, H. *et al.* Developmental regulation of diacylglycerol acyltransferase family gene expression in tung tree tissues. *PLoS One* **8**, e76946 (2013).
22. Pan, X., Siloto, R. M., Wickramaratna, A. D., Mietkiewska, E. & Weselake, R. J. Identification of a pair of phospholipid:diacylglycerol acyltransferases from developing flax (*Linum usitatissimum* L.) seed catalyzing the selective production of trilinolenin. *J Biol Chem* **288**, 24173–24188 (2013).
23. Zhou, X. R., Shrestha, P., Yin, F., Petrie, J. R. & Singh, S. P. AtDGAT2 is a functional acyl-CoA:diacylglycerol acyltransferase and displays different acyl-CoA substrate preferences than AtDGAT1. *FEBS Lett* **587**, 2371–2376 (2013).
24. Burgal, J. *et al.* Metabolic engineering of hydroxy fatty acid production in plants: RcDGAT2 drives dramatic increases in ricinoleate levels in seed oil. *Plant Biotechnol J* **6**, 819–831 (2008).
25. Kroon, J. T., Wei, W., Simon, W. J. & Slabas, A. R. Identification and functional expression of a type 2 acyl-CoA:diacylglycerol acyltransferase (DGAT2) in developing castor bean seeds which has high homology to the major triglyceride biosynthetic enzyme of fungi and animals. *Phytochemistry* **67**, 2541–2549 (2006).
26. Li, R. *et al.* Vernonia DGATs can complement the disrupted oil and protein metabolism in epoxygenase-expressing soybean seeds. *Metab Eng* **14**, 29–38 (2012).
27. Lardizabal, K. *et al.* Expression of *Umbelopsis ramanniana* DGAT2A in seed increases oil in soybean. *Plant Physiol* **148**, 89–96 (2008).
28. Cases, S. *et al.* Cloning of DGAT2, a second mammalian diacylglycerol acyltransferase, and related family members. *J Biol Chem* **276**, 38870–38876 (2001).
29. Yen, C.-L., Stone, S. J., Koliwad, S., Harris, C. & Farese, R. V. Thematic review series: glycerolipids. DGAT enzymes and triacylglycerol biosynthesis. *J Lipid Res* **49**, 2283–2300 (2008).
30. Liu, Q., Siloto, R. M., Snyder, C. L. & Weselake, R. J. Functional and topological analysis of yeast acyl-CoA:diacylglycerol acyltransferase 2, an endoplasmic reticulum enzyme essential for triacylglycerol biosynthesis. *J Biol Chem* **286**, 13115–13126 (2011).
31. Wang, H. W., Zhang, J. S., Gai, J. Y. & Chen, S. Y. Cloning and comparative analysis of the gene encoding diacylglycerol acyltransferase from wild type and cultivated soybean. *Theor Appl Genet* **112**, 1086–1097 (2006).
32. Eskandari, M., Cober, E. R. & Rajcan, I. Using the candidate gene approach for detecting genes underlying seed oil concentration and yield in soybean. *Theor Appl Genet* **126**, 1839–1850 (2013).
33. Roudier, F. *et al.* Very-long-chain fatty acids are involved in polar auxin transport and developmental patterning in Arabidopsis. *Plant Cell* **22**, 364–375 (2010).
34. Sandager, L. *et al.* Storage lipid synthesis is non-essential in yeast. *J Biol Chem* **277**, 6478–6482 (2002).
35. Siloto, R. M., Truksa, M., He, X., McKeon, T. & Weselake, R. J. Simple methods to detect triacylglycerol biosynthesis in a yeast-based recombinant system. *Lipids* **44**, 963–973 (2009).
36. Guan, R. *et al.* Salinity tolerance in soybean is modulated by natural variation in GmSALT3. *Plant J* **80**, 937–950 (2014).
37. Lu, C., Xin, Z., Ren, Z., Miquel, M. & Browse, J. An enzyme regulating triacylglycerol composition is encoded by the ROD1 gene of Arabidopsis. *Proc Natl Acad Sci USA* **106**, 18837–18842 (2009).
38. Oakes, J. *et al.* Expression of fungal diacylglycerol acyltransferase 2 genes to increase kernel oil in maize. *Plant Physiol* **155**, 1146–1157 (2011).
39. McFie, P. J., Stone, S. L., Banman, S. L. & Stone, S. J. Topological orientation of acyl-CoA:diacylglycerol acyltransferase-1 (DGAT1) and identification of a putative active site histidine and the role of the N terminus in dimer/tetramer formation. *J Biol Chem* **285**, 37377–37387 (2010).
40. Zheng, P. *et al.* A phenylalanine in DGAT is a key determinant of oil content and composition in maize. *Nat Genet* **40**, 367–372 (2008).
41. Santos-Mendoza, M. *et al.* Deciphering gene regulatory networks that control seed development and maturation in Arabidopsis. *Plant J* **54**, 608–620 (2008).
42. Iyer, V. V. *et al.* Metabolic flux maps comparing the effect of temperature on protein and oil biosynthesis in developing soybean cotyledons. *Plant Cell Environ* **31**, 506–517 (2008).
43. Tjellström, H., Strawsine, M. & Ohlrogge, J. B. Tracking synthesis and turnover of triacylglycerol in leaves. *J Exp Bot* **66**, 1453–1461 (2015).
44. Figueroa, P. & Browse, J. Male sterility in Arabidopsis induced by overexpression of a MYC5-SRDX chimeric repressor. *Plant J* **81**, 849–860 (2015).
45. Stintzi, A. & Browse, J. The Arabidopsis male-sterile mutant, opr3, lacks the 12-oxophytodienoic acid reductase required for jasmonate synthesis. *Proc Natl Acad Sci USA* **99**, 8512–8517 (2000).

46. Guelette, B. S., Benning, U. F. & Hoffmann-Benning, S. Identification of lipids and lipid-binding proteins in phloem exudates from *Arabidopsis thaliana*. *J Exp Bot* **63**, 3603–3616 (2012).
47. Li, Q. *et al.* Understanding the biochemical basis of temperature-induced lipid pathway adjustments in plants. *Plant Cell* **27**, 86–103 (2015).
48. Sharma, N. *et al.* Transgenic increases in seed oil content are associated with the differential expression of novel Brassica-specific transcripts. *BMC Genomics* **9**, 619 (2008).
49. Zhao, J. *et al.* AtCHX13 is a plasma membrane K⁺ transporter. *Plant Physiol* **148**, 796–807 (2008).
50. Zhao, J. *et al.* MATE2 mediates vacuolar sequestration of flavonoid glycosides and glycoside malonates in *Medicago truncatula*. *Plant Cell* **23**, 1536–1555 (2011).
51. Wagner, M. *et al.* Identification and characterization of an acyl-CoA:diacylglycerol acyltransferase 2 (DGAT2) gene from the microalga *O. tauri*. *Plant Physiol Biochem* **48**, 407–416 (2010).
52. Wang, Z. & Benning, C. Arabidopsis thaliana polar glycerolipid profiling by thin layer chromatography (TLC) coupled with gas-liquid chromatography (GLC). *J Vis Exp* **49**, 2518 (2011).
53. Browse, J., McCourt, P. J. & Somerville, C. R. Fatty acid composition of leaf lipids determined after combined digestion and fatty acid methyl ester formation from fresh tissue. *Anal Biochem* **152**, 141–145 (1986).
54. Christie, W. W. Preparation of lipid extracts from tissue in *Advances in Lipid Methodology-II* (ed. Christie, W. W.) 195–213 (The Oily Press, 1993).

Acknowledgements

We thank Dr. Edgar Carhoon at University of Nebraska-Lincoln for providing yeast strain H1246, Dr. John Browse at Washington State University for providing *rod1* mutant seeds. Illumina data for the gene expression analysis during soybean seed development was carried out by Shanghai Aoxiang Scientific Company. This project is supported in part by Major State Basic Research Development Program of China (973 Program, Grant 2013CB127001) and the Huazhong Agricultural University Scientific & Technological Self-innovation Foundation (2012RC007) to J.Z.

Author Contributions

J.Z. and B.C. designed the research experiments, wrote the main manuscript text. B.C. did the most of experiments and analyzed data. G.Z., J.W., J.L. and S.M. did part of experiments (gene cloning, plant transformation, yeast complementation, hairy root transformation, lipid analysis, DNA extraction, and data analysis). H.H. provided facilities for Arabidopsis growth and transformation. J.Z., B.C. and H.H. edited the manuscript. All authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Chen, B. *et al.* Two types of soybean diacylglycerol acyltransferases are differentially involved in triacylglycerol biosynthesis and response to environmental stresses and hormones. *Sci. Rep.* **6**, 28541; doi: 10.1038/srep28541 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>