Social inequalities in female mortality by region and by selected causes of death, England and Wales, 2001–03

Ann Langford, Brian Johnson and
Alaa Al-Hamad
Office for National Statistics

This article reports on social inequalities in female mortality by region and cause of death for women aged 25–59 years in England and Wales in the period 2001–03. It is the first official compilation of detailed mortality statistics for women based on the National Statistics Socio-economic Classification (NS-SEC). It is part of a series of articles measuring inequalities in adult mortality.

The results demonstrate a strong socio-economic effect on the mortality of women in all regions. This pattern remained consistent between regions.

There were marked differences in the socio-economic gradient by cause. Mortality rates for women in the least and most advantaged NS-SEC classes were similar for breast cancer. In contrast, compared to women in the most advantaged class, mortality rates were three times as high for the least advantaged women for lung cancer and cerebrovascular disease, around five times as high for ischaemic heart disease and all digestive diseases, and six times as high for respiratory diseases.

Introduction

This article describes social inequalities in adult female mortality rates for selected causes of death and across the Government Office Regions of England, and Wales. It is the fifth in a series of articles reporting mortality using the final version of the National Statistics Socio-economic Classification (NS-SEC). The first three articles covered social inequalities by NS-SEC for men,¹ by cause of death,² and by Government Office Region.³ The fourth article examined inequalities for women in all-cause mortality,⁴ and presented results for two classifications, one based on a woman's own occupation, and another on a 'combined' classification based on the most advantaged NS-SEC class of the woman or her husband.

This study focuses on women aged 25–59 in the period 2001–03, and uses the 'combined' NS-SEC classification to analyse mortality rates by region and selected causes of death.

Background

There is a long history of the study of health inequalities by socio-economic classification in England and Wales. The influential Black report⁵ showed that there had been a striking lack of improvement in the health experience of the less advantaged social classes up to the 1970s. The Acheson Report⁶ in 1998 highlighted widening differences between the expectation of life of the most advantaged and most disadvantaged groups in society. The Government strategy *Tackling Health Inequalities: A programme for action*⁷ aspired to 'address the inequalities that are found across different geographical areas, between genders... and between different social and economic groups'. Four years later, however, the *2007 Status Report on the Programme for Action*⁸ reported that 'the gap has not narrowed for life expectancy in disadvantaged areas; indeed, the gap has widened, particularly for women.'

The interest in health inequalities has led to a large volume of literature on the analysis of mortality by socio-economic classification^{9,10,11} but relatively little on mortality in women. This is due, in part, to a number of well-known difficulties inherent in any analysis of female mortality by a classification based on occupation.¹² There are conceptual difficulties because many women have weaker ties to the labour market than men, which reduces the potential relevance of occupation-based indicators of social class. There are also practical difficulties, since the occupation of a substantial minority of women is inadequately described at death registration and, in many cases, is not recorded at all.

The previous article in this series⁴ examined two methods of classification for women. One was based on the woman's own occupation, and the other on a 'combined' measure which also took into account the husband's classification, where available and if reflecting more advantage than the woman's own occupational class. The concept behind the latter classification is similar to the 'dominance' approach first suggested by Erikson¹³ who maintained that the life-chances of individuals in a family unit are more likely to be aligned with those of the most advantaged individual in that unit. The idea was summarised as follows: 'A secretary who is married to an executive may have life chances closer to those of executives than to those of other secretaries.'¹⁴ The previous article in this series⁴ found that the 'combined' measure was a better discriminator of female mortality, with the mortality rate among those assigned to the least advantaged class 2.6 times that of those in the most advantaged class.

Previous authors^{11,15} have studied inequalities in the mortality of women by cause using the Registrar General's Social Class (RGSC).¹⁵ Others have studied inequalities in the mortality of women by region and cause in the early 1990s¹⁶ and covering the period of this study¹⁷ but both these studies analysed by deprivation of area of residence rather than individual socio-economic characteristics. All have found evidence of inequalities that vary by cause and region. However this article is the first to present an analysis of the mortality of women by selected causes of death and region using the new measure of social class – the NS-SEC – which replaced the RGSC in 2001.

Methods

In this article the measure of social class used is the National Statistics Socio-economic Classification (NS-SEC), which is described below, along with the definition of the regions and the selected causes analysed. The sources of data, calculation methods and outcomes are also described in this section.

The classification of women by the National Statistics Socio-economic Classification (NS-SEC)

The National Statistics Socio-economic Classification (NS-SEC) was developed in order to replace the Registrar General's Social Class, which had been criticised as lacking a coherent theoretical basis and becoming increasingly irrelevant to the changing patterns of industry and employment in modern economies.¹⁸

The conceptual basis of the NS-SEC is the structure of employment relations operating in modern developed economies.¹⁹ Occupations are differentiated in terms of reward mechanisms, promotion prospects, autonomy and job security. The most advantaged NS-SEC classes (for example higher managerial and professional occupations), typically exhibit personalised reward structures, have good opportunities for advancement, have relatively high levels of autonomy within the job, and are relatively secure. These attributes tend to be reversed for the most disadvantaged classes (for example routine occupations). Box One shows the NS-SEC analytical class breakdowns used in this analysis, and provides examples of the occupations included in each class.

Further information on the rationale, derivation and application of the NS-SEC is available on the Office for National Statistics (ONS) website.²⁰

This analysis uses the 'combined' classification method, whereby a non-married woman is assigned an NS-SEC class on the basis of her own occupation and employment status, and a married woman is assigned the most advantaged NS-SEC class of either herself or her husband.

Regions and selected causes of death

This article focuses on mortality in the Government Office Regions of England and in Wales. Government Office Regions are the largest level statistical sub-divisions in England (Box Two). Wales is not sub-divided in this analysis. For convenience, use of the term 'region' throughout this article includes Wales.

Further information can be found in the guide to UK geography on the ONS website.²¹

Box one

National Statistics Socio-economic Classification – analytic classes

Analytic class	Examples of occupations included
1 Higher managerial and professional occupations	Directors and chief executives of major organisations, civil engineers, medical practitioners, IT strategy and planning professionals, legal professionals, architects, senior officials in national and local government
2 Lower managerial and professional occupations	Teachers in primary and secondary schools, quantity surveyors, public service administrative professionals, social workers, nurses, IT technicians
3 Intermediate occupations	Graphic designers, medical and dental technicians, Civil Service administrative officers and local government clerical officers, counter clerks, school and company secretaries
4 Small employers and own account workers	Hairdressing and beauty salon proprietors, shopkeepers, dispensing opticians in private practice, farmers, self-employed decorators
5 Lower supervisory and technical occupations	Bakers and flour confectioners, catering supervisor, head waitress, postal supervisor, sales assistant supervising others
6 Semi-routine occupations	Retail assistants, catering assistants, clothing cutters, dressmaker, traffic wardens, veterinary nurses and assistants, shelf fillers
7 Routine occupations	Hairdressing employees, floral arrangers, sewing machinists, bar staff, cleaners and domestics
Other	Full-time students, never worked, long-term unemployed, inadequately described, not classifiable for other reasons

Source: NS-SEC User Manual, Office for National Statistics

Box two

The Government Office Regions of England

North East (NE)
 North West (NW)
 Yorkshire and The Humber (YH)
 East Midlands (EM)
 West Midlands (WM)
 East of England (E)
 London (L)
 South East (SE)
 South West (SW)

The causes of death examined in this article are listed in Box Three, along with the number and percentage of all deaths represented in the study population. These causes of death account for approximately 80 per cent of all deaths among women aged 25–59. Causes of death were grouped to ensure sufficiently large numbers for robust statistical analysis when broken down by region and NS-SEC class. Thus the mortality rates reported in this article are for all deaths, four major groups of causes and four specific causes of death. Deaths were coded to the tenth revision of the International Classification of Diseases (ICD).²²

Box three

Causes of death to women 25–59 included in the analysis

Cause of death	ICD-10 codes	Number of deaths	Percentage of deaths
All causes	A00–R99, V00–Y89	65,276	100
All cancers	C00–C97	31,639	48
Trachea, bronchus and lung	C33–C34	4,607	7
Breast cancer	C50	9,313	14
All circulatory diseases	I00–I99	11,505	18
Ischemic heart disease	I20–I25	4,616	7
Cerebrovascular disease	I60–I69	3,559	5
All respiratory diseases	J00–J99	3,794	6
All digestive diseases	K00–K93	5,322	8

Sources of data

Numerators

The routine collection of data at death registration provided the number of deaths in each age-group for each NS-SEC class in each region for the period 2001–03. These data were used as the basis of numerators for the mortality rates.

The numbers of deaths were subject to two adjustments. The first reapportioned some deaths from NS-SEC analytical classes 3 to class 2. In the first study of this series¹ examination of the ONS Longitudinal Study (LS), a one per cent sample of linked census and death records, revealed that a number of men had been allocated at death registration to NS-SEC class 3 on the basis that they had no supervisory duties. Examination of their 2001 linked Census records revealed, however, that they had in fact been supervisors in 2001, and thus were more appropriately assigned to NS-SEC class 2. A similar phenomenon was observed on examination of

the records of women in the ONS Longitudinal Study, and appropriate adjustment factors were calculated to correct for this bias.⁴

The second adjustment was to correct for the under-reporting of the occupation of women at death. The previous article⁴ reported that for 19 per cent of deaths insufficient occupational details were available to allow classification by 'combined' NS-SEC. In a sample of 158 women who were 'unclassified at death', the NS-SEC classification could be determined by reference to their census records. The distribution of this sample (Table 1) was used to reallocate the unclassified women across NS-SEC classes in this study.

Appendix A contains Tables A1 and A2, showing the numbers of deaths after these adjustments, by five-year age-band and 'combined' NS-SEC, for region and selected causes respectively.

Table 1

NS-SEC class¹ at Census for those female members of the LS who died 2001–05 and were not classified or inadequately described at death

England and Wales		Numbers and percentages								
		NS-SEC analytic class								Total
		1	2	3	4	5	6	7	Other and FTS ²	
Numbers		6	12	19	7	14	32	22	46	158
Percentages		4	8	12	4	9	20	14	29	100

1 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification.

2 Other and FTS includes never worked, long-term unemployed, inadequately described, not classifiable for other reasons and full-time students.

Source: ONS Longitudinal Study

Denominators

All denominators were based on the 'optimised population estimates' for mortality analysis presented in the previous article.⁴ To obtain regional denominators, for each age group and NS-SEC combination, the percentage distribution across the regions was calculated from 2001 Census data. The resulting percentages were applied to the optimised population estimates in order to obtain an estimate of person-years at risk by five-year age group and NS-SEC for each region.

Appendix A also contains Table A3, showing the resulting population estimates by region.

Outcome measures

Mortality rates per 100,000 person years, age-standardised to the European standard population, were calculated for each NS-SEC class in each Government Office Region, and for each selected cause of death. As in previous articles in this series^{1,2,3,4} the ratio between the mortality rates of the least and most advantaged NS-SEC classes is also presented. For convenience, this mortality rate ratio is referred to as the 'socio-economic gradient'.

Results

Mortality by region

Age-standardised mortality rates by NS-SEC for Government Office Regions and Wales are presented in Table 2. Shaded areas denote rates that are statistically significantly different from the corresponding rate for England and Wales as a whole. The results are illustrated graphically in Figure 1. Figure 2 displays the socio-economic gradient, that is the ratio of mortality rates of those assigned to routine occupations relative to those assigned to higher managerial and professional occupations, for each region.

The highest gradients are in London and the North West, where women classified to routine occupations have a mortality rate 3.0

Figure 1

Age-standardised mortality rates¹ by NS-SEC,² women aged 25–59, 2001–03

Government Office Regions of England, Wales

1 Directly age-standardised rate using the European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods

Table 2 Age-standardised mortality rates¹ by NS-SEC,² women aged 25–59, 2001–03

Government Office Regions of England, Wales Rates per 100,000

	NS-SEC analytic class							Socio-economic gradient ³
	1	2	3	4	5	6	7	
North East	126	153 H	167	201 H	237	237	329	2.6
North West	123	156 H	169	195 H	244 H	260 H	367 H	3.0
Yorkshire and The Humber	118	138	141	171	192	208	285	2.4
East Midlands	120	142	141	169	182 L	209	258 L	2.1
West Midlands	116	137	138	164	217	216	290	2.5
East of England	118	123 L	141	152	179 L	202	253 L	2.1
London	114	133	164	187	235	258 H	341	3.0
South East	111	128 L	137	133 L	204	201	282	2.5
South West	112	127 L	130 L	145 L	193	189 L	264	2.4
Wales	127	148	170	179	218	215	340	2.7
England	116	136	147	164	209	221	298	2.6
England and Wales	118	137	149	165	210	221	302	2.6

1 Directly standardised rate using the European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

3 Ratio of mortality rates of NS-SEC class 7 to the mortality rate of NS-SEC class 1.

H Rate is statistically significantly higher than that for England and Wales.

L Rate is statistically significantly lower than that for England and Wales.

Figure 2 Socio-economic gradients,¹ women aged 25–59, 2001–03

Government Office Regions of England, Wales

1 Ratio of mortality rates of NS-SEC class 7 to the mortality rate of NS-SEC Class 1 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Figure 3 Age-standardised mortality rates¹ by NS-SEC analytic class,² women aged 25–59, 2001–03

Government Office Regions of England, Wales

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

times that of women classified to higher managerial and professional occupations. This gradient is not however statistically significantly higher than the gradient for England and Wales as a whole (2.6). The lowest gradients are in the East of England and the East Midlands (2.1), and these gradients are statistically significantly lower than those for England and Wales.

Figure 3 demonstrates that there is relatively little difference in mortality rates between regions for the more advantaged classes, but more substantial differences for the more disadvantaged classes.

In detail, the results show that age-adjusted mortality rates in the North West for most NS-SEC classes are statistically significantly higher than the corresponding rates for England and Wales. In both the North West and the North East, mortality rates for each NS-SEC class are consistently higher than the corresponding rate for England and Wales.

Most mortality rates in the South West are statistically significantly lower than the corresponding NS-SEC class rates for England and Wales. In both the South West and the South East, mortality rates for each NS-SEC class are consistently lower than the corresponding rate for England and Wales.

Mortality by selected causes

Cancer

Table 3 presents the age-standardised rates by NS-SEC for all cancers; trachea, bronchus and lung cancer, and breast cancer. The results are displayed in Figure 4.

The mortality rates for all cancers displayed a distinct socio-economic pattern. The socio-economic gradient, that is the ratio of mortality rate of NS-SEC class 7 to NS-SEC class 1, was approximately 1.5. This implies that women in the least advantaged class had a mortality rate approximately one-and-a-half times that of women in the most advantaged class.

Table 3

Age-standardised mortality rates¹ from selected malignant neoplasms by NS-SEC², women aged 25–59, 2001–03

England and Wales										Rate per 100,000
NS-SEC analytic class	All cancers			Trachea, bronchus and lung			Breast cancer			
	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval	
1 Higher managerial and professional	75	72	78	7	6	8	26	25	28	
2 Lower managerial and professional	77	75	80	9	9	10	25	24	26	
3 Intermediate	73	69	77	9	8	10	22	21	24	
4 Small employers and own account workers	87	82	91	14	12	15	23	22	25	
5 Lower supervisory and technical	102	96	108	17	15	19	27	25	29	
6 Semi-routine	90	86	95	16	15	18	24	22	25	
7 Routine	116	108	123	23	21	25	29	26	31	
Ratio of classes 7:1	1.5	1.4	1.7	3.3	2.9	3.7	1.1	1.0	1.2	

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Figure 4

Age-standardised mortality rates¹ for all cancers, lung cancer and breast cancer by NS-SEC², women aged 25–59, 2001–03

England and Wales

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

The mortality rates for breast cancer, however, were fairly consistent over the socio-economic classes, with a gradient of only 1.1. The rate for the most advantaged class was not statistically significantly different from that of the least advantaged class. The pattern for lung cancer was quite different. Most classes had a statistically significantly higher mortality rate than the previous class, and mortality rates for women in the least advantaged class were more than three times that of women in the most advantaged class.

Circulatory diseases

Table 4 presents the age-standardised rates by NS-SEC for all circulatory diseases, and for ischaemic heart disease and cerebrovascular diseases separately. The results are displayed in Figure 5.

The mortality rates for all circulatory diseases displayed a more pronounced socio-economic pattern than those for cancer. Most classes had significantly higher mortality rates than the previous class. The exception was between NS-SEC classes 5 and 6 (lower supervisory and technical, and semi-routine). The mortality rate for NS-SEC class 7 (routine workers) was particularly high. The socio-economic gradient, that is the ratio of mortality rate of NS-SEC class 1 to NS-SEC class 7, was approximately 4.2. This implies that women in the least advantaged class had a mortality rate approximately four times that of women in the most advantaged class.

The socio-economic pattern was very similar for ischaemic heart disease. Women in the least advantaged class had a mortality rate 5.3 times that of women in the most advantaged class.

The socio-economic pattern for cerebrovascular disease was rather different to that of ischaemic heart disease. (Figure 5) The differences in mortality rates between adjacent classes were more consistent and the mortality rate for NS-SEC class 7 was not markedly higher. The socio-economic gradient was still high at 3.4, but was statistically significantly lower than the gradient for all circulatory diseases.

Respiratory and digestive diseases

The results for these two groups of diseases are shown in Table 5 and displayed in Figure 6. The socio-economic pattern of mortality rates was similar to that for ischaemic heart disease. Mortality rates rose gradually between NS-SEC classes 1 and 4. However, the difference in mortality rates between NS-SEC classes 4 and 5, and between classes 6 and 7 were statistically significant. The socio-economic gradients were 6.2 for respiratory diseases and 5.2 for digestive diseases.

Table 4

Age-standardised mortality rates¹ from selected circulatory diseases by NS-SEC,² women aged 25–59, 2001–03

England and Wales

Rate per 100,000

NS-SEC analytic class	All circulatory diseases			Ischaemic heart disease			Cerebrovascular diseases		
	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval
1 Higher managerial and professional	15	13	16	5	5	6	5	5	6
2 Lower managerial and professional	21	20	22	7	7	8	7	7	8
3 Intermediate	24	22	27	9	8	10	8	7	9
4 Small employers and own account workers	30	27	33	11	10	12	10	9	11
5 Lower supervisory and technical	41	37	45	18	16	20	12	11	14
6 Semi-routine	45	41	48	19	17	20	13	12	14
7 Routine	64	58	69	29	27	32	17	15	19
Ratio of classes 7:1	4.2	3.7	4.8	5.4	4.6	6.3	3.4	2.9	3.9

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Figure 5

Age-standardised mortality rates¹ for all circulatory diseases, ischaemic heart disease and cerebrovascular diseases by NS-SEC,² women aged 25–59, 2001–03

England and Wales

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Table 5

Age-standardised mortality rates¹ from all respiratory and all digestive diseases by NS-SEC,² women aged 25–59, 2001–03

England and Wales

Rate per 100,000

NS-SEC analytic class	All respiratory diseases			All digestive diseases		
	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval
1 Higher managerial and professional	4	3	5	6	5	7
2 Lower managerial and professional	6	5	6	9	9	10
3 Intermediate	7	6	9	12	11	14
4 Small employers and own account workers	9	7	10	12	10	14
5 Lower supervisory and technical	15	13	17	18	16	21
6 Semi-routine	16	14	17	20	18	22
7 Routine	24	21	27	32	28	35
Ratio of classes 7:1	6.2	5.0	7.5	5.2	4.3	6.2

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Figure 6**Age-standardised mortality rates¹ for all respiratory diseases and all digestive diseases by NS-SEC,² women aged 25–59, 2001–03**

England and Wales

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Comparison of gradients by cause

Figure 7 shows the socio-economic gradients for each of the disease categories. The smallest gradients were for all cancers combined and for breast cancer. The highest gradients were for respiratory diseases, digestive diseases and ischaemic heart disease, for which women in the least advantaged classes had mortality rates more than five times that of women in the most advantaged classes.

For all diseases studied, mortality rates for those classed as small employers and own account workers (NS-SEC class 4) were significantly lower than those classed as lower supervisory and technical (NS-SEC class 5). There were no significant differences between the rates for class 5 and the rates for those classed as semi-routine workers (NS-SEC class 6). Rates for those classed as routine workers (NS-SEC class 7) were statistically significantly higher than those classed as semi-routine workers (NS-SEC class 6) for all causes of death studied.

Figure 7**Socio-economic gradients¹ by selected causes, women 25–59, 2001–03**

England and Wales

1 Ratio of mortality rates of NS-SEC class 7 to the mortality rate of NS-SEC Class 1. NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Mortality by selected region and cause

The socio-economic variation of mortality by cause across regions was examined only for the North West and the South West, the regions with the highest and lowest all-cause mortality rates for most NS-SEC classes. Owing to the lack of statistically significant variation between regions for all-cause mortality, only mortality rates for the two major causes of death 'All cancers' (48 per cent of deaths) and 'All circulatory diseases' (18 per cent of deaths) are displayed in Table 6 and illustrated in Figure 8.

The estimated mortality rates for the North West are generally higher than those in the South West and statistically significantly so for NS-SEC classes 2, 3, 4 and 6 for all cancers combined, and for all classes except NS-SEC class 3 for circulatory diseases. By contrast, the socio-economic gradients are not statistically significantly different between the two regions for either major cause of death.

Table 6**Age-standardised mortality rates¹ for selected causes and regions by NS-SEC,² women aged 25–59, 2001–03**

NS-SEC analytic class	Rates per 100,000			
	All cancers		All circulatory diseases	
	North West	South West	North West	South West
1 Higher managerial and professional	69	76	18	11
2 Lower managerial and professional	86	74	25	18
3 Intermediate	76	65	29	22
4 Small employers and own account workers	97	83	35	23
5 Lower supervisory and technical	107	98	51	38
6 Semi-routine	104	78	51	38
7 Routine	125	112	78	49
Ratio of classes 7:1	1.8	1.5	4.4	4.4

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Figure 8

Age-standardised mortality rates¹ for selected causes and regions by NS-SEC,² women aged 25–59, 2001–03

England and Wales

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

Discussion

Regional mortality

The results show variation in mortality rates between regions, with rates in the North West and North East consistently higher for all classes than rates in the South West and South East. A 'north-south' divide was also reported by authors considering female mortality by deprivation of area of residence both in the 1990s¹⁶ and in the period covered in this study.¹⁷ A 'north west-south east divide' in self-reported health²³ was also reported where the authors found that each of the seven NS-SEC classes had 'higher rates of poor health in Wales, the North East and the North West regions of England than elsewhere'.

This analysis also found that regional differences in mortality were smaller for the most advantaged classes and greater for the least advantaged classes. The comparative insensitivity of the most advantaged classes to regional effects has also been found by authors working on local area deprivation.¹⁷ That study found that 'those living in the least deprived areas had similar mortality rates, independent of region'. A similar pattern was reported in the third article in the current series which considered male socio-economic inequalities by region using NS-SEC.³ On the other hand, for the least advantaged, differences in mortality were more apparent between regions. The North West had mortality rate of 367 all-cause deaths per 100,000 person-years while the East of England had a rate of 253 all-cause deaths per 100,000 person years. A similar effect was found in the analysis of men,³ however the effect for men appeared to be stronger, since the mortality rate for the least advantaged class varied by approximately 80 per cent across regions, compared with a variation of approximately 45 per cent for women.

The current results do not indicate large regional variation in socio-economic gradients. The highest and lowest gradients were 2.1 and 3.0 respectively, but only two regions had significantly different socio-economic gradients to those for England and Wales as a whole. In the corresponding analysis of men³ the gradients were similar, but there were significant differences in seven of the ten regions. This may be because relative differences between the sexes were smaller, or because the methodology adopted in the current analysis was not sufficiently sensitive to detect smaller differences.

The crucial aspect of the methodology that could affect the ability of this study to detect smaller differences involves the adjustments that have been made to the data to correct for the under-recording of occupation at death. These adjustments were made to correct for bias, but have necessarily resulted in larger confidence intervals for the estimates. Consequently there

were difficulties in the detection of differences in smaller sub-samples, such as the smaller regions and also in smaller causes of death.

However, authors considering mortality by deprivation^{16,17} have also found that although mortality rates increase with deprivation for both sexes, the relationship was generally stronger for males. Figures published by authors studying self-reported health²³ have indicated that relative differences between rates of those reporting 'not good' general health between the least and most advantaged NS-SEC classes for England were of the order of 2.7 for men, but only 2.2 for women. Hence there is some evidence in the literature that relative differences in regional socio-economic inequalities were smaller for women.

Many reasons have been suggested as to why there may be inequalities between regions, particularly in the most disadvantaged classes.³ Theoretical explanations include suggestions that the regional differences in mortality are associated with regional differences in unemployment risk, selective migration, differential concentrations of deprived areas and material disadvantage, or differing geographic patterns in health-related behaviours.

It is well established that there is a higher prevalence of ill-health and excess mortality in men and women who are unemployed.^{24,25} In addition, the authors suggested that job insecurity itself may constitute a psychosocial hazard to health. As unemployment and the risk of unemployment is greater in the north,²⁶ this implies greater psychosocial stress on both men and women, particularly on those in routine or semi-routine occupations. Women may be affected directly, but possibly also through their husbands' employment position, potentially leading to increased stress and consequent mortality for women in the north, and particularly for the more disadvantaged.

Another potential explanation of regional inequalities is that of selective migration.²⁷ This explanation suggests that those with good long-term health are more likely to move (for example to find work) while those with serious health problems are less likely to move and will have higher mortality rates. This is the so-called 'healthy migrant effect' which would lead to varying mortality rates between advantaged and disadvantaged areas. Some authors²⁸ however have questioned whether this is a significant effect at the regional level. There are also questions as to how relevant this factor is for female mortality, although others have found that gender made little difference to the likelihood of migrants moving to more advantaged areas.²⁹

A further explanation of regional differences may be that regions in the north have a higher concentration of deprived neighbourhoods, whose

effect is not fully captured by the use of an occupational based socio-economic measure such as NS-SEC. A study³⁰ on cancer survival trends of males and females in England, for instance, found that survival was consistently lower for patients in the deprived Spearhead Primary Care Trust (PCT) Areas than for those resident in the rest of England. Most of these PCT areas are in the North, while there are none in the South West or South East. The disproportionate concentration of local deprivation may therefore be part of the explanation of the regional differences observed in the current study.

A related explanation is that regional differences may be directly related to material deprivation. Materialist explanations of inequalities suggest that individual incomes determine living conditions, including diet and housing quality, which consequently affect health and mortality.²⁴ Both income inequality and mean income have indeed been found to be associated with mortality³¹ and it has been reported that in 2003–04 average incomes in the North East and the North West were among the smallest in England.³²

Another explanation of the regional differences is that there may be regional differences in health-related behaviour between regions. Lifestyles have been recognised as contributing strongly to inequalities in mortality³³ and differences in the prevalence of smoking, in particular, have been found to account for much of the variation in mortality between areas.³⁴ More recently it has been shown in a study looking at self-reported health in women that the strongest independent effect on health was from smoking.³⁵ It is known that levels of smoking vary quite markedly over both regions and social class. The General Household Survey found that in England in 2001, 30 per cent of women living in households classified to the manual group smoked cigarettes, compared with 20 per cent of those in households classified as non-manual. The GHS also reported on variation by region with 29 per cent of women smoking in the North West in 2001, but only 22 per cent in the South West.³⁶ This would also seem to be a plausible explanation as to why more disadvantaged women in the north have a higher mortality rate than women in a similar position in the south.

Mortality by cause

Malignant neoplasms

The results presented here show a significant, but relatively small, socio-economic gradient for all cancers combined (1.5), a marked gradient for lung cancer (3.3) but no significant gradient for breast cancer.

Many authors have also found strong socio-economic effects in lung cancer mortality.^{11,15} In addition, an international study³⁷ comparing mortality rates for 1981–85 with 1991–95, found that in four western European countries rates of mortality from lung cancer in women had risen over the decade. The authors also found that this rise had been greater for women with lower educational levels in three of the countries studied. It has been estimated³⁸ that eight in ten lung cancer deaths in women are smoking-attributable, so the socio-economic gradient observed in the current study may well reflect the differential smoking patterns in socio-economic groups.³⁶ Many authors^{39,40} go further, postulating that ‘the combination of the greatly increased mortality of smokers with the now much lower prevalence of smoking among the more affluent is the major contributor to the widening health inequalities’³⁹ and that ‘scope for reducing health inequalities related to social position is limited unless many smokers in lower social positions stop smoking’.³⁹

In contrast, a flat or inverse socio-economic gradient in breast cancer mortality has been well documented. A study of mortality covering the 1970s and 1980s, which based socio-economic status on tenure and car-ownership, found that breast cancer did not appear to differ significantly between socio-economic groups.⁴¹ Later findings¹¹ using the Registrar General’s Social Class (RGSC) covering the late 1990s found an inverse gradient in breast cancer mortality and concluded that this inverse

gradient had increased over the period 1986–2000. An international study⁴² also covering the 1990s found that in eight of the 11 European populations studied, women with a higher education level had a greater risk of dying from breast cancer than women of a lower educational level. It has been suggested that this pattern may be related to the rise in incidence of later childbearing, which is more prominent in women in more advantaged classes.^{24,43}

The comparatively flat socio-economic pattern in breast cancer mortality found in this study is probably attributable to the fact that while breast cancer incidence is higher among more affluent women, survival is also higher among these groups. Breast cancer incidence has been rising steadily, both in the age range invited for breast screening (50–69 years) and at other ages.^{44,45} Survival has also been rising, however.^{46,47} Thus although overall five-year survival rose from 68 per cent for women diagnosed in the late 1980s to 80 per cent for women diagnosed in the late 1990s, the gap in survival between the most affluent and the most deprived was stable at around 6 per cent.

The results shown here for all cancers combined can be compared with those from an international study⁴⁸ of women aged 20–74 in seven countries that used educational qualifications as a proxy for socio-economic status. That study found an inverse gradient for all neoplasms in two eastern European countries, a small positive gradient in two Scandinavian countries and no significant gradient in the other three countries. The results presented here are more compatible with the Scandinavian results.

Circulatory diseases

This study found higher inequalities for all circulatory diseases and particularly for ischaemic heart disease, the socio-economic gradients were 4.3 and 5.4 respectively. Earlier authors looking at manual/non-manual mortality rate ratios for the 1980s¹⁵ and 1990s¹¹ also found that ischaemic heart disease had a ratio more than twice that of breast cancer. An international comparison⁴⁸ also found stronger socio-economic gradients for cardiovascular disease than for all cancer amongst women.

The explanations that have been put forward for the existence of a socio-economic gradient for circulatory diseases have included differential behavioural factors, material circumstances and exposure to psychosocial hazards at work. Some⁴⁸ have suggested that inequalities in ischaemic heart disease could be due to a strong social patterning of behavioural risk factors (dietary factors, lack of physical activity, obesity) among women. Indeed, it has been found that in 2001, 30 per cent of women in routine occupations were obese compared with 16 per cent in higher managerial and professional occupations.⁴⁹ The authors⁴⁸ suggest that obesity may be a response of women to material disadvantage and/or psychosocial stressors. Support for psychosocial explanations comes from authors who found that high job strain and effort-reward imbalance seemed to increase the risk of cardiovascular mortality in a study that included both sexes.⁵⁰ Others⁵¹ have also claimed that much of the difference in incidence of coronary heart disease can be explained by differences in the psychosocial work environment, with additional contributions from behavioural factors such as smoking and from some life-course factors. The psychosocial explanation is more problematic for this study, as the combined indicator allocates a substantial minority of women to their husband’s occupation, rather than their own.

Respiratory and digestive diseases

This study found that mortality rates from these diseases exhibited a similar pattern to that of mortality rates from IHD. It has been estimated³⁸ that over 80 per cent of deaths from chronic obstructive lung disease can be attributed to smoking. As this disease accounts for over 70 per cent of respiratory disease deaths, it seems that the prevalence of smoking among differing social classes would again be an important factor in explaining socio-economic gradients observed.

Approximately three-quarters of deaths from all digestive diseases are associated with liver disease. The relationship between alcohol consumption and liver disease is well founded^{52,53} but the relationship between alcohol consumption and socio-economic position is complex. Surveys on alcohol consumption have reported that women in managerial and professional households are more likely than other women to have drunk alcohol in the last week,^{36,54} and more likely to have drunk more in the week than other women³⁶. However, it has also been found⁵⁴ that among women, those living in semi-routine and routine households were the most likely to indulge in binge drinking, that is drinking that exceeds twice the recommended daily limits. This later study also indicated that this drinking behaviour was far more common at younger ages. As other authors⁵⁵ have found that, for younger women, those in the manual classes were more likely to die from alcohol-related causes, binge drinking may be an important factor in explaining the socio-economic mortality gradient found in this study.

Another possible explanation may be related to the individual occupations included in the more disadvantaged classes. A recent study of alcohol-related mortality by occupation⁵⁶ found that, for women aged 20–64, the seven occupations with the highest alcohol-related mortality included bar staff, waitresses, hairdressers and elementary office occupations. Most of these occupations would primarily be classified as semi-routine or routine, and may therefore also partly explain the socio-economic gradient found in this study.

Mortality by selected region and cause

Only the two regions with highest and lowest all-cause mortality rates were compared for all cancers and all circulatory diseases. There was evidence of differences in mortality rates between the two regions in all classes but the socio-economic gradients, as measured by the ratio of the mortality rates between the least and most advantaged classes, were not statistically significant. This may be partly due to methodological difficulties in detecting small differences noted earlier. However, other authors¹⁷ have also found relatively small differences between inequalities for these two diseases and regions. For instance, they found that the ratio of mortality rates between the first and fifth deprivation quintile for all cancers was 1.5 for the North West and 1.4 for the South West. For all circulatory diseases the ratio was 2.4 for both the North West and the South East.

Comparison with male mortality

The second article in this series described the socio-economic inequalities by cause of death for men aged 25–64.² However, mortality rates for the different diseases between the two articles are not directly comparable for three main reasons. The first is that since women in 2001 retired at 60, the working-age range was chosen to be 25–59, and so the age range is not comparable. The second reason is that it is well known that the effect of disease varies considerably between sexes. The final reason is that the male analysis was conducted using the man's 'own' NS-SEC, but the female analysis was done on a 'combined' quasi-household measure.

Given these provisos, although the mortality rates themselves are not comparable, the inequalities, that is the socio-economic gradients, may be compared. Table 7 presents the socio-economic gradients from the study of men alongside those presented above.

There are similarities in the patterns – inequalities in the diseases studied for both sexes are smallest for all cancers, but higher for lung cancer. In this study the gradients for lung cancer appear similar, other studies have also found that the ratio between most and least deprived were similar for males and females.¹⁷ However, others^{15,57} have found that women have an apparently greater relative risk than men of lung cancer.

The socio-economic gradient for ischaemic heart disease for women is markedly higher than that observed for men. A very similar pattern was observed in a study by area deprivation.^{16,17} The inequalities for women

Table 7

Socio-economic gradients for males and females by selected causes

	Men ¹	Women ²
All Cancer	1.8	1.5 (1.4, 1.7)
<i>Cancer of Trachea, bronchus and lung</i>	3.7	3.3 (2.9, 3.7)
All circulatory diseases	2.8	4.2 (3.7, 4.8)
<i>Ischemic heart disease</i>	2.9	5.4 (4.6, 6.3)
<i>Cerebrovascular disease</i>	2.9	3.4 (2.9, 3.9)
All respiratory diseases	4.9	6.2 (5.0, 7.5)
All digestive diseases	3.5	5.2 (4.3, 6.2)

1. From previous article in series¹

2. 95% confidence interval in brackets

seem also to be higher for respiratory and digestive diseases, but given the provisos mentioned earlier and the breadth of the confidence intervals it is difficult to draw any firm conclusion.

Limitations of the analysis

Owing to the very sparse recording of women's occupations at death after normal retirement age, it was necessary to restrict the analysis to women aged 25–59. Since only 8 per cent of adult women died age 59 or lower in the years 2001–03, this analysis is focused only on a minority of 'premature' deaths.

There are conceptual problems with the use of a 'combined' NS-SEC. NS-SEC is theoretically based on the employment relations of the individual, using the 'most advantaged' NS-SEC of a married couple implies that one can use the most advantaged NS-SEC of marriage partners as a proxy for the life-chances of each partner. This difficulty is fully discussed in the article on female all cause mortality.⁴

The results are sensitive to the adjustment to the deaths not classified to an occupied NS-SEC class. This adjustment was, of necessity, based on a relatively small sample (only 158 deaths) and resulted in much larger confidence intervals for the estimates presented than for the unadjusted figures. The unadjusted estimates are available for comparison in Appendix B, Table B1–B5. The size of the confidence intervals makes it much more difficult to detect differences in smaller sub-samples, and may partly explain why no significant differences between regional socio-economic gradients within causes were detected.

The outcome measure used throughout this series of articles, referred to above as the socio-economic gradient, has a number of limitations. As it is an age-standardised mortality ratio, it does not take account of the size of each class, nor what happens in the intervening classes. More complex outcome measures, such as the slope index of inequality or the GINI coefficient,⁵⁸ could potentially be used in future studies.

The death registers during the period of this study did not recognise civil partnerships, and thus women in such partnerships have been treated as 'single'. The identical definition was used to obtain the census populations so the results presented above are consistent. However there is an argument for treating women in partnerships in the same way as married women.

Conclusions

This analysis has estimated standardised mortality rates by 'combined' NS-SEC for women aged 25–59 in the period 2001–03 by region and cause of death.

A clear social gradient is evident for each region. Regional differences in absolute mortality rates were smaller for the most advantaged class, no region had a mortality rate statistically significantly different from

Key findings

- There were socio-economic differences in mortality for women across Wales and all the English regions for the period 2001–03, with higher mortality rates in the more disadvantaged classes
- Regional differences in mortality were small for the most advantaged classes and greatest for the least advantaged
- The socio-economic gradient did not vary considerably from region to region for all-cause mortality – only two regions had gradients that were significantly different from that of England and Wales as a whole
- Among the causes of death analysed there were marked socio-economic differences in all respiratory diseases, all digestive diseases including liver diseases, ischaemic heart disease, all circulatory diseases, cerebrovascular disease and lung cancer
- There was no significant difference between mortality rates for the most and least advantaged for breast cancer

the England and Wales rate. Differences between regions were more pronounced for the least advantaged, but only three regions had rates that were statistically significantly different from the England and Wales rate.

Mortality rates were significantly higher for the North West region than for England and Wales as a whole for most NS-SEC classes, while mortality rates were lower in the South West region for most NS-SEC classes. None of the regional socio-economic gradients (that is the ratio between mortality rates for the least and most advantaged) were statistically significantly higher than that for England and Wales, as a whole.

A social gradient was also evident for all causes of death studied except for breast cancer. For breast cancer the mortality rates for the least and most advantaged were not statistically significantly different. For the other diseases studied there were marked socio-economic differences. The most disadvantaged women had approximately three times the mortality rate of more advantaged women for lung cancer and cerebrovascular disease, around five times as high for ischaemic heart disease and digestive diseases, and six times as high for respiratory diseases.

Mortality rates for the regions with generally the lowest and highest mortality rates were compared for the two largest causes studied, 'all cancers' and 'all circulatory diseases'. There were statistically significant differences between the mortality rates of the two regions in most NS-SEC classes, but no statistically significant differences in the socio-economic gradient between the two regions.

In comparison to men, there were similarities in the pattern of inequalities between diseases. Inequalities were smallest for all cancer and largest for respiratory and digestive diseases. Inequalities for ischaemic heart disease were particularly high for women.

References

1. White C, Glickman M, Johnson B and Corbin T (2007) 'Social inequalities in adult male mortality by the National Statistics Socio-Economic Classification, England and Wales, 2001–03', *Health Statistics Quarterly* **36**, 6–23. Available on the Office for National Statistics website at:

www.statistics.gov.uk/statbase/Product.asp?vlnk=6725

2. White C, Edgar G and Siegler V (2008) 'Social inequalities in male mortality for selected causes of death by the National Statistics Socio-Economic Classification, England and Wales, 2001–03', *Health Statistics Quarterly* **38**, 19–31. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
3. Siegler V, Langford A and Johnson B (2008) 'Regional differences in male mortality using the National Statistics Socio-economic Classification, England and Wales, 2001–03', *Health Statistics Quarterly* **40**, 6–18. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
4. Langford A and Johnson B (2009) 'Social inequalities in female mortality by the National Statistics Socio-economic Classification, England and Wales, 2001–03', *Health Statistics Quarterly* **42**, 6–21. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
5. Black D, Morris J, Smith C and Townsend P (1980) *Inequalities in health: report of a research working group*. Department of Health and Social Security: London.
6. Acheson D (1998) *Independent inquiry into inequalities in health*, The Stationery Office: London.
7. Department of Health (2003) *Tackling Health Inequalities: A programme for action*, Department of Health: London.
8. Department of Health (2007) *Tackling Health Inequalities: 2007 Status Report on the Programme for Action*, Department of Health: London.
9. Goldblatt P (ed) (1990), *Longitudinal Study: Mortality and Social Organisation*, Series LS No.6, Her Majesty's Stationery Office: London.
10. Drever F and Whitehead M (eds) (1997), *Health Inequalities: Decennial Supplement*, Series DS no. 15, The Stationery Office: London.
11. White C, van Galen F and Chow Y H (2003) 'Trends in social class differences in mortality by cause, 1986 to 2000', *Health Statistics Quarterly* **20**, 25–37. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
12. Moser K, Goldblatt P and Pugh H (1990) 'Occupational Mortality of Women in Employment' in Goldblatt P (ed), *Longitudinal Study: Mortality and Social Organisation*, Series LS no.6, Her Majesty's Stationery Office: London.
13. Erikson R (1984) 'Social class of men, women and families', *Sociology*, **18**, 500–14.
14. Erikson R (2006) 'Social class assignment and mortality in Sweden', *Social Science in Medicine* **62** (9) 2151–60.
15. Harding S, Bethune A, Maxwell R, and Brown J (1997) 'Mortality trends using the Longitudinal study' in Drever F and Whitehead M (eds), *Health Inequalities: Decennial Supplement*, Series DS no. 15, The Stationery Office: London, 143–55.
16. Uren Z and Fitzpatrick J (2001) in Griffiths C and Fitzpatrick J (eds), *Geographic Variations in Health*, Series DS no 16 (Chapter 11), 326–36. The Stationery Office: London.
17. Romeri E, Baker A and Griffiths C (2006) 'Mortality by deprivation and cause of death in England and Wales, 1999–2003', *Health Statistics Quarterly* **32**, 19–34. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
18. Goldthorpe J (2000) 'Social class and the differentiation of employment contracts' in Goldthorpe J (ed), *On Sociology: Numbers, Narratives and the Integration of Research and Theory*, Oxford University Press: Oxford.
19. Rose D and Pevalin D (eds) (2003) *A Researcher's Guide to the National Statistics Socio-economic Classification*, Sage: London.
20. Office for National Statistics, The National Statistics Socio-economic

- Classification on-line edition. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=13561
21. Office for National Statistics, Geography User Guide. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=7428&Pos=2&ColRank=1&Rank=240 per cent20(geog)
 22. Rooney C and Smith S (2000) 'Implementation of ICD-10 for mortality data in England and Wales from January 2001', *Health Statistics Quarterly* **8**, 41–51. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
 23. Doran T, Drever F and Whitehead M (2004) 'Is there a north-south divide in social class inequalities in health in Great Britain? Cross sectional study using data from the 2001 Census', *British Medical Journal*, **328**, 1043–45.
 24. Bartley M (2004) *Health Inequality: an introduction to theories, concepts and methods*, Polity Press: Cambridge.
 25. Bartley M, Ferrie J and Montgomery S (1999) 'Living in a high-unemployment economy: understanding the health consequences' in Marmot M and Wilkinson R G (eds) *Social Determinants of Health*, Oxford University Press: Oxford.
 26. Summerfield C and Babb P (eds) (2003) 'Unemployment rates: by region, 2002', *Social Trends* **33**, 83. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/ssdataset.asp?vlnk=6280
 27. Griffiths C, Goldblatt P and Fitzpatrick J (2001) 'Geographic variations in health: main findings and implications for the future' in Griffiths C and Fitzpatrick J (eds), *Geographic Variations in Health*, Series DS no **16** (Chapter 11), 360–68. The Stationery Office: London.
 28. Brimblecombe N, Dorling D and Shaw M (1999) 'Mortality and migration in Britain, first results from the British Household Panel Survey', *Social Science and Medicine* **49**, 981–88.
 29. Connolly S, O'Reilly D and Rosato M (2007) 'Increasing inequalities in health: Is it an artefact caused by the selective movement of people?', *Social Science & Medicine* **64**, 2008–15.
 30. Ellis L, Rachet B, Shah A *et al* (2009) 'Trends in cancer survival in Spearhead Primary Care Trusts in England, 1998–2004', *Health Statistics Quarterly* **41**, 7–12. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
 31. Stanistreet D, Scott-Samuel A and Bellis M A (1999) 'Income inequality and mortality in England', *Journal of Public Health Medicine* **21**, 205–07.
 32. Her Majesty's Revenue and Customs. Income Distribution. Accessed 27 October 2009, available at:
www.hmrc.gov.uk/stats/income_distribution/3_13_apr06.PDF
 33. Balia S and Jones A (2008) 'Mortality, lifestyle and socio-economic status', *Journal of Health Economics* **27**, 1–26.
 34. Law M and Morris J K (1998) 'Why is mortality higher in poorer areas and in more northern areas of England and Wales?', *Journal of Epidemiology and Community Health* **52**, 344–52.
 35. Sacker A, Bartley M, Firth D *et al* (2001) 'Dimensions of social inequality in the health of women in England: occupational, material and behavioural pathways', *Social Science & Medicine* **52**, 763–81.
 36. Office for National Statistics (2002) *Living in Britain 2001*. Available on the Office for National Statistics website at:
www.statistics.gov.uk/lib2001/Section3536.html
 37. Mackenbach J P, Bos V, Andersen O *et al* (2003) 'Widening socio-economic inequalities in mortality in six Western European countries' *International Journal of Epidemiology* **32**, 830–37.
 38. Twigg L, Moon G and Walker S (2004) *The Smoking Epidemic in England*, Health Development Agency: London.
 39. Gruer L, Hart C L, Gordon D S *et al* (2009) 'Effect of tobacco smoking on survival of men and women by social position: a 28 year cohort study', *British Medical Journal* **338**, 480.
 40. Jarvis M J and Wardle J (1999) 'Social patterning of individual health behaviours: the case of cigarette smoking' in Marmot M and Wilkinson R G (eds) *Social Determinants of Health*, Oxford University Press: Oxford
 41. Smith J and Harding S (1997) 'Mortality of women and men using alternative social classifications' in Drever F and Whitehead M (eds), *Health Inequalities: Decennial Supplement*, Series DS no. **15**, The Stationery Office: London, 143–55.
 42. Strand B H, Kunst A, Huisman M *et al* (2007) 'The reversed social gradient: Higher breast cancer mortality in the higher educated compared to the lower educated. A comparison of 11 European populations during the 1990s', *European Journal of Cancer* **42**, 1200–07.
 43. Brown J, Harding S, Bethune A *et al* (1997) 'Incidence of Health of the Nation cancers by social class', *Population Trends* **90**, 40–48.
 44. Westlake S and Cooper N (2008) 'Cancer incidence and mortality: trends in the United Kingdom and constituent countries 1993 to 2004' *Health Statistics Quarterly* **38**, 33–43. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
 45. Rowan S (2007) 'Trends in cancer incidence by deprivation, England and Wales, 1990–2002', *Health Statistics Quarterly* **36**, 24–35. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
 46. Quinn M J, Cooper N, Rachet B *et al* (2008) 'Survival from cancer of the breast in women in England and Wales up to 2001', *British Journal of Cancer* **99** (Suppl 1), 53–55.
 47. Coleman M P, Rachet B, Woods L M *et al* (2004) 'Trends in socio-economic inequalities in cancer survival in England and Wales up to 2001', *British Journal of Cancer* **90**, 1367–73.
 48. Mackenbach J P, Kunst A E, Groenhouf F *et al* (1999) 'Socio-economic Inequalities in Mortality Among Women and Among Men: An International Study', *American Journal of Public Health* **89**, 1800–06.
 49. Office for National Statistics (2004) 'Obesity among adults: by sex and NS-SEC 2001', *Social Trends* **34**, 111–12. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/ssdataset.asp?vlnk=7447
 50. Kivimäki M, Leino-Arjas P, Luukkainen R *et al* (2002) 'Work stress and risk of cardiovascular mortality: prospective cohort study of industrial employees' *British Medical Journal* **325**, 857.
 51. Marmot M G, Bosma H, Hemingway H *et al* (1997) 'Contribution of job control and other risk factors to social variations in coronary heart disease incidence' *The Lancet* **350**, 235–39.
 52. Brunn K, Edwards G, Lumio M *et al* (1975) *Alcohol Control Policies in Public Health Perspective*, Finnish Foundation for Alcohol Studies: Helsinki.
 53. Anderson P (1995) 'Alcohol and the risk of physical harm' in Holder H and Edwards G (eds), *Alcohol and Public Policy: evidence and Issues*, 82–113.
 54. Erens B (2003) 'Alcohol Consumption' in Sproston K and Primates P (eds), *Health Survey for England 2003*, Department of Health: London.
 55. Harrison L and Gardiner E (2002) 'Do the rich really die young? Alcohol related mortality and social class in Great Britain 1988–94', *Addiction* **94**:12, 1871–80.
 56. Romeri E, Baker A and Griffiths C (2007) 'Alcohol-related deaths by occupation, England and Wales, 2001–05', *Health Statistics Quarterly* **35**, 6–12. Available on the Office for National Statistics website at:
www.statistics.gov.uk/statbase/Product.asp?vlnk=6725
 57. Payne S (2001) 'Smoke like a man, die like a man?: A review of the relationship between gender, sex and lung cancer', *Social Science and Medicine* **53**, 1067–68.
 58. South East Public Health Observatory (2005). 'The Public Health Observatory Handbook of Health Inequalities Measurement'. Accessed on 27 October 2009, available at: www.sepho.org.uk/extras/rch_handbook.aspx

Appendix A

Table A1

Number of deaths by NS-SEC¹ and age, women aged 25–59, 2001–03

Government Office Regions of England, Wales

Numbers

	Higher managerial and professional	Lower managerial and professional	Intermediate	Small employers and own account workers	Lower supervisory and technical	Semi-routine	Routine	Others ²	Total
North West									
25–29	13	42	48	12	17	51	32	42	258
30–34	39	65	65	21	35	97	63	55	440
35–39	60	120	90	41	59	127	85	64	647
40–44	85	199	130	72	89	187	148	76	986
45–49	128	347	177	121	153	287	227	114	1,554
50–54	197	562	280	218	263	442	351	121	2,434
55–59	296	722	358	336	420	600	532	151	3,414
Total	818	2,058	1,148	820	1,035	1,793	1,438	623	9,733
Percentage	8.4	21.1	11.8	8.4	10.6	18.4	14.8	6.4	
North East									
25–29	3	13	15	4	5	21	11	13	86
30–34	9	20	21	7	14	33	21	19	143
35–39	14	42	20	16	22	50	28	22	213
40–44	18	80	46	27	40	66	50	31	358
45–49	53	123	69	41	59	92	94	40	570
50–54	67	221	104	72	122	160	161	53	960
55–59	93	228	130	78	186	249	239	65	1,269
Total	258	727	405	245	447	671	604	243	3,599
Percentage	7.2	20.2	11.2	6.8	12.4	18.6	16.8	6.8	
Yorkshire and The Humber									
25–29	11	29	25	7	8	48	28	26	182
30–34	23	62	40	19	21	48	32	27	273
35–39	42	101	42	36	36	86	67	37	448
40–44	53	140	70	50	58	142	101	40	654
45–49	105	216	112	80	104	168	136	56	977
50–54	134	343	155	146	151	253	208	58	1,446
55–59	184	428	229	214	280	397	353	85	2,170
Total	552	1,319	673	551	657	1,142	924	330	6,150
Percentage	9.0	21.5	11.0	9.0	10.7	18.6	15.0	5.4	
East Midlands									
25–29	9	24	17	8	6	22	22	18	125
30–34	24	44	29	17	22	53	30	26	245
35–39	36	85	45	28	30	67	38	29	359
40–44	59	124	54	41	44	103	72	33	531
45–49	90	165	91	67	84	144	98	41	780
50–54	148	320	144	116	128	219	185	56	1,318
55–59	184	431	191	212	235	332	313	78	1,976
Total	550	1,194	572	488	550	942	757	281	5,334
Percentage	10.3	22.4	10.7	9.2	10.3	17.7	14.2	5.3	
West Midlands									
25–29	14	19	26	8	13	32	23	24	158
30–34	18	52	33	16	22	63	41	30	275
35–39	34	86	52	29	43	66	54	33	397
40–44	88	150	75	44	66	118	97	38	676
45–49	91	198	93	83	102	207	136	52	962
50–54	161	381	187	147	206	323	244	77	1,726
55–59	239	501	230	229	323	452	342	87	2,404
Total	646	1,386	695	555	776	1,262	937	341	6,598
Percentage	9.8	21.0	10.5	8.4	11.8	19.1	14.2	5.2	
East of England									
25–29	15	46	27	5	9	31	14	18	166
30–34	32	62	47	20	20	46	30	30	287
35–39	53	103	62	33	29	91	47	35	453
40–44	81	126	80	56	48	101	61	33	586
45–49	136	221	126	88	90	161	95	49	965
50–54	199	396	209	161	146	253	149	60	1,574
55–59	308	543	284	246	217	354	198	80	2,231
Total	824	1,498	835	609	559	1,037	594	305	6,262
Percentage	13.2	23.9	13.3	9.7	8.9	16.6	9.5	4.9	
London									
25–29	31	60	37	17	15	53	27	48	288
30–34	47	107	74	25	30	88	48	74	494
35–39	88	180	115	50	50	114	67	99	763
40–44	121	246	167	68	64	152	100	96	1,014
45–49	150	305	171	92	98	218	114	108	1,256
50–54	187	455	278	176	126	322	189	123	1,857
55–59	248	572	375	269	222	429	307	143	2,565
Total	873	1,925	1,216	698	606	1,377	850	692	8,237
Percentage	10.6	23.4	14.8	8.5	7.4	16.7	10.3	8.4	

¹ NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

² Includes full-time students, never worked, long-term unemployed, inadequately described and not classified for other reasons.

**Table A1
continued****Number of deaths by NS-SEC¹ and age, women aged 25–59, 2001–03**

Government Office Regions of England, Wales

Numbers

	Higher managerial and professional	Lower managerial and professional	Intermediate	Small employers and own account workers	Lower supervisory and technical	Semi-routine	Routine	Others ²	Total
South East									
25–29	29	54	34	8	14	46	28	32	246
30–34	61	108	57	22	30	70	44	39	431
35–39	102	177	87	36	61	90	63	57	673
40–44	155	226	129	78	82	150	82	59	961
45–49	217	334	170	112	116	197	115	70	1,330
50–54	345	643	306	192	206	298	187	103	2,280
55–59	456	871	412	304	339	479	310	127	3,297
Total	1,365	2,414	1,195	752	848	1,329	828	486	9,218
Percentage	14.8	26.2	13.0	8.2	9.2	14.4	9.0	5.3	
South West									
25–29	10	20	18	6	10	26	20	18	128
30–34	22	59	40	14	11	38	28	17	229
35–39	49	87	51	30	45	58	44	28	393
40–44	76	127	64	45	48	107	52	38	559
45–49	107	232	109	91	92	158	102	48	939
50–54	170	391	160	166	146	236	142	61	1,472
55–59	217	518	218	285	225	318	212	78	2,071
Total	651	1,434	660	638	578	940	601	289	5,791
Percentage	11.2	24.8	11.4	11.0	10.0	16.2	10.4	5.0	
Wales									
25–29	6	14	12	5	5	17	13	13	84
30–34	9	40	25	13	15	28	19	24	174
35–39	16	55	30	12	21	51	38	33	255
40–44	34	92	49	40	30	61	64	35	405
45–49	49	149	65	59	71	107	88	45	632
50–54	86	213	92	85	113	158	140	49	936
55–59	126	312	166	161	191	239	233	71	1,499
Total	326	873	439	375	445	661	596	270	3,985
Percentage	8.2	21.9	11.0	9.4	11.2	16.6	14.9	6.8	

1 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

2 Includes full-time students, never worked, long-term unemployed, inadequately described and not classified for other reasons.

Table A2**Number of deaths by NS-SEC,¹ cause² and age, women aged 25–59, 2001–03**

England and Wales

Numbers

	Higher managerial and professional	Lower managerial and professional	Intermediate	Small employers and own account workers	Lower supervisory and technical	Semi-routine	Routine	Others ³	Total
All malignant neoplasms									
25–29	54	99	58	21	30	64	32	28	387
30–34	138	270	137	60	65	123	83	44	921
35–39	270	477	218	133	156	248	145	79	1,727
40–44	472	774	366	260	228	423	270	105	2,897
45–49	730	1,289	574	439	468	721	430	182	4,832
50–54	1,142	2,363	1,035	857	867	1,217	843	245	8,568
55–59	1,589	3,110	1,439	1,316	1,371	1,805	1,349	327	12,307
Total	4,395	8,383	3,827	3,086	3,186	4,600	3,150	623	31,639
Percentage	13.9	26.5	12.1	9.8	10.1	14.5	10.0	2.0	
Breast cancer									
25–29	12	15	9	5	5	5	5	2	59
30–34	47	79	43	17	15	30	24	10	266
35–39	129	197	93	45	51	102	50	24	691
40–44	198	321	143	90	85	136	92	34	1,098
45–49	292	455	192	120	155	207	121	59	1,601
50–54	410	738	305	229	241	341	201	61	2,526
55–59	484	881	380	314	286	372	278	76	3,072
Total	1,572	2,687	1,165	821	839	1,194	771	265	9,313
Percentage	16.9	28.9	12.5	8.8	9.0	12.8	8.3	2.8	

1 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

2 Younger age bands have been amalgamated where necessary due to avoid statistical disclosure issues.

3 Includes full-time students, never worked, long-term unemployed, inadequately described and not classified for other reasons.

**Table A2
continued****Number of deaths by NS-SEC,¹ cause² and age, women aged 25–59, 2001–03**

England and Wales

Numbers

	Higher managerial and professional	Lower managerial and professional	Intermediate	Small employers and own account workers	Lower supervisory and technical	Semi-routine	Routine	Others ³	Total
Cancer of trachea, bronchus and lung									
25–39	10	22	11	8	12	14	21	8	107
40–44	24	55	29	23	23	45	28	10	238
45–49	67	137	69	72	72	130	72	27	646
50–54	121	326	155	145	155	229	183	46	1,359
55–59	182	462	232	255	280	420	349	76	2,257
Total	405	1,003	496	503	541	838	653	168	4,607
Percentage	8.8	21.8	10.8	10.9	11.8	18.2	14.2	3.6	
All circulatory diseases									
25–29	15	38	23	6	8	34	17	22	162
30–34	19	57	54	22	37	65	39	39	333
35–39	57	136	85	53	59	129	78	65	663
40–44	92	216	134	85	113	221	168	83	1,111
45–49	144	385	207	148	195	376	252	119	1,826
50–54	188	590	316	270	298	558	424	171	2,816
55–59	355	840	457	483	577	895	753	235	4,594
Total	870	2,261	1,276	1,068	1,287	2,277	1,730	735	11,505
Percentage	7.6	19.7	11.1	9.3	11.2	19.8	15.0	6.4	
Ischaemic heart disease									
25–34	4	13	7	3	10	16	10	9	72
35–39	15	23	21	17	16	37	21	16	166
40–44	26	54	42	19	37	72	65	30	345
45–49	41	114	70	54	81	154	112	48	675
50–54	66	220	120	103	132	238	203	77	1,160
55–59	160	374	217	213	285	433	394	122	2,198
Total	311	798	477	409	562	951	806	303	4,616
Percentage	6.7	17.3	10.3	8.9	12.2	20.6	17.5	6.6	
Cerebrovascular disease									
25–29	5	9	10	3	3	16	4	8	57
30–34	5	27	23	7	13	19	13	11	118
35–39	20	57	31	18	18	40	28	19	230
40–44	38	89	46	32	41	78	49	24	396
45–49	56	168	71	49	61	119	76	32	632
50–54	73	201	104	94	92	166	122	43	894
55–59	106	239	129	152	155	224	173	54	1,232
Total	303	789	413	355	383	661	465	190	3,559
Percentage	8.5	22.2	11.6	10.0	10.8	18.6	13.1	5.4	
All respiratory diseases									
25–29	3	10	16	5	5	20	9	11	79
30–34	9	19	16	11	16	30	15	16	132
35–39	13	30	20	10	18	37	26	28	183
40–44	21	58	30	23	33	64	45	29	303
45–49	24	64	50	42	41	97	74	50	443
50–54	65	159	88	74	123	198	185	76	968
55–59	91	281	172	139	224	354	309	115	1,686
Total	226	621	393	306	460	800	663	325	3,794
Percentage	6.0	16.4	10.3	8.1	12.1	21.1	17.5	8.6	
All digestive diseases									
25–29	5	6	12	4	6	19	7	12	72
30–34	16	37	34	11	17	58	36	35	244
35–39	30	75	63	24	39	78	68	54	430
40–44	40	139	93	50	71	146	116	79	733
45–49	81	196	119	80	102	180	188	98	1,045
50–54	91	292	165	105	137	243	188	91	1,311
55–59	103	290	168	151	192	274	225	84	1,487
Total	367	1,035	653	424	564	998	827	454	5,322
Percentage	6.9	19.5	12.3	8.0	10.6	18.8	15.5	8.5	

1 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

2 Younger age bands have been amalgamated where necessary due to avoid statistical disclosure issues.

3 Includes full-time students, never worked, long-term unemployed, inadequately described and not classified for other reasons.

Table A3

Population¹ by NS-SEC² and age, women aged 25–59, 2001–03

Government Office Regions of England, Wales

Thousands

	Higher managerial and professional	Lower managerial and professional	Intermediate	Small employers and own account workers	Lower supervisory and technical	Semi-routine	Routine	Others ³	Total
North West									
25–29	69	170	109	22	40	98	50	55	611
30–34	110	209	116	41	54	116	61	42	749
35–39	127	227	115	55	61	117	62	34	798
40–44	118	214	103	57	60	101	54	27	733
45–49	103	195	89	54	56	84	47	22	650
50–54	96	188	93	64	59	93	53	19	666
55–59	74	157	87	62	60	95	58	18	612
Total	697	1,361	712	354	389	704	384	218	4,819
Percentage	14.5	28.2	14.8	7.4	8.1	14.6	8.0	4.5	
North East									
25–29	20	55	39	7	15	39	21	21	218
30–34	31	70	42	13	23	48	27	16	270
35–39	38	81	44	18	27	50	29	13	300
40–44	38	80	41	18	28	45	26	11	287
45–49	33	74	35	17	26	36	23	9	253
50–54	30	69	33	19	27	39	25	8	249
55–59	23	55	29	17	25	36	26	8	218
Total	212	485	263	107	172	293	178	85	1,795
Percentage	11.8	27.0	14.7	6.0	9.6	16.3	9.9	4.7	
Yorkshire and The Humber									
25–29	51	123	75	18	32	74	44	38	455
30–34	77	150	79	35	45	91	53	28	558
35–39	88	162	78	44	51	91	52	21	586
40–44	83	156	72	44	50	80	45	17	546
45–49	73	142	62	42	45	65	39	14	481
50–54	68	138	66	49	48	75	43	12	500
55–59	52	112	62	46	47	78	47	12	455
Total	492	983	494	278	317	554	323	141	3,581
Percentage	13.7	27.5	13.8	7.8	8.9	15.5	9.0	3.9	
East Midlands									
25–29	46	107	61	15	28	57	35	26	374
30–34	75	133	66	29	40	70	45	20	478
35–39	88	142	64	37	43	70	44	16	504
40–44	79	132	59	38	42	62	39	12	465
45–49	69	122	53	37	38	55	34	9	418
50–54	66	119	56	44	42	61	39	8	435
55–59	52	102	54	43	42	66	44	9	413
Total	476	857	413	243	277	440	280	100	3,087
Percentage	15.4	27.8	13.4	7.9	9.0	14.3	9.1	3.2	
West Midlands									
25–29	55	134	79	19	35	78	41	39	479
30–34	87	163	87	35	46	94	51	29	591
35–39	101	171	83	44	50	93	48	23	614
40–44	94	160	72	44	49	82	42	18	561
45–49	83	148	65	44	45	72	38	15	509
50–54	78	140	66	51	49	77	43	13	517
55–59	66	125	69	50	52	86	51	12	512
Total	565	1,041	520	286	325	581	313	149	3,781
Percentage	15.0	27.5	13.7	7.6	8.6	15.4	8.3	3.9	
East of England									
25–29	74	158	87	20	33	64	33	28	497
30–34	118	189	92	39	41	73	37	21	609
35–39	138	201	88	52	45	75	36	17	653
40–44	123	186	80	52	43	68	30	13	595
45–49	106	174	75	50	40	61	27	11	544
50–54	101	175	84	61	44	70	32	9	577
55–59	79	151	83	59	45	80	37	10	543
Total	739	1,234	588	334	291	491	233	110	4,020
Percentage	18.4	30.7	14.6	8.3	7.2	12.2	5.8	2.7	
London									
25–29	209	349	165	32	39	90	43	106	1,033
30–34	236	332	153	51	41	96	44	72	1,026
35–39	206	303	138	64	43	94	43	57	946
40–44	151	243	114	61	40	84	37	41	771
45–49	114	205	96	54	34	68	31	29	630
50–54	96	188	95	54	33	67	31	21	585
55–59	72	158	91	47	35	71	34	17	525
Total	1,084	1,777	852	363	263	571	263	342	5,516
Percentage	19.7	32.2	15.5	6.6	4.8	10.3	4.8	6.2	

1 Population optimised for mortality analysis as described in Methods.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

3 Includes full-time students, never worked, long-term unemployed, inadequately described and not classified for other reasons.

**Table A3
continued****Population¹ by NS-SEC² and age, women aged 25–59, 2001–03**

Government Office Regions of England, Wales

Thousands

	Higher managerial and professional	Lower managerial and professional	Intermediate	Small employers and own account workers	Lower supervisory and technical	Semi-routine	Routine	Others ³	Total
South East									
25–29	126	239	124	28	44	84	45	43	734
30–34	207	287	132	55	55	94	45	30	905
35–39	243	309	129	75	61	97	47	25	986
40–44	222	288	116	75	58	88	40	19	906
45–49	188	265	108	72	52	78	34	15	810
50–54	174	268	123	85	57	89	37	13	847
55–59	137	238	124	83	60	102	45	13	802
Total	1,298	1,894	856	474	386	632	292	158	5,989
Percentage	21.7	31.6	14.3	7.9	6.5	10.6	4.9	2.6	
South West									
25–29	51	126	68	18	32	62	31	25	413
30–34	84	161	75	38	41	71	36	19	526
35–39	101	175	74	50	45	73	35	16	570
40–44	98	168	68	53	42	67	30	12	539
45–49	86	159	65	54	38	60	26	10	500
50–54	84	162	72	67	41	68	30	9	532
55–59	68	144	71	68	42	78	36	10	518
Total	573	1,096	494	348	283	480	224	101	3,598
Percentage	15.9	30.5	13.7	9.7	7.9	13.3	6.2	2.8	
Wales									
25–29	23	67	40	9	19	41	22	22	245
30–34	37	85	43	19	27	49	27	17	303
35–39	44	93	42	25	30	50	28	14	326
40–44	42	92	39	26	29	45	23	11	307
45–49	39	86	35	27	26	39	21	10	282
50–54	38	85	35	33	28	41	24	9	295
55–59	31	73	33	32	28	43	25	10	275
Total	254	581	267	172	187	308	171	93	2,033
Percentage	12.5	28.6	13.1	8.5	9.2	15.2	8.4	4.6	

1 Population optimised for mortality analysis as described in Methods.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have been applied as described in Methods.

3 Includes full-time students, never worked, long-term unemployed, inadequately described and not classified for other reasons.

Appendix B

Table B1

Age Standardised mortality rates¹ by NS-SEC² without adjustments for under-recording of occupation at death, women aged 25–59, 2001–03

Government Office Regions of England, Wales

Rate per 100,000

	NS-SEC analytic class							Socio-economic gradient ³	
	1	2	3	4	5	6	7	Without adjustments	With adjustment
North East	112	139 H	133	171	200 H	183	271 H	2.4	2.6
North West	112	143 H	136 H	171 H	200 H	201 H	294 H	2.6	3.0
Yorkshire and The Humber	109	128	117	155	163	168	238	2.2	2.4
East Midlands	113	133	117	154	155 L	169	215 L	1.9	2.1
West Midlands	109	127	114 L	147	188	177	240	2.2	2.5
East of England	113	116 L	123	139	150 L	163	194 L	1.7	2.1
London	105	122	135 H	164 H	168	185 H	233	2.2	3.0
South East	107	121 L	118	120 L	170	153 L	208 L	2.0	2.5
South West	106	120 L	110 L	133 L	165	150 L	206 L	1.9	2.4
Wales	115	136	136	160	183	162	276 H	2.4	2.7
England and Wales	110	128	124	148	175	172	239	2.2	2.6

1 Directly standardised rate using the European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have NOT been applied as described in Methods.

3 Ratio of mortality rates of NS-SEC class 7 to the mortality rate of NS-SEC class 1.

H Rate is statistically significantly higher than that for England and Wales.

L Rate is statistically significantly lower than that for England and Wales.

Table B2

Age-standardised mortality rates¹ from selected malignant neoplasms by NS-SEC,² women aged 25–59, 2001–03

England and Wales

Rate per 100,000

NS-SEC analytic class	All malignant neoplasms			Trachea, bronchus and lung			Breast cancer		
	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval
1 Higher managerial and professional	73	71	75	7	6	7	26	25	27
2 Lower managerial and professional	75	73	76	9	8	10	24	23	25
3 Intermediate	66	63	68	8	7	9	20	19	22
4 Small employers and own account workers	82	79	85	13	12	14	22	21	24
5 Lower supervisory and technical	92	89	96	15	14	17	25	23	26
6 Semi-routine	77	75	80	14	13	15	20	19	22
7 Routine	98	95	102	21	19	22	24	22	26
Ratio 7:1	1.4	1.3	1.4	3.0	2.7	3.4	0.9	0.9	Rate

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have NOT been applied.

Table B3**Age-standardised mortality rates¹ from selected circulatory diseases by NS-SEC,² women aged 25–59, 2001–03**

England and Wales

Rate per 100,000

NS-SEC analytic class	All circulatory diseases			Ischaemic heart disease			Cerebrovascular diseases		
	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval
1 Higher managerial and professional	13	12	14	5	4	5	5	4	5
2 Lower managerial and professional	19	18	20	7	6	7	7	6	7
3 Intermediate	19	18	20	7	6	8	7	6	7
4 Small employers and own account workers	27	25	28	10	9	11	9	8	10
5 Lower supervisory and technical	34	32	36	15	13	16	11	9	12
6 Semi-routine	35	34	37	15	13	16	11	10	12
7 Routine	51	48	54	24	22	26	14	13	15
Ratio 7:1	3.8	3.5	4.1	5.0	4.4	5.7	3.0	2.6	3.4

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have NOT been applied.

Table B4**Age-standardised mortality rates¹ from all respiratory and all digestive diseases by NS-SEC,² women aged 25–59, 2001–03**

England and Wales

Rate per 100,000

NS-SEC analytic class	All respiratory diseases			All digestive diseases		
	Rate	Lower 95% confidence interval	Upper 95% confidence interval	Rate	Lower 95% confidence interval	Upper 95% confidence interval
1 Higher managerial and professional	3	3	4	5	5	6
2 Lower managerial and professional	5	4	5	8	8	9
3 Intermediate	5	4	6	9	8	10
4 Small employers and own account workers	7	6	8	10	9	11
5 Lower supervisory and technical	11	10	13	14	13	15
6 Semi-routine	11	10	12	14	13	15
7 Routine	18	17	20	24	22	26
Ratio 7:1	5.8	4.9	6.8	4.5	4.0	5.2

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have NOT been applied.

Table B5**Age-standardised mortality rates¹ for selected causes and regions by NS-SEC,² women aged 25–59, 2001–03**

Rate per 100,000

NS-SEC analytic class	All cancers		All circulatory diseases	
	North West	South West	North West	South West
1 Higher managerial and professional	66	74	15	10
2 Lower managerial and professional	82	72	23	16
3 Intermediate	68	59	22	19
4 Small employers and own account workers	91	80	31	21
5 Lower supervisory and technical	95	90	43	33
6 Semi-routine	88	67	40	31
7 Routine	105	96	64	39
Ratio 7:1	1.6	1.3	4.1	3.8

1 Directly age-standardised using European standard population.

2 NS-SEC assigned by the 'combined' method whereby if married the most advantaged class of either the woman or her husband is used to represent the woman's classification. Adjustments for the under-recording of death have NOT been applied.