The *Eucalyptus* canker pathogen *Holocryphia eucalypti* on *Eucalyptus* in New Zealand

M. Gryzenhout^{A,D}, M. Vermeulen^B, M. Dick^C and M. J. Wingfield^A

^AForestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002, South Africa.

^BDepartment of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002, South Africa.

^CScion Research, New Zealand Forest Institute Ltd, Rotorua, New Zealand.

^DCorresponding author. Email: Marieka.Gryzenhout@fabi.up.ac.za

Abstract. *Holocryphia eucalypti* is an opportunistic canker pathogen of *Eucalyptus* and *Corymbia* spp. (Myrtaceae, Myrtales) in Australia and South Africa. It is also known in Australia on *Tibouchina* trees (Melastomataceae, Myrtales). Using DNA sequence comparisons and morphological characterisation, we show for the first time that *H. eucalypti* is present in New Zealand on *Eucalyptus* spp.

Holocryphia eucalypti (= *Cryphonectria eucalypti*) causes die-back, and stem and branch cankers of several *Eucalyptus* and *Corymbia* spp. in natural forests and commercial plantations in Australia and South Africa (Walker *et al.* 1985; Van der Westhuizen *et al.* 1993; Carnegie 2007). The pathogen has also recently been reported on *Tibouchina urvilleana* (Lasiandra) planted as ornamentals in Australia (Heath *et al.* 2007). The canker disease caused by *H. eucalypti* is closely associated with stress conditions such as drought and cankers can be relatively mild or lethal where host trees are susceptible and stress is severe (Old *et al.* 1986; Wardlaw 1999; Gryzenhout *et al.* 2003). Because predisposition plays such an important role in disease development, *H. eucalypti* is not considered as severe (Yuan and Mohammed 2000) as other closely related *Eucalyptus* pathogens such as species of *Chrysosporthe* (Gryzenhout *et al.* 2009).

Canker caused by *H. eucalypti* is common on many *Eucalyptus* and *Corymbia* species in forests and plantations of eastern Australia including Tasmania, and it also occurs in Western Australia (Old *et al.* 1986; Davison and Coates 1991; Wardlaw 1999; Yuan and Mohammed 2000; Carnegie 2007). In South Africa, it occurs on commercially propagated, non-native *Eucalyptus* spp. in plantations (Van der Westhuizen *et al.* 1993; Gryzenhout *et al.* 2003). Population genetic studies employing microsatellite data have shown that *H. eucalypti* was most likely introduced into South Africa, although evidence for its origin in Western Australia was less convincing (Nakabonge *et al.* 2008).

During the course of the past 24 years, isolates of a fungus in the Cryphonectriaceae have been collected from various *Eucalyptus* species in New Zealand. Most locations were in the southern half of the North Island with one from the north of the South Island (Table 1). Based on cultural characteristics and the morphology of conidia produced in culture, the isolates were

identified as *Endothiella*-like, which represent the anamorph state of some species in the Cryphonectriaceae (Gryzenhout *et al.* 2009). However, in the absence of sexual fruiting bodies or well-defined fruiting bodies on host tissue and in culture, a definitive identification could not be made. The aim of this study was thus to characterise the isolates from *Eucalyptus* stems in New Zealand based on DNA sequence comparisons for the β -Tubulin gene region.

Original isolates curated in NZFS (National Forestry Culture Collection) were obtained from discoloured yellowish wood, from branch cankers and die-back. These isolates are maintained at 4°C in the culture collection (NZFS) of Scion, New Zealand Forest Research Institute, Rotorua, New Zealand (Table 1). Duplicates of the cultures, as well as an isolate (CMW12723) obtained from asexual fruiting bodies from a later collection, are housed in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa (Table 1). The representative specimen of the original bark material from which the culture CMW12723 was made has been deposited in the National Collection of Fungi (PREM), Pretoria, South Africa (Table 1).

For DNA sequence comparisons, isolates were grown in 2% Malt Extract broth (100 μ L in 1.5 μ L Eppendorf tubes), the mycelium harvested and the DNA extracted following Gryzenhout *et al.* (2006). The DNA was used in polymerase chain reactions (PCR) of the β -Tubulin gene regions 1 and 2 and sequenced using the same protocols as those described by Gryzenhout *et al.* (2006). The sequences were included in a data matrix containing sequences of *H. eucalypti* isolates previously published (Gryzenhout *et al.* 2006; Heath *et al.* 2007). Sequences of isolates CMW10010 and CMW10011 were obtained from the study of Myburg (2003). Species of

Table 1. Isolates and specimens used in this study

Species identity	Isolate ^A	Host	Origin	Collector	Date	GenBank accession ^B
<i>Holcynophia eucalypti</i>	CMW7036/CBS119478	<i>Eucalyptus</i> sp.	Mpumalanga South Africa	I van der Westhuizen	n/a	AF 368341, AF 368340
	CMW7037/CBS119477	<i>Eucalyptus delegatensis</i>	NSW, Australia	KM Old	n/a	AF 368343, AF 368342
	CMW7038/CBS119475	<i>Eucalyptus globulus</i>	WA, Australia	MJ Wingfield	September 1997	AF 368345, AF 368344
	CMW14545/CBS115852	<i>Eucalyptus</i> sp.	South Africa	I van der Westhuizen	n/a	DQ 368730, DQ 368731
	CMW8541/CBS115720	<i>Eucalyptus</i> sp.	Nseleni, KwaZulu-Natal, South Africa	J Roux	July 2007	GQ 465234
	CMW7033/CBS115842	<i>Eucalyptus grandis</i>	South Africa	M Venter		DQ 368728, DQ 368729
	CMW6246	<i>Tibouchina urvilleana</i>	Melbourne, Australia	MJ Wingfield	September 2000	EF 127995, EF 128000
	CMW6249	<i>T. urvilleana</i>	Melbourne, Australia	MJ Wingfield	September 2000	EF 127996, EF 128001
	CMW6244	<i>T. urvilleana</i>	Melbourne, Australia	MJ Wingfield	September 2000	EF 127997, EF 128002
	CMW10729	<i>T. urvilleana</i>	Brisbane, Australia	MJ Wingfield	2000	EF 127998, EF 128003
	CMW10010/NZFS79/E	<i>Eucalyptus fastigata</i>	Castleridge, Wellington, New Zealand	BJ Rogan	1997	AY 214236, AY 214272
	CMW10011/NZFS79/F	<i>Eucalyptus</i> sp.	Central Park, Wellington, New Zealand	BJ Rogan	1998	AY 214237, AY 214273
<i>Cryphonectria parasitica</i> <i>Endothia gyrosa</i> <i>Microthia havanensis</i>	CMW10015/NZFS79.15	<i>Eucalyptus fastigata</i>	Jackson Road, Genmarie Blk., Bay of Plenty, New Zealand	RJ van Boven	1999	GQ 427071
	CMW10016/NZFS79.17	<i>Eucalyptus nitens</i>	Omataroa, Bay of Plenty, New Zealand	MR Twaddle	1999	GQ 427068
	CMW10017/NZFS372	<i>Eucalyptus ficifolia</i>	Lambton Quay, Wellington, New Zealand	L Renney	2000	GQ 427069
	CMW10021/NZFS833	<i>E. globulus</i>	Westport, Buller, New Zealand	PM Bradbury	2001	GQ 427072
	CMW10797/NZFS79	<i>Eucalyptus regnans</i>	Wanganui, New Zealand	MA Dick	1985	AY 214238, AY 214274
	CMW12723 ^C	<i>E. globulus</i>	Auckland, New Zealand	MJ Wingfield	February 2003	GQ 427070
	CMW7048/ATCC48198	<i>Quercus virginiana</i>	Virginia, USA	FF Lombard	May 1960	AF 273076, AF 273470
	CMW2091/ATCC48192	<i>Quercus palustris</i>	Virginia, USA	RJ Stipes	n/a	AF 368337, AF 368336
	CMW14550/CBS115855	<i>Eucalyptus saligna</i>	Tabasco State, Mexico	CS Hodges	February 1998	DQ 368741, DQ 368742

^ACMW = Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria; NZFS = National Forestry Culture Collection, Scion, New Zealand, Forest Research Institute Ltd.^BAccession numbers refer to the combined sequence data of the β -tubulin 1 and 2 regions.^COriginal bark material deposited as PREM 60237 in the National Collection of Fungi (PREM), Pretoria, South Africa.

other genera in the Cryphonectriaceae (*Endothia gyrosa*, *Cryphonectria parasitica*, *Microthia havanensis*) were defined as outgroups. The dataset consisted of 20 taxa and the sequences were aligned with the CLUSTAL function of the program MEGA ver. 4 (Tamura *et al.* 2007) and verified manually.

Phylogenetic analyses were run with PAUP (Phylogenetic Analysis Using Parsimony) version 4.0b10 (Swofford 2002). Phylogenetic trees were obtained with maximum parsimony (uninformative sites excluded, heuristic search with 100 random sequence additions and tree-bisection-reconnection branch swapping, MULTREES off, base pairs re-weighted according to the consistency index). The strength of branches was tested with a 70% bootstrap analysis (1000 replicates).

Of the 928 characters in the DNA matrix, 165 were uninformative, 645 were constant and 118 were informative. Two trees that were identical topologically but differed slightly based on length of some branches (Tree Length=170.3, Consistency Index=0.882, Retention Index=0.895, g1 value=-2.62), were obtained in the analysis. The phylogenetic trees (Fig. 1) clearly showed that the isolates from New Zealand grouped with those of *H. eucalypti* from South Africa and Australia (bootstrap support 100%). The isolates from New Zealand formed a group with those from *T. urvilleana* in Australia (bootstrap support 87%). Two other groups, including

Fig. 1. Most parsimonious tree based on DNA sequences of the β -tubulin gene. Confidence levels based on a 70% bootstrap analysis are indicated on the branches, with isolates sequenced in this study in bold typeface. *Cryphonectria parasitica*, *Endothia gyrosa* and *Microthia havanensis* were defined as outgroups.

South African and Australian isolates, were also evident (bootstrap support 100% and below 50%, respectively).

This study reports the presence of the *Eucalyptus* pathogen *H. eucalypti* in New Zealand for the first time. This report further contributes to the databases and checklists of pathogens and fungi in New Zealand (<http://nzfungi.landcareresearch.co.nz/html/mycology.asp> database; Dingley 1969; Pennycook 1989; McKenzie *et al.* 2000; Pennycook and Galloway 2004). Due to the opportunistic nature of *H. eucalypti* on *Eucalyptus* and *Tibouchina*, the occurrence of the fungus in New Zealand is most likely not of economic significance, although its presence deserves to be monitored.

Relatively little is known regarding the Cryphonectriaceae in New Zealand. Although there are reports of *Endothia gyrosa*, a previous name used for *H. eucalypti* (Gryzenhout *et al.* 2006), in the country (Gryzenhout *et al.* 2009), these reports are from *Nothofagus* (McKenzie *et al.* 2000; Pennycook and Galloway 2004), *Myrsine salicina* and dead wood of an unknown host (<http://nzfungi.landcareresearch.co.nz/html/mycology.asp> database) and need to be verified based on recent taxonomic changes to the group (Gryzenhout *et al.* 2009). Reports of other Cryphonectriaceae that could represent incorrectly identified *H. eucalypti* specimens in New Zealand, include those of *Rostrareum longirostris* on *Nothofagus* (Nothofagaceae, Fagales), *Amphilogia gyrosa* and *Amphilogia major* on *Elaeocarpus* spp. (Elaeocarpaceae, Oxalidales), an *Endothiella* sp. and *Cryphonectria radicalis* (<http://nzfungi.landcareresearch.co.nz/html/mycology.asp> database; Dingley 1969; Pennycook 1989; McKenzie *et al.* 2000; Pennycook and Galloway 2004; Gryzenhout *et al.* 2005). In the NZFS collection, cultures of an *Endothiella* sp. identified only on the basis of culture characteristics, originate from *Myrsine chathamica* and *Leptospermum scoparium* as well as various species of eucalypts. A member of the Cryphonectriaceae has also been found on a *Coriaria* sp. (Coriariaceae, Cucurbitales) in New Zealand (PDD28477, Waiomu, Thames, Auckland, J.M. Dingley, August 1958) but the identity of this collection relies only on a herbarium specimen and cannot be confirmed (M. Gryzenhout, unpubl. data). Thus, only the identities of *A. gyrosa* and *A. major* have been verified based on recent phylogenies (Gryzenhout *et al.* 2005).

The relative proximity of New Zealand to Australia, as well as the common occurrence of *H. eucalypti* in the eastern part of Australia indicates a possibility that the fungus originated in the latter country. However, it is also possible that *H. eucalypti* occurs on native Myrtaceae in New Zealand. It may also occur on other non-native shrubs such as *Tibouchina* spp. (Melastomataceae, Myrtales), which are hosts of this fungus (Heath *et al.* 2007) and many other members of the Cryphonectriaceae (Gryzenhout *et al.* 2009). Answers to these intriguing questions will require further surveys and population genetic studies.

Acknowledgements

Funding from the National Research Foundation (NRF), members of the Tree Protection Co-operative Program (TPCP), the THRIP support program of the Department of Trade and Industry, and the Department of Science and Technology/National Research Foundation Centre of Excellence in Tree Health Biotechnology, South Africa made this study possible.

References

- Carnegie AJ (2007) Forest health condition in New South Wales, Australia, 1996–2005. II. Fungal damage recorded in eucalypt plantations during forest health surveys and their management. *Australasian Plant Pathology* **36**, 225–239. doi:10.1071/AP07021
- Davison EM, Coates DJ (1991) Identification of *Cryphonectria cubensis* and *Endothia gyrosa* from eucalypts in Western Australia using isozyme analysis. *Australasian Plant Pathology* **20**, 157–160. doi:10.1071/AP09910157
- Dingley JM (1969) Records of plant diseases in New Zealand. *New Zealand Department of Scientific and Industrial Research Bulletin* **192**, 120.
- Gryzenhout M, Eisenberg BE, Coutinho TA, Wingfield BD, Wingfield MJ (2003) Pathogenicity of *Cryphonectria eucalypti* to *Eucalyptus* clones in South Africa. *Forest Ecology and Management* **176**, 427–437. doi:10.1016/S0378-1127(02)00309-2
- Gryzenhout M, Glen HF, Wingfield BD, Wingfield MJ (2005) *Amphilogia* gen. nov. for *Cryphonectria*-like fungi from *Elaeocarpus* spp. in New Zealand and Sri Lanka. *Taxon* **54**, 1009–1021.
- Gryzenhout M, Myburg H, Hodges CS, Wingfield BD, Wingfield MJ (2006) *Microthia*, *Holocryphia* and *Ursicollum*, three new genera on *Eucalyptus* and *Coccoloba* for fungi previously known as *Cryphonectria*. *Studies in Mycology* **55**, 35–52. doi:10.3114/sim.55.1.35
- Gryzenhout M, Wingfield BD, Wingfield MJ (2009) Taxonomy, phylogeny, and ecology of bark-infecting and tree killing fungi in the Cryphonectriaceae. APS Press, St. Paul.
- Heath RN, Roux J, Gryzenhout M, Carnegie AJ, Smith IW, Wingfield MJ (2007) *Holocryphia eucalypti* on *Tibouchina urvilleana* in Australia. *Australasian Plant Pathology* **36**, 560–564. doi:10.1071/AP07059
- McKenzie EHC, Buchanan PK, Johnston PR (2000) Checklist of fungi on *Nothofagus* species in New Zealand. *New Zealand Journal of Botany* **38**, 635–720.
- Myburg H (2003) Genera and species in the *Cryphonectria/Endothia* complex and their placement in the Diaporthales: A molecular and morphological synopsis. In 'Molecular Phylogenetic Studies on Species of *Cryphonectria* and Related Fungi'. pp. 203–273. Ph.D. Thesis. (University of Pretoria: Pretoria)
- Nakabonge G, Burgess T, Gryzenhout M, Roux J, Wingfield BD, Wingfield MJ (2008) Population structure of the fungal pathogen *Holocryphia eucalypti* in Australia and South Africa. *Australasian Plant Pathology* **37**, 154–161. doi:10.1071/AP08004
- Old KM, Murray DIL, Kile GA, Simpson J, Malafant KWJ (1986) The pathology of fungi isolated from eucalypt cankers in south-eastern Australia. *Australian Forest Research* **16**, 21–36.
- Pennycook SR (1989) Part II. Fungal plant diseases. Plant Diseases Recorded in New Zealand 2. Plant Diseases Division, Auckland.
- Pennycook SR, Galloway DJ (2004) Checklist of New Zealand "fungi". In 'Introduction to the Fungi of New Zealand'. (Ed. E McKenzie) pp. 401–488. (Fungal Diversity Press: Hong Kong)
- Swofford DL (2002) PAUP. Phylogenetic Analysis Using Parsimony. Version 4.0b10. Sinauer Associates, Sunderland.
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* **24**, 1596–1599. doi:10.1093/molbev/msm092
- Van der Westhuizen IP, Wingfield MJ, Kemp GHJ, Swart WJ (1993) First report of the canker pathogen *Endothia gyrosa* on *Eucalyptus* in South Africa. *Plant Pathology* **42**, 661–663. doi:10.1111/j.1365-3059.1993.tb01548.x
- Walker J, Old KM, Murray DIL (1985) *Endothia gyrosa* on *Eucalyptus* in Australia with notes on some other species of *Endothia* and *Cryphonectria*. *Mycotaxon* **23**, 353–370.
- Wardlaw TJ (1999) *Endothia gyrosa* associated with severe stem cankers on plantation grown *Eucalyptus nitens* in Tasmania, Australia. *European Journal of Forest Pathology* **29**, 199–208. doi:10.1046/j.1439-0329.1999.00143.x
- Yuan ZQ, Mohammed C (2000) The pathogenicity of isolates of *Endothia gyrosa* to *Eucalyptus nitens* and *E. globulus*. *Australasian Plant Pathology* **29**, 29–35. doi:10.1071/AP00005

Manuscript received 26 October 2009, accepted 7 December 2009