

## Corrigendum

The publisher wishes to draw attention to an error in the following article:

Li L, Kayukawa Y, Imai M *et al.* Association of sleep-disordered breathing with hypertension in Japanese industrial workers. *Sleep Biol. Rhyth.* 2003; 1: 221–7.

On page 223, table 1 should have been as follows:

**Table 1** Sociodemographic characteristics, sleep, lifestyle and health variables associated with sleep-disordered breathing (SDB)

Variables	Total (n = 2512)	HS (n = 243)	OSAS (n = 61)	Non-SDB (n = 2238)
Age (years)	40.5±11.8	44.8±10.2***	46.5±9.7***	40.0±11.9
Gender (%)				
Male	85.7	93.0**	88.5	84.9
Female	14.3	7.0**	11.5	15.1
BMI (kg/m <sup>2</sup> )	22.4±3.0	24.3±3.1***	23.7±3.1***	22.2±2.9
ESS score	5.9±3.8	6.8±4.0***	8.4±3.3***	5.8±3.7
Alcohol consumption				
Non-drinkers	42.4	35.4	29.5	43.3
Light drinkers	45.9	49.4	45.9	45.5
Heavy drinkers	11.7	15.2*	24.6**	11.2
Hypertension	22.4	35.0***	29.5	21.3
Systolic BP (mmHg)	125.9±17.4	133.1±19.5***	129.4±20.4*	125.1±17.0
Diastolic BP (mmHg)	76.4±12.1	81.5±12.6***	79.3±11.3*	75.8±12.1

BMI, body mass index; BP, blood pressure; ESS, Epworth Sleepiness Scale; HS, habitual snoring; OSAS, obstructive sleep apnea syndrome. Data are mean ± SD or number of subjects (%). \**P*<0.05, \*\**P*<0.001, \*\*\**P*<0.001 compared with non-SDB.

The publisher wishes to apologise for any inconvenience caused by this error.