Erratum: “New Polyamides with Main-Chain Cyanine Chromophores” [Polymer Science, Ser. A, 53 (6), 457 (2011)]

M. Ya. Goikhman^a, N. P. Yevlampieva^b, N. V. Kamanina^c, I. V. Podeshvo^a,
I. V. Gofman^a, S. A. Mil'tsov^d, A. P. Khurchak^b, and A. V. Yakimanskii^a

^a Institute of Macromolecular Compounds, Russian Academy of Sciences,
Bol'shoi pr. 31, St. Petersburg, 199004 Russia

^b Fock Institute of Physics, St. Petersburg State University,
Ul'yanovskaya ul. 1, Petrodvorets, St. Petersburg, 198504 Russia

^c NPK Vavilov State Optical Institute,
Birzhevaya liniya 12, St. Petersburg, 199034 Russia

^d St. Petersburg State University, Universitetskii pr. 26, Petrodvorets, St. Petersburg, 198504 Russia
e-mail: mgoikhman@yandex.ru

Received October 25, 2010;

Revised Manuscript Received February 1, 2011

DOI: 10.1134/S0965545X11130013

In the paper “New Polyamides with Main-Chain Cyanine Chromophores” by M.Ya. Goikhman, N.P. Yevlampieva, N.V. Kamanina, I.V. Podeshvo, I.V. Gofman, S.A. Mil'tsov, A.P. Khurchak, and A.V. Yakimanskii published in issue 6, pp. 457–468, 2011, Table 5 was missed by the authors. Therefore, Table 5 should be read instead of Table 4 (p. 467, left column).

Table 5. Nonlinear Optical Properties of Films of Chromophore-Containing Polymers

Copolymer	Film thickness, μm	Δn_i	$n_2 \times 10^{10}, \text{cm}^2 \text{W}^{-1}$	$\chi^{(3)} \times 10^9, \text{cm}^3 \text{erg}^{-1}$
X	3.5 ± 0.1	0.0430	1.440	4.49
XIII	7.0	0.0045	0.150	0.46
XIII	16.0	0.0013	0.044	0.14