
ERRATA

**Erratum: “About Convex 4-Isosceles 7-Point Sets”
[*Mathematical Notes* 85 (5–6), 760–763 (2009)]**

Xianglin Wei^{1*}, Wenhua Lan^{2}, and Ren Ding^{2***}**

¹*College of Science, Hebei University of Science and Technology, China*

²*College of Mathematics, Hebei Normal University, China*

Received June 21, 2009

DOI: 10.1134/S0001434609110352

The following corrections must be made in our paper.

- Theorem 1 must be formulated as follows.

Theorem 1. *Suppose that P is the set R_7 minus one point. Then there exist two convex 4-isosceles 7-point sets containing P .*

- On p. 761 on lines 21–22 from above, the following part of the sentence:

“If x lies on \widehat{ab} (with the exception of the endpoint b), then the set $\{3, 5, 6, x\}$ is not 4-isosceles;”

must be replaced by

“If x lies on \widehat{ab} (with the exception of the endpoint b), then it must be a point q such that $d(1, q) = d(6, q)$, because, otherwise, the set $\{3, 5, 6, x\}$ is not 4-isosceles and we can easily see that $\{1, 2, 3, 4, 5, 6, q\}$ is a 4-isosceles set;”

- The caption for Fig. 1 must be the following:

“The convex 4-isosceles 7-point sets containing the set R_7 minus one point”.

- The last-but-one paragraph in the proof of Theorem 1 must be replaced by

“Thus, there exist two convex 4-isosceles 7-point sets containing the set R_7 minus one point.”

ACKNOWLEDGMENTS

This research was supported by the Science Foundation of Hebei University of Science and Technology (grant no. XL200818).

*E-mail: wxlhebtu@126.com

**E-mail: lanandshuo@126.com

***E-mail: rending@hebtu.edu.cn